

DOCTORAL THESIS

Developments to established dose-finding methodologies for application in trials with complex and innovative designs

Author:

Amit PATEL

Supervisor:

Prof. Lucinda BILLINGHAM

Dr. Kristian BROCK

*A thesis submitted in fulfillment of the requirements
for the degree of Doctor of Philosophy*

Institute of Cancer and Genomic Sciences

College of Medical and Dental Sciences

University of Birmingham

February 21, 2021

Abstract

**Developments to established dose-finding methodologies for application
in trials with complex and innovative designs**

by Amit PATEL

Insert abstract here...

Acknowledgements

Acknowledge people here ...

Contents

Abstract	iii
Acknowledgements	v
1 Introduction	1
2 Implementing the PO-TITE-CRM trial design into ADePT-DDR	3
2.1 Introduction	3
2.2 The PO-TITE-CRM Design	9
2.3 PO-TITE-CRM in ADePT-DDR	13
2.3.1 Partial Ordering in Practice	14
2.3.2 The TITE component	18
2.3.3 Stopping Rules	20
2.3.4 Operating Characteristics	21
2.4 Exploring other designs	27
2.5 Discussion	35
2.6 Conclusion	38
Bibliography	41

List of Figures

2.1	Example dose levels to illustrate partial ordering.	7
2.2	ADePT-DDR dose levels across dose and duration.	16
2.3	Weight function across the follow-up period.	19
2.4	Illustration of true DLT rates used in simulations.	28
2.5	Plot of simulations comparing designs for ordering 1.	33
2.6	Plot of simulations comparing designs for ordering 2.	34

List of Tables

2.1	Example drug combinations with two agents.	9
2.2	ADePT-DDR dose-levels.	15
2.3	Operating Characteristics for ordering 1.	24
2.4	Operating Characteristics for ordering 2.	25
2.5	Summary of simulated patient numbers for each scenario. . . .	26
2.6	Alternative designs selection probabilities for ordering 1.	31
2.7	Alternative designs selection probabilities for ordering 2.	32

Chapter 1

Introduction

Chapter 2

Implementing the PO-TITE-CRM trial design into ADePT-DDR

2.1 Introduction

Worldwide there are approximately 600,000 new cases of Head and Neck Squamous Cell Carcinoma (HNSCC) each year [1]. Of which, 12,000 occur in the UK with the most common forms of treatment being surgery, radiotherapy and/or chemotherapy [2]. Radiotherapy is essential for the treatment of cancer. It has been estimated that more than 40% of patients will receive radiotherapy at some point in their treatment [3]. However, despite recent advancements in radiation techniques and the use of concomitant chemo radiotherapy, patients with solid tumours such as head and neck cancer have suboptimal cure rates [2], [4]. For those with advanced HNSCC primary radiotherapy with concurrent chemotherapy is often offered but, it has not been shown to improve survival in patients aged over 70 compared to radiotherapy alone [5]. Therefore, any strategy to improve the efficacy of radiotherapy without increasing toxicity would have a significant impact on patient outcomes.

DNA damage repair (DDR) inhibition is a potential technique which could

be utilised as it potentiates the therapeutic effects of ionising radiation in cancer cells [6]. Combining radiotherapy with DDR inhibition could improve clinical outcomes for these patients [7].

The ADePT-DDR trial is a platform trial which aims to evaluate the safety and efficacy of different DDR agents, or different immunotherapy agents and/or DDR and immunotherapy combinations, together with radiotherapy in patients with HNSCC. The initial component of this trial is a single-arm dose-finding trial investigating the ataxia telangiectasis and Rad3-related (ATR) inhibitor AZD6738 in combination with radiotherapy. ATR inhibitors not only stop DNA repair but impair the mechanism that allows for repairs to take place. Preclinical models have shown this double blocking to be effective in killing cancer cells [8].

Traditionally dose-finding trials aim to determine the maximum tolerated dose (MTD) of a treatment based on the cytotoxic assumption that the most toxic dose is the most efficacious. Rule-based or 'up and down' designs achieve this by escalating and de-escalating doses dependent on the observation of severe toxicity due to the drug, commonly referred to as a dose-limiting toxicity (DLT). In the case of the 3+3 design escalation continues until at least two patients in a cohort of three or six experience a DLT. More explicitly, the MTD is the dose level below the dose at which $\geq 33\%$ of patients experience a DLT [9]. Model-based designs such as the continual reassessment method (CRM) [10] work on the assumption that the probability of toxicity increases monotonically with increases in dose levels. The CRM aims to find the MTD which is a dose with specified target toxicity level.

Due to the historical use of rule-based designs [11], [12], the majority of the terminology used to describe them, and the ambiguity they raise, have been inherited by modern designs such as the CRM. The MTD in the context of a CRM is not the 'maximum' dose patients could tolerate but rather a dose in

which there would be an acceptable target probability of a DLT occurring. For example, if the target is set at 25% the MTD would be the dose at which there is a 25% probability of experiencing a DLT. Rather than using the term MTD, the dose to be found will be referred to as the target dose (TD%%, where the %'s are replaced by the target probability), i.e. TD25 would be the dose expected to be toxic in 25% of patients.

The investigation of multiple-agent treatments, where the monotonicity assumption may not hold, is increasing in early phase trials. Finding the TD in combinations of treatments, compared to single-agents, presents methodological challenges. Each drug individually may obey the monotonicity assumption, however, when combined, the ordering of doses in terms of toxicity may not be fully apparent. An order may be identified for a subset of the doses which would result in a partial order. Without a fully understood ordering it is uncertain which dose should be chosen in decisions of escalation and de-escalation and ultimately as the TD. This issue is not exclusively reserved for trials with multiple-agents. The monotonicity assumption may not hold for certain drugs in single-agent studies leading to partial orders of dose toxicity. For example, when dose and frequency of administration vary between dose levels. Monotonicity is a very strong assumption. It requires that probability of toxicity always increases - staying the same is not enough. At high enough doses, this assumption is almost surely violated for all interventions when the event probability reaches its maximum. Thus, even when total ordering is possible, the monotonicity assumption could be violated [13]. This can occur in scenarios where multiple parameters of the treatment schedule are altered for each dose level. For example, either dose or treatment duration could be increased and even if patients receive an equal dose it would remain unclear as to if prolonged exposure to a lower dose is more toxic than short exposure to a higher dose, which implies a partial ordering of toxicity probabilities.

Further methodological challenges revolve around the issue of late-onset toxicities. Typically, early phase trials implement a short window to observe DLTs. This works well in situations where toxicities are likely to occur rapidly after treatment. However, this is not optimal for treatments that could cause late-onset toxicities such as radiotherapy. The aim here would be to incorporate a larger observation window to account for potential late-onset toxicities whilst also minimising the trial duration.

Cheung and Chappel [14] introduced an extension to the CRM to deal with the issues of treatments that may cause late-onset toxicity. This design referred to as the time-to-event CRM (TITE-CRM), uses a weighted dose-response model to incorporate the time it takes for a DLT to occur in a patient. There have also been published trial designs to deal with the issues that arise from investigating combinations of treatments. Thall et al. [15] proposed an adaptive two-stage Bayesian design which utilises a parametric model of toxicity as a function of two doses. Yin and Yuan [16] present a Bayesian design that uses a copula regression model to evaluate the joint toxicity probabilities of combined drugs. The continual reassessment method for partial orders (PO-CRM) developed by Wages et al. [17] extends the CRM design by relaxing the assumption of monotonicity and by modelling different potential orders. Figure 2.1 shows a simple example of partial ordering where the order of two out of the four dose levels are unknown.

Wages et al. [17], [18] further developed their work on the PO-CRM to deal with late-onset toxicities by implementing a TITE component. This trial design, referred to as the time-to-event continual reassessment method in the presence of partial orders (PO-TITE-CRM) by the authors, was chosen to be used in ADePT-DDR. A search of PubMed, conducted on the 25th of July 2020, found six articles that had cited the PO-TITE-CRM design by Wages et al. [18].

FIGURE 2.1: Example dose levels to illustrate partial ordering.



Of these six articles non actually implement the design into a trial. The following paragraphs provide more details.

Five of these papers were methodological in nature, two of which only include the PO-TITE-CRM design in a brief introduction to current methodology before going on to present new Bayesian trial designs [19], [20]. The other three papers were authored by Wages. The first of which details practical considerations and specifications for the PO-CRM design, the TITE variant is only cited as the source of an example which is being used [21]. One paper presents an R package ‘pocrm’ [22], [23]. The package is only capable of analysing the PO-CRM design. The TITE variant is only referenced here as it illustrates the issue of partial ordering. The last methodological paper by Wages et al. [24] presents three different methods for phase I studies of drug combinations one of which is the PO-CRM however, PO-TITE-CRM is only mentioned as an extension to this design. A key message in this paper is the fact that novel methodologies

are constantly emerging but are rarely implemented in practice.

The last paper is a protocol paper for a phase I/II study, OLA-TMZ-RTE-01 [25]. The phase I component of the study aims to determine the recommended phase II dose (RP2D) of olaparib combined with a standard schedule of radiotherapy and temozolomide (TMZ) as first line treatment for patients with unresectable glioblastoma (GBM). The treatment schedule is divided into a radiotherapy and maintenance period. They propose to conduct two sequential dose-escalations of seven different olaparib dose-levels. Patients in the first escalation will be allocated to a dose level of olaparib for 10 weeks including radiotherapy for six weeks with TMZ given each day during radiotherapy and then for six cycles four weeks post radiotherapy during the maintenance period. They state the MTD1 will be determined using a TITE-CRM. Patients in the second escalation olaparib at the MTD1 during the radiotherapy period along with the same schedule of radiotherapy and TMZ. Those patients will then be allocated to one of the seven dose levels of olaparib during the maintenance period. Again, it is stated that the MTD2 will be determined using TITE-CRM modelling. The RP2D is the MTD1 and MTD2 during the radiotherapy and maintenance period respectively. Even though a combination of treatments is being investigated only olaparib is being escalated and doses for other treatments are fixed for all patients. Furthermore, the dose-levels for olaparib increase consistently in either amount or duration meaning there are no issues of partial ordering which would warrant the use of PO-TITE-CRM. The authors reference the TITE-CRM methodology with two papers. One of them being the paper detailing the PO-TITE-CRM design and the other being a paper by Huang and Kuan [26] which proposes an adaptive weight function that incorporates cyclical data of treatment into the TITE-CRM. It is unclear as to why the PO-TITE-CRM is cited as its methodology is not mentioned anywhere in methods.

This is just a brief review of the current literature but it seems that the PO-TITE-CRM has rarely been used or discussed since its inception.

This chapter provides novel insight into the methodology of PO-TITE-CRM through application in a real-world scenario. Section 2.2 will detail how the PO-TITE-CRM works. Section 2.3 discusses the justification for implementing the design into the ADePT-DDR trial and our experiences doing so. Section 2.4 explores other alternative designs which could have been implemented and assess how they perform in comparison to the PO-TITE-CRM. We provide some discussion in Section 2.5 and finally some conclusions in Section 2.6.

2.2 The PO-TITE-CRM Design

Wages et al. [18] introduced the PO-TITE-CRM design which builds directly upon the PO-CRM design by incorporating a TITE component into the dose toxicity model. The aim of which is to determine the target dose for combinations of drugs where the monotonicity assumption does not hold, in a setting where late-onset toxicities are possible.

TABLE 2.1: Example of drug combinations for a trial investigating two agents.

Agent	Drug combinations					
	d_1	d_2	d_3	d_4	d_5	d_6
A (mg/day)	0.25	0.5	1.0	0.25	0.5	1.0
B (mg/day)	1.0	1.0	1.0	1.5	1.5	1.5

To help understand partial ordering consider an example of an early phase trial investigating the combination of two agents. Drug A which consists of three doses (0.25, 0.5, 1.0 mg/day) and drug B which consists of two doses (1.0, 1.5 mg/day), for a total of six drug combinations d_1, \dots, d_6 (Table 2.1). For each drug independently we assume they have a monotonic dose-toxicity curve however, the ordering of toxicity probabilities for some of the treatment

combinations is unknown. Specifically, we can say d_1 is less toxic than d_2 as the dose of drug A increased whilst the dose of drug B stayed the same. This is also the case for d_2 and d_3 . So, d_1 can always be considered less toxic than d_2 which is always less toxic than d_3 . The same can be said for doses d_4 , d_5 and d_6 , these three doses are can also all be considered more toxic than d_1 as well. The order between d_4 and d_5 in comparison to d_3 is not known because the dose of drug A decreases whilst the dose of drug B increases. Similarly the order between d_2 and d_4 is unknown. Assessing all these potential order toxicity relationships leaves five possible orderings.

1. $d_1 \rightarrow d_2 \rightarrow d_3 \rightarrow d_4 \rightarrow d_5 \rightarrow d_6$
2. $d_1 \rightarrow d_2 \rightarrow d_4 \rightarrow d_3 \rightarrow d_5 \rightarrow d_6$
3. $d_1 \rightarrow d_2 \rightarrow d_4 \rightarrow d_5 \rightarrow d_3 \rightarrow d_6$
4. $d_1 \rightarrow d_4 \rightarrow d_2 \rightarrow d_3 \rightarrow d_5 \rightarrow d_6$
5. $d_1 \rightarrow d_4 \rightarrow d_2 \rightarrow d_5 \rightarrow d_3 \rightarrow d_6$

Using the notation of Wages et al. [17], [18], let M denote the number of possible orders and Y be an indicator of a toxicity event. Then for a trial investigating k combinations, d_1, \dots, d_k , the dose for the j th patient, X_j , $j = 1, \dots, n$ can be thought of as random $x_j \in (d_1, \dots, d_k)$. For a specific ordering m , $m = 1, \dots, M$ the toxicity probability $R(d_i)$ is modelled by

$$R(d_i) = \phi_m(d_i, w, \beta) = w\psi_m(d_i, \beta) \quad i = 1, \dots, k; \quad m = 1, \dots, M \quad (2.1)$$

for a weighted dose response model $\phi_m(d_i, w, \beta)$ where $\beta \in (-\infty, \infty)$. The weight, w as defined by Cheung and Chappel [14], is a function of the time-to-event of each patient and is incorporated linearly with the dose toxicity model

ψ so that $0 \leq w \leq 1$. Each patient is followed for a fixed amount of time T . Let U_j represent the time-to-toxicity of patient j . Then for $u \leq T$,

$$P(U_j \leq u) = P(U_j \leq u | U_j \leq T)P(U_j \leq T) \equiv w(u; T)\psi_m(d_i, \beta). \quad (2.2)$$

For simplicity we will refer to the weight function $w(u; T)$ as w . The weight function will have to be decided upon by the trials team, dependent on the scenario, a simple linear function or a more complex adaptive weights function could be utilised. There are also several working dose models which could be used for ψ , Wages et al. [18] present their design with the power parameter model given by

$$\psi_m(d_i, \beta) = \alpha_{mi}^{\exp(\beta)} \quad i = 1, \dots, k; \quad m = 1, \dots, M. \quad (2.3)$$

Here $0 < \alpha_{m1} < \dots < \alpha_{mk} < 1$ are the prior estimates of toxicity probabilities, or skeleton, for each potential ordering. Furthermore, prior probabilities are assigned to each order M to account for any prior information regarding the plausibility of each model such that, $p(m) = \{p(1), \dots, p(M)\}$, where $p(m) \geq 0$ and $\sum_m p(m) = 1$. When all orders are equally likely or there is no prior information available on possible orderings the prior is discretely uniform and would be $p(m) = 1/M$.

A Bayesian framework is used and a prior probability distribution $g(\beta)$ is assigned to the parameter β . The ordering with the largest prior probability is selected as the starting ordering, in the scenario where all priors are equal an ordering is selected at random, subsequently a starting dose is also chosen. After j patients have been entered into the trial data is collected in the form of $\Omega_j = \{x_1, y_1, \dots, x_j, y_j\}$. A weighted likelihood for the parameter β is used to establish running probabilities of toxicity for each treatment combination. The

weighted likelihood under ordering m , is given by

$$\tilde{L}_m(\beta|\Omega_j) = \prod_{l=1}^j \phi_m^{y_l}(x_l, w_l, \beta) \{1 - \phi_m(x_l, w_l, \beta)\}^{(1-y_l)} \quad (2.4)$$

which can be used to generate a summary value $\hat{\beta}_{mj}$ for each ordering. With the likelihood and the data Ω_j , the posterior density for β can be calculated using

$$\tilde{f}_m(\beta|\Omega_j) = \frac{\tilde{L}_m(\beta|\Omega_j)g(\beta)}{\int_{\beta} \tilde{L}_m(\beta|\Omega_j)g(\beta)d\beta} \quad (2.5)$$

This can then be used to establish posterior probabilities of the orderings given the data as

$$\tilde{\pi}(m|\Omega_j) = \frac{p(m) \int_{\beta} \tilde{L}_m(\beta|\Omega_j)g(\beta)d\beta}{\sum_{m=1}^M p(m) \int_{\beta} \tilde{L}_m(\beta|\Omega_j)g(\beta)d\beta}. \quad (2.6)$$

We select the single ordering, h , with the largest posterior probability along with its associated working model $\psi_h(d_i, \beta)$ and generate toxicity probabilities for each dose level. Once the j th patient has been included the posterior probability of DLT can be calculated for d_i so that

$$\hat{R}(d_i) = \psi_h(d_i, \hat{\beta}_{hj}); \hat{\beta}_h = \int_{\beta} \beta \tilde{f}_h(\beta|\Omega_j)d\beta. \quad (2.7)$$

In turn, the dose level $x_j \in \{d_1, \dots, d_k\}$ assigned to the $(j+1)$ th patient is the dose, d_i , which minimises

$$\triangle(\hat{R}(d_i), \theta) = |\hat{R}(d_i) - \theta|, \quad i = 1, \dots, k \quad (2.8)$$

where θ is the target toxicity rate. Similarly, once all patients have been recruited and observed and the trial ends, the target dose (TD θ) is the dose, d_i , which minimises (2.8).

2.3 PO-TITE-CRM in ADePT-DDR

The decision to implement PO-TITE-CRM into ADePT-DDR was made by Piers Gaunt (PG) after discussions with other statisticians Kristian Brock (KB) and Daniel Slade (DS), as well as the chief investigator and other co-investigators. The design was chosen as the toxicity probabilities of the dose levels weren't monotonically increasing which restricts the use of most early phase designs such as the CRM. Additionally, the design also handles late-onset toxicities which would be an issue in ADePT-DDR due to the treatment involving radiotherapy. The availability of software to conduct the trial was also a factor that was considered. The R package 'pocrm' [22] only provides a means for implementing the PO-CRM design but the easy accessibility to this code meant that it could be extended to include the TITE component.

The intended use of this design is for dose-finding in combinations of therapies, as this is the source of the partial ordering issue. ADePT-DDR however, is a unique implementation of the design as even though it involves a combination of therapies (radiotherapy and AZD6783) the dose of radiotherapy is fixed and dose-finding is only planned for AZD6783. PO-TITE-CRM is still applicable in this case as the design includes combinations of dose and duration for AZD6783 which are partially ordered.

A two-stage PO-TITE-CRM will be used to find the TD25 of AZD6783. This will be determined by dose-limiting toxicities evaluated by Common Terminology Criteria for Adverse Events (CTCAE) v5.0 and Radiation Therapy Oncology Group (RTOG) late toxicity score. The binary DLT events are pre-defined by a variety of grade 3-4 adverse events notably, haematological, cardiovascular and gastrointestinal/hepatic toxicities as well as significant non-haematological events and specific treatment-related toxicities. DLTs will be monitored for the duration of treatment (seven weeks) and throughout the

follow-up period. The total follow-up period post treatment is 52 weeks, so patients will spend a total of 59 weeks in the trial.

A maximum of 60 patients will be recruited for the dose-finding aspect of this trial and up to 20 patients as controls. Controls will be utilised to make comparisons for secondary outcomes such as survival and efficacy. Control patients will only be receiving radiotherapy, the dose of which is fixed at 70Gy/35F. Cohorts of three patients will be recruited and assigned to dose levels chosen by the PO-TITE-CRM. Controls will be recruited in the interim period between the recruitment of the third patient in a cohort and the completion of the minimum follow-up period.

2.3.1 Partial Ordering in Practice

Each patient entered into ADePT-DDR will receive fixed dose radiation, totalling 70 Gy in 35 fractions over seven weeks. For the dose-finding aspect we investigate six doses of AZD6783 detailed in table 2.2. Treatment dose and duration to be selected for dose level 3 will be determined based on a combination of data observed, adverse events and compliance. The issue of partial ordering is illustrated in Figure 2.2 inspired from plots by Wages et al. [18]. The doses to be used in this trial are detailed in their appropriate box. Additionally, each dot represents a potential dose combination which theoretically could be investigated. The combinations are colour coordinated to indicate where partial ordering exists in this dose combination space. Doses across the same colour (each diagonal) cannot be distinguished from each other in terms of probability of toxicity. However, it forms a hierarchy in which doses of the same colour can be thought of as less/more toxic than doses in another colour i.e the red dose levels would have a higher probability of toxicity than the yellow dose levels. It is clear that dose levels 2a and 2b would be considered more toxic

than dose level 1 due to the increase in treatment duration and treatment dose respectively. When comparing 2a and 2b it is unknown whether the increase in dose or duration will be more toxic. Hence there are two possible orderings for ADePT-DDR.

$$1. d_{-1} \rightarrow d_0 \rightarrow d_1 \rightarrow d_{2a} \rightarrow d_{2b} \rightarrow d_3$$


$$2. d_{-1} \rightarrow d_0 \rightarrow d_1 \rightarrow d_{2b} \rightarrow d_{2a} \rightarrow d_3$$

TABLE 2.2: ADePT-DDR dose-levels and duration of treatment for AZD6783.

Dose Level	AZD6783 Daily dose (mg BD)	Weeks	Duration (days)	Radiotherapy
-1	20	1	5	70 Gy/ 35 F
0	20	1&4	10	70 Gy/ 35 F
1	40	1&4	10	70 Gy/ 35 F
2a	40	1,2,4&5	20	70 Gy/ 35 F
2b	80	1&4	10	70 Gy/ 35 F
3	120	1&4	10	70 Gy/ 35 F
	80	1,2,4&5	20	70 Gy/ 35 F

Traditionally, dose-finding trials for combinations would select dose levels to form a 'path' through the dose combination space such that each subsequent dose level was logically more toxic. This avoids the issue of partial ordering but means doses of interest or effective dose combinations may be missed or not investigated. Specifically, for ADePT-DDR this allows two 'paths' from dose level 1 extending to 2a and 2b. In terms of dose level 3 only one of the doses in that tier will be investigated, it was unclear as to which dose level would be best due to a lack of historical data. Even though dose level 3 is not yet specified in terms of modelling and simulations it was treated as singular dose. This was done as clinicians thought that it would be unlikely that we'd reach these doses and that the probability of toxicity between them would be similar.

FIGURE 2.2: ADePT-DDR dose levels across dose and duration.



Preliminary designs of the trial included only five dose levels and planned to use dose level 0 as the starting dose. During the trial design phase it was decided a new lower dose (dose level -1) would be introduced to allow for de-escalation if the initial dose was found to be too toxic. Dose escalation/de-escalation for subsequent cohorts would be determined from the two-stage

PO-TITE-CRM. A two-stage design allows for escalation according to a pre-defined escalation scheme similar to a '3+3' design. The first stage dictates that if no DLT's are observed in the current cohort the dose allocated to the next cohort is the following dose in the escalation scheme. Dose levels continue to be incremented in this fashion until the first DLT is observed. In stage two, dose levels are determined by the PO-TITE-CRM.

Typically CRM designs begin by testing the first patient, or cohort, at the prior guess of TD or at a lower dose to be safe. However, clinicians may have safety concerns beginning the trial at higher dose levels as well as escalating to higher dose levels without testing lower ones. Investigators in ADePT-DDR expressed similar concerns as such a two-stage design was adopted. The escalation scheme used in stage one of ADePT-DDR will follow that of the first ordering ($d_{-1} \rightarrow d_0 \rightarrow d_1 \rightarrow d_{2a} \rightarrow d_{2b} \rightarrow d_3$). If patients in the first cohort (assigned to dose level 0) don't experience a DLT the next cohort will be allocated to dose level 1 and then if no DLTs are observed again the third cohort will be allocated to dose level 2a and so on and so forth. The dose escalation scheme was determined based on the prior probabilities of toxicity generated for each dose level.

Information elicited from the investigators helped generate prior probabilities of toxicity for each dose level. They believed that dose level 2b would be the TD25 with 2a being less toxic. This was used in conjunction with the `getprior` function from the `dfcrm` R package [27] which yielded priors of 0.012, 0.036, 0.084, 0.157, 0.25 and 0.355 for dose levels -1, 0, 1, 2a, 2b and 3 respectively. The half-width of the indifference interval was set at 0.05. The indifference interval is an interval in which the toxicity probability of the selected dose will eventually fall. Prior probabilities are also required for the plausibility of each model and even though the clinicians think that 2b will be more toxic than 2a there is no clear evidence and still a lot of uncertainty. As such it is

sensible to assume a plausibility probability of 0.5 for each ordering, implying both orders are equally likely to be the true ordering of these dose levels.

2.3.2 The TITE component


The observation window for this trial will be up to a year post-treatment as the combination of radiotherapy with AZD6738 is anticipated to cause late-onset toxicity. The Acute DLT observation period is 12 weeks (84 days) post radiotherapy end with a minimum of 8 weeks (56 days) for the last patient of each cohort. However, patients will continuously be monitored for occurrence of DLT for at least 12 weeks (84 days), i.e. at least 12 weeks (84 days) from the end of radiotherapy. The full window will last for 52 weeks (365 days) post-treatment.

The TITE component incorporates a weighting contribution for each patient dependent on how long that patient has been evaluable in the study. This allows a patient to be evaluated once they have been observed for the minimum DLT period of 8 weeks (56 days). The weighting at this point is 60% rising to 80% at 12 weeks (84 days). A patient will not contribute fully to the model until they have completed 52 weeks (365 days) follow up (or have experienced a DLT at any stage in which case they will be weighted as a whole contribution). Linear weighting functions will be employed for any patient with a length of follow up between these three time points. One weight function to calculate weights between 8-12 weeks and another for weights between 12-52 weeks. For the weighting function $w(u; t_1, t_2, t_3)$ where u is the time-to-toxicity of patient j and t_1, t_2, t_3 is the time time period with values 8, 12 and 52 respectively. Then for $t_1 \leq u \leq t_3$

$$w(u; t_1, t_2, t_3) = 0.6 + 0.2 \frac{\min(0, \min(u, t_2) - t_1)}{t_2 - t_1} + 0.2 \frac{\max(0, u - t_2)}{t_3 - t_2}. \quad (2.9)$$

All patients will have a minimum weight of 60% as that is the prescribed weighting to the minimum follow up period before dose escalation/de-escalation decisions can be made. For each additional week the patient is observed, without a DLT occurring, between weeks 8 and 12 their weighting increases by 5%. Similarly for each week between 12 and 52 weeks, without a DLT, weighting increases by 0.5%. Figure 2.3 illustrates the weight function and how the weight changes for patients dependent on how long they have been followed-up.

FIGURE 2.3: Weights of patients who have not experienced a DLT across the observation window.



The TITE-CRM originally presented by Cheung and Chappel [14] did not incorporate a minimum follow-up period and their design allowed for the continual recruitment of patients whenever they became available. There are some practical considerations which make this infeasible in ADePT-DDR. The model

would need to be run each time a new patient entered the study which requires statistical input hence the introduction of cohorts. Clinicians may also have safety concerns if we see rapid recruitment at the start of the trial and the model keeps escalating so we impose a minimum follow-up period. Initially this was set at 12 weeks (at 80% weighting) however, statisticians AP and PG pointed out that dose escalation/de-escalation decisions would have to take place 19 weeks (7 weeks treatment and 12 weeks follow-up) after recruitment of the third patient in the cohort. Dependent on the recruitment rates this could extend the duration of the trial and negates the benefits of using a TITE design. The investigators also agreed this was too long and settled on lowering this period to 8 weeks (at 60% weighting) whilst also including the original 12 week weighting of 80%.

2.3.3 Stopping Rules

A practical modification was included to allow for early stopping of the trial if there is sufficient evidence that the TD has been reached. Sufficient evidence is achieved once 15 patients (five cohorts) have been treated at the same dose level and the model allocates that dose level again to a sixth cohort. This rule evolved from the original designs of the trial which involved 30 patients with a dose expansion cohort to ensure at least 15 patients were treated at the TD.

Initial simulations highlighted the inadequacy of these design parameters as operating characteristics for various scenarios were poor, specifically in terms of correct TD selection. Clinicians explained the inclusion of the dose expansion cohort was to ensure the dose-finding aspect of the trial did not take a large amount of time whilst also allowing safety to be assessed at the TD. In order to ensure that a reasonable amount of patients would be treated at the

TD, the trial wouldn't take longer than necessary and operating characteristics improved, the sample size was increase and this rule was introduced.

A rule was also implemented to allow for early termination of the trial in the case of excess toxicity at the lowest dose. If the probability of DLT at the lowest dose is higher than 0.35 with a probability of 80% and has been tested the trials safety committee will be alerted and will recommend if the trial should be stopped. As the trial starts at dose level 0, which is not the lowest dose, it is possible for the trial to recommend terminating without ever allocating patients to the lowest dose level. As such it was decided early termination would only occur once at least 3 patients (1 cohort) have been allocated dose level -1.

An approximate estimate of the variance was calculated using methodology presented by O'Quigley and Shen [28]. The observed information matrix is obtained by taking the second derivative of the likelihood (eq. 2.4) which is then used to calculate the variance $v(\hat{\beta}_j)$, for estimate β_j which becomes more accurate with larger sample sizes. After each cohort, we sample many times from a normal distribution with parameters based on the estimate of β_j and its variance. These samples are then plugged into our dose-toxicity model to ascertain the probability of toxicity at the lowest dose. The trial will be recommended to stop if it breaks the rule based on the criteria above.

2.3.4 Operating Characteristics

Simulations were continually utilised during the design process of the trial to assess how various changes impact the overall performance. These changes to design features such as the sample size, weight function and stopping rules helped inform decisions which led to the final design.

Functions from pocrm package in R [22], [23] were modified in order to perform simulations and conduct the trial. The majority of work involved integrating the TITE component and the stopping rules into the code. In standard CRM designs a binary outcome for toxicity is generated for each patient based on a pre-specified true DLT rates for the dose they are assigned. Adding the TITE component means the time the toxicity occurs also has to be generated, the simulation must also track this time and incorporate this information into the PO-TITE-CRM model when it needs to make dose allocation decisions for the next cohort. We defined multiple scenarios to reflect various real life possibilities in order to assess the designs performance.

Standard scenarios to run involve adjusting the true DLT rates to reflect each dose being the TD25. For each of these we calculate the probability of selecting each dose as the TD25. It would be expected the dose with the highest probability of being selected has its true DLT rate set at 25% to match the target rate. A high probability of selection for the correct dose implies the design works well in the specified scenario. Additional characteristics such as the average number of patients at each dose level are also investigated. This can be used to look at how many patients may potentially be allocated to a toxic dose. It is also necessary to consider performance when all doses are too toxic, here we would want the design to recommend stopping early. Usually the true DLT rates used to define these scenarios abide by the monotonicity assumption. Due to the partial ordering we consider scenarios in which the true DLT rates follow both orders. For trials with a large amount of orders it may be unfeasible to run so many simulations. However, as ADePT-DDR only has two orders we explored all scenarios for each ordering.

We simulated 10000 trials for each scenario using the finalised design detailed in section 2.3. Simulations were based on the assumption that the trial would recruit one patient per month. The occurrence of DLT's were randomly

generated for patients in each cohort using a Bernoulli distribution with the probability set at the true DLT rate for the cohorts assigned dose level in the specific scenario. For patients who had a DLT occur, the time at which the DLT occurred was randomly generated using a uniform distribution which spanned the start of treatment to the end of follow-up. The simulations presented in Tables 2.3 and 2.4 took approximately 5 hours and 53 minutes to run. The Monte Carlo standard error for probabilities estimated by 10000 simulations is $\sqrt{0.5 \times 0.5 / 10000} = 0.5\%$.

Table 2.3 details simulations for eight scenarios to test the performance of the PO-TITE-CRM design using true DLT rates which reflect the first ordering. We analyse scenarios where each dose is the TD25 (scenarios 1-6) and when all doses are too toxic (scenario 8). Additionally, we also investigate performance under conditions where the probability of DLT is fairly similar between doses (scenario 7). This is a notoriously difficult circumstance for CRM designs to deal with as the limited number of patients and events at each dose make it hard to accurately estimate toxicity probabilities if they are similar. Simulation results for ordering 2 are shown in Table 2.4 where dose level 2a is considered more toxic than 2b. This is achieved by altering the true DLT rates so 2b has a lower probability of DLT compared to 2a.

Ideally we want the probability of selection for the dose allocated at TD25 to be as high as possible and greater than other dose levels. For scenarios 1-7 the TD25 is highlighted in bold along with results from the simulations. However, for scenario 8 where all doses are too toxic we expect the trial to terminate early, here 'stop' should be selected and its associated probability of stopping is shown in bold.

In scenarios 1 - 6 (Table 2.3), this design correctly selects the TD25 with probabilities between 43% and 78%, under the assumption 2b is more toxic than 2a. Likewise, for the ordering where 2a is more toxic than 2b, scenarios

TABLE 2.3: Operating Characteristics of the two-stage PO-TITE-CRM (with true DLT rates that imply 2b is more toxic than 2a) based on 10000 simulated trials. Definitions: DLT: Dose-limiting toxicity. P(select): Probability of selecting a dose as the TD25.

Scenario	Prior DLT	Dose Levels						Stop
		-1	0	1	2a	2b	3	
1: TD25 @-1	True DLT rate	0.25	0.4	0.45	0.5	0.55	0.6	0.0849
	P(select)	0.68	0.18	0.05	0.01	0	0	
	% of patients	39	32	20	6	3	0	
	Mean number of patients	10.17	8.46	5.33	1.67	0.69	0.07	
2: TD25 @0	True DLT rate	0.12	0.25	0.4	0.45	0.5	0.55	0.0079
	P(select)	0.23	0.51	0.2	0.03	0.02	0	
	% of patients	17	35	29	11	6	1	
	Mean number of patients	5.24	10.48	8.75	3.4	1.83	0.26	
3: TD25 @1	True DLT rate	0.09	0.12	0.25	0.4	0.45	0.5	4e-04
	P(select)	0.02	0.2	0.55	0.14	0.09	0.01	
	% of patients	4	20	34	23	16	3	
	Mean number of patients	1.22	6.41	10.97	7.23	5.14	1.02	
4: TD25 @2a	True DLT rate	0.06	0.09	0.12	0.25	0.4	0.45	1e-04
	P(select)	0	0.02	0.22	0.48	0.23	0.05	
	% of patients	1	12	20	31	25	11	
	Mean number of patients	0.47	3.88	6.74	10.43	8.2	3.5	
5: TD25 @2b	True DLT rate	0.03	0.06	0.09	0.12	0.25	0.4	0
	P(select)	0	0	0.02	0.3	0.43	0.25	
	% of patients	1	10	12	24	28	25	
	Mean number of patients	0.25	3.36	4.15	8.17	9.33	8.33	
6: TD25 @3	True DLT rate	0.01	0.03	0.06	0.09	0.12	0.25	0
	P(select)	0	0	0	0.09	0.13	0.78	
	% of patients	0	10	11	18	18	42	
	Mean number of patients	0.1	3.13	3.49	5.46	5.6	13.14	
7: Equal steps in DLT rate	True DLT rate	0.05	0.1	0.15	0.2	0.25	0.3	3e-04
	P(select)	0	0.03	0.12	0.31	0.28	0.26	
	% of patients	2	13	18	26	23	19	
	Mean number of patients	0.55	4.03	5.72	8.32	7.15	5.96	
8: All toxic	True DLT rate	0.5	0.6	0.65	0.7	0.75	0.8	0.7404
	P(select)	0.26	0	0	0	0	0	
	% of patients	56	26	15	2	0	0	
	Mean number of patients	9.05	4.27	2.4	0.37	0.04	0	

9-14 (Table 2.4) have probabilities between 43% and 78% of correctly selecting the TD25. Correct selection probabilities are generally higher when the TD25 is at the first and last dose levels compared to dose levels 2a and 2b. However, these dose levels are still chosen with the highest probability as the TD25 in their given scenarios. For scenarios 7 and 15, the probabilities of toxicity are equally spaced, approximately 5% apart. This is a relatively difficult scenario for dose-finding studies to handle. The probability of selecting the TD25 is 28%

TABLE 2.4: Operating Characteristics of the two-stage PO-TITE-CRM (with true DLT rates that imply 2a is more toxic than 2b) based on 10000 simulated trials. Definitions: DLT: Dose-limiting toxicity. P(select): Probability of selecting a dose as the TD25.

Scenario	Prior DLT	Dose Levels						Stop
		-1	0	1	2a	2b	3	
		0.01	0.04	0.08	0.16	0.25	0.35	
9: TD25 @-1	True DLT rate	0.25	0.4	0.45	0.55	0.5	0.6	0.0827
	P(select)	0.67	0.19	0.05	0	0.01	0	
	% of patients	39	32	20	6	3	0	
	Mean number of patients	10.19	8.43	5.27	1.6	0.68	0.07	
10: TD25 @0	True DLT rate	0.12	0.25	0.4	0.5	0.45	0.55	0.0074
	P(select)	0.23	0.52	0.2	0.02	0.02	0	
	% of patients	18	36	29	11	6	1	
	Mean number of patients	5.24	10.64	8.82	3.16	1.85	0.24	
11: TD25 @1	True DLT rate	0.09	0.12	0.25	0.45	0.4	0.5	4e-04
	P(select)	0.02	0.2	0.55	0.09	0.14	0.01	
	% of patients	4	20	34	21	17	3	
	Mean number of patients	1.16	6.43	11.07	6.83	5.6	1.07	
12: TD25 @2a	True DLT rate	0.06	0.09	0.12	0.25	0.15	0.45	2e-04
	P(select)	0	0.01	0.08	0.44	0.33	0.14	
	% of patients	1	11	16	30	24	18	
	Mean number of patients	0.48	3.78	5.24	10.1	7.9	6.07	
13: TD25 @2b	True DLT rate	0.03	0.06	0.09	0.35	0.25	0.4	0
	P(select)	0	0	0.15	0.31	0.43	0.11	
	% of patients	1	11	18	30	28	14	
	Mean number of patients	0.25	3.5	5.9	9.82	9.14	4.54	
14: TD25 @3	True DLT rate	0.01	0.03	0.06	0.12	0.09	0.25	0
	P(select)	0	0	0	0.13	0.09	0.78	
	% of patients	0	10	11	19	16	43	
	Mean number of patients	0.1	3.13	3.51	5.88	5.06	13.13	
15: Equal steps in DLT rate	True DLT rate	0.05	0.1	0.15	0.25	0.2	0.3	2e-04
	P(select)	0	0.02	0.12	0.32	0.27	0.26	
	% of patients	2	13	19	27	22	18	
	Mean number of patients	0.54	4.02	5.93	8.56	6.89	5.75	
16: All toxic	True DLT rate	0.5	0.6	0.65	0.75	0.7	0.8	0.7273
	P(select)	0.27	0	0	0	0	0	
	% of patients	56	27	15	2	0	0	
	Mean number of patients	9.01	4.28	2.39	0.38	0.05	0	

and 32% for orderings 1 and 2 respectively and even if the performance is poor the correct dose is still likely to be selected. In scenarios 8 and 16, where all the doses are too toxic, the design very seldom allocates patients higher than the first three doses and there is a high chance (74% and 73% respectively) that the trial will recommend early stopping.

Additionally, we assess designs based on how doses are allocated to patients. Designs may correctly select the TD however, this could be undesirable

and unethical if the majority of patients are over dosed at the more toxic dose levels. The average number and the percentage of patients at each dose level, for each scenario, is recorded in Tables 2.3 and 2.4.

The percentage of patients treated at the TD25 ranges between 23% and 43% for each scenario under both orderings. The design also allocates the most patients on average to the TD25 apart from in scenario 7. In this case more patients were allocated to the next lowest dose, we have already discussed the difficulties of this scenario so this characteristic is not too concerning. The mean number of patients recruited for scenarios 1-6 is 26, 30, 32, 33, 34 and 31 respectively. Similarly for scenarios 9-14 its 26, 30, 32, 34, 33 and 31. Even though we allow for up to 60 patients the majority of trials terminate early based on the pre-defined rules for selecting the TD25. This information is presented in Table 2.5 which also shows how often the max sample size is reached from the 10000 trials for each scenario.

TABLE 2.5: Summary of simulated patient numbers for each scenario.

Scenario	Max no. of patients	No. of times max reached	Mean no. of patients
1: TD25 @-1	60	21	26.38
2: TD25 @0	60	8	29.97
3: TD25 @1	60	5	32.01
4: TD25 @2a	60	12	33.22
5: TD25 @2b	60	6	33.60
6: TD25 @3	60	2	30.92
7: Equal steps	60	1	31.74
8: All toxic	54	1	16.14
9: TD25 @-1	60	17	26.23
10: TD25 @0	60	11	29.95
11: TD25 @1	60	6	32.15
12: TD25 @2a	60	7	33.56
13: TD25 @2b	60	3	33.16
14: TD25 @3	60	8	30.81
15: Equal steps	60	2	31.69
16: All toxic	51	1	16.11

Overall, the simulation results show the specification of this design performs relatively well in a number of scenarios. We have shown there is a high

probability of the trial stopping early if all dose-levels are too toxic. We have also shown the design behaves in an appropriate manner when there is a lack of disparity between dose-levels in terms of toxicity. Finally, we have demonstrated that regardless of the ordering we observe the PO-TITE-CRM has a high probability of selecting the correct dose. There are a number of limitations to the operating characteristics presented here which are due to the specification of the simulations and trial design. Section 2.4 explores and discusses these limitations in more detail.


2.4 Exploring other designs

The operating characteristics presented in section 2.3.4 provide an insight into how the trial design operates and its effectiveness at selecting the TD25. However, several factors impact the results seen here. These factors can be grouped into two main categories, limitations with the simulations performed or the trial design.

To simulate various scenarios the true DLT rates are adjusted to reflect the TD25 being at different dose levels. There is no formal process to select these values as such their selection is fairly arbitrary. We set one dose level as the TD25 with lower and higher dose levels set at lower and higher DLT rates respectively. Figure 2.4 illustrates the dose levels for scenarios in Table 2.3 where dose level 2b is more toxic than 2a. The DLT rates cover some possible scenarios and account for a range of plausible values. However, these true DLT rates may not accurately reflect what we observe once the trial begins. Also, the relationship between the rates and the dose levels may not be similar to what we use in the simulations. Multiple other scenarios could be investigated but it would still be impossible to account for all possible variations which may occur. Hence when evaluating the performance of a design it is important to

note the scenario in which it is being evaluated and whether or not the design performs as expected and to an adequate level. For ADePT-DDR, the design produces reasonable operating characteristics under each scenario.

FIGURE 2.4: True DLT rates used for each of the scenarios where dose level 2b is more toxic than 2a. The dotted red line represents the target dlt rate of 25% (TD25).



The original methodological papers by Wages et al. [17], [18] only provide simulations for their examples using true DLT rates that are monotonically increasing which represented one of their possible orderings. We cannot tell how their design would perform under scenarios from different orderings. It is unclear as to why this wasn't examined. Initially, in ADePT-DDR we followed suit and only produced simulations under a monotonically increasing DLT rate (order where 2b is more toxic than 2a, Table 2.3). However, as we are unclear on the ordering of 2a and 2b there is a possibility that 2b is less toxic. So those initial simulations would not provide an accurate assessment of the design

in that circumstance. This was the main motivation for running scenarios in Table 2.4, in which we see the design performs at a similar level regardless of the partial ordering. ADePT-DDR is a simple case of partial ordering as there are only two possible orderings and six dose-levels. For trials with higher numbers of orderings or dose-levels the number of scenarios that would have to be evaluated would increase which may be infeasible. Here it may be more beneficial to choose a handful of scenarios from multiple different orderings to cover a wider range of possible outcomes for the trial to assess the design.

There are various features in this trial design that impact how it performs. The partial ordering caused by dose levels 2a and 2b adds complexity to design. If one of these dose-levels were to be removed or a normal ordering was assumed a standard TITE-CRM design could be used instead. However, this would take away from what the trial is trying to discover. This trial also has a long follow-up period due to potential late-onset toxicities and in turn, will have a long duration. The TITE component will allow for the duration to be a lot shorter than it would be otherwise. TITE-CRM designs allow for patients to be recruited sequentially and allocated a dose based on available information from patients already in the trial. The design for ADePT-DDR uses cohorts of three and a minimum follow-up period. The dose-escalation decisions will only be made every third patient after a specific amount of time. This is done for safety and practicality reasons but means that some patients may not be able to enter the trial and it also loses some of the benefits of the TITE-CRM. We also have a sample size of 60 patients but include a stopping rule for when a consensus is reached which means we often don't recruit the maximum sample size. Further simulations were produced to investigate how these features affected the trial. Tables 2.6 and 2.7 compares selection probabilities from the ADePT-DDR trial design with five alternative designs based on the two different orderings. Figures 2.5 and 2.6 visualise the result from each of these tables

respectively. 10000 trials were simulated for each scenario which took 62 hours and 22 minutes to complete.

1. TITE-CRM design. This design assumes that partial ordering doesn't exist and that dose-level 2b is more toxic than 2a. A TITE-CRM is used instead of a PO-TITE-CRM. All other stopping rules and details remain the same.
2. PO (Partial Ordering design). This design removes the TITE component and uses a PO-CRM as detailed by [17]. This requires the removal of the minimum follow-up period so all dose allocation decisions are made once all 3 patients in a cohort have been observed for the full follow-up period of one year. All other stopping rules and details remain the same.
3. $N = 30$. This design uses a fixed sample size of 30 patients and removes the stopping rule for reaching consensus. The analysis is still conducted using the PO-TITE-CRM. All other stopping rules and details remain the same.
4. $N = 60$. This design uses a fixed sample size of 60 patients and removes the stopping rule for reaching consensus. The analysis is still conducted using the PO-TITE-CRM. All other stopping rules and details remain the same.
5. $CS = 1$. This design uses a cohort size (CS) of one. All other stopping rules remain the same.

The TITE-CRM performs similarly to our original design for scenarios 1-8 where 2a is assumed less toxic than 2b (Table 2.3). We see increases in probability selection for scenarios 4 and 5 where the target dose is at 2a and 2b respectively. This increase in performance can be attributed to the fact that the


TABLE 2.6: Selection probabilities of the TD25 and expected trial duration (in months) for the PO-TITE, TITE and PO-CRM designs as well as modified PO-TITE-CRM designs for scenarios 1-8 (where 2b is considered more toxic than 2a) based on 10000 simulated trials.

Scenario	CRM details	Prior DLT	Dose Levels						Stop	Duration	Mean N
			-1	0	1	2a	2b	3			
			0.01	0.04	0.08	0.16	0.25	0.35			
1: TD25 @-1		True DLT rate	0.25	0.4	0.45	0.5	0.55	0.6			
	PO-TITE	P(select)	0.68	0.18	0.05	0.01	0	0	0.08	57.61	26.38
	TITE	P(select)	0.7	0.21	0.05	0.01	0	0	0.03	59.55	27.46
	PO	P(select)	0.59	0.18	0.04	0.01	0	0	0.19	132.19	22.11
	N = 30	P(select)	0.67	0.19	0.05	0.01	0	0	0.08	60.02	27.72
	N = 60	P(select)	0.78	0.12	0.01	0	0	0	0.09	106.61	53.62
	CS = 1	P(select)	0.63	0.17	0.04	0.01	0.01	0	0.15	87.62	22.44
2: TD25 @0		True DLT rate	0.12	0.25	0.4	0.45	0.5	0.55			
	PO-TITE	P(select)	0.23	0.51	0.2	0.03	0.02	0	0.01	64.06	29.97
	TITE	P(select)	0.22	0.54	0.2	0.04	0.01	0		63.5	29.65
	PO	P(select)	0.2	0.54	0.21	0.02	0.01	0	0.02	163.53	27.79
	N = 30	P(select)	0.23	0.5	0.21	0.03	0.02	0	0.01	63.26	29.52
	N = 60	P(select)	0.15	0.69	0.14	0.01	0	0	0.01	116.29	59
	CS = 1	P(select)	0.25	0.46	0.19	0.03	0.02	0.01	0.04	105.41	27.59
3: TD25 @1		True DLT rate	0.09	0.12	0.25	0.4	0.45	0.5			
	PO-TITE	P(select)	0.02	0.2	0.55	0.14	0.09	0.01		67.74	32.01
	TITE	P(select)	0.01	0.2	0.54	0.2	0.04	0		64.44	30.18
	PO	P(select)	0.02	0.16	0.59	0.14	0.07	0.01		178.12	30.43
	N = 30	P(select)	0.01	0.19	0.52	0.15	0.12	0.01		64.04	29.96
	N = 60	P(select)	0	0.14	0.7	0.1	0.06	0		117.89	59.89
	CS = 1	P(select)	0.05	0.18	0.51	0.14	0.09	0.02	0.01	114.18	30.13
4: TD25 @2a		True DLT rate	0.06	0.09	0.12	0.25	0.4	0.45			
	PO-TITE	P(select)	0	0.02	0.22	0.48	0.23	0.05		69.91	33.22
	TITE	P(select)	0	0.02	0.21	0.56	0.18	0.03		66.99	31.59
	PO	P(select)	0	0.02	0.22	0.49	0.22	0.05		189.69	32.52
	N = 30	P(select)	0	0.01	0.22	0.45	0.27	0.05		64.1	29.99
	N = 60	P(select)	0	0	0.18	0.64	0.18	0.01		118.01	59.96
	CS = 1	P(select)	0.03	0.02	0.22	0.45	0.2	0.08		113.77	30.01
5: TD25 @2b		True DLT rate	0.03	0.06	0.09	0.12	0.25	0.4			
	PO-TITE	P(select)	0	0	0.02	0.3	0.43	0.25		70.6	33.6
	TITE	P(select)	0	0	0.02	0.22	0.56	0.19		68.34	32.34
	PO	P(select)	0	0	0.03	0.26	0.44	0.27		194.41	33.38
	N = 30	P(select)	0	0	0.03	0.3	0.43	0.24		64.11	29.99
	N = 60	P(select)	0	0	0	0.24	0.61	0.14		118.08	59.99
	CS = 1	P(select)	0.01	0	0.02	0.26	0.4	0.3		110.26	29
6: TD25 @3		True DLT rate	0.01	0.03	0.06	0.09	0.12	0.25			
	PO-TITE	P(select)	0	0	0	0.09	0.13	0.78		65.78	30.92
	TITE	P(select)	0	0	0	0.03	0.23	0.73		65.6	30.82
	PO	P(select)	0	0	0	0.08	0.12	0.8		183.48	31.4
	N = 30	P(select)	0	0	0	0.1	0.14	0.75		64.12	30
	N = 60	P(select)	0	0	0	0.06	0.1	0.85		118.09	60
	CS = 1	P(select)	0.01	0	0	0.06	0.1	0.83		92.92	23.98
7: Equal steps		True DLT rate	0.05	0.1	0.15	0.2	0.25	0.3			
	PO-TITE	P(select)	0	0.03	0.12	0.31	0.28	0.26		67.25	31.74
	TITE	P(select)	0	0.03	0.14	0.3	0.32	0.21		65.22	30.61
	PO	P(select)	0.01	0.03	0.14	0.3	0.27	0.25		186.6	31.96
	N = 30	P(select)	0	0.02	0.12	0.3	0.3	0.25		64.07	29.97
	N = 60	P(select)	0	0	0.07	0.33	0.35	0.25		118.04	59.97
	CS = 1	P(select)	0.03	0.02	0.09	0.24	0.23	0.39		101.81	26.55
8: All toxic		True DLT rate	0.5	0.6	0.65	0.7	0.75	0.8			
	PO-TITE	P(select)	0.26	0	0	0	0	0	0.74	39.19	16.14
	TITE	P(select)	0.37	0	0	0	0	0	0.63	44.65	19.17
	PO	P(select)	0.13	0	0	0	0	0	0.86	70.38	10.92
	N = 30	P(select)	0.23	0	0	0	0	0	0.77	41.35	17.34
	N = 60	P(select)	0.13	0	0	0	0	0	0.87	47.35	20.68
	CS = 1	P(select)	0.33	0	0	0	0	0	0.67	46.43	10.51

TABLE 2.7: Selection probabilities of the TD25 and expected trial duration (in months) for the PO-TITE, TITE and PO-CRM designs as well as modified PO-TITE-CRM designs for scenarios 9-16 (where 2a is considered more toxic than 2b) based on 10000 simulated trials.

Scenario	CRM details	Prior DLT	Dose Levels						Stop	Duration	Mean N
			-1	0	1	2a	2b	3			
			0.01	0.04	0.08	0.16	0.25	0.35			
9: TD25 @-1		True DLT rate	0.25	0.4	0.45	0.55	0.5	0.6			
	PO-TITE	P(select)	0.67	0.19	0.05	0	0.01	0	0.08	57.35	26.23
	TITE	P(select)	0.7	0.21	0.05	0	0	0	0.03	59.34	27.34
	PO	P(select)	0.58	0.17	0.04	0	0	0	0.2	131.46	21.98
	N = 30	P(select)	0.68	0.18	0.04	0	0.01	0	0.09	59.85	27.62
	N = 60	P(select)	0.78	0.13	0.01	0	0	0	0.09	106.5	53.56
	CS = 1	P(select)	0.62	0.16	0.05	0	0.01	0	0.15	87.6	22.43
10: TD25 @0		True DLT rate	0.12	0.25	0.4	0.5	0.45	0.55			
	PO-TITE	P(select)	0.23	0.52	0.2	0.02	0.02	0	0.01	64.02	29.95
	TITE	P(select)	0.21	0.56	0.19	0.02	0.01	0		63.57	29.69
	PO	P(select)	0.2	0.54	0.2	0.01	0.02	0	0.02	163.06	27.7
	N = 30	P(select)	0.24	0.51	0.2	0.02	0.03	0	0.01	63.35	29.57
	N = 60	P(select)	0.15	0.7	0.13	0	0.01	0	0.01	116.22	58.96
	CS = 1	P(select)	0.25	0.47	0.19	0.02	0.03	0.01	0.04	105.04	27.49
11: TD25 @1		True DLT rate	0.09	0.12	0.25	0.45	0.4	0.5			
	PO-TITE	P(select)	0.02	0.2	0.55	0.09	0.14	0.01		68	32.15
	TITE	P(select)	0.01	0.22	0.58	0.14	0.04	0		64.86	30.41
	PO	P(select)	0.03	0.17	0.59	0.09	0.12	0.01		177.68	30.35
	N = 30	P(select)	0.02	0.2	0.51	0.1	0.16	0.01		64.03	29.95
	N = 60	P(select)	0	0.14	0.71	0.05	0.1	0		117.88	59.88
	CS = 1	P(select)	0.05	0.17	0.53	0.09	0.13	0.02	0.01	113.98	30.07
12: TD25 @2a		True DLT rate	0.06	0.09	0.12	0.25	0.15	0.45			
	PO-TITE	P(select)	0	0.01	0.08	0.44	0.33	0.14		70.52	33.56
	TITE	P(select)	0	0.02	0.14	0.26	0.45	0.14		67.23	31.73
	PO	P(select)	0.01	0.02	0.09	0.45	0.28	0.15		194.42	33.38
	N = 30	P(select)	0	0.01	0.08	0.42	0.35	0.14		64.09	29.98
	N = 60	P(select)	0	0	0.03	0.57	0.33	0.07		117.98	59.94
	CS = 1	P(select)	0.02	0.01	0.07	0.4	0.32	0.18		113.02	29.8
13: TD25 @2b		True DLT rate	0.03	0.06	0.09	0.35	0.25	0.4			
	PO-TITE	P(select)	0	0	0.15	0.31	0.43	0.11		69.8	33.16
	TITE	P(select)	0	0.01	0.26	0.36	0.28	0.1		65.95	31.02
	PO	P(select)	0	0.01	0.14	0.34	0.43	0.1		190.68	32.7
	N = 30	P(select)	0	0	0.15	0.3	0.44	0.11		64.12	30
	N = 60	P(select)	0	0	0.1	0.29	0.57	0.04		118.07	59.99
	CS = 1	P(select)	0.02	0	0.14	0.27	0.4	0.17		112.27	29.58
14: TD25 @3		True DLT rate	0.01	0.03	0.06	0.12	0.09	0.25			
	PO-TITE	P(select)	0	0	0	0.13	0.09	0.78		65.58	30.81
	TITE	P(select)	0	0	0	0.04	0.21	0.74		65.66	30.86
	PO	P(select)	0	0	0	0.13	0.07	0.8		183.9	31.47
	N = 30	P(select)	0	0	0	0.13	0.11	0.75		64.12	30
	N = 60	P(select)	0	0	0	0.09	0.06	0.85		118.09	60
	CS = 1	P(select)	0.01	0	0	0.1	0.07	0.82		93.1	24.03
15: Equal steps		True DLT rate	0.05	0.1	0.15	0.25	0.2	0.3			
	PO-TITE	P(select)	0	0.02	0.12	0.32	0.27	0.26		67.17	31.69
	TITE	P(select)	0	0.03	0.19	0.27	0.28	0.24		65.01	30.49
	PO	P(select)	0	0.03	0.16	0.32	0.26	0.23		186.35	31.92
	N = 30	P(select)	0	0.02	0.13	0.3	0.29	0.25		64.1	29.99
	N = 60	P(select)	0	0	0.07	0.36	0.31	0.25		117.97	59.94
	CS = 1	P(select)	0.03	0.02	0.09	0.23	0.23	0.4		102.22	26.67
16: All toxic		True DLT rate	0.5	0.6	0.65	0.75	0.7	0.8			
	PO-TITE	P(select)	0.27	0	0	0	0	0	0.73	39.14	16.11
	TITE	P(select)	0.37	0	0	0	0	0	0.63	44.23	18.94
	PO	P(select)	0.13	0	0	0	0	0	0.87	69.79	10.81
	N = 30	P(select)	0.23	0	0	0	0	0	0.77	41.18	17.24
	N = 60	P(select)	0.13	0	0	0	0	0	0.87	47.29	20.64
	CS = 1	P(select)	0.33	0	0	0	0	0	0.67	45.76	10.31

FIGURE 2.5: Plot of the simulation results presented in Table 2.6 detailing selection probabilities for multiple designs across scenarios 1-8 (where 2b is considered more toxic than 2a).



partial ordering no longer exists as we have assumed an ordering. The lower selection probabilities for the PO-TITE-CRM can be seen as the price to pay for the uncertainty of not knowing the order of 2a and 2b. However, the TITE-CRM underperforms in scenarios 9-16 where 2a is assumed more toxic than 2b. Specifically, scenarios 12 and 13 where it fails to identify the TD25 the majority of the time.

The PO-CRM design without a TITE component also performs similarly except for scenarios 1, 8, 9 and 16 where the trial stops more regularly for excess toxicity at the lowest dose. This would be because patients complete the full follow-up window before the next dose allocation decision is made. In a TITE setting a new cohort could be recruited before patients in previous cohorts experience a DLT. The main difference between these designs is the

FIGURE 2.6: Plot of the simulation results presented in Table 2.7 detailing selection probabilities for multiple designs across scenarios 9-16 (where 2a is considered more toxic than 2b).



trial duration. Without the TITE component the trial duration is significantly longer, with the average length ranging from 70 to 195 months compared to 39 to 71 months for PO-TITE-CRM.

The design with a fixed sample size of 30 performs is comparable to our design with the sample size of 60 and the consensus stopping rule. With a sample size of 30 selection probabilities are only 2-5% lower. For the design with 60 patients, we see much improved operating characteristics with selection probabilities ranging from 31% to 85% for the various scenarios. Even though our original design specifies a sample size of 60 we rarely ever reach it as we often stop for consensus hence why this design performs better. The trade-off here is trial duration. Recruitment and follow-up under the constraints of these simulations will take much longer compared to our specification which is not ideal

for an early-phase trial. Originally our design for a sample size of 30 but as the clinicians wanted a dose expansion cohort we opted to use the consensus rule to ensure a minimum number of patients would be treated at the TD25.

For the design with no cohorts, or cohort size of one, we see somewhat comparable performance to that of the PO-TITE-CRM design. The design performs similarly for scenarios where the TD25 is at the lowest or highest dose level but underperforms for the more complex scenarios in terms of selection probabilities. This discrepancy in performance may be related to how the simulations recruit patients into the trial and the large DLT follow-up period. Meaning more frequent dose allocation decisions are being made each with less available information. This also leads to the no cohorts design having a longer duration. Patients entered into the trial in cohorts of three won't have to wait the full minimum follow-up period between patients within the cohort.

2.5 Discussion

The PO-CRM and PO-TITE-CRM designs offer solutions to the issue of partial ordering where the order of the treatments is only partially known. The original methodology details that this issue commonly arises in trials of multiple agents, where each drug individually may follow the monotonicity assumption but when combined at certain dose levels this may not hold. This issue is typically dealt with by fixing the dose of one of the agents and escalating in the other or escalating in both agents simultaneously. This means certain drug combinations that are clinically relevant may not be investigated or even considered.

Here we have shown that these issues can also arise in other situations. Even though the ADePT-DDR trial uses multiple agents the issue of partial ordering still occurs due to the varying treatment dose and schedule for one

of its agents AZD6738. Implementing the PO-TITE-CRM design allowed us to deal with this issue effectively. There may be other factors or variables in single-agent dose-finding trials that would lead to the issue of partial ordering and would warrant the use of either PO-CRM or PO-TITE-CRM. The small literature review conducted highlighted this may be the first instance of the PO-TITE-CRM design being applied. It is important to note that although this methodology takes into account all the various orderings the main aim is to identify the TD and does not attempt to identify the most correct order.

Compared to other CRM based designs only a few additional pieces of information are required to implement the PO-CRM design. Most importantly is the number of toxicity orderings and prior probabilities for the orders. Dependent on how many dose combinations are available it may not be feasible to investigate all combinations and all orderings. Careful thought and consideration should be given to the combinations and orderings selected which would require input from all relevant investigators. In terms of priors for orderings if no prior information is available all orders should be treated as equally likely to occur. Extending this design to the PO-TITE-CRM requires a fit for purpose weight function and is applied similarly to the TITE-CRM methodology. There is an R package available with functions that can be used to run and simulate a PO-CRM trial. These functions were extended to include weighted dose toxicity models as described in this chapter to implement PO-TITE-CRM into ADePT-DDR. The lack of available software for PO-TITE-CRM specifically may be one of the reasons for its lack of use.

In terms of ADePT-DDR, dose combinations were decided upon by the clinical investigators. The issue of partial ordering was due to the dose-levels 2a and 2b as such this methodology was employed to deal with that scenario. Meaning that this is a very simple example of partial ordering as we only have two possible orderings and six dose levels. The necessity of implementing this

methodology was discussed and whether or not adopting an easier solution by simply altering the dose levels would have been better. Ultimately, the dose levels selected by the clinicians were deemed the most relevant with the TD25 likely to be one of these doses.

Simulations and operating characteristics were the main tools used to assess the designs performance as well as help understand the impact of sample size and stopping rules. This was an iterative process that involved running multiple iterations of simulations under various scenarios until the design was finalised. A key point is that scenarios from simulations should account for each of the possible orderings. ADePT-DDR only has two orderings, we ran scenarios for both. For a trial with a greater number of orderings, this may be unfeasible but at least some scenarios should be assessed to ensure the design is behaving as expected. Overall, the design operating characteristics performed reasonably well even in difficult scenarios.

One limitation of the simulations is how the time-to-event data is generated. The time of DLTs is sampled from a uniform distribution $U(0, 413)$, where the time of the DLT can occur at any time between the patient beginning treatment and the end of follow-up (413 days). Using this uniform distribution implies that a DLT has an equal probability of occurring at any time-point in the observation window. This may not be an accurate representation of what happens in the actual trial. Similar comments can be made about the accrual rate used in the simulations. Here we specified the recruitment of one patient per month which is in no way guaranteed for the actual trial. Wages et al. [18], when presenting this methodology investigated four different applications of the PO-TITE-CRM which used different models to enroll patients and allocate DLTs. Results across these four applications were comparable.

The simulations are also able to instantaneously determine dose-levels for incoming cohorts with all available information. This does not fully reflect the

process in which dose-escalation decisions would be made during the actual running of the trial. The analysis would require a data snapshot and time would have to be spent cleaning the data and determining the next dose-level. Meaning any data from the point of the snapshot would not be included in any dose escalation/de-escalation decisions.

Similarly, there may also be limitations with some of the design choices made concerning to cohort size and sample size. These were investigated alongside a variety of other trial designs that could have been implemented. This was done to validate the choices we made with the design and highlight the differences in operating characteristics due to the varying assumptions and components in the designs. The standard PO-CRM had a much longer average duration due to the lack of TITE component whereas a standard TITE-CRM overall performs better but assumes the ordering of toxicity is known.

2.6 Conclusion

The monotonicity assumption may not hold in some dose-finding trials leading to the issue of partial ordering. This could be due to multiple-agents being investigated or varying factors in single-agent treatment. PO-CRM and PO-TITE-CRM are important trial designs as they address this core issue. ADePT-DDR is a platform trial with its initial component being a dose-finding trial investigating AZD6738 in combination with radiotherapy, of which the toxicity ordering for two of the dose levels for investigation is unknown. The PO-TITE-CRM design allows for us to deal with this issue of partial ordering as well as account for potential late-onset toxicities due to radiotherapy with its TITE component.

We detail the issue of partial ordering and how we implemented the trial design, in what we believe is the first real-world application of this design. A

large amount of simulation work is required to assess the performance of the design. This is often an iterative process to refine decisions that were made and often requires input from both clinical and statistical investigators. We recommend running several varied scenarios for each potential ordering that will be investigated. Finally, we also compared the implementation of PO-TITE-CRM to various other designs.

Bibliography

- [1] N. Stransky, A. M. Egloff, A. D. Tward, *et al.*, “The Mutational Landscape of Head and Neck Squamous Cell Carcinoma,” *Science (New York, N.Y.)*, vol. 333, no. 6046, pp. 1157–1160, Aug. 26, 2011, ISSN: 0036-8075. DOI: 10.1126/science.1208130. pmid: 21798893. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3415217/>.
- [2] Cancer Reaserch UK. (). “Head and neck cancers statistics,” [Online]. Available: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/head-and-neck-cancers>.
- [3] C. E. Round, M. V. Williams, T. Mee, *et al.*, “Radiotherapy Demand and Activity in England 2006–2020,” *Clinical Oncology*, vol. 25, no. 9, pp. 522–530, Sep. 1, 2013, ISSN: 0936-6555. DOI: 10.1016/j.clon.2013.05.005. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/S0936655513002343>.
- [4] D. M. Cognetti, R. S. Weber, and S. Y. Lai, “Head and Neck Cancer: An Evolving Treatment Paradigm,” *Cancer*, vol. 113, pp. 1911–1932, 7 Oct. 1, 2008, ISSN: 0008-543X. DOI: 10.1002/cncr.23654. pmid: 18798532. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2751600/>.
- [5] J. Pignon, J Bourhis, C Domenge, *et al.*, “Chemotherapy added to locoregional treatment for head and neck squamous-cell carcinoma: Three

- meta-analyses of updated individual data," *The Lancet*, vol. 355, no. 9208, pp. 949–955, Mar. 18, 2000, ISSN: 0140-6736. DOI: 10.1016/S0140-6736(00)90011-4. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/S0140673600900114>.
- [6] M. J. O'Connor, "Targeting the DNA Damage Response in Cancer," *Molecular Cell*, vol. 60, no. 4, pp. 547–560, Nov. 19, 2015, ISSN: 1097-2765. DOI: 10.1016/j.molcel.2015.10.040. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/S109727651500831X>.
- [7] A. Chalmers, "Science in Focus: Combining Radiotherapy with Inhibitors of the DNA Damage Response," *Clinical Oncology*, vol. 28, no. 5, pp. 279–282, May 2016, ISSN: 09366555. DOI: 10.1016/j.clon.2016.01.035. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S0936655516000649>.
- [8] L. Mei, J. Zhang, K. He, *et al.*, "Ataxia telangiectasia and Rad3-related inhibitors and cancer therapy: Where we stand," *Journal of Hematology & Oncology*, vol. 12, no. 1, p. 43, Apr. 24, 2019, ISSN: 1756-8722. DOI: 10.1186/s13045-019-0733-6. [Online]. Available: <https://doi.org/10.1186/s13045-019-0733-6>.
- [9] C. Le Tourneau, J. J. Lee, and L. L. Siu, "Dose Escalation Methods in Phase I Cancer Clinical Trials," *JNCI: Journal of the National Cancer Institute*, vol. 101, no. 10, pp. 708–720, May 20, 2009, ISSN: 0027-8874. DOI: 10.1093/jnci/djp079. [Online]. Available: <https://academic.oup.com/jnci/article/101/10/708/969691>.
- [10] J. O'Quigley, M. Pepe, and L. Fisher, "Continual Reassessment Method: A Practical Design for Phase 1 Clinical Trials in Cancer," *Biometrics*, vol. 46, no. 1, pp. 33–48, 1990, ISSN: 0006-341X. DOI: 10.2307/2531628. JSTOR: 2531628.

- [11] A. Rogatko, D. Schoeneck, W. Jonas, *et al.*, "Translation of Innovative Designs Into Phase I Trials," *Journal of Clinical Oncology*, vol. 25, no. 31, pp. 4982–4986, Nov. 1, 2007, ISSN: 0732-183X. DOI: [10.1200/JCO.2007.12.1012](https://doi.org/10.1200/JCO.2007.12.1012). [Online]. Available: <https://ascopubs.org/doi/full/10.1200/JCO.2007.12.1012>.
- [12] C. Chiuzan, J. Shtaynberger, G. A. Manji, *et al.*, "Dose-finding designs for trials of molecularly targeted agents and immunotherapies," *Journal of Biopharmaceutical Statistics*, vol. 27, no. 3, pp. 477–494, May 4, 2017, ISSN: 1054-3406. DOI: [10.1080/10543406.2017.1289952](https://doi.org/10.1080/10543406.2017.1289952). pmid: 28166468. [Online]. Available: <https://doi.org/10.1080/10543406.2017.1289952>.
- [13] K. Brock, V. Homer, G. Soul, *et al.*, "Is more better? An analysis of toxicity and response outcomes from dose-finding clinical trials in cancer," *medRxiv*, p. 2020.08.18.20177212, Aug. 21, 2020. DOI: [10.1101/2020.08.18.20177212](https://doi.org/10.1101/2020.08.18.20177212). [Online]. Available: <https://www.medrxiv.org/content/10.1101/2020.08.18.20177212v1>.
- [14] Y. K. Cheung and R. Chappell, "Sequential Designs for Phase I Clinical Trials with Late-Onset Toxicities," *Biometrics*, vol. 56, no. 4, pp. 1177–1182, 2000, ISSN: 1541-0420. DOI: [10.1111/j.0006-341X.2000.01177.x](https://doi.org/10.1111/j.0006-341X.2000.01177.x). [Online]. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.0006-341X.2000.01177.x>.
- [15] P. F. Thall, R. E. Millikan, P. Mueller, *et al.*, "Dose-Finding with Two Agents in Phase I Oncology Trials," *Biometrics*, vol. 59, no. 3, pp. 487–496, 2003, ISSN: 1541-0420. DOI: [10.1111/1541-0420.00058](https://doi.org/10.1111/1541-0420.00058). [Online]. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/1541-0420.00058>.
- [16] G. Yin and Y. Yuan, "Bayesian dose finding in oncology for drug combinations by copula regression," *Journal of the Royal Statistical Society:*

- Series C (Applied Statistics)*, vol. 58, no. 2, pp. 211–224, 2009, ISSN: 1467-9876. DOI: [10.1111/j.1467-9876.2009.00649.x](https://doi.org/10.1111/j.1467-9876.2009.00649.x). [Online]. Available: <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9876.2009.00649.x>.
- [17] N. A. Wages, M. R. Conaway, and J. O’Quigley, “Continual reassessment method for partial ordering,” *Biometrics*, vol. 67, no. 4, pp. 1555–1563, Dec. 2011, ISSN: 1541-0420. DOI: [10.1111/j.1541-0420.2011.01560.x](https://doi.org/10.1111/j.1541-0420.2011.01560.x). pmid: [21361888](https://pubmed.ncbi.nlm.nih.gov/21361888/).
- [18] —, “Using the time-to-event continual reassessment method in the presence of partial orders,” *Statistics in Medicine*, vol. 32, no. 1, pp. 131–141, Jan. 15, 2013, ISSN: 1097-0258. DOI: [10.1002/sim.5491](https://doi.org/10.1002/sim.5491). pmid: [22806898](https://pubmed.ncbi.nlm.nih.gov/22806898/).
- [19] S. Liu, G. Yin, and Y. Yuan, “BAYESIAN DATA AUGMENTATION DOSE FINDING WITH CONTINUAL REASSESSMENT METHOD AND DELAYED TOXICITY,” *The annals of applied statistics*, vol. 7, no. 4, pp. 1837–2457, Dec. 1, 2013, ISSN: 1932-6157. DOI: [10.1214/13-AOAS661](https://doi.org/10.1214/13-AOAS661). pmid: [24707327](https://pubmed.ncbi.nlm.nih.gov/24707327/). [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3972824/>.
- [20] G. M. Wheeler, M. J. Sweeting, and A. P. Mander, “A Bayesian model-free approach to combination therapy phase I trials using censored time-to-toxicity data,” *Journal of the Royal Statistical Society. Series C, Applied statistics*, vol. 68, no. 2, pp. 309–329, Feb. 2019, ISSN: 0035-9254. DOI: [10.1111/rssc.12323](https://doi.org/10.1111/rssc.12323). pmid: [30880843](https://pubmed.ncbi.nlm.nih.gov/30880843/). [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6420054/>.
- [21] N. A. Wages and M. R. Conaway, “Specifications of a continual reassessment method design for phase I trials of combined drugs,” *Pharmaceutical statistics*, vol. 12, no. 4, 2013, ISSN: 1539-1604. DOI: [10.1002/pst.1575](https://doi.org/10.1002/pst.1575).

- pmid: 23729323. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3771354/>.
- [22] N. A. Wages and N. Varhegyi, "Pocrm: An R-package for phase I trials of combinations of agents," *Computer Methods and Programs in Biomedicine*, vol. 112, no. 1, pp. 211–218, Oct. 2013, ISSN: 1872-7565. DOI: 10.1016/j.cmpb.2013.05.020. pmid: 23871691.
- [23] N. A. Wages, *Pocrm: Dose Finding in Drug Combination Phase I Trials Using PO-CRM*, version 0.12, Mar. 15, 2019. [Online]. Available: <https://CRAN.R-project.org/package=pocrm>.
- [24] N. A. Wages, A. Ivanova, and O. Marchenko, "Practical designs for phase I combination studies in oncology," *Journal of biopharmaceutical statistics*, vol. 26, no. 1, pp. 150–166, 2016, ISSN: 1054-3406. DOI: 10.1080/10543406.2015.1092029. pmid: 26379085. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4720553/>.
- [25] P. Lesueur, J. Lequesne, J.-M. Grellard, *et al.*, "Phase I/IIa study of concomitant radiotherapy with olaparib and temozolomide in unresectable or partially resectable glioblastoma: OLA-TMZ-RTE-01 trial protocol," *BMC Cancer*, vol. 19, Mar. 4, 2019, ISSN: 1471-2407. DOI: 10.1186/s12885-019-5413-y. pmid: 30832617. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6399862/>.
- [26] B. Huang and P. F. Kuan, "Time-to-event continual reassessment method incorporating treatment cycle information with application to an oncology phase I trial," *Biometrical Journal. Biometrische Zeitschrift*, vol. 56, no. 6, pp. 933–946, Nov. 2014, ISSN: 1521-4036. DOI: 10.1002/bimj.201300261. pmid: 24895140.

-
- [27] K. Cheung, *Dfcrm: Dose-Finding by the Continual Reassessment Method*, version 0.2-2.1, Jan. 26, 2019. [Online]. Available: <https://CRAN.R-project.org/package=dfcrm>.
- [28] J. O'Quigley and L. Z. Shen, "Continual Reassessment Method: A Likelihood Approach," *Biometrics*, vol. 52, no. 2, pp. 673–684, 1996, ISSN: 0006-341X. DOI: [10.2307/2532905](https://doi.org/10.2307/2532905). JSTOR: 2532905.