DOCTORAL THESIS

Developments to established dose-finding methodologies for application in trials with complex and innovative designs

Author: Supervisor:

Amit Patel Prof. Lucinda Billingham

Dr. Kristian BROCK

A thesis submitted in fulfillment of the requirements for the degree of Doctor of Philosophy

Institute of Cancer and Genomic Sciences

College of Medical and Dental Sciences

University of Birmingham

July 21, 2020

Abstract

Developments to established dose-finding methodologies for application in trials with complex and innovative designs

by Amit Patel

Insert abstract here...

Acknowledgements

Acknowledge people here ...

Contents

Ał	ostra	ct Control of the Con	iii
Ac	knov	wledgements	v
1	Imp	elementing the PO-TITE-CRM trial design into ADePT-DDR	1
	1.1	Draft Structure	1
	1.2	Introduction	2
		1.2.1 Subsection 1	3
		1.2.2 Subsection 2	3
	1.3	Main Section 2	3
A	App	pendix Title Here	5
Bi	bliog	raphy	7

List of Figures

List of Tables

Chapter 1

Implementing the PO-TITE-CRM trial design into ADePT-DDR

1.1 Draft Structure

- Introduction
 - Basic biological background
 - Main objective of the trial
 - Introduce the new TD notation link to previous trial designs
 - Paragraph on traditional dose finding trial designs 3+3, CRM etc.
 - Methodological issues which arise due to investigating combination of drugs/varying parameters (new concept by Piers)
 - Necessity of time-to-event components for DLTs which may occur later
 - Other possible methodologies in this area which may be of use to solve this problem
 - Mini literature search will do a citation search for both methodology papers (potentially use a table/figure to summarise)
 - Detail whats to come in the chapter

- The PO-TITE-CRM Design
- PO-TITE-CRM in ADePT-DDR
- Modifications to the specification to improve operating characteristics

1.2 Introduction

Worldwide there are approximately 600,000 new cases of Head and Neck Squamous Cell Carcinoma (HNSCC) each year [1]. Of which, 12,000 occur in the UK with the most common forms of treatment being surgery, radiotherapy and/or [2]. Radiotherapy is essential for the treatment of cancer, it has been estimated that more than 40% of patients will receive radiotherapy at some point in their treatment [3]. However, despite recent advancements in radiation techniques and the use of of concomitant chemo radiotherapy, patients with solid tumours such as head and neck cancer have suboptimal cure rates [2], [4]. For those with advance HNSCC primary radiotherapy with concurrent chemotherapy is often offered but, it has not been shown to improve survival in patients aged over 70 compared to radiotherapy alone [5]. Therefore, any strategy to improve the efficacy of radiotherapy without increasing toxicity to normal tissue would have a significant impact for patients. DNA damage repair (DDR) inhibition is a potential technique which could be utilised as it potentiates the therapeutic effects of ionising radiation in cancer cells without a substantial increase in acute and late toxicity. Combining radiotherapy with DDR inhibition could improve clinical outcomes for these patients [6].

1.3. Main Section 2

1.2.1 Subsection 1

Nunc posuere quam at lectus tristique eu ultrices augue venenatis. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Aliquam erat volutpat. Vivamus sodales tortor eget quam adipiscing in vulputate ante ullamcorper. Sed eros ante, lacinia et sollicitudin et, aliquam sit amet augue. In hac habitasse platea dictumst.

1.2.2 Subsection 2

Morbi rutrum odio eget arcu adipiscing sodales. Aenean et purus a est pulvinar pellentesque. Cras in elit neque, quis varius elit. Phasellus fringilla, nibh eu tempus venenatis, dolor elit posuere quam, quis adipiscing urna leo nec orci. Sed nec nulla auctor odio aliquet consequat. Ut nec nulla in ante ullam-corper aliquam at sed dolor. Phasellus fermentum magna in augue gravida cursus. Cras sed pretium lorem. Pellentesque eget ornare odio. Proin accumsan, massa viverra cursus pharetra, ipsum nisi lobortis velit, a malesuada dolor lorem eu neque.

1.3 Main Section 2

Sed ullamcorper quam eu nisl interdum at interdum enim egestas. Aliquam placerat justo sed lectus lobortis ut porta nisl porttitor. Vestibulum mi dolor, lacinia molestie gravida at, tempus vitae ligula. Donec eget quam sapien, in viverra eros. Donec pellentesque justo a massa fringilla non vestibulum metus vestibulum. Vestibulum in orci quis felis tempor lacinia. Vivamus ornare ultrices facilisis. Ut hendrerit volutpat vulputate. Morbi condimentum venenatis augue, id porta ipsum vulputate in. Curabitur luctus tempus justo. Vestibulum risus lectus, adipiscing nec condimentum quis, condimentum nec

nisl. Aliquam dictum sagittis velit sed iaculis. Morbi tristique augue sit amet nulla pulvinar id facilisis ligula mollis. Nam elit libero, tincidunt ut aliquam at, molestie in quam. Aenean rhoncus vehicula hendrerit.

Appendix A

Appendix Title Here

Write your Appendix content here.

Bibliography

- [1] N. Stransky, A. M. Egloff, A. D. Tward, A. D. Kostic, K. Cibulskis, A. Sivachenko, G. V. Kryukov, M. Lawrence, C. Sougnez, A. McKenna, E. Shefler, A. H. Ramos, P. Stojanov, S. L. Carter, D. Voet, M. L. Cortés, D. Auclair, M. F. Berger, G. Saksena, C. Guiducci, R. Onofrio, M. Parkin, M. Romkes, J. L. Weissfeld, R. R. Seethala, L. Wang, C. Rangel-Escareño, J. C. Fernandez-Lopez, A. Hidalgo-Miranda, J. Melendez-Zajgla, W. Winckler, K. Ardlie, S. B. Gabriel, M. Meyerson, E. S. Lander, G. Getz, T. R. Golub, L. A. Garraway, and J. R. Grandis, "The mutational landscape of head and neck squamous cell carcinoma," *Science (New York, N.Y.)*, vol. 333, no. 6046, pp. 1157–1160, Aug. 26, 2011, ISSN: 0036-8075. DOI: 10.1126/science.1208130. [Online]. Available: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3415217/ (visited on 07/20/2020).
- [2] C. R. UK. (Feb. 8, 2017). "Head and neck cancers statistics," Cancer Research UK. Library Catalog: www.cancerresearchuk.org, [Online]. Available: https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/head-and-neck-cancers (visited on 07/21/2020).
- [3] C. E. Round, M. V. Williams, T. Mee, N. F. Kirkby, T. Cooper, P. Hoskin, and R. Jena, "Radiotherapy demand and activity in england 2006–2020," *Clinical Oncology*, vol. 25, no. 9, pp. 522–530, Sep. 1, 2013, ISSN: 0936-6555.

 DOI: 10.1016/j.clon.2013.05.005. [Online]. Available: http://www.

8 Bibliography

sciencedirect.com/science/article/pii/S0936655513002343 (visited on 07/21/2020).

- [4] D. M. Cognetti, R. S. Weber, and S. Y. Lai, "Head and neck cancer: An evolving treatment paradigm," *Cancer*, vol. 113, no. 7, pp. 1911–1932, Oct. 1, 2008, ISSN: 0008-543X. DOI: 10.1002/cncr.23654. [Online]. Available: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2751600/ (visited on 07/21/2020).
- [5] J. Pignon, J Bourhis, C Domenge, and L Designé, "Chemotherapy added to locoregional treatment for head and neck squamous-cell carcinoma: Three meta-analyses of updated individual data," *The Lancet*, vol. 355, no. 9208, pp. 949–955, Mar. 18, 2000, ISSN: 0140-6736. DOI: 10.1016/S0140-6736(00)90011-4. [Online]. Available: http://www.sciencedirect.com/science/article/pii/S0140673600900114 (visited on 07/20/2020).
- [6] A. Chalmers, "Science in focus: Combining radiotherapy with inhibitors of the DNA damage response," *Clinical Oncology*, vol. 28, no. 5, pp. 279–282, May 2016, ISSN: 09366555. DOI: 10.1016/j.clon.2016.01.035. [Online]. Available: https://linkinghub.elsevier.com/retrieve/pii/S0936655516000649 (visited on 07/21/2020).