

UUID: B25D2818-B3C6-49F0-81E6-224488C1E926
TCGA-5M-AAT6-01A-PR Redacted



Date:

SPECIMEN: Terminal ileum, cecum, appendix, ascending and transverse colon
PROCEDURE: Right hemicolectomy

MACROSCOPY

- Product of right hemicolectomy.
- Segment of terminal ileum, cecum, segment of ascending colon and appendix.
- The serosa is focally irregular, and displays multiple nodules in addition to a perforation 2.0 x 1.5 cm large, which is continuous to an ulcerative, vegetative luminal lesion measuring 8.0 x 8.0 cm, located 8.0 cm from the nearest surgical margin.
- The lesion penetrates the colon wall to the serosa.
- 11 lymph nodes were dissected from the pericolic fat; sizes ranging from 0.3 to 0.6 cm.

MICROSCOPIC EXAMINATION

- Moderately differentiated tubular adenocarcinoma of the ascending colon;
- Microscopic tumor extension: invasion of pericolic soft tissues;
- Perineural invasion: present;
- Lymph-vascular invasion: present;
- Moderate desmoplastic stromal reaction;
- Surgical margins free of neoplastic involvement;
- Number of lymph nodes examined: 11;
- Number of lymph nodes involved: 10;
- Stage: pT4c/d, pN2

PATHOLOGISTS

CLUE ICD-O-3
path Adenocarcinoma NOS 8140/3
Adenocarcinoma, tubular 8211/3
Site: Ascending colon C18.2
p 2/1/14

Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HIPAA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Noted		☒
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	BRH	Date Reviewed: 1/24/14