


Theoretical task 3.

Recommendations: all solutions should be short, mathematically strict (unless qualitative explanation is needed), precise with respect to the stated question and clearly written. Solutions may be submitted in any readable format, including images

1. Consider linear regression task in one-dimensional space. $(x_i, y_i)_{i=1}^N$ is a training dataset and for object $i : x_i \in [-1, 1]$ is a feature, y_i is an answer we want to predict, $y_i^* = kx_i + b$ is our prediction. At the picture below you can see three different plots of the prediction error $(y - y^*)$ against x . Which of these plots cannot be obtained if least squares method is used to train a regression model?



2. Ridge regression – model like usual linear regression, but with add l_2 regularization. Denote $X : n \times m$ matrix of feature descriptions of objects, $y : n$ – vector of target variable, β – vector of weights (model's parameters), λ – hyperparameter of regularization. Show, that optimal β is expressed like

$$\beta = (X^T X + \lambda I)^{-1} X^T y,$$

where I - identity matrix size of $m \times m$

You can assume that

$$\nabla_{\beta} L = -2X^T y + 2X^T X \beta$$

and

$$\frac{dx^T A x}{dx} = (A + A^T)x, \text{ if } A = A^T \rightarrow \frac{dx^T A x}{dx} = 2Ax$$

3. For linear regression task with one variable $\hat{y} = \beta_0 + \beta_1 x$ show, that

$$\beta_1 = \frac{\sum_{i=1}^n x_i y_i - \frac{\sum_{i=1}^n y_i \sum_{i=1}^n x_i}{n}}{\sum_{i=1}^n x_i^2 - \frac{\sum_{i=1}^n x_i \sum_{i=1}^n x_i}{n}} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

where $\bar{x} = \frac{1}{n} \sum_i x_i$

4. While fit linear regression we solve this optimal task::

$$\|Xw - y\|^2 \rightarrow \min_w.$$

From lectures we know analytical solution of this task:

$$w^* = (X^T X)^{-1} X^T y$$

Proof that solution don't use derivatives and optimization methods. Proof a linear regression solution based on geometric considerations.. To do this, use the fact that Xw defines a linear space spanned by column vectors of X .

Hint: the shortest distance between a point and a plane is the length of a perpendicular dropped from a point to a plane.

5. Consider the dataset with the following features: height, age(from 10 to 45), sex(=1 for females, =0 for males). You would like to build a regression model to estimate person's height based on other features. Write down the regression model that would consider all the following facts:

- on average the height of males and females is different,
- at the age of 25 human rate of growth dramatically drops down, but still assumed to be linear.

Explain your answer.