

## INSTRUCTIONS FOR USING THE KEPLER FITS v2.0 FILE VStar PLUG IN

This contains instructions for first time use. After that you can skip directly to the URL for the Kepler Data Search and Retrieval form in the Mikulsi Archive for Space Telescopes (MAST).

The inputs to the Kepler data Plug in are Kepler light curve .FITS files. It does not use complete Kepler data records or the pixel files. Here's how you get a Kepler FITS file:


1. Go the Kepler Official Site Home page: <http://kepler.nasa.gov/>




2. Click on the Science tab and select “for Scientists” in the drop down




3. Select MAST data archive



- Click on the highlighted text Mikulski Data Archive for Space Telescopes (MAST), which takes you to the MAST Kepler home page (not back to the Kepler home page)



- Select the Data Search tab which brings up the search input form



MAST data is linked to Simbad. Therefore, you can use most if not all names for an object found in



Simbad. You can also use coordinates as the input or Kepler ID numbers and a bunch of other fields and qualifiers to narrow the search. Usually you will just use name or coordinates. You can also customize the information in the output table provided by the search, but since what you are after for input into VStar is the light curve, unless there is specific information about the object that you want in the output table, you will normally use the defaults. Notice in particular the form default will retrieve longs (30 minutes) and short (1 minute) cadence data. These are the time periods of the individual 6 and 2/3rds second individual images that were summed on-board into a single equivalent exposure.

- When you click on the search button at the top of the form or at the bottom (not shown in the image) you will get a summary output table that looks like the following for the eclipsing binary V 579 Lyr:

Kepler Data Search Results

Object name **V 579 Lyr** resolved by **Simbad** (via **SANTIA**) to **V\* V579 Lyr (EB\*WU3a)**  
**RA:** 18 50 52.37 **Dec:** 43 40 12.06 (**J2000**)

number of rows returned = 18  
*note: reload page if no results are shown*

Click on top column headers to sort the table on the column contents.  
Click on bottom column headers for more information about the data in that column.  
Click on Ref entries to display list of published papers.  
Click links under Kepler ID to see available Simbad information.  
Click on Condition Flag entries for more information on flag definitions.  
Click on Dataset Name entries to preview information on data set.


Mark	Kepler ID	Investigation ID	Dataset Name	Quarter	RA (J2000)	Dec (J2000)	Target Type	Archive Class	Ref	Actual Start Time	Actual End Time	Release Date	R Mag	J Mag	K Mag	2MASS ID	2MASS conflict flag	Tet	Log G	Metallicity	E(B-V)	Radius	Total PM	G-R color	Module
u	781200	EX	KPLR00781200-2009131105131	0	18 50 52.371	+43 40 12.07	LC	CLC	2	2009-05-02 00:54:56	2009-05-11 17:51:31	2010-06-15 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	2
u	781200	EX	KPLR00781200-2009160043257	1	18 50 52.371	+43 40 12.07	LC	CLC	2	2009-05-13 00:15:49	2009-06-15 11:32:57	2010-06-15 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	2
u	781200	EX	KPLR00781200-200915160929	2	18 50 52.371	+43 40 12.07	LC	CLC	1	2009-06-20 00:21:09	2009-09-16 23:09:29	2011-02-02 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	16
u	781200	EX	KPLR00781200-200915015506	3	18 50 52.371	+43 40 12.07	LC	CLC	0	2009-09-18 17:19:58	2009-12-16 23:55:06	2011-09-23 09:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	24
u	781200	EX	KPLR00781200-2010070809131	4	18 50 52.371	+43 40 12.07	LC	CLC	0	2009-12-19 21:03:56	2010-03-19 16:53:31	2012-01-07 12:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	10
u	781200	EX	KPLR00781200-2010070809206	5	18 50 52.371	+43 40 12.07	LC	CLC	0	2010-03-20 23:47:15	2010-06-23 15:50:26	2012-01-07 12:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	2
u	781200	EX	KPLR00781200-2010061421224	6	18 50 52.371	+43 40 12.07	LC	CLC	0	2010-06-24 22:44:09	2010-09-22 19:01:09	2012-04-19 12:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	16
u	781200	EX	KPLR00781200-2010033117224	7	18 50 52.371	+43 40 12.07	LC	CLC	0	2010-09-25 16:08:24	2010-12-21 01:10:41	2012-07-28 12:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	24
u	781200	EX	KPLR00781200-201101133229	8	18 50 52.371	+43 40 12.07	LC	CLC	0	2011-01-06 20:45:08	2011-03-14 20:18:16	2012-07-28 12:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	10
u	781200	EX	KPLR00781200-201117082412	9	18 50 52.371	+43 40 12.07	LC	CLC	0	2011-03-21 00:24:25	2011-06-26 10:10:29	2012-11-19 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	2
u	781200	EX	KPLR00781200-201117111324	10	18 50 52.371	+43 40 12.07	LC	CLC	0	2011-06-27 08:14:32	2011-09-28 18:22:51	2012-10-28 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	16
u	781200	EX	KPLR00781200-2012040120808	11	18 50 52.371	+43 40 12.07	LC	CLC	0	2011-09-29 16:56:19	2012-01-04 19:50:25	2012-10-28 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	24
u	781200	EX	KPLR00781200-2012080547206	12	18 50 52.371	+43 40 12.07	LC	CLC	0	2012-01-05 21:49:54	2012-03-28 12:32:43	2012-10-28 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	10
u	781200	EX	KPLR00781200-2012120603101	13	18 50 52.371	+43 40 12.07	LC	CLC	1	2012-03-29 05:42:32	2012-06-27 13:18:20	2012-10-28 00:00:00	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	2
u	781200	EX	KPLR00781200-20121217125493	14	18 50 52.371	+43 40 12.07	LC	CLC	0	2012-06-28 15:17:47	2012-10-03 19:40:10	2012-02-06 03:51:51	12.783	11.548	12.799	18505237+4340120	0	4644	4.534	-0.271	0.034	0.801	0.000	0.847	16

- Choose the dataset of interest. By default they are listed in increasing order of the quarter in which they were taken. Start and end times of the datasets are shown in other columns. Choose the dataset by clicking on the dataset name. DO NOT USE THE CHECK BOXES IN THE MOST LEFTHAND COLUMN. The column footers (not shown in the image) link to explanations of the terms used in the various columns. For example the Target Type column in the displayed rows tells you that the datasets shown contain long cadence data.


When you click on the dataset name a preview page opens similar to the following for V 579 Lyr:



8. Now comes the easy part. Select New Star From Kepler FITS v2.0 from the file drop down list



9. Then select the file you saved from Kepler in the normal file browse and select dialog. If you have eliminated the .txt extension, you will have to select all files as the file type. The default shows .txt files but not .fits files.



10. If you want to add Kepler Data files together you can use the “Add to Current Check box at the top right of the open file dialog.

Kepler data is NOT restricted to Kepler exoplanet objects of interest. You can find data on essentially any star in the field, certainly any star that was in the Kepler input file. Also ASAS imaged most of the field for about 6 months in 1999 in preparation for the Kepler mission. You can cross index ASAS and Kepler IDs for objects in the field.

There is a wealth of information about the Kepler data sets on the Kepler and MAST websites. If you are going to work with Kepler data two documents that are invaluable are Kepler Data Field Descriptions and the Kepler Archive Manual. Keep in mind that Kepler satellite data is in units of flux. The archive includes flux and magnitude data from follow up sources but caution is needed because the sources have different offsets and are not transformed to a common scale.

Notes:

1. **PDCSAP\_FLUX** [32-bit floating point] – The flux contained in the optimal aperture in electrons per second after the PDC module has applied its detrending algorithm to the PA light curve. To better understand how PDC manipulated the light curve, read Section 2.3.1.1 [of the Kepler Archive Manual] and see the PDCSAPFL keyword in the header.

Brad Walter, WBY

Rev E

20140610

**Revision History**

Rev	Date	Description
A	6/7/2014	Initial Release
B	6/8/2014	Added item 10
C	6/9/2014	Added Footnote 1, changed text in item 10 describing locations of the “Search” buttons and added revision history
D	6/9/2014	Correct item 10
E	6/10/2014	Correct item 7 & add page numbers