



## Statistics for the SDGs - indicators for national priorities



| Name of the indicator          | 2.7.a Prevalence of obesity among adults  |
|--------------------------------|---|
|                                |   |
| Sustainable Development Goal   | Goal 2. Zero hunger   |
| Priority                       | Promotion of healthy lifestyle among adults and children (decrease in proportion of persons with overweight and obesity)  |
| Definition                     | Percentage of population aged 18 years and more with a BMI equal 30 or more.  |
| Unit                           | percent [%]   |
| Available dimentions           | sex   |
| Methodological<br>explanations | Obesity is defined as abnormal or excessive fat accumulation that presents a risk to health. A crude population measure of obesity is the body mass index (BMI). A person with a BMI of 30 or more is generally considered obese.  Body mass index (BMI) is defined as a person's weight in kilograms divided by the square of the person's height in metres (kg/m2). For adults over 20 years old, BMI falls into one of the following categories:  - <18.5 - underweight, - 18.5 - 24.9 - normal weight, - 25.0 - 29.9 - pre-obesity, - 30.0 - 34.9 - obesity class II, - >=40 - obesity class III.  The BMI ranges are based on the effect excessive body fat has on disease and death and are reasonably well related to adiposity. BMI was developed as a risk indicator of disease; as BMI increases, so does the risk for some diseases. Some common conditions related to overweight and obesity include: premature death, cardiovascular diseases, high blood pressure, osteoarthritis, some cancers and diabetes. |
| Data source                    | World Health Organization   |
| Data availability              | Annual data; since 2010   |
| Notes                          |   |

Last update: 12-05-2020, 11:15