


Chapter 6: Strings in Background Fields

Note this chapter is specific to *closed oriented strings*. As such, we will not consider the effects of the boundary.

0. This is not a required problem but it certainly should be ¹. Let's *calculate the β -functions of the nonlinear sigma model*. Here, I will borrow diagrams from the very nice set of TASI lecture notes of Callan and Thorlacius

First, it is worth using a normal coordinate system for the X^μ (one in which all of the Γ symbols vanish and all higher symmetrized Γ symbols also vanish). We want to look at radiative corrections to $\langle T_{++} \rangle$, since they have integrals that are easier to handle than those for $\langle T_{+-} \rangle$. From conservation this will give us the trace anomaly for $\langle T_{+-} \rangle$. We will first look at how G , B affect the trace on a flat worldsheet.


For the graviton contribution to β^G , we have only one diagram



This contributes an anomalous trace of

$$\langle T_{+-} \rangle = \frac{1}{4} R_{\mu\nu} \partial_a X_0^\mu \partial^a X_0^\nu$$

For the B contribution to β^B , we have two such diagrams:



These contribute anomalous traces of:

$$-\frac{1}{16} H_{\mu\rho\sigma} H_{\nu\rho\sigma} \partial_a X_0^\mu \partial^a X_0^\nu, \quad \frac{1}{8} \nabla^\lambda H_{\mu\nu\lambda} \epsilon^{ab} \partial_a X_0^\mu \partial_b X_0^\nu$$

respectively.

The dilaton contribution *also* affects the trace on the flat world sheet (even though it does not couple at $R = 0$), by affecting the stress energy tensor as it is defined by varying the action w.r.t. the metric. Kiritsis has worked this out before and shown that the dilaton contributes $(\partial_a \partial_b - g_{ab} \square) \Phi$ to the stress energy tensor, from which we get a dilaton contribution of $\square_\xi \Phi(X(\xi))$ to the trace. Using covariant expressions for the D'Alambertian we arrive at a contribution

$$\nabla_\mu \nabla_\nu \Phi(X_0) \partial_a X_0^\mu \partial^a X_0^\nu - \frac{1}{2} \nabla^\lambda \Phi(X_0) H_{\mu\nu\lambda}(X_0) \partial_a X_0^\mu \partial_b X_0^\nu \epsilon^{ab}$$

Combining all of this together, we see that we will get the β -functions:

$$\beta^G = R_{\mu\nu} - \frac{1}{4} H_{\mu\rho\sigma} H_{\nu}^{\rho\sigma} + 2 \nabla_\mu \nabla_\nu \Phi, \quad \beta^B = -\frac{1}{2} \nabla^\lambda H_{\lambda\mu\nu} + \nabla^\lambda \Phi H_{\lambda\mu\nu}.$$

As pointed out, these are not quite the RG beta functions, but around the fixed point, they capture the correct first order behavior. In particular their vanishing will mean that we have no Weyl anomaly.

Now we need to account for the effects of a curved worldsheet geometry. We can account for this by looking at a $\langle T_{+-} T_{+-} \rangle$ correlator:

$$\frac{\delta}{\delta \phi(\xi)} \langle T_{+-}(0) \rangle_{e^\phi \delta_{ab}} = -\frac{1}{4\pi} \langle T_{+-}(\xi) T_{+-}(0) \rangle_{\delta_{ab}} \quad (1)$$

Again we can get this by first looking at $\langle T_{++} T_{++} \rangle$ and appealing to conservation. The Weyl anomaly comes from this diagram:

¹After seeing the details of this calculation, I can understand why it was omitted.




This gives $\langle T_{+-} T_{+-} \rangle = \frac{\pi D}{12} \square \delta^{(2)}(\xi)$. Here we have a factor of D coming from each degree of freedom. This can be used to integrate equation (1) to yield:

$$\langle T_{+-} \rangle = -\frac{D}{48} \square \phi = \frac{D}{24} \sqrt{\gamma} R$$

Note that the ghosts (which are otherwise decoupled) will here contribute their factor of -26 .

We also now need to consider *two-loop* contributions of G, B to the TT correlator. The following diagrams contribute:



The calculations here are very involved, but will precisely give us

$$\frac{\alpha}{8} \left(-R + \frac{H^2}{12} \right)$$

Finally, the dilaton both modifies the energy-momentum tensor, giving rise to a tree-level propagator contribution to the two-point function:



This contributes $\langle T_{+-}^{dil} T_{+-}^{dil} \rangle = \pi \alpha' (\nabla \Phi)^2 \square \delta^{(2)}(\xi)$ which will integrate to give a factor of $\frac{\alpha'}{2} (\nabla \Phi)^2 \sqrt{\gamma} R$.

Also, the dilaton gives a loop-contribution to the unmodified energy-momentum tensor:



Which contributes the term $\langle T_{+-}^{dil} T_{+-}^{dil} \rangle = -\pi \alpha' \square \Phi \square \delta^{(2)}(\xi)$ which will integrate to give a factor of $-\frac{\alpha'}{2} \square \Phi \sqrt{\gamma} R$.

Altogether this gives:

$$\beta^\Phi = D - 26 + \frac{3}{2} \alpha' \left[4(\nabla \Phi)^2 - 4 \square \Phi - R + \frac{1}{12} H^2 \right].$$

as required.

1. Each β -function of a coupling constant G, B, Φ as given in **6.1.5**, **6.1.6**, **6.1.7** is $\frac{\delta}{\delta\phi}$ of that coupling constant, since our scaling $\mu = e^\phi \Rightarrow \log \mu = \phi$. Since

$$T_a^a = \frac{\beta^\Phi}{12} R^{(2)} + \frac{1}{2\ell_s^2} (\beta_{\mu\nu}^G g^{\alpha\beta} + \beta_{\mu\nu}^B \varepsilon^{\alpha\beta}) \partial_\alpha X \partial_\beta X$$

The change in effective action under an infinitesimal Weyl transformation $\delta g^{\alpha\beta} = -g^{\alpha\beta} \delta\phi$ is

$$\delta \log Z = -\delta S = \frac{1}{4\pi} \int d^2\xi \sqrt{g} T_a^a \delta\phi = \frac{1}{4\pi} \int d^2\xi \left[\frac{\beta^\Phi}{12} \sqrt{g} R^{(2)} + \frac{1}{2\ell_s^2} (\beta_{\mu\nu}^G \sqrt{g} g^{ab} + \beta_{\mu\nu}^B \varepsilon^{ab}) \partial_a X \partial_b X \right] \delta\phi$$

We can integrate this to get the change after a finite conformal transformation:

$$\frac{1}{4\pi} \int d^2\xi \left[\frac{\beta^\Phi}{12} \sqrt{g} \left(R^{(2)} \phi + \frac{1}{2} g^{ab} \nabla_a \phi \nabla_b \phi \right) + \frac{\phi}{2\ell_s^2} (\beta_{\mu\nu}^G \sqrt{g} g^{ab} + \beta_{\mu\nu}^B \varepsilon^{ab}) \partial_a X \partial_b X \right]$$

this vanishes, of course, when all beta functions are zero. When β^G, β^B are zero we can show (exercise 3) that β^Φ is a constant, and we recover the Liouville action from before.

2. First write G explicitly in the action:

$$S = \frac{1}{2\kappa^2} \int d^D x \sqrt{-\det G} e^{-2\Phi} \left[R + 4G^{\alpha\beta} \nabla_\alpha \Phi \nabla_\beta \Phi - \frac{1}{12} G^{\alpha\delta} G^{\beta\epsilon} G^{\gamma\zeta} H_{\alpha\beta\gamma} H_{\delta\epsilon\zeta} + 2 \frac{26-D}{3\ell_s^2} \right]$$

The classical equations of motion from varying the action with respect to G give

$$\begin{aligned} 0 &= \overbrace{R_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi - 4\nabla_\mu \Phi \nabla_\nu \Phi - 2G_{\mu\nu} \square \Phi + 4G_{\mu\nu} (\nabla \Phi)^2}^{R \text{ variation}} + \overbrace{4\nabla_\mu \Phi \nabla_\nu \Phi}^{(\nabla \Phi)^2 \text{ variation}} \\ &\quad \underbrace{-\frac{1}{4} H_{\mu\rho\sigma} H_\nu^{\rho\sigma}}_{H^2 \text{ variation}} - \underbrace{\frac{1}{2} G_{\mu\nu} \left(R + 4(\nabla \Phi)^2 - \frac{1}{12} H^2 + \frac{2}{3\ell_s^2} (26-D) \right)}_{\sqrt{-\det G} \text{ variation}} \\ &= \underbrace{R_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi - \frac{1}{4} H_{\mu\rho\sigma} H_\nu^{\rho\sigma}}_{:=\beta_{\mu\nu}^G} - \frac{1}{2} G_{\mu\nu} \left(R - 4(\nabla \Phi)^2 + 4\square \Phi - \frac{1}{12} H^2 + 2 \frac{26-D}{3\ell_s^2} \right) \end{aligned} \quad (2)$$

With respect to B we get:

$$-\frac{1}{12} e^{-2\Phi} (2(\delta_{B\mu\nu} (\partial_\alpha B_{\beta\gamma} + 2 \text{ perms.})) H^{\alpha\beta\gamma}) \xrightarrow{IBP} \frac{2 \times 3}{12} \nabla^\alpha (e^{-2\Phi} H_{\alpha\mu\nu}) \Rightarrow \underbrace{-\frac{1}{2} \nabla^\alpha (e^{-2\Phi} H_{\alpha\mu\nu})}_{:=\beta_{\mu\nu}^B} = 0$$

Finally, with respect to Φ we get:

$$0 = -2 \left(R + 4(\nabla \Phi)^2 - \frac{1}{12} H^2 + 2 \frac{26-D}{3\ell_s^2} \right) - 8\square \Phi + 16(\nabla \Phi)^2 = -2 \underbrace{\left(R - 4(\nabla \Phi)^2 + 4\square \Phi - \frac{1}{12} H^2 + 2 \frac{26-D}{3\ell_s^2} \right)}_{:= -\frac{2}{3\ell_s^2} \beta^\Phi}$$

The term in parentheses is the same as the term in parentheses the bottom line of (2). This agrees with **Polchinski 3.7.21** (with appropriate conventions adopted)

$$\delta S = -\frac{1}{2\kappa^2} \int d^D x \sqrt{-\det G} e^{-2\Phi} \left[\delta G^{\mu\nu} \left(\beta_{\mu\nu}^G - \frac{1}{2} G_{\mu\nu} \frac{2}{3\ell_s^2} \beta^\Phi \right) + \delta B^{\mu\nu} \beta_{\mu\nu}^B + 2\delta \Phi \frac{2}{3\ell_s^2} \beta^\Phi \right]$$

3. Let's look at $\frac{2}{3\ell_s^2}\nabla\beta^\Phi$. We get:

$$8\nabla_\nu\Phi\nabla_\mu\nabla^\nu\Phi - 4\Box\nabla_\mu\Phi - \nabla_\mu R + \frac{1}{6}(\nabla_\mu H_{\alpha\beta\gamma})H^{\alpha\beta\gamma}$$

The contracted Bianchi identity $\nabla_\mu R = 2\nabla^\nu R_{\mu\nu}$ together with the vanishing of $\beta_{\mu\nu}^G$ gives:

$$\nabla_\mu R = 2\nabla^\nu R_{\mu\nu} = \frac{1}{2}\nabla^\nu(H_{\mu\rho\sigma}H_\nu^{\rho\sigma}) - 4\Box\nabla_\mu\Phi$$

which in turn gives

$$8\nabla_\nu\Phi\nabla_\mu\nabla^\nu\Phi - \frac{1}{2}\nabla^\nu(H_{\mu\rho\sigma}H_\nu^{\rho\sigma}) + \frac{1}{6}(\nabla_\mu H_{\alpha\beta\gamma})H^{\alpha\beta\gamma}$$

The fact that H is exact gives us $dH = 0$ so $\partial_{[\alpha}H_{\beta\gamma\delta]} = 0$. The symmetry properties of H imply that summing over the four cyclic permutations of this gives zero. Contracting with the metric then implies a contracted Bianchi-type identity for H , namely that $\nabla^\alpha H_{\alpha\beta\gamma} = 0$.

Using $\beta^B = 0$ together with the Bianchi identity, we have $0 = \nabla^\rho H_{\mu\nu\rho} = 2\nabla^\rho\Phi H_{\mu\nu\rho}$. So we have that H is divergence-free, and $\nabla^\rho\Phi$ dotted with any component of H is zero. This lets us rewrite:

$$\begin{aligned} -\frac{1}{2}\nabla^\nu(H_{\mu\rho\sigma}H_\nu^{\rho\sigma}) &= -\frac{1}{2}H^{\nu\rho\sigma}\nabla_\nu H_{\mu\rho\sigma} \\ \frac{1}{6}\nabla_\mu(H_{\alpha\beta\gamma})H^{\alpha\beta\gamma} &= -\frac{1}{6}H^{\alpha\beta\gamma}(\nabla_\alpha H_{\beta\gamma\mu} - \nabla_\beta H_{\gamma\alpha\mu} + \nabla_\gamma H_{\alpha\beta\mu}) = -\frac{1}{6}H^{\nu\rho\sigma}\nabla_\nu H_{\mu\rho\sigma} \\ \Rightarrow \frac{1}{12\ell_s^2}\nabla_\mu\beta^\Phi &= \nabla_\nu\Phi\nabla_\mu\nabla^\nu\Phi - \frac{1}{12}\nabla^\nu(H_{\mu\rho\sigma}H_\nu^{\rho\sigma}) = -\frac{1}{2}\nabla^\nu\Phi R_{\mu\nu} - \frac{1}{12}\nabla^\nu H_{\mu\nu} \end{aligned}$$

One last step. I am missing something.

This gives that $\nabla_\mu\beta^\Phi = 0$ as required. So $\beta^\Phi = c$ is a constant.

4. We get a linear dilaton giving rise to a Liouville action with $Q = 0$. This is our familiar free massless boson in $2D$ with $1D$ target space. So we get a string propagating in a single dimension.
5. Note that the only relevant parameters are ℓ_s , with units of length, and whatever length scales there are on the manifold, all of which depend on its volume (since its compact) as $V^{1/D}$. In particular $c = \beta^\Phi$ depends on ℓ_s as

$$c = D + O(\ell_s^2/V^{2/D}).$$

I think this is correct, though it is different from Kiristis' equation.

6. Note that a nonzero total flux of H over any closed 3-manifold is incompatible with $H = dB$ for a single-valued B . We can write:

$$e^{\frac{i}{2\pi\ell_s^2}\int_M B} = e^{\frac{i}{2\pi\ell_s^2}\int_N H}$$

where M is the 2D manifold corresponding to the embedding of the world-sheet into the target space and N is any manifold whose boundary is M . We need this to be independent of N , so for any three-cycle M_3 we need:

$$\frac{1}{2\pi\ell_s^2}\int_{M_3} H \in 2\pi\mathbb{Z} \Rightarrow \frac{1}{4\pi^2\ell_s^2}\int_{M_3} H \in \mathbb{Z}$$

7. (a) We have

$$H = 2R^2 \sin^2 \psi \sin \theta \, d\psi \wedge d\theta \wedge d\phi \Rightarrow \int_{S^3} H = 4\pi^2 R^2 \Rightarrow \frac{1}{4\pi^2 \ell_s^2} \int_{S^3} H = \frac{R^2}{\ell_s^2} \in \mathbb{Z}$$

(b) The dilaton is $\Phi = 0$. Using Mathematica, the Ricci tensor is:

$$R_{\mu\nu} = \text{diag}(2, 2 \sin^2 \psi, 2 \sin^2 \psi \sin^2 \theta)$$

Which gives a Ricci scalar of $6/R^2$. From the previous part, $H_{123} = 2R^2 \sin^2 \psi \sin \theta$. From the metric being diagonal, we get that $H_{\mu\nu}^2 := H_{\mu\rho\sigma} H_{\nu}^{\rho\sigma}$ is diagonal. We have

$$H_{\mu\nu}^2 = \text{diag}(8, 8 \sin^2 \psi, 8 \sin^2 \psi \sin^2 \theta) \Rightarrow \beta^G = R_{\mu\nu} - \frac{1}{4} H_{\mu\nu}^2 = 0$$

as desired. Next, $\beta_{\mu\nu}^B = -\frac{1}{2} \nabla^\alpha (H_{\mu\nu\alpha})$. To take a contravariant divergence we divide by the volume element and differentiate, but the volume element is $\sin^2 \psi \sin \theta$ which will give $H/\sqrt{g}$ is a constant, so $\beta_{\mu\nu}$ will vanish.

Lastly, $H^2 = 3!(2R^2)^2/R^6 = 24/R^2$ so that $-R + \frac{1}{12} H^2 = -\frac{4}{R^2}$. Ignoring ghosts, this gives a central charge of:

$$D - 6 \frac{\ell_s^2}{R^2} + O(\ell_s^4) = D - \frac{6}{k} + O(\ell_s^4)$$

as desired.

(c) Without using coordinates, the isometry of S^3 is $G = \text{SO}(4) = [\text{SU}(2) \times \text{SU}(2)]/\mathbb{Z}_2$. To see that equivalence, think of S^3 as the unit quaternions, and take $\text{SU}(2) \times \text{SU}(2)$ act as unit quaternions on the left and right. We get a right G -action by: $x \rightarrow a^{-1}xb$. Note the kernel is the set of $(a, b) \in G$ $ax = xb$ for all x . In particular, for $x = 1$ we get $a = b$ so the kernel lies in the diagonal subgroup. To act trivially on all quaternions, a must be in the center, and for the unit quaternions this is exactly ± 1 . So this is an injection $\varphi : [\text{SU}(2) \times \text{SU}(2)]/\mathbb{Z}_2 \rightarrow \text{SO}(4)$. Since $\text{SO}(4)$ is compact and connected, it is generated by the image of exponentiating $\mathfrak{so}(4)$, and so surjectivity of φ at the level of the Lie algebras (which is true by dimension-counting) implies surjectivity and hence equivalence at the level of Lie groups.

So we see that $\mathfrak{so}(4)$ acting on S^3 is just a simultaneous left and right copy of $\mathfrak{su}(2)$ acting on $\text{SU}(2)$. Thus, we view this as the CFT of a nonlinear sigma model with target space $G = \text{SU}(2)$ and the left, right copies of the $\mathfrak{su}(2)$ action correspond to currents $J = g^{-1}\partial g$ and $\bar{J} = \bar{\partial} g g^{-1}$

We indeed get the central charge $c = \frac{3k}{k+2}$ which has the large k expansion $3 - 6/k + O(1/k^2)$. Since k is a non-negative integer in WZW models, except for the case $k = 0$ corresponding to the trivial CFT, we must have $k \geq 1$, where we get $R \geq \ell_s$.

8. Here the metric has three degrees of freedom and $B_{\mu\nu}, \Phi$ both have only one degree of freedom (which can be spatially varying). H , being a 3-index antisymmetric tensor, must vanish in 1+1D, and so we will always have $\beta^B = 0$. The other two constraints become:

$$0 = \beta_{\mu\nu}^G = \frac{1}{2} R g_{\mu\nu} + 2 \nabla_\mu \nabla_\nu \Phi, \quad 0 = \beta^\Phi = -24 + \frac{3}{2} \ell_s^2 [4(\nabla\Phi)^2 - 4\Box\Phi - R]$$

Translational isometry implies that R, g depend on only the time variable t . The x variable can therefore parameterize either S^1 or $\mathbb{R}$ endowed with constant metric.

Now taking the trace of the first equation implies $R(t) = -2\Box\Phi(x, t)$. Then the second equation will give:

$$\frac{16}{\ell_s^2} = 4(\nabla\Phi(x, t))^2 - 2(\Box\Phi)(t)$$

The only way for this to work is for $R = \Box\Phi = 0$ so that $\nabla\Phi$ can be a constant. We then have $\Phi = \alpha x + \beta t$ so that $\alpha^2 + \beta^2 = 4/\ell_s^2$, and g is Ricci flat everywhere (so we can pick it to be constant). In the case of either $\alpha, \beta = 0$, we can also safely take x, t respectively to be periodic without having Φ be multi-valued.

9. We still have $\beta^B = 0$, but $\beta^G = R_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi$ while $\beta^\Phi = D - 26 + \frac{3}{2}\ell_s^2(4(\nabla\Phi)^2 - 4\Box\Phi - R)$

This can be recast in terms of a new 4D *Ricci flat* metric $ds^2 = F(\phi)d\phi^2 + \phi R^2 d\Omega_3^2$.

Using Mathematica again to take the trace of this gives R_{ij} for $i = j \geq 1$ proportional to $R^2\phi F'(\phi) + 8\phi F(\phi)^2 - R^2 F(\phi)$. Solving this differential equation for F gives

$$F(\phi) = \frac{R^2\phi}{4\phi^2 + R^2 c_1}$$

Setting $c_1 = 0$, $F(\phi) = R^2/4\phi$ will also make R_{00} vanish. Then we can take the dilaton to be zero $\Phi(\phi) = 0$.

10. As stated in the problem, upon gauging the adapted compact $U(1) : \theta \rightarrow \theta + \epsilon$, which has radius 2π , we modify our derivative operator to act as $\partial_\alpha \theta \rightarrow \partial_\alpha \theta + A_\alpha$, where A_α gives our connection on the $U(1)$ principal bundle associated with gauging the Killing symmetry. The action gets modified:

$$S \supseteq \frac{R^2}{4\pi\ell_s^2} \int |\partial\theta|^2 \rightarrow \frac{R^2}{4\pi\ell_s^2} \int |\partial\theta + A|^2$$

This is a new theory, but we can *return to the old one* by enforcing that A be pure gauge as follows: introduce an auxiliary field ϕ and add to S the term

$$\frac{i}{2\pi} \int \phi \epsilon^{\alpha\beta} \partial_\alpha A_\beta = -\frac{i}{2\pi} \int d\phi \wedge A.$$

Integrating out ϕ gives exactly a δ -function enforcing $\epsilon^{\alpha\beta} \partial_\alpha A_\beta = 0$. This gives that A is closed, but it need not be exact if our manifold has nontrivial topology. Going around any cycle, $\int A$ can pick up a factor of $2\pi n$.

For a closed, genus g Riemann surface, there are $2g$ cycles labeled by a_i, b_i , $1 \leq i \leq g$ coming from viewing it as a $2g$ -gon. we have *Riemann's bilinear identity*, namely for two closed 1-forms ω_1, ω_2 ,

$$\int_\Sigma \omega_1 \wedge \omega_2 = \sum_{i=1}^g \left(\int_{a_i} \omega_1 \int_{b_i} \omega_2 - \int_{a_i} \omega_2 \int_{b_i} \omega_1 \right) \quad (3)$$

Now take $\omega_1 = A$, $\omega_2 = d\phi$. Now (3) gives us that $\frac{1}{2\pi} \int d\phi \wedge A$ will not be zero in general, but in the path integral, it suffices to have it be an integral multiple of 2π , since then the nontrivial holonomies will have no contribution to the action. We have that A can have winding $2\pi\mathbb{Z}$, so the only solution is to have ϕ have winding $2\pi\mathbb{Z}$. This will exactly leave over a factor of $2\pi\mathbb{Z}$. So we return to our original action by introducing the field ϕ of period 2π . (NB if I had kept the fields dimensionful, then ϕ would have period $2\pi/R$ when θ has period $2\pi R$)

In this new, equivalent action, we can gauge-fix $\theta = 0$ (do I need ghosts? No because this is abelian $U(1)$) and integrate out A . We get:

$$\frac{\ell_s^4/R^2}{4\pi\ell_s^2} \int d^2\xi (\partial\phi)^2$$

so we have obtained the same action but now on a circle of radius ℓ_s^2/R instead of R .

In doing this path integral we get a determinant factor of $\sqrt{4\pi^2\ell_s^2/R^2} = 2\pi\ell_s/R$ for each mode. Using zeta function regularization this is equal to $\sqrt{R/2\pi\ell_s}$ which we can understand as adding a $-\log(R/\ell_s)$ term to the action that will couple to the curvature R (**Show why**), this shifting the dilaton as required.

11. We can simplify things by using the conventions of the next problem to do this one. Here, we have a *single* compact coordinate θ . In our convention:

$$\hat{G}_{\mu\nu} = \begin{pmatrix} G_{00} & G_{00}A_j \\ G_{00}A_i & g_{ij} + G_{00}A_iA_j \end{pmatrix}, \quad B = (d\theta + A_i dx^i) \wedge B_j dx^j + b_{ij} dx^i \wedge dx^j, \quad \phi = \Phi - \frac{1}{4} \log G_{00}$$

From formula **F.3** specialized to this case, we get that the metric and dilaton terms become

$$\int d^D x \sqrt{-\det \hat{G}_{\mu\nu}} e^{-2\Phi} \left[\hat{R} + 4(\partial_\mu \Phi)^2 \right] = \int d^{D-1} x \sqrt{-\det g} e^{-2\phi} \left[R + 4(\partial_\mu \phi)^2 + \frac{1}{4} \partial_\mu G_{00} \partial^\mu G^{00} - \frac{1}{4} G_{00} (F_{\mu\nu}^A)^2 \right] \quad (4)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and $\hat{R}$ corresponds to the original $\hat{G}_{\mu\nu}$ while R corresponds to g_{ij} . Further $G^{00} = 1/G_{00}$.


From **F.6-F.9**, the antisymmetric tensor changes as:

$$-\frac{1}{12} \int d^D x \sqrt{-\det \hat{G}} e^{-2\Phi} \hat{H}_{ijk} \hat{H}^{ijk} = - \int d^{D-1} x \sqrt{-\det g} e^{-2\phi} \left[\frac{1}{12} H_{ijk} H^{ijk} + \frac{1}{4} \hat{H}_{ij0} \hat{H}^{ij0} \right] \quad (5)$$

Here where $H_{ij0} = \hat{H}_{ij0}$ and $H_{ijk} = \hat{H}_{ijk} - (A_i H_{0jk} + \text{cyclic})$. Here H_{ijk} is defined so that it is invariant under T -duality (**TYSM Kiritsis for pre-organizing these terms for me**). Further, under T -duality

$$\begin{aligned} G_{00} &\rightarrow G_{00}^{-1} = G^{00} \Rightarrow \partial_\mu G_{00} \partial^\mu G^{00} \text{ invariant} \\ g_{ij} &\rightarrow g_{ij} \Rightarrow R \text{ invariant} \\ A_i &\rightarrow B_i \\ B_i &\rightarrow A_i \\ \Phi &\rightarrow \Phi - \frac{1}{2} \log G_{00} \Rightarrow \phi \rightarrow \phi \Rightarrow (\partial_\mu \phi) \text{ invariant.} \end{aligned} \quad (6)$$

We see that $\sqrt{-\det g} e^{-2\phi}$ as well as the first three terms of equation (4) are invariant. We have that $F_{\mu\nu}^A \rightarrow \partial_\mu B_\nu - \partial_\nu B_\mu =: F_{\mu\nu}^B$ and $F_{ij}^B = H_{ij0}$. The last term of (4) will therefore swap with the last term of (5) and we are done.



12. This one is quick. We have

$$ds^2 = G_{00} d\theta^2 + 2G_{00} A_i dx^i dx^0 + G_{ij} dx^i dx^j, \quad B = B_j d\theta \wedge dx^j + (b_{ij} + A_i B_j) dx^i \wedge dx^j$$

Certainly we have $\tilde{G}_{00} = 1/G_{00}$, $\tilde{B}_i = G_{00} A_i / G_{00}$. Then $\tilde{A}_i = B_i$ is consistent both for the $i, 0$ components of the line element and the $dx^i \wedge dx^j$ components of the B -field as long as we keep $\tilde{b}_{ij} = b_{ij}$ and $\tilde{g}_{ij} = g_{ij}$. Finally, the dilaton must be shifted by $\tilde{\Phi} = \Phi - \frac{1}{2} \log G_{00}$.

13. The N commuting isometries correspond to a fibration by N -dimensional tori over each point in the base space. As we have seen before (for strings valued in a N -dimensional torus target space), we have that modes are described by two momenta p_L, p_R that lie on an integral lattice. Naively, we can rotate p_L, p_R by any $GL(N)$ transformation, but the integrality condition restricts us to $GL(N, \mathbb{Z})$. Now $GL(N)$ acts separately on the left and the right momenta, but we are allowed to exchange between these two by applying T -duality, which still preserves our Lorentzian norm, so the T -duality group gets enhanced to $O(N, N, \mathbb{Z})$.

14. This is clear, since orientation reversal acts trivially on $g^{ab} G_{\mu\nu} \partial_a X^\mu \partial_b X^\nu$ while it acts with a minus sign on $\epsilon^{ab} B_{\mu\nu} \partial_a X^\mu \partial_b X^\nu$. The corresponding vertex operators are:

$$: \partial X^\mu \bar{\partial} X_\mu e^{ikX} :, \quad : \zeta_{\mu\nu} \partial X^\mu \bar{\partial} X^\nu e^{ikX} :, \quad R^{(2)} : e^{ikX} :$$

If we assume the tachyon $: e^{ikX} :$ is negative under parity then so are the dilaton and graviton.

This is incompatible with **6.1.10**, as then parity will flip the sign of the dilaton in the exponential, substantially changing the action of the theory.