



Probabilistic Graphical Models

Daphne Koller, Kevin Murphy
Winter 2011-2012

[Home](#)
[Quizzes](#)
[Theory Problems](#)
[Assignments](#)
[Assignment Questions](#)
[Video Lectures](#)
[Discussion Forums](#)
[Course Wiki](#)
[Lecture Slides](#)
[Course Schedule](#)
[Course Logistics](#)
[Course Information](#)
[Course Staff](#)
[Octave Installation](#)

Feedback — PA6 Quiz

You achieved a score of 35.00 out of 35.00

Question 1

We have provided an instantiated influence diagram FullI (complete with a decision rule for D) in the file FullI.mat. What is the expected utility for this influence diagram? Please round to the nearest tenth (i.e., 1 decimal place), do not include commas, and do not write the number in scientific notation.


-685.9

Your Answer		Score	Explanation
-685.9	✓	2.00	This is the output of SimpleCalcExpectedUtility(FullI).
Total		2.00	

Question 2


Run ObserveEvidence.m on FullI to account for the following: We have been informed that variable 3 in

the model, which models an overall genetic risk for ARVD, has value 2 (indicating the presence of genetic risk factors). Then run SimpleCalcExpectedUtility on the modified influence diagram. What happened to the expected utility? (Hint -- ObserveEvidence does not re-normalize the factors so that they are again valid CPDs unless the normalize flag is set to 1. -- If you do not use the normalize flag, you can use NormalizeCPDFactors.m to do the normalization.)

Your Answer		Score	Explanation
<input checked="" type="radio"/> It substantially decreased.		3.00	It decreased from -685.9 to -729.3
Total		3.00	


Question 3

Why can we explicitly enumerate all the possible decision rules while we often cannot enumerate over all possible CPDs?

Your Answer		Score	Explanation
<input checked="" type="radio"/> All choices have a probability of either 0 or 1, where in a general CPD, choices could take on any value in [0, 1].		3.00	Because each choice is restricted to a finite set of probabilities, there is a finite number of possible decision rules, so we can enumerate over all of them.
Total		3.00	


Question 4

Let a decision node D take on d possible values. Let it have m parents that can each take on n possible values. How many possible decision rules δ_D are there?

Your Answer	Score	Explanation
<input checked="" type="radio"/> $d^{(n^m)}$	 3.00	Since each of the m parents can take on n values, there are n^m joint assignments over the parents. Since there are d values for the decision node, and all combinations of decisions over all joint assignments of parents to values are possible decision rules, there are $d^{(n^m)}$ possible decision rules.
Total	3.00	


Question 5

Consider an influence diagram with 1 decision node D that can take on d values. Let D have m parents that can each take on n values. Assume that running sum-product inference takes $O(S)$ time. What is the run-time complexity of running OptimizeMEU on this influence diagram?

Your Answer	Score	Explanation
<input checked="" type="radio"/> $O(S + dn^m)$	 4.00	Sum-product inference can be run only once, and it requires $O(S)$ time. The results of sum-product inference can be used to construct a table that is the number of instantiations to the parents, which is n^m , by the number of decision values, which is d . To choose the optimal decision, it is sufficient to scan through the table once. Thus, the total run-time is $O(S + dn^m)$.
Total	4.00	

Question 6

In which of the following situations does it make sense to use `OptimizeWithJointUtility` instead of `OptimizeLinearExpectations`?

Your Answer	Score	Explanation
<input checked="" type="radio"/> When there are large factors in the random-variables part of the influence diagram, making inference over the network slow, and there are only a few utility factors, each involving a small number of variables.	 5.00	If there are only a few utility factors, each involving a small number of variables, combining them into a joint factor won't result in too big a factor. Since the size of the utility factor is not in running inference, and since inference is expensive and we don't want to run it once for each utility factor, it makes sense to create a joint utility factor.
Total	5.00	

Question 7

In the field below, enter the dollar value of each of the tests, starting with T1, then T2 and T3, separated by commas and rounded to the nearest cent (e.g., "1.23, 4.54, 3.21" means that you would pay \$1.23 for the first test; any more than that, and your net utility will be lower than if you didn't perform any test). Do not precede with the amounts with dollar signs.

155.97, 2.82, 846.15

Your Answer		Score	Explanation
155.97	✓	5.00	
2.82	✓	5.00	
846.15	✓	5.00	
Total		15.00	