

# 19-Regression-III-More-Linear

November 15, 2016

## 1 More on Linear Regression

Today, we'll look at some additional aspects of Linear Regression.

Let's look at a standard regression.

Here are the pieces of a standard linear model.

```
In [2]: nsample = 100
        x = np.linspace(0, 10, 100)
        #print(x)
        X = np.column_stack((x, x**2))
        print(X)
        beta = np.array([1, 0.1, 10])
        e = np.random.normal(size=nsample)
```

```
[[ 0.00000000e+00  0.00000000e+00]
 [ 1.01010101e-01  1.02030405e-02]
 [ 2.02020202e-01  4.08121620e-02]
 [ 3.03030303e-01  9.18273646e-02]
 [ 4.04040404e-01  1.63248648e-01]
 [ 5.05050505e-01  2.55076013e-01]
 [ 6.06060606e-01  3.67309458e-01]
 [ 7.07070707e-01  4.99948985e-01]
 [ 8.08080808e-01  6.52994592e-01]
 [ 9.09090909e-01  8.26446281e-01]
 [ 1.01010101e+00  1.02030405e+00]
 [ 1.11111111e+00  1.23456790e+00]
 [ 1.21212121e+00  1.46923783e+00]
 [ 1.31313131e+00  1.72431385e+00]
 [ 1.41414141e+00  1.99979594e+00]
 [ 1.51515152e+00  2.29568411e+00]
 [ 1.61616162e+00  2.61197837e+00]
 [ 1.71717172e+00  2.94867871e+00]
 [ 1.81818182e+00  3.30578512e+00]
 [ 1.91919192e+00  3.68329762e+00]
 [ 2.02020202e+00  4.08121620e+00]
 [ 2.12121212e+00  4.49954086e+00]
 [ 2.22222222e+00  4.93827160e+00]
```

[ 2.32323232e+00	5.39740843e+00]
[ 2.42424242e+00	5.87695133e+00]
[ 2.52525253e+00	6.37690032e+00]
[ 2.62626263e+00	6.89725538e+00]
[ 2.72727273e+00	7.43801653e+00]
[ 2.82828283e+00	7.99918376e+00]
[ 2.92929293e+00	8.58075707e+00]
[ 3.03030303e+00	9.18273646e+00]
[ 3.13131313e+00	9.80512193e+00]
[ 3.23232323e+00	1.04479135e+01]
[ 3.33333333e+00	1.11111111e+01]
[ 3.43434343e+00	1.17947148e+01]
[ 3.53535354e+00	1.24987246e+01]
[ 3.63636364e+00	1.32231405e+01]
[ 3.73737374e+00	1.39679625e+01]
[ 3.83838384e+00	1.47331905e+01]
[ 3.93939394e+00	1.55188246e+01]
[ 4.04040404e+00	1.63248648e+01]
[ 4.14141414e+00	1.71513111e+01]
[ 4.24242424e+00	1.79981635e+01]
[ 4.34343434e+00	1.88654219e+01]
[ 4.44444444e+00	1.97530864e+01]
[ 4.54545455e+00	2.06611570e+01]
[ 4.64646465e+00	2.15896337e+01]
[ 4.74747475e+00	2.25385165e+01]
[ 4.84848485e+00	2.35078053e+01]
[ 4.94949495e+00	2.44975003e+01]
[ 5.05050505e+00	2.55076013e+01]
[ 5.15151515e+00	2.65381084e+01]
[ 5.25252525e+00	2.75890215e+01]
[ 5.35353535e+00	2.86603408e+01]
[ 5.45454545e+00	2.97520661e+01]
[ 5.55555556e+00	3.08641975e+01]
[ 5.65656566e+00	3.19967350e+01]
[ 5.75757576e+00	3.31496786e+01]
[ 5.85858586e+00	3.43230283e+01]
[ 5.95959596e+00	3.55167840e+01]
[ 6.06060606e+00	3.67309458e+01]
[ 6.16161616e+00	3.79655137e+01]
[ 6.26262626e+00	3.92204877e+01]
[ 6.36363636e+00	4.04958678e+01]
[ 6.46464646e+00	4.17916539e+01]
[ 6.56565657e+00	4.31078461e+01]
[ 6.66666667e+00	4.44444444e+01]
[ 6.76767677e+00	4.58014488e+01]
[ 6.86868687e+00	4.71788593e+01]
[ 6.96969697e+00	4.85766758e+01]
[ 7.07070707e+00	4.99948985e+01]

```
[ 7.17171717e+00  5.14335272e+01]
[ 7.27272727e+00  5.28925620e+01]
[ 7.37373737e+00  5.43720029e+01]
[ 7.47474747e+00  5.58718498e+01]
[ 7.57575758e+00  5.73921028e+01]
[ 7.67676768e+00  5.89327620e+01]
[ 7.77777778e+00  6.04938272e+01]
[ 7.87878788e+00  6.20752984e+01]
[ 7.97979798e+00  6.36771758e+01]
[ 8.08080808e+00  6.52994592e+01]
[ 8.18181818e+00  6.69421488e+01]
[ 8.28282828e+00  6.86052444e+01]
[ 8.38383838e+00  7.02887460e+01]
[ 8.48484848e+00  7.19926538e+01]
[ 8.58585859e+00  7.37169677e+01]
[ 8.68686869e+00  7.54616876e+01]
[ 8.78787879e+00  7.72268136e+01]
[ 8.88888889e+00  7.90123457e+01]
[ 8.98989899e+00  8.08182838e+01]
[ 9.09090909e+00  8.26446281e+01]
[ 9.19191919e+00  8.44913784e+01]
[ 9.29292929e+00  8.63585348e+01]
[ 9.39393939e+00  8.82460973e+01]
[ 9.49494949e+00  9.01540659e+01]
[ 9.59595960e+00  9.20824406e+01]
[ 9.69696970e+00  9.40312213e+01]
[ 9.79797980e+00  9.60004081e+01]
[ 9.89898990e+00  9.79900010e+01]
[ 1.00000000e+01  1.00000000e+02]]
```

We add one more column of ones (1) so that we can estimate the intercept.  
We then go on to create data that comes from a particular model.

```
In [3]: X = sm.add_constant(X)
        y = np.dot(X, beta) + e
```

Now we have the explanatory variable X (predictor, regressor, input variable) and the response variable y (predicted, regressand, output).

We are ready to fit the model.

```
In [4]: model = sm.OLS(y, X)
        results = model.fit()
        print(results.summary())
```

```

                                OLS Regression Results
=====
Dep. Variable:                  y      R-squared:                1.000
Model:                        OLS      Adj. R-squared:           1.000
```

```

Method:                Least Squares      F-statistic:                4.020e+06
Date:                  Tue, 15 Nov 2016    Prob (F-statistic):         2.83e-239
Time:                  10:16:58           Log-Likelihood:             -146.51
No. Observations:      100               AIC:                        299.0
Df Residuals:          97               BIC:                        306.8
Df Model:              2
Covariance Type:       nonrobust

```

```

=====
              coef      std err          t      P>|t|      [95.0% Conf. Int.]
-----
const         1.3423      0.313        4.292      0.000        0.722      1.963
x1            -0.0402      0.145       -0.278      0.781       -0.327      0.247
x2            10.0103      0.014       715.745      0.000        9.982     10.038
=====

Omnibus:                2.042    Durbin-Watson:                2.274
Prob(Omnibus):           0.360    Jarque-Bera (JB):           1.875
Skew:                    0.234    Prob(JB):                   0.392
Kurtosis:                2.519    Cond. No.                   144.
=====

```

Warnings:

```
[1] Standard Errors assume that the covariance matrix of the errors is correctly sp
```

Let's look at the various aspects of the fitted model.

```

In [5]: #dir(results)
        print('Parameters: ', results.params)
        print('R2: ', results.rsquared)
        print('Predicted values: ', results.predict())

```

```
Parameters: [ 1.34233516 -0.04024948 10.01025357]
```

```
R2: 0.999987936503
```

```

Predicted values: [ 1.34233516  1.44040458  1.74274404  2.24935355
 3.87538271  4.99480235  6.31849204  7.84645178  9.57868156
11.51518139 13.65595126 16.00099118 18.55030114 21.30388115
24.2617312  27.4238513  30.79024145 34.36090164 38.13583187
42.11503215 46.29850248 50.68624285 55.27825327 60.07453373
65.07508424 70.27990479 75.68899539 81.30235603 87.11998672
93.14188746 99.36805824 105.79849906 112.43320993 119.27219085
126.31544181 133.56296282 141.01475387 148.67081497 156.53114611
164.5957473  172.86461853 181.33775981 190.01517113 198.8968525
207.98280392 217.27302538 226.76751688 236.46627843 246.36931003
256.47661167 266.78818336 277.30402509 288.02413687 298.94851869
310.07717056 321.41009247 332.94728443 344.68874643 356.63447848
368.78448058 381.13875272 393.69729491 406.46010714 419.42718941
432.59854174 445.9741641  459.55405651 473.33821897 487.32665148
501.51935402 515.91632662 530.51756926 545.32308194 560.33286467

```

```

575.54691745    590.96524027    606.58783314    622.41469605    638.445829
654.68123201    671.12090505    687.76484815    704.61306128    721.66554447
738.9222977     756.38332097    774.04861429    791.91817766    809.99201107
828.27011452    846.75248802    865.43913157    884.33004516    903.4252288
922.72468248    942.22840621    961.93639998    981.8486638     1001.96519767]

```

Now we'll load a standard dataset as an example.

The Longley dataset contains various US macroeconomic variables that are known to be highly collinear. It has been used to appraise the accuracy of least squares routines.

```

In [6]: from statsmodels.datasets.longley import load_pandas
        y = load_pandas().endog
        X = load_pandas().exog
        X = sm.add_constant(X)

```

```

In [7]: ols_model = sm.OLS(y, X)
        ols_results = ols_model.fit()
        print(ols_results.summary())

```

```

                                OLS Regression Results
=====
Dep. Variable:                  TOTEMP      R-squared:                0.995
Model:                            OLS      Adj. R-squared:            0.992
Method:                 Least Squares      F-statistic:                330.3
Date:                  Tue, 15 Nov 2016      Prob (F-statistic):          4.98e-10
Time:                      10:16:58      Log-Likelihood:              -109.62
No. Observations:                  16      AIC:                        233.2
Df Residuals:                      9      BIC:                        238.6
Df Model:                          6
Covariance Type:                  nonrobust
=====

```

	coef	std err	t	P> t	[95.0% Conf. Int.]	
const	-3.482e+06	8.9e+05	-3.911	0.004	-5.5e+06	-1.47e+06
GNPDEFL	15.0619	84.915	0.177	0.863	-177.029	207.153
GNP	-0.0358	0.033	-1.070	0.313	-0.112	0.040
UNEMP	-2.0202	0.488	-4.136	0.003	-3.125	-0.915
ARMED	-1.0332	0.214	-4.822	0.001	-1.518	-0.549
POP	-0.0511	0.226	-0.226	0.826	-0.563	0.460
YEAR	1829.1515	455.478	4.016	0.003	798.788	2859.515

```

=====
Omnibus:                        0.749      Durbin-Watson:                2.559
Prob(Omnibus):                  0.688      Jarque-Bera (JB):              0.684
Skew:                          0.420      Prob(JB):                      0.710
Kurtosis:                      2.434      Cond. No.:                     4.86e+09
=====

```

Warnings:

```
[1] Standard Errors assume that the covariance matrix of the errors is correctly sp
[2] The condition number is large, 4.86e+09. This might indicate that there are
strong multicollinearity or other numerical problems.
```

```
/Users/crovella/anaconda3/lib/python3.5/site-packages/scipy/stats/stats.py:1327: Us
"anyway, n=%i" % int(n))
```

What does this mean?

In statistics, multicollinearity (also collinearity) is a phenomenon in which two or more predictor variables in a multiple regression model are highly correlated, meaning that one can be linearly predicted from the others with a substantial degree of accuracy.

(Wikipedia)

The condition number being referred to is the condition number of the design matrix.

That is the  $X$  in  $X\beta = y$ .

If  $X$  did not have full column rank, then there would be multiple solutions to the normal equations:

$$X^T X \beta = X^T y.$$

The solutions are in a sense all equivalent in that they yield the same value of  $\|X\beta - y\|$ .

However, the actual values of  $\beta$  can vary tremendously and so it is not clear how best to interpret the case when  $X$  does not have full column rank.

This happens when the columns of  $X$  are not linearly independent, ie, one column can be expressed as a linear combination of the other columns.

Condition number is a measure of whether  $X$  is **nearly** lacking full column rank.

In other words, whether some column is **close to** being a linear combination of the other columns.

Even in this case, the actual values of  $\beta$  can vary a lot due to the limitations of numerical precision in the computer.

This does not happen too often in practice, but there are some things that can be done if it does happen.

We can try to find one of the offending predictor variables, and leave it out of the regression.

## 1.1 Influence

Also, it can happen that dropping a single observation can have a dramatic effect on the coefficient estimates.

Restricting ourselves to the first 14 observations:

```
In [8]: ols_results2 = sm.OLS(y.ix[:14], X.ix[:14]).fit()
```

Let us see how much each parameter changes if we compare to all 16 observations:

```
In [9]: print("Percentage change %4.2f%%\n"*7 % tuple([i for i in (ols_results2.par
```

```

Percentage change -13.35%
Percentage change -236.18%
Percentage change -23.69%
Percentage change -3.36%
Percentage change -7.26%
Percentage change -200.46%
Percentage change -13.34%

```

Formal statistics for this such as the DFBETAS – a standardized measure of how much each coefficient changes when that observation is left out.

An observation is considered troublesome if its DFBETA is greater than  $2/\sqrt{\text{number of observations}}$ .

```

In [10]: infl = ols_results.get_influence()
          ## significant influence
          2./len(X)**.5

```

```

Out[10]: 0.5

```

```

In [11]: print(infl.summary_frame().filter(regex="dfb"))

```

	dfb_const	dfb_GNPDEFL	dfb_GNP	dfb_UNEMP	dfb_ARMED	dfb_POP	dfb_YEAR
0	-0.016406	-0.234566	-0.045095	-0.121513	-0.149026	0.211057	0.013388
1	-0.020608	-0.289091	0.124453	0.156964	0.287700	-0.161890	0.025958
2	-0.008382	0.007161	-0.016799	0.009575	0.002227	0.014871	0.008103
3	0.018093	0.907968	-0.500022	-0.495996	0.089996	0.711142	-0.040056
4	1.871260	-0.219351	1.611418	1.561520	1.169337	-1.081513	-1.864186
5	-0.321373	-0.077045	-0.198129	-0.192961	-0.430626	0.079916	0.323275
6	0.315945	-0.241983	0.438146	0.471797	-0.019546	-0.448515	-0.307517
7	0.015816	-0.002742	0.018591	0.005064	-0.031320	-0.015823	-0.015583
8	-0.004019	-0.045687	0.023708	0.018125	0.013683	-0.034770	0.005116
9	-1.018242	-0.282131	-0.412621	-0.663904	-0.715020	-0.229501	1.035723
10	0.030947	-0.024781	0.029480	0.035361	0.034508	-0.014194	-0.030805
11	0.005987	-0.079727	0.030276	-0.008883	-0.006854	-0.010693	-0.005323
12	-0.135883	0.092325	-0.253027	-0.211465	0.094720	0.331351	0.129120
13	0.032736	-0.024249	0.017510	0.033242	0.090655	0.007634	-0.033114
14	0.305868	0.148070	0.001428	0.169314	0.253431	0.342982	-0.318031
15	-0.538323	0.432004	-0.261262	-0.143444	-0.360890	-0.467296	0.552421

```


In [12]: sns.heatmap(infl.summary_frame().filter(regex="dfb") > 0.5 )

```

```

Out[12]: <matplotlib.axes._subplots.AxesSubplot at 0x11aa379e8>

```



## 1.2 Flexible Modeling

The Guerry dataset is a collection of historical data used in support of Andre-Michel Guerry's 1833 "Essay on the Moral Statistics of France."

Andre-Michel Guerry's (1833) *Essai sur la Statistique Morale de la France* was one of the foundation studies of modern social science. Guerry assembled data on crimes, suicides, literacy and other "moral statistics," and used tables and maps to analyze a variety of social issues in perhaps the first comprehensive study relating such variables.

Wikipedia

Guerry's results were startling for two reasons. First he showed that rates of crime and suicide remained remarkably stable over time, when broken down by age, sex, region of France and even season of the year; yet these numbers varied systematically across departements of France. This regularity of social numbers created the possibility to conceive, for the first time, that human actions in the social world were governed by social laws, just as inanimate objects were governed by laws of the physical world.

Source: "A.-M. Guerry's Moral Statistics of France: Challenges for Multivariable Spatial Analysis", Michael Friendly. *Statistical Science* 2007, Vol. 22, No. 3, 368–399.



```
In [13]: # Lottery is per-capital wager on Royal Lottery
df = sm.datasets.get_rdataset("Guerry", "HistData").data
df = df[['Lottery', 'Literacy', 'Wealth', 'Region']].dropna()
df.head()
```

```
Out[13]:
```

	Lottery	Literacy	Wealth	Region
0	41	37	73	E
1	38	51	22	N
2	66	13	61	C
3	80	46	76	E
4	79	69	83	E

We can use another version of the module that can directly type formulas and expressions in the functions of the models.

We can specify the name of the columns to be used to predict another column, remove columns, etc.

```
In [14]: mod = smf.ols(formula='Lottery ~ Literacy + Wealth + Region', data=df)
res = mod.fit()
print(res.summary())
```

```

OLS Regression Results
=====
Dep. Variable:          Lottery    R-squared:                0.338
Model:                  OLS        Adj. R-squared:           0.287
Method:                 Least Squares    F-statistic:             6.636
Date:                  Tue, 15 Nov 2016    Prob (F-statistic):      1.07e-05
Time:                  10:16:59    Log-Likelihood:          -375.30
No. Observations:      85    AIC:                     764.6
Df Residuals:          78    BIC:                     781.7
Df Model:               6
Covariance Type:       nonrobust
=====

```

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	38.6517	9.456	4.087	0.000	19.826 57.478
Region[T.E]	-15.4278	9.727	-1.586	0.117	-34.793 3.938
Region[T.N]	-10.0170	9.260	-1.082	0.283	-28.453 8.419
Region[T.S]	-4.5483	7.279	-0.625	0.534	-19.039 9.943
Region[T.W]	-10.0913	7.196	-1.402	0.165	-24.418 4.235
Literacy	-0.1858	0.210	-0.886	0.378	-0.603 0.232
Wealth	0.4515	0.103	4.390	0.000	0.247 0.656

```

=====
Omnibus:                 3.049    Durbin-Watson:           1.785
Prob(Omnibus):           0.218    Jarque-Bera (JB):        2.694
Skew:                    -0.340    Prob(JB):                0.260
Kurtosis:                2.454    Cond. No.:               371.
=====

```

Warnings:

```
[1] Standard Errors assume that the covariance matrix of the errors is correctly sp
```

### Categorical variables

Patsy is the name of the interpreter that parses the formulas.

Looking at the summary printed above, notice that patsy determined that elements of Region were text strings, so it treated Region as a categorical variable. patsy's default is also to include an intercept, so we automatically dropped one of the Region categories.

### Removing variables

The "-" sign can be used to remove columns/variables. For instance, we can remove the intercept from a model by:

```
In [15]: res = smf.ols(formula='Lottery ~ Literacy + Wealth + C(Region) -1 ', data=
print(res.summary())
```

```

                                OLS Regression Results
=====
Dep. Variable:                  Lottery    R-squared:                0.338
Model:                            OLS      Adj. R-squared:            0.287
Method:                 Least Squares    F-statistic:                6.636
Date:                Tue, 15 Nov 2016    Prob (F-statistic):        1.07e-05
Time:                  10:17:00    Log-Likelihood:            -375.30
No. Observations:                85      AIC:                       764.6
Df Residuals:                    78      BIC:                       781.7
Df Model:                        6
Covariance Type:                nonrobust
=====

```

	coef	std err	t	P> t	[95.0% Conf. Int.]
C(Region) [C]	38.6517	9.456	4.087	0.000	19.826 57.478
C(Region) [E]	23.2239	14.931	1.555	0.124	-6.501 52.949
C(Region) [N]	28.6347	13.127	2.181	0.032	2.501 54.769
C(Region) [S]	34.1034	10.370	3.289	0.002	13.459 54.748
C(Region) [W]	28.5604	10.018	2.851	0.006	8.616 48.505
Literacy	-0.1858	0.210	-0.886	0.378	-0.603 0.232
Wealth	0.4515	0.103	4.390	0.000	0.247 0.656

```
=====
Omnibus:                3.049    Durbin-Watson:            1.785
Prob(Omnibus):           0.218    Jarque-Bera (JB):         2.694
Skew:                   -0.340    Prob(JB):                 0.260
Kurtosis:               2.454    Cond. No.:                653.
=====
```

Warnings:

```
[1] Standard Errors assume that the covariance matrix of the errors is correctly sp
```

### Functions

We can also apply vectorized functions to the variables in our model:

```
In [16]: res = smf.ols(formula='Lottery ~ np.log(Literacy)', data=df).fit()
         print(res.summary())
```

```

                        OLS Regression Results
=====
Dep. Variable:          Lottery      R-squared:                0.161
Model:                  OLS         Adj. R-squared:            0.151
Method:                 Least Squares   F-statistic:              15.89
Date:                  Tue, 15 Nov 2016   Prob (F-statistic):       0.000144
Time:                  10:17:00         Log-Likelihood:           -385.38
No. Observations:      85              AIC:                     774.8
Df Residuals:          83              BIC:                     779.7
Df Model:               1
Covariance Type:       nonrobust
=====
                        coef      std err          t      P>|t|      [95.0% Conf. Int.]
-----
Intercept              115.6091      18.374        6.292      0.000       79.064   152.15
np.log(Literacy)      -20.3940       5.116       -3.986      0.000      -30.570  -10.21
=====
Omnibus:               8.907      Durbin-Watson:           2.019
Prob(Omnibus):         0.012      Jarque-Bera (JB):         3.299
Skew:                 0.108      Prob(JB):                 0.192
Kurtosis:             2.059      Cond. No.                 28.7
=====
```

Warnings:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

### 1.3 Understanding Problematic Observations

```
In [17]: from statsmodels.formula.api import ols
         from statsmodels.compat import lzip
         prestige = sm.datasets.get_rdataset("Duncan", "car", cache=True).data
```

This is data on the “prestige” and other characteristics of 45 U. S. occupations in 1950.

- type: Type of occupation. A factor with the following levels: prof, professional and managerial; wc, white-collar; bc, blue-collar.
- income: Percent of males in occupation earning USD 3500 or more in 1950.
- education: Percent of males in occupation in 1950 who were high-school graduates.
- prestige: Percent of raters in NORC study rating occupation as excellent or good in prestige.

```
In [18]: prestige.head()
```

```
Out[18]:
```

	type	income	education	prestige
accountant	prof	62	86	82
pilot	prof	72	76	83
architect	prof	75	92	90
author	prof	55	90	76
chemist	prof	64	86	90

```
In [19]: prestige_model = ols("prestige ~ income + education", data=prestige).fit()
print(prestige_model.summary())
```

```

OLS Regression Results
=====
Dep. Variable:          prestige    R-squared:                0.828
Model:                  OLS        Adj. R-squared:           0.820
Method:                 Least Squares    F-statistic:             101.2
Date:                  Tue, 15 Nov 2016    Prob (F-statistic):      8.65e-17
Time:                  10:17:00          Log-Likelihood:          -178.98
No. Observations:      45              AIC:                    364.0
Df Residuals:          42              BIC:                    369.4
Df Model:               2
Covariance Type:       nonrobust
=====
               coef      std err          t      P>|t|      [95.0% Conf. Int.]
-----
Intercept    -6.0647      4.272      -1.420      0.163      -14.686      2.556
income        0.5987      0.120       5.003      0.000       0.357      0.840
education     0.5458      0.098       5.555      0.000       0.348      0.744
=====
Omnibus:            1.279    Durbin-Watson:           1.458
Prob(Omnibus):      0.528    Jarque-Bera (JB):         0.520
Skew:               0.155    Prob(JB):                 0.771
Kurtosis:           3.426    Cond. No.                  163.
=====

```

Warnings:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

### Influence plots

To ask about the impact of individual observations on the overall model, we can calculate the influence of each observation.

leverage is a measure of how far away the independent variable values of an observation are from those of the other observations.

The residual is the error obtained when predicting the observation.

Influence plots show the studentized residuals vs. the leverage of each observation as measured by the projection matrix.

Externally studentized residuals are residuals that are scaled by their standard deviation where

$$\text{var}(\hat{\epsilon}_i) = \hat{\sigma}^2(1 - h_{ii}).$$

$h_{ii}$  is the  $i$ -th diagonal element of the projection matrix

$$H = X(X^T X)^{-1} X^T$$


where  $X$  is the design matrix.

It can be shown that

$$h_{ii} = \frac{\partial \hat{y}_i}{\partial y_i}.$$

The influence of each point can be visualized by the criterion keyword argument. Options are Cook's distance and DFFITS, two measures of influence. Basically these methods combine leverage and residual.

```
In [20]: fig, ax = plt.subplots(figsize=(10,6))
         fig = sm.graphics.influence_plot(prestige_model, ax=ax, criterion="cooks")
```




As you can see there are a few worrisome observations. Both contractor and reporter have low leverage but a large residual. RR.engineer has small residual and large leverage. Conductor and minister have both high leverage and large residuals, and, therefore, large influence.

**Partial Regression Plots** are a principled way of looking at individual independent variables versus the dependent variable, for visualization.

In a partial regression plot, to discern the relationship between the response variable and the  $k$ -th variable, we compute the residuals by regressing the response variable versus the independent variables excluding  $X_k$ . We can denote this by  $X_{\sim k}$ . We then compute the residuals by regressing  $X_k$  on  $X_{\sim k}$ . The partial regression plot is the plot of the former versus the latter residuals.

```
In [21]: fix, ax = plt.subplots(figsize=(12,14))
         fig = sm.graphics.plot_partregress("prestige", "income", ["education"], da
```



As you can see the partial regression plot confirms the influence of conductor, minister, and RR.engineer on the partial relationship between income and prestige. The cases greatly decrease the effect of income on prestige. Dropping these cases confirms this.

```
In [22]: subset = ~prestige.index.isin(["conductor", "RR.engineer", "minister"])
prestige_model2 = ols("prestige ~ income + education", data=prestige, subset=subset)
print(prestige_model2.summary())
```

OLS Regression Results

=====

```

Dep. Variable:          prestige    R-squared:          0.876
Model:                  OLS         Adj. R-squared:      0.870
Method:                 Least Squares    F-statistic:        138.1
Date:                  Tue, 15 Nov 2016    Prob (F-statistic):  2.02e-18
Time:                  10:17:01         Log-Likelihood:     -160.59
No. Observations:      42             AIC:               327.2
Df Residuals:          39             BIC:               332.4
Df Model:               2
Covariance Type:       nonrobust

```

```

=====
              coef      std err          t      P>|t|      [95.0% Conf. Int.]
-----
Intercept    -6.3174      3.680      -1.717      0.094      -13.760      1.125
income        0.9307      0.154       6.053      0.000       0.620      1.242
education     0.2846      0.121       2.345      0.024       0.039      0.530
=====
Omnibus:            3.811    Durbin-Watson:          1.468
Prob(Omnibus):      0.149    Jarque-Bera (JB):          2.802
Skew:              -0.614    Prob(JB):              0.246
Kurtosis:           3.303    Cond. No.              158.
=====

```

Warnings:

[1] Standard Errors assume that the covariance matrix of the errors is correctly specified

```

In [23]: fig = plt.figure(figsize=(12,8))
         fig = sm.graphics.plot_partregress_grid(prestige_model, fig=fig)

```

