

# Infinitely Repeated Games

Game Theory

Vincent Knight

$$\begin{pmatrix} (2, 2) & (0, 3) \\ (3, 0) & (1, 1) \end{pmatrix}$$

$$\begin{pmatrix} (2, 2) & (0, 3) \\ (3, 0) & (1, 1) \end{pmatrix}$$

- ▶  $s_C$ : cooperate at every stage
- ▶  $s_D$ : defect at every stage

$$\begin{pmatrix} (2, 2) & (0, 3) \\ (3, 0) & (1, 1) \end{pmatrix}$$

- ▶  $s_C$ : cooperate at every stage
- ▶  $s_D$ : defect at every stage

$$u_1(s_C, s_C) = \sum_{i=1}^{\infty} 2 > \infty$$

$$\begin{pmatrix} (2, 2) & (0, 3) \\ (3, 0) & (1, 1) \end{pmatrix}$$

- ▶  $s_C$ : cooperate at every stage
- ▶  $s_D$ : defect at every stage

$$u_1(s_C, s_C) = \sum_{i=1}^{\infty} \delta^i 2 < \infty \text{ if } |\delta| < 1$$

$$\begin{pmatrix} (2, 2) & (0, 3) \\ (3, 0) & (1, 1) \end{pmatrix}$$


- ▶  $s_C$ : cooperate at every stage
- ▶  $s_D$ : defect at every stage

$$u_1(s_C, s_C) = \sum_{i=1}^{\infty} \delta^i 2 < \infty \text{ if } |\delta| < 1$$


Possible interpretation of  $\delta$ : probability of game ending at any stage.


$$\bar{T} = \frac{1}{1 - \delta}$$

$$\frac{1}{\bar{T}} U_i(r, s) = (1 - \delta) U_i(r, s)$$









## Folk Theorem

Let  $(u_1^*, u_2^*)$  be a pair of Nash equilibrium payoffs for a stage game. For every individually rational pair  $(v_1, v_2)$  there exists  $\bar{\delta}$  such that for all  $1 > \delta > \bar{\delta} > 0$  there is a subgame perfect Nash equilibrium with payoffs  $(v_1, v_2)$ .