SBML Model Report

Model name: "Bornheimer2004_GTPaseCycle"



May 6, 2016

1 General Overview

This is a document in SBML Level 2 Version 1 format. This model was created by Enuo He¹ at November 29th 2006 at 5:40 p. m. and last time modified at July fifth 2012 at 2:49 p. m. Table 1 provides an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	17
events	0	constraints	0
reactions	24	function definitions	0
global parameters	2	unit definitions	0
rules	2	initial assignments	0

Model Notes

This model is according to the paper *Computational modeling reveals how interplay between components of a GTPase-cycle module regulates signal transduction* by Bornheimer et al 2004. The figure 3 is reproduced by Copasi 4.0.19 (development). It is three-dimensional logarithmic plots show the output of simulations of Z and v at various concentrations of R and GAP.

¹University of Oxford, enuo.he@wolfson.ox.ac.uk

To the extent possible under law, all copyright and related or neighbouring rights to this encoded model have been dedicated to the public domain worldwide. Please refer to CCO Public Domain Dedication for more information.

In summary, you are entitled to use this encoded model in absolutely any manner you deem suitable, verbatim, or with modification, alone or embedded it in a larger context, redistribute it, commercially or not, in a restricted way or not.

To cite BioModels Database, please use: Li C, Donizelli M, Rodriguez N, Dharuri H, Endler L, Chelliah V, Li L, He E, Henry A, Stefan MI, Snoep JL, Hucka M, Le Novre N, Laibe C (2010) BioModels Database: An enhanced, curated and annotated resource for published quantitative kinetic models. BMC Syst Biol., 4:92.

2 Unit Definitions

This is an overview of five unit definitions which are all predefined by SBML and not mentioned in the model.

2.1 Unit substance

Notes Mole is the predefined SBML unit for substance.

Definition mol

2.2 Unit volume

Notes Litre is the predefined SBML unit for volume.

Definition 1

2.3 Unit area

Notes Square metre is the predefined SBML unit for area since SBML Level 2 Version 1.

Definition m²

2.4 Unit length

Notes Metre is the predefined SBML unit for length since SBML Level 2 Version 1.

Definition m

2.5 Unit time

Notes Second is the predefined SBML unit for time.

Definition s

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
compartment_0	cell		3	1	litre	Z	

3.1 Compartment compartment_0

This is a three dimensional compartment with a constant size of one litre.

Name cell

4 Species

This model contains 17 species. Section 8 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condi- tion
species_0	A	${\tt compartment_0}$	$\operatorname{mol} \cdot 1^{-1}$		
species_1	G	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_2	GA	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_3	T	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_4	R	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_5	G*T	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_6	GD	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_7	Pi	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_8	D	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_9	RG	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_10	RG*T	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_11	G*AT	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_12	GAD	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_13	RGD	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_14	RGA	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		
species_15	RG*AT	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		\Box
species_16	RGAD	${\tt compartment_0}$	$\text{mol} \cdot l^{-1}$		

5 Parameters

This model contains two global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO Value Unit	Constant
parameter_0	Z	0.0	
$parameter_1$	V	0.0	

6 Rules

This is an overview of two rules.

6.1 Rule parameter_0

Rule parameter_0 is an assignment rule for parameter parameter_0:

$$parameter_0 = \frac{[species_5] + [species_10] + [species_15] + [species_11]}{1.0E - 8}$$
 (1)

Notes The [Gtot]=the initial value of [G]=10e-9 M

6.2 Rule parameter_1

Rule parameter_1 is an assignment rule for parameter parameter_1:

$$parameter_{1} = \frac{25 \cdot [species_{1}5] + 25 \cdot [species_{1}1] + 0.013 \cdot [species_{1}0] + 0.013 \cdot [species_{5}]}{1.0E - 8}$$
(2)

Notes The [Gtot]=the initial value of [G]=10e-9 M

7 Reactions

This model contains 24 reactions. All reactions are listed in the following table and are subsequently described in detail. If a reaction is affected by a modifier, the identifier of this species is written above the reaction arrow.

Table 5: Overview of all reactions

Nº	Id	Name	Reaction Equation	SBO
1	reaction_0	G protein binding GAP	$species_1 + species_0 \Longrightarrow species_2$	
2	${\tt reaction_1}$	G protein binding GTP	$species_1 + species_3 \Longrightarrow species_5$	
3	${\tt reaction_2}$	G binding Receptor	$species_1 + species_4 \Longrightarrow species_9$	
4	$reaction_3$	G*T hydrolysis	$species_5 \Longrightarrow species_6 + species_7$	
5	${\tt reaction_4}$	GD dissociation	$species_6 \Longrightarrow species_1 + species_8$	
6	${\tt reaction_5}$	RG binding GTP	$species_9 + species_3 \Longrightarrow species_10$	
7	${\tt reaction_6}$	G*T binding Receptor	$species_5 + species_4 \Longrightarrow species_10$	
8	${\tt reaction_7}$	G*T binding GAP	$species_5 + species_0 \Longrightarrow species_11$	
9	${\tt reaction_8}$	GD binding GAP	$species_6 + species_0 \Longrightarrow species_{12}$	
10	${\tt reaction_9}$	GD binding Receptor	$species_6 + species_4 \Longrightarrow species_13$	
11	${\tt reaction_10}$	RG*T hydrolysis	$species_10 \Longrightarrow species_13 + species_7$	
12	${\tt reaction_11}$	RGD dissociation	$species_13 \Longrightarrow species_9 + species_8$	
13	${\tt reaction_12}$	GA binding GTP	$species_2 + species_3 \Longrightarrow species_11$	
14	$reaction_13$	G*AT hydrolysis	$species_11 \Longrightarrow species_12 + species_7$	
15	${\tt reaction_14}$	GAD dissociation	$species_12 \Longrightarrow species_2 + species_8$	
16	${\tt reaction_15}$	RG binding GAP	$species_9 + species_0 \Longrightarrow species_14$	
17	${\tt reaction_16}$	GA binding Receptor	$species_2 + species_4 \Longrightarrow species_14$	
18	${\tt reaction_17}$	RGA binding GTP	$species_14 + species_3 \Longrightarrow species_15$	
19	${\tt reaction_18}$	G*AT binding Receptor	$species_11 + species_4 \Longrightarrow species_15$	
20	${\tt reaction_19}$	RG*T binding GAP	$species_10 + species_20 \Longrightarrow species_15$	
21	$reaction_20$	RG*AT hydrolysis	$species_15 \Longrightarrow species_16 + species_7$	
22	$reaction_21$	RGD binding GAP	$species_13 + species_0 \Longrightarrow species_16$	
23	${\tt reaction_22}$	GAD binding Receptor	$species_12 + species_4 \Longrightarrow species_16$	

Nº Id	Name	Reaction Equation	SBO
24 reaction_23	RGAD dissociation	species_16 ⇒ species_14 + species_8	

7.1 Reaction reaction_0

This is a reversible reaction of two reactants forming one product.

Name G protein binding GAP

Reaction equation

$$species_1 + species_0 \Longrightarrow species_2$$
 (3)

Reactants

Table 6: Properties of each reactant.

Id	Name	SBO
species_1	G	
${\tt species_0}$	A	

Product

Table 7: Properties of each product.

Id	Name	SBO
species_2	GA	

Kinetic Law

Derived unit contains undeclared units

$$v_1 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_1}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_2}])$$
 (4)

Table 8: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			8780000.0		$ \mathbf{Z} $
k2			8.0		Ø

7.2 Reaction reaction_1

This is a reversible reaction of two reactants forming one product.

Name G protein binding GTP

Reaction equation

$$species_1 + species_3 \Longrightarrow species_5$$
 (5)

Reactants

Table 9: Properties of each reactant.

Id	Name	SBO
species_1 species_3	G T	

Product

Table 10: Properties of each product.

Id	Name	SBO
species_5	G*T	

Kinetic Law

Derived unit contains undeclared units

$$v_2 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_1}] \cdot [\text{species_3}] - \text{k2} \cdot [\text{species_5}])$$
 (6)

Table 11: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			529000.000)	
k2			$8.38 \cdot 10^{-6}$	Ó	$\overline{\mathbf{Z}}$

7.3 Reaction reaction_2

This is a reversible reaction of two reactants forming one product.

Name G binding Receptor

Reaction equation

$$species_1 + species_4 \Longrightarrow species_9$$
 (7)

Reactants

Table 12: Properties of each reactant.

Id	Name	SBO
species_1 species_4	G R	

Product

Table 13: Properties of each product.

Id	Name	SBO
species_9	RG	

Kinetic Law

Derived unit contains undeclared units

$$v_3 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_1}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_9}])$$
 (8)

Table 14: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			$6.36 \cdot 10^{8}$		lacksquare
k2			0.018		\square

7.4 Reaction reaction_3

This is a reversible reaction of one reactant forming two products.

Name G*T hydrolysis

Reaction equation

$$species_5 \Longrightarrow species_6 + species_7 \tag{9}$$

Reactant

Table 15: Properties of each reactant.

Id	Name	SBO
species_5	G*T	

Products

Table 16: Properties of each product.

Id	Name	SBO
species_6		
species_7	Pi	

Kinetic Law

Derived unit contains undeclared units

$$v_4 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_5}] - \text{k2} \cdot [\text{species_6}] \cdot [\text{species_7}])$$
 (10)

Table 17: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			0.013		
k2			$9.03 \cdot 10^{-7}$		

7.5 Reaction reaction_4

This is a reversible reaction of one reactant forming two products.

Name GD dissociation

Reaction equation

$$species_6 \Longrightarrow species_1 + species_8$$
 (11)

Reactant

Table 18: Properties of each reactant.

Id	Name	SBO
species_6	GD	

Products

Table 19: Properties of each product.

Id	Name	SBO
species_1	G	
species_8	D	

Kinetic Law

Derived unit contains undeclared units

$$v_5 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_6}] - \text{k2} \cdot [\text{species_1}] \cdot [\text{species_8}])$$
 (12)

Table 20: Properties of each parameter.

Id	Name	SBO Value Unit	Constant
k1		10^{-4}	lacksquare
k2		62.300	

7.6 Reaction reaction_5

This is a reversible reaction of two reactants forming one product.

Name RG binding GTP

Reaction equation

$$species_9 + species_3 \Longrightarrow species_{10}$$
 (13)

Reactants

Table 21: Properties of each reactant.

Id	Name	SBO
species_9 species_3	RG T	

Product

Table 22: Properties of each product.

Id	Name	SBO
species_10	RG*T	

Derived unit contains undeclared units

$$v_6 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_9}] \cdot [\text{species_3}] - \text{k2} \cdot [\text{species_10}])$$
 (14)

Table 23: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			853000.000)	\overline{Z}
k2			0.005	;	

7.7 Reaction reaction_6

This is a reversible reaction of two reactants forming one product.

Name G*T binding Receptor

Reaction equation

$$species_5 + species_4 \Longrightarrow species_{10}$$
 (15)

Reactants

Table 24: Properties of each reactant.

Id	Name	SBO
species_5	G*T	
${\tt species_4}$	R	

Product

Table 25: Properties of each product.

Id	Name	SBO
species_10	RG*T	

Derived unit contains undeclared units

$$v_7 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_5}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_10}])$$
 (16)

Table 26: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			$1.32\cdot 10^8$		
k2			1.280		

7.8 Reaction reaction_7

This is a reversible reaction of two reactants forming one product.

Name G*T binding GAP

Reaction equation

$$species_5 + species_0 \Longrightarrow species_{11}$$
 (17)

Reactants

Table 27: Properties of each reactant.

Id	Name	SBO
species_5	G*T	
species_0	A	

Product

Table 28: Properties of each product.

Id	Name	SBO
species_11	G*AT	

Kinetic Law

Derived unit contains undeclared units

$$v_8 = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_5}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_11}])$$
 (18)

Table 29: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			386000.000)	\square
k2			0.041		\mathbf{Z}

7.9 Reaction reaction_8

This is a reversible reaction of two reactants forming one product.

Name GD binding GAP

Reaction equation

$$species_6 + species_0 \Longrightarrow species_{12}$$
 (19)

Reactants

Table 30: Properties of each reactant.

Id	Name	SBO
species_6	GD	
species_0	A	

Product

Table 31: Properties of each product.

Id	Name	SBO
species_12	GAD	

Kinetic Law

Derived unit contains undeclared units

$$v_9 = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_6}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_12}])$$
 (20)

Table 32: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			64100.00		lacksquare
k2			0.95		

7.10 Reaction reaction_9

This is a reversible reaction of two reactants forming one product.

Name GD binding Receptor

Reaction equation

$$species_6 + species_4 \Longrightarrow species_{13}$$
 (21)

Reactants

Table 33: Properties of each reactant.

Id	Name	SBO
species_6	GD	
species_4	R	

Product

Table 34: Properties of each product.

Id	Name	SBO
species_13	RGD	

Kinetic Law

Derived unit contains undeclared units

$$v_{10} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_6}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_13}])$$
 (22)

Table 35: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			$9.47 \cdot 10^7$		
k2			0.002		✓

7.11 Reaction reaction_10

This is a reversible reaction of one reactant forming two products.

Name RG*T hydrolysis

Reaction equation

$$species_10 \Longrightarrow species_13 + species_7$$
 (23)

Reactant

Table 36: Properties of each reactant.

Id	Name	SBO
species_10	RG*T	

Products

Table 37: Properties of each product.

Id	Name	SBO
species_13 species_7	RGD Pi	

Kinetic Law

Derived unit contains undeclared units

$$v_{11} = \text{vol} \left(\text{compartment_0} \right) \cdot \left(\text{k1} \cdot [\text{species_10}] - \text{k2} \cdot [\text{species_13}] \cdot [\text{species_7}] \right)$$
 (24)

Table 38: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			0.013		
k2			$2.22 \cdot 10^{-9}$)	$ \overline{\mathbf{Z}} $

7.12 Reaction reaction_11

This is a reversible reaction of one reactant forming two products.

Name RGD dissociation

Reaction equation

$$species_13 \Longrightarrow species_9 + species_8$$
 (25)

Reactant

Table 39: Properties of each reactant.

Id	Name	SBO
species_13	RGD	

Products

Table 40: Properties of each product.

Id	Name	SBO
species_9 species_8		

Kinetic Law

Derived unit contains undeclared units

$$v_{12} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_13}] - \text{k2} \cdot [\text{species_9}] \cdot [\text{species_8}])$$
 (26)

Table 41: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			2.0		\overline{Z}
k2			1470000.0		

7.13 Reaction reaction_12

This is a reversible reaction of two reactants forming one product.

Name GA binding GTP

Reaction equation

$$species_2 + species_3 \Longrightarrow species_{11}$$
 (27)

Reactants

Table 42: Properties of each reactant.

Id	Name	SBO
species_2	GA T	

Product

Table 43: Properties of each product.

Id	Name	SBO
species_11	G*AT	

Kinetic Law

Derived unit contains undeclared units

$$v_{13} = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_2}] \cdot [\text{species_3}] - \text{k2} \cdot [\text{species_11}])$$
 (28)

Table 44: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			4700.000		Ø
k2		8	$.32 \cdot 10^{-8}$		\mathbf{Z}

7.14 Reaction reaction_13

This is a reversible reaction of one reactant forming two products.

Name G*AT hydrolysis

Reaction equation

$$species_11 \Longrightarrow species_12 + species_7$$
 (29)

Reactant

Table 45: Properties of each reactant.

Id	Name	SBO
species_11	G*AT	

Products

Table 46: Properties of each product.

Id	Name	SBO
species_12	GAD	
species_7	Pi	

Kinetic Law

Derived unit contains undeclared units

$$v_{14} = \text{vol} \left(\text{compartment_0} \right) \cdot \left(\text{k1} \cdot [\text{species_11}] - \text{k2} \cdot [\text{species_12}] \cdot [\text{species_7}] \right)$$
 (30)

Table 47: Properties of each parameter.

Id	Name	SBO Value Unit	Constant
k1		25.000	
k2		0.244	

7.15 Reaction reaction_14

This is a reversible reaction of one reactant forming two products.

Name GAD dissociation

Reaction equation

$$species_12 \rightleftharpoons species_2 + species_8$$
 (31)

Reactant

Table 48: Properties of each reactant.

Id	Name	SBO
species_12	GAD	

Products

Table 49: Properties of each product.

Id	Name	SBO
species_2 species_8	GA D	

Derived unit contains undeclared units

$$v_{15} = \text{vol} \left(\text{compartment_0} \right) \cdot \left(\text{k1} \cdot [\text{species_12}] - \text{k2} \cdot [\text{species_2}] \cdot [\text{species_8}] \right)$$
 (32)

Table 50: Properties of each parameter.

Id	Name	SBO Value Unit	Constant
k1		10^{-4}	
k2		3.830	

7.16 Reaction reaction_15

This is a reversible reaction of two reactants forming one product.

Name RG binding GAP

Reaction equation

$$species_9 + species_0 \Longrightarrow species_{14}$$
 (33)

Reactants

Table 51: Properties of each reactant.

Id	Name	SBO
species_9	RG	
${\tt species_0}$	A	

Product

Table 52: Properties of each product.

Id	Name	SBO
species_14	RGA	

Derived unit contains undeclared units

$$v_{16} = \text{vol} (\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_9}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_14}])$$
 (34)

Table 53: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			74300.000		\overline{Z}
k2			0.006		\mathbf{Z}

7.17 Reaction reaction_16

This is a reversible reaction of two reactants forming one product.

Name GA binding Receptor

Reaction equation

$$species_2 + species_4 \Longrightarrow species_{14}$$
 (35)

Reactants

Table 54: Properties of each reactant.

Id	Name	SBO
species_2	GA	
species_4	R	

Product

Table 55: Properties of each product.

Id	Name	SBO
species_14	RGA	

Derived unit contains undeclared units

$$v_{17} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_2}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_14}])$$
 (36)

Table 56: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			$2.28\cdot 10^7$		
k2			$5.43 \cdot 10^{-5}$		

7.18 Reaction reaction_17

This is a reversible reaction of two reactants forming one product.

Name RGA binding GTP

Reaction equation

$$species_14 + species_3 \Longrightarrow species_15$$
 (37)

Reactants

Table 57: Properties of each reactant.

Id	Name	SBO
species_14	RGA	
species_3	T	

Product

Table 58: Properties of each product.

Id	Name	SBO
species_15	RG*AT	

Kinetic Law

Derived unit contains undeclared units

$$v_{18} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_14}] \cdot [\text{species_3}] - \text{k2} \cdot [\text{species_15}])$$
 (38)

Table 59: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			1620000.000)	
k2			0.009)	

7.19 Reaction reaction_18

This is a reversible reaction of two reactants forming one product.

Name G*AT binding Receptor

Reaction equation

$$species_11 + species_4 \Longrightarrow species_15$$
 (39)

Reactants

Table 60: Properties of each reactant.

Id	Name	SBO
species_11	G*AT	
species_4	R	

Product

Table 61: Properties of each product.

Id	Name	SBO
species_15	RG*AT	

Kinetic Law

Derived unit contains undeclared units

$$v_{19} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_11}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_15}])$$
 (40)

Table 62: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			6200000.000)	\overline{Z}
k2			0.043	3	\checkmark

7.20 Reaction reaction_19

This is a reversible reaction of two reactants forming one product.

Name RG*T binding GAP

Reaction equation

$$species_10 + species_20 \Longrightarrow species_15$$
 (41)

Reactants

Table 63: Properties of each reactant.

Id	Name	SBO
species_10 species_0	RG*T A	

Product

Table 64: Properties of each product.

Id	Name	SBO
species_15	RG*AT	

Kinetic Law

Derived unit contains undeclared units

$$v_{20} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_10}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_15}])$$
 (42)

Table 65: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			6300000.000)	$ \mathcal{Q} $
k2			0.478	3	\square

7.21 Reaction reaction_20

This is a reversible reaction of one reactant forming two products.

Name RG*AT hydrolysis

Reaction equation

$$species_15 \Longrightarrow species_16 + species_7$$
 (43)

Reactant

Table 66: Properties of each reactant.

Id	Name	SBO
species_15	RG*AT	

Products

Table 67: Properties of each product.

Id	Name	SBO
species_16 species_7	RGAD Pi	

Kinetic Law

Derived unit contains undeclared units

$$v_{21} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_15}] - \text{k2} \cdot [\text{species_16}] \cdot [\text{species_7}])$$
 (44)

Table 68: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			25.000		\square
k2			0.003		\square

7.22 Reaction reaction_21

This is a reversible reaction of two reactants forming one product.

Name RGD binding GAP

Reaction equation

$$species_13 + species_20 \Longrightarrow species_16$$
 (45)

Reactants

Table 69: Properties of each reactant.

Id	Name	SBO
species_13	RGD	
species_0	A	

Product

Table 70: Properties of each product.

Id	Name	SBO
species_16	RGAD	

Kinetic Law

Derived unit contains undeclared units

$$v_{22} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_13}] \cdot [\text{species_0}] - \text{k2} \cdot [\text{species_16}])$$
 (46)

Table 71: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			13000.000		
k2			0.685		

7.23 Reaction reaction_22

This is a reversible reaction of two reactants forming one product.

Name GAD binding Receptor

Reaction equation

$$species_12 + species_4 \Longrightarrow species_16$$
 (47)

Reactants

Table 72: Properties of each reactant.

Id	Name	SBO
species_12	GAD	
species_4	R	

Product

Table 73: Properties of each product.

Id	Name	SBO
species_16	RGAD	

Kinetic Law

Derived unit contains undeclared units

$$v_{23} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_12}] \cdot [\text{species_4}] - \text{k2} \cdot [\text{species_16}])$$
 (48)

Table 74: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			$4.94 \cdot 10^7$		
k2		0.004			

7.24 Reaction reaction_23

This is a reversible reaction of one reactant forming two products.

Name RGAD dissociation

Reaction equation

$$species_16 \rightleftharpoons species_14 + species_8$$
 (49)

Reactant

Table 75: Properties of each reactant.

Id	Name	SBO
species_16	RGAD	

Products

Table 76: Properties of each product.

Id	Name	SBO
species_14	RGA	
species_8	D	

Kinetic Law

Derived unit contains undeclared units

$$v_{24} = \text{vol}(\text{compartment_0}) \cdot (\text{k1} \cdot [\text{species_16}] - \text{k2} \cdot [\text{species_14}] \cdot [\text{species_8}])$$
 (50)

Table 77: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1			2.75		
k2			2940.00		

8 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

Identifiers for kinetic laws highlighted in gray cannot be verified to evaluate to units of SBML substance per time. As a result, some SBML interpreters may not be able to verify the consistency of the units on quantities in the model. Please check if

- parameters without an unit definition are involved or
- volume correction is necessary because the hasOnlySubstanceUnits flag may be set to false and spacialDimensions> 0 for certain species.

8.1 Species species_0

Name A

Notes GAP

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in six reactions (as a reactant in reaction_0, reaction_7, reaction_8, reaction_15, reaction_19, reaction_21).

$$\frac{d}{dt} \text{species}_{0} = -|v_{1}| - |v_{8}| - |v_{9}| - |v_{16}| - |v_{20}| - |v_{22}|$$
(51)

8.2 Species species_1

Name G

Notes inactive G protein

Initial concentration $10^{-8} \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_0, reaction_1, reaction_2 and as a product in reaction_4).

$$\frac{d}{dt} \text{species}_{-1} = |v_5| - |v_1| - |v_2| - |v_3| \tag{52}$$

8.3 Species species_2

Name GA

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_12, reaction_16 and as a product in reaction_0, reaction_14).

$$\frac{d}{dt} \text{species} 2 = |v_1| + |v_{15}| - |v_{13}| - |v_{17}|$$
(53)

8.4 Species species_3

Name T

Notes GTP

Initial concentration $4.68 \cdot 10^{-4} \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_1, reaction_5, reaction_12, reaction_17).

$$\frac{d}{dt} \text{species}_{3} = -|v_{2}| - |v_{6}| - |v_{13}| - |v_{18}| \tag{54}$$

8.5 Species species_4

Name R

Notes Receptor

Initial concentration $10^{-6} \text{ mol} \cdot l^{-1}$

This species takes part in six reactions (as a reactant in reaction_2, reaction_6, reaction_9, reaction_16, reaction_18, reaction_22).

$$\frac{d}{dt} \text{species} = 4 = -|v_3| - |v_7| - |v_{10}| - |v_{17}| - |v_{19}| - |v_{23}|$$
(55)

8.6 Species species_5

Name G*T

Notes active G protein

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_3, reaction_6, reaction_7 and as a product in reaction_1).

$$\frac{d}{dt} \text{species}_{5} = |v_{2}| - |v_{4}| - |v_{7}| - |v_{8}| \tag{56}$$

8.7 Species species_6

Name GD

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_4, reaction_8, reaction_9 and as a product in reaction_3).

$$\frac{d}{dt} \text{species}_{.6} = |v_4| - |v_5| - |v_9| - |v_{10}| \tag{57}$$

8.8 Species species_7

Name Pi

Initial concentration $0.0044 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a product in reaction_3, reaction_10, reaction_13, reaction_20).

$$\frac{d}{dt} \text{species}_{7} = |v_{4}| + |v_{11}| + |v_{14}| + |v_{21}|$$
(58)

8.9 Species species_8

Name D

Notes GDP

Initial concentration $1.49 \cdot 10^{-4} \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a product in reaction_4, reaction_11, reaction_14, reaction_23).

$$\frac{d}{dt} \text{species_8} = |v_5| + |v_{12}| + |v_{15}| + |v_{24}| \tag{59}$$

8.10 Species species_9

Name RG

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_5, reaction_15 and as a product in reaction_2, reaction_11).

$$\frac{d}{dt} \text{species}_{9} = |v_{3}| + |v_{12}| - |v_{6}| - |v_{16}| \tag{60}$$

8.11 Species species_10

Name RG*T

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_10, reaction_19 and as a product in reaction_5, reaction_6).

$$\frac{d}{dt} \text{species}_{10} = |v_6| + |v_7| - |v_{11}| - |v_{20}|$$
(61)

8.12 Species species_11

Name G*AT

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_13, reaction_18 and as a product in reaction_7, reaction_12).

$$\frac{d}{dt} \text{species}_{11} = v_8 + v_{13} - v_{14} - v_{19}$$
 (62)

8.13 Species species_12

Name GAD

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_14, reaction_22 and as a product in reaction_8, reaction_13).

$$\frac{d}{dt} \text{species}_{12} = |v_9| + |v_{14}| - |v_{15}| - |v_{23}| \tag{63}$$

8.14 Species species_13

Name RGD

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_11, reaction_21 and as a product in reaction_9, reaction_10).

$$\frac{d}{dt} \text{species}_{-13} = |v_{10}| + |v_{11}| - |v_{12}| - |v_{22}| \tag{64}$$

8.15 Species species_14

Name RGA

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_17 and as a product in reaction_15, reaction_16, reaction_23).

$$\frac{d}{dt} \text{species}_{-}14 = |v_{16}| + |v_{17}| + |v_{24}| - |v_{18}| \tag{65}$$

8.16 Species species_15

Name RG*AT

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_20 and as a product in reaction_17, reaction_18, reaction_19).

$$\frac{d}{dt} \text{species}_{-15} = |v_{18}| + |v_{19}| + |v_{20}| - |v_{21}|$$
(66)

8.17 Species species_16

Name RGAD

Initial concentration $0 \text{ mol} \cdot l^{-1}$

This species takes part in four reactions (as a reactant in reaction_23 and as a product in reaction_20, reaction_21, reaction_22).

$$\frac{d}{dt} \text{species}_{-16} = |v_{21}| + |v_{22}| + |v_{23}| - |v_{24}|$$
(67)

 $\mathfrak{BML2}^{lAT}$ EX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany