


A


nFeature_RNA
12000
9000
6000
3000
0
24h−blastoids
48h−blastoids
72h−blastoids
Identity
nCount_RNA
4e+05
3e+05
2e+05
1e+05
0e+00
pct_counts_mt
80
60
40
20
0
pct_counts_hb
3
2
1
0
pct_counts_ribo
0.050
0.025
0.000
−0.025
−0.050


B


nCount_RNA distribution per sample
72h−blastoids
48h−blastoids
24h−blastoids
0e+00
1e+05
2e+05
3e+05
4e+05
nCount_RNA

C


nFeature_RNA distribution per sample
72h−blastoids
48h−blastoids
24h−blastoids
0
4000
8000
12000
nFeature_RNA

D


log10(nFeature_RNA) vs log10(nCount_RNA)
Exponential saturation fit: y = Asym + (R0 − Asym) * exp(−exp(lrc) * x)
24h−blastoids
48h−blastoids
72h−blastoids
log10_nFeature
4.5
4.0
3.5
3.0
2.5
2.0
3
4
5
log10_nCount
24h−blastoids: Asym = 5.080, R0 = −4.362, lrc = −0.759 | 48h−blastoids: Asym = 4.638, R0 = −6.525, lrc = −0.499 | 72h−blastoids: Asym = 4.627, R0 = −5.873, lrc = −0.522