

## 13. Douglas-Rachford method and ADMM

- Douglas-Rachford splitting method
- examples
- alternating direction method of multipliers
- image deblurring example
- convergence

# Douglas-Rachford splitting algorithm

$$\text{minimize } f(x) + g(x)$$

$f$  and  $g$  are closed convex functions

**Douglas-Rachford iteration:** start at any  $y^{(0)}$  and repeat for  $k = 1, 2, \dots$ ,

$$x^{(k)} = \text{prox}_f(y^{(k-1)})$$

$$y^{(k)} = y^{(k-1)} + \text{prox}_g(2x^{(k)} - y^{(k-1)}) - x^{(k)}$$

- useful when  $f$  and  $g$  have inexpensive prox-operators
- $x^{(k)}$  converges to a solution of  $0 \in \partial f(x) + \partial g(x)$  (if a solution exists)
- not symmetric in  $f$  and  $g$

# Douglas-Rachford iteration as fixed-point iteration

- iteration on page 13-2 can be written as fixed-point iteration

$$y^{(k)} = F(y^{(k-1)})$$

where

$$F(y) = y + \text{prox}_g(2\text{prox}_f(y) - y) - \text{prox}_f(y)$$

- $y$  is a fixed point of  $F$  if and only if  $x = \text{prox}_f(y)$  satisfies  $0 \in \partial f(x) + \partial g(x)$ :

$$y = F(y)$$

$$\Updownarrow$$

$$0 \in \partial f(\text{prox}_f(y)) + \partial g(\text{prox}_f(y))$$

(proof on next page)

*Proof.*

$$x = \text{prox}_f(y), \quad y = F(y)$$

$$\Updownarrow$$

$$x = \text{prox}_f(y), \quad x = \text{prox}_g(2x - y)$$

$$\Updownarrow$$

$$y - x \in \partial f(x), \quad x - y \in \partial g(x)$$

- therefore, if  $y = F(y)$ , then  $x = \text{prox}_f(y)$  satisfies

$$0 = (y - x) + (x - y) \in \partial f(x) + \partial g(x)$$

- conversely, if  $-z \in \partial f(x)$  and  $z \in \partial g(x)$ , then  $y = x - z$  is a fixed point of  $F$

## Equivalent form of DR algorithm

- start iteration on page 13-2 at  $y$ -update and renumber iterates

$$y^{(k)} = y^{(k-1)} + \text{prox}_g(2x^{(k-1)} - y^{(k-1)}) - x^{(k-1)}$$

$$x^{(k)} = \text{prox}_f(y^{(k)})$$

- switch  $y$ - and  $x$ -updates

$$u^{(k)} = \text{prox}_g(2x^{(k-1)} - y^{(k-1)})$$

$$x^{(k)} = \text{prox}_f(y^{(k-1)} + u^{(k)} - x^{(k-1)})$$

$$y^{(k)} = y^{(k-1)} + u^{(k)} - x^{(k-1)}$$

- make change of variables  $w^{(k)} = x^{(k)} - y^{(k)}$

$$u^{(k)} = \text{prox}_g(x^{(k-1)} + w^{(k-1)})$$

$$x^{(k)} = \text{prox}_f(u^{(k)} - w^{(k-1)})$$

$$w^{(k)} = w^{(k-1)} + x^{(k)} - u^{(k)}$$

# Scaling

algorithm applied to cost function scaled by  $t > 0$

$$\text{minimize } tf(x) + tg(x)$$

- algorithm of page 13-2

$$\begin{aligned}x^{(k)} &= \text{prox}_{tf}(y^{(k-1)}) \\y^{(k)} &= y^{(k-1)} + \text{prox}_{tg}(2x^{(k)} - y^{(k-1)}) - x^{(k)}\end{aligned}$$

- algorithm of page 13-5

$$\begin{aligned}u^{(k)} &= \text{prox}_{tg}(x^{(k-1)} + w^{(k-1)}) \\x^{(k)} &= \text{prox}_{tf}(u^{(k)} - w^{(k-1)}) \\w^{(k)} &= w^{(k-1)} + x^{(k)} - u^{(k)}\end{aligned}$$

- the algorithm is not invariant with respect to scaling
- in theory,  $t$  can be any positive constant; several heuristics exist for adapting  $t$

# Douglas-Rachford iteration with relaxation

- fixed-point iteration with relaxation

$$y^{(k)} = y^{(k-1)} + \rho_k (F(y^{(k-1)}) - y^{(k-1)})$$

$1 < \rho_k < 2$  is overrelaxation,  $0 < \rho_k < 1$  underrelaxation

- algorithm of page 13-2 with relaxation

$$\begin{aligned}x^{(k)} &= \text{prox}_f(y^{(k-1)}) \\y^{(k)} &= y^{(k-1)} + \rho_k (\text{prox}_g(2x^{(k)} - y^{(k-1)}) - x^{(k)})\end{aligned}$$

- algorithm of page 13-5

$$\begin{aligned}u^+ &= \text{prox}_g(x + w) \\x^+ &= \text{prox}_f(x + \rho(u^+ - x) - w) \\w^+ &= w + x^+ - x + \rho(x - u^+)\end{aligned}$$

# Primal-dual formulation

primal:      minimize    $f(x) + g(x)$

dual:        maximize    $-g^*(z) - f^*(-z)$

- use Moreau decomposition to simplify step 2 of DR iteration (page 13-2):

$$x^{(k)} = \text{prox}_f(y^{(k-1)})$$

$$y^{(k)} = x^{(k)} - \text{prox}_{g^*}(2x^{(k)} - y^{(k-1)})$$

- make change of variables  $z^{(k)} = x^{(k)} - y^{(k)}$ :

$$x^{(k)} = \text{prox}_f(x^{(k-1)} - z^{(k-1)})$$

$$z^{(k)} = \text{prox}_{g^*}(z^{(k-1)} + 2x^{(k)} - x^{(k-1)})$$



# Outline

- Douglas-Rachford splitting method
- **examples**
- alternating direction method of multipliers
- image deblurring example
- convergence

# Sparse inverse covariance selection

$$\text{minimize} \quad \text{tr}(CX) - \log \det X + \gamma \sum_{i>j} |X_{ij}|$$

variable is  $X \in \mathbf{S}^n$ ; parameters  $C \in \mathbf{S}_{++}^n$  and  $\gamma > 0$  are given

## Douglas-Rachford splitting

$$f(X) = \text{tr}(CX) - \log \det X, \quad g(X) = \gamma \sum_{i>j} |X_{ij}|$$

- $X = \text{prox}_{tf}(\hat{X})$  is positive solution of  $C - X^{-1} + (1/t)(X - \hat{X}) = 0$   
easily solved via eigenvalue decomposition of  $\hat{X} - tC$  (see homework)
- $X = \text{prox}_{tg}(\hat{X})$  is soft-thresholding

# Spingarn's method of partial inverses

**Equality constrained convex problem** ( $f$  closed and convex;  $V$  a subspace)

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & x \in V\end{array}$$

**Spingarn's method:** Douglas-Rachford splitting with  $g = \delta_V$  (indicator of  $V$ )

$$\begin{aligned}x^{(k)} &= \text{prox}_{tf}(y^{(k-1)}) \\ y^{(k)} &= y^{(k-1)} + P_V(2x^{(k)} - y^{(k-1)}) - x^{(k)}\end{aligned}$$

**Primal-dual form** (algorithm of page 13-8):

$$\begin{aligned}x^{(k)} &= \text{prox}_{tf}(x^{(k-1)} - z^{(k-1)}) \\ z^{(k)} &= P_{V^\perp}(z^{(k-1)} + 2x^{(k)} - x^{(k-1)})\end{aligned}$$

# Application to composite optimization problem

$$\text{minimize } f_1(x) + f_2(Ax)$$

$f_1$  and  $f_2$  have simple prox-operators

- problem is equivalent to minimizing  $f(x_1, x_2)$  over subspace  $V$  where

$$f(x_1, x_2) = f_1(x_1) + f_2(x_2), \quad V = \{(x_1, x_2) \mid x_2 = Ax_1\}$$

- $\text{prox}_{tf}$  is separable:

$$\text{prox}_{tf}(x_1, x_2) = (\text{prox}_{tf_1}(x_1), \text{prox}_{tf_2}(x_2))$$

- projection of  $(x_1, x_2)$  on  $V$  reduces to linear equation:

$$\begin{aligned} P_V(x_1, x_2) &= \begin{bmatrix} I \\ A \end{bmatrix} (I + A^T A)^{-1} (x_1 + A^T x_2) \\ &= \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} A^T \\ -I \end{bmatrix} (I + AA^T)^{-1} (x_2 - Ax_1) \end{aligned}$$

# Decomposition of separable problems

$$\text{minimize} \quad \sum_{j=1}^n f_j(x_j) + \sum_{i=1}^m g_i(A_{i1}x_1 + \cdots + A_{in}x_n)$$

- same problem as page 12-17, but without strong convexity assumption
- we assume the functions  $f_j$  and  $g_i$  have inexpensive prox-operators

## Equivalent formulation

$$\begin{aligned} &\text{minimize} \quad \sum_{j=1}^n f_j(x_j) + \sum_{i=1}^m g_i(y_{i1} + \cdots + y_{in}) \\ &\text{subject to} \quad y_{ij} = A_{ij}x_j, \quad i = 1, \dots, m, \quad j = 1, \dots, n \end{aligned}$$

- prox-operator of first term requires evaluations of  $\text{prox}_{tf_j}$  for  $j = 1, \dots, n$
- prox-operator of 2nd term requires  $\text{prox}_{ntg_i}$  for  $i = 1, \dots, m$  (see page 8-8)
- projection on constraint set reduces to  $n$  independent linear equations

# Decomposition of separable problems

**Second equivalent formulation:** introduce extra splitting variables  $x_{ij}$

$$\begin{aligned} &\text{minimize} && \sum_{j=1}^n f_j(x_j) + \sum_{i=1}^m g_i(y_{i1} + \cdots + y_{in}) \\ &\text{subject to} && x_{ij} = x_j, \quad i = 1, \dots, m, \quad j = 1, \dots, n \\ &&& y_{ij} = A_{ij}x_{ij}, \quad i = 1, \dots, m, \quad j = 1, \dots, n \end{aligned}$$

- make first set of constraints part of domain of  $f_j$ :

$$\tilde{f}_j(x_j, x_{1j}, \dots, x_{mj}) = \begin{cases} f_j(x_j) & x_{ij} = x_j, \quad i = 1, \dots, m \\ +\infty & \text{otherwise} \end{cases}$$

prox-operator of  $\tilde{f}_j$  reduces to prox-operator of  $f_j$

- projection on other constraints involves  $mn$  independent linear equations

# Outline

- Douglas-Rachford splitting method
- examples
- **alternating direction method of multipliers**
- image deblurring example
- convergence

# Dual application of Douglas-Rachford method

## Separable convex problem

$$\begin{array}{ll}\text{minimize} & f_1(x_1) + f_2(x_2) \\ \text{subject to} & A_1x_1 + A_2x_2 = b\end{array}$$

## Dual problem

$$\text{maximize} \quad -b^T z - f_1^*(-A_1^T z) - f_2^*(-A_2^T z)$$

we apply the Douglas-Rachford method (page 13-5) to minimize

$$\underbrace{b^T z + f_1^*(-A_1^T z)}_{g(z)} + \underbrace{f_2^*(-A_2^T z)}_{f(z)}$$



## Douglas Rachford applied to the dual

$$u^+ = \text{prox}_{tg}(z + w), \quad z^+ = \text{prox}_{tf}(u^+ - w), \quad w^+ = w + z^+ - u^+$$

**First line:** use result on page 10-7 to compute  $u^+ = \text{prox}_{tg}(z + w)$

$$\begin{aligned} \hat{x}_1 &= \underset{x_1}{\text{argmin}} (f_1(x_1) + z^T(A_1x_1 - b) + \frac{t}{2}\|A_1x_1 - b + w/t\|_2^2) \\ u^+ &= z + w + t(A_1\hat{x}_1 - b) \end{aligned}$$

**Second line:** similarly, compute  $z^+ = \text{prox}_{tf}(z + t(A_1\hat{x}_1 - b))$

$$\begin{aligned} \hat{x}_2 &= \underset{x_2}{\text{argmin}} (f_2(x_2) + z^TA_2x_2 + \frac{t}{2}\|A_1\hat{x}_1 + A_2x_2 - b\|_2^2) \\ z^+ &= z + t(A_1\hat{x}_1 + A_2\hat{x}_2 - b) \end{aligned}$$

**Third line** reduces to  $w^+ = tA_2\hat{x}_2$

# Alternating direction method of multipliers (ADMM)

1. minimize augmented Lagrangian over  $x_1$

$$x_1^{(k)} = \underset{x_1}{\operatorname{argmin}} \left( f_1(x_1) + (z^{(k-1)})^T A_1 x_1 + \frac{t}{2} \|A_1 x_1 + A_2 x_2^{(k-1)} - b\|_2^2 \right)$$

2. minimize augmented Lagrangian over  $x_2$

$$x_2^{(k)} = \underset{x_2}{\operatorname{argmin}} \left( f_2(x_2) + (z^{(k-1)})^T A_2 x_2 + \frac{t}{2} \|A_1 x_1^{(k)} + A_2 x_2 - b\|_2^2 \right)$$

3. dual update

$$z^{(k)} = z^{(k-1)} + t(A_1 x_1^{(k)} + A_2 x_2^{(k)} - b)$$

this the alternating direction method of multipliers or split Bregman method

# Comparison with other multiplier methods

**Alternating minimization method** (page 12-22) with  $g(y) = \delta_{\{b\}}(y)$

- same dual update, same update for  $x_2$
- $x_1$ -update in alternating minimization method is simpler:

$$x_1^{(k)} = \underset{x_1}{\operatorname{argmin}} (f_1(x_1) + (z^{(k-1)})^T A_1 x_1)$$

- ADMM does not require strong convexity of  $f_1$
- in theory, parameter  $t$  in ADMM can be any positive constant

**Augmented Lagrangian method** (page 12-23) with  $g(y) = \delta_{\{b\}}(y)$

- same dual update
- AL method requires joint minimization of the augmented Lagrangian

$$f_1(x_1) + f_2(x_2) + (z^{(k-1)})^T (A_1 x_1 + A_2 x_2) + \frac{t}{2} \|A_1 x_1 + A_2 x_2 - b\|_2^2$$

# Application to composite optimization (method 1)

$$\text{minimize } f_1(x) + f_2(Ax)$$

- apply ADMM to

$$\begin{array}{ll} \text{minimize} & f_1(x_1) + f_2(x_2) \\ \text{subject to} & Ax_1 = x_2 \end{array}$$

- augmented Lagrangian is

$$f_1(x_1) + f_2(x_2) + \frac{t}{2} \|Ax_1 - x_2 + z/t\|_2^2$$

- $x_1$ -update requires (possibly nontrivial) minimization of

$$f_1(x_1) + \frac{t}{2} \|Ax_1 - x_2 + z/t\|_2^2$$

- $x_2$ -update is evaluation of  $\text{prox}_{t^{-1}f_2}$

## Application to composite optimization (method 2)

introduce an extra ‘splitting’ variable  $x_3$

$$\begin{array}{ll} \text{minimize} & f_1(x_3) + f_2(x_2) \\ \text{subject to} & \begin{bmatrix} A \\ I \end{bmatrix} x_1 = \begin{bmatrix} x_2 \\ x_3 \end{bmatrix} \end{array}$$

- alternate minimization of augmented Lagrangian over  $x_1$  and  $(x_2, x_3)$

$$f_1(x_3) + f_2(x_2) + \frac{t}{2} (\|Ax_1 - x_2 + z_1/t\|_2^2 + \|x_1 - x_3 + z_2/t\|_2^2)$$

- $x_1$ -update: linear equation with coefficient  $I + A^T A$
- $(x_2, x_3)$ -update: decoupled evaluations of  $\text{prox}_{t-1} f_1$  and  $\text{prox}_{t-1} f_2$

# Outline

- Douglas-Rachford splitting method
- examples
- alternating direction method of multipliers
- **image deblurring example**
- convergence

# Image blurring model

$$b = Kx_t + w$$

- $x_t$  is unknown image
- $b$  is observed (blurred and noisy) image;  $w$  is noise
- $N \times N$ -images are stored in column-major order as vectors of length  $N^2$

## Blurring matrix $K$

- represents 2D convolution with space-invariant point spread function
- with periodic boundary conditions, block-circulant with circulant blocks
- can be diagonalized by multiplication with unitary 2D DFT matrix  $W$ :

$$K = W^H \mathbf{diag}(\lambda) W$$

equations with coefficient  $I + K^T K$  can be solved in  $O(N^2 \log N)$  time

# Total variation deblurring with 1-norm

$$\begin{aligned} & \text{minimize} && \|Kx - b\|_1 + \gamma \|Dx\|_{\text{tv}} \\ & \text{subject to} && 0 \leq x \leq \mathbf{1} \end{aligned}$$

second term in objective is *total variation* penalty

- $Dx$  is discretized first derivative in vertical and horizontal direction

$$D = \begin{bmatrix} I \otimes D_1 \\ D_1 \otimes I \end{bmatrix}, \quad D_1 = \begin{bmatrix} -1 & 0 & 0 & \cdots & 0 & 0 & 1 \\ 1 & -1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 1 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & -1 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 1 & -1 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 & -1 \end{bmatrix}$$

- $\|\cdot\|_{\text{tv}}$  is a sum of Euclidean norms:  $\|(u, v)\|_{\text{tv}} = \sum_{i=1}^n \sqrt{u_i^2 + v_i^2}$



# Solution via Douglas-Rachford method

an example of a composite optimization problem

$$\text{minimize } f_1(x) + f_2(Ax)$$

with  $f_1$  the indicator of  $[0, 1]^n$  and

$$A = \begin{bmatrix} K \\ D \end{bmatrix}, \quad f_2(u, v) = \|u\|_1 + \gamma \|v\|_{\text{tv}}$$

**Primal DR method** (page 13-11) and **ADMM** (page 13-19) require:


- decoupled prox-evaluations of  $\|u\|_1$  and  $\gamma \|v\|_{\text{tv}}$ , and projections on  $C$
- solution of linear equations with coefficient matrix

$$I + K^T K + D^T D$$

solvable in  $O(N^2 \log N)$  time

# Example

- $1024 \times 1024$  image, periodic boundary conditions
- Gaussian blur
- salt-and-pepper noise (50% pixels randomly changed to 0/1)



original




noisy/blurred



restored

# Convergence



cost per iteration is dominated by 2D FFTs

# Outline

- Douglas-Rachford splitting method
- examples
- alternating direction method of multipliers
- image deblurring example
- **convergence**

# Douglas-Rachford iteration mappings

define iteration map  $F$  and negative step  $G$  (in notation of page 13-7)

$$F(y) = y + \operatorname{prox}_g(2\operatorname{prox}_f(y) - y) - \operatorname{prox}_f(y)$$

$$G(y) = y - F(y)$$

$$= \operatorname{prox}_f(y) - \operatorname{prox}_g(2\operatorname{prox}_f(y) - y)$$

- $F$  is firmly nonexpansive (co-coercive with parameter 1)

$$(F(y) - F(\hat{y}))^T (y - \hat{y}) \geq \|F(y) - F(\hat{y})\|_2^2 \quad \forall y, \hat{y}$$

- this implies that  $G$  is firmly nonexpansive:

$$\begin{aligned} & (G(y) - G(\hat{y}))^T (y - \hat{y}) \\ &= \|G(y) - G(\hat{y})\|_2^2 + (F(y) - F(\hat{y}))^T (y - \hat{y}) - \|F(y) - F(\hat{y})\|_2^2 \\ &\geq \|G(y) - G(\hat{y})\|_2^2 \end{aligned}$$

*Proof* (of firm nonexpansiveness of  $F$ ).

- define  $x = \text{prox}_f(y)$ ,  $\hat{x} = \text{prox}_f(\hat{y})$ , and

$$v = \text{prox}_g(2x - y), \quad \hat{v} = \text{prox}_g(2\hat{x} - \hat{y})$$

- substitute expressions  $F(y) = y + v - x$  and  $F(\hat{y}) = \hat{y} + \hat{v} - \hat{x}$ :

$$\begin{aligned} & (F(y) - F(\hat{y}))^T (y - \hat{y}) \\ & \geq (y + v - x - \hat{y} - \hat{v} + \hat{x})^T (y - \hat{y}) - (x - \hat{x})^T (y - \hat{y}) + \|x - \hat{x}\|_2^2 \\ & = (v - \hat{v})^T (y - \hat{y}) + \|y - x - \hat{y} + \hat{x}\|_2^2 \\ & = (v - \hat{v})^T (2x - y - 2\hat{x} + \hat{y}) - \|v - \hat{v}\|_2^2 + \|F(y) - F(\hat{y})\|_2^2 \\ & \geq \|F(y) - F(\hat{y})\|_2^2 \end{aligned}$$

inequalities use firm nonexpansiveness of  $\text{prox}_f$  and  $\text{prox}_g$  (page 6-9):

$$(x - \hat{x})^T (y - \hat{y}) \geq \|x - \hat{x}\|_2^2, \quad (2x - y - 2\hat{x} + \hat{y})^T (v - \hat{v}) \geq \|v - \hat{v}\|_2^2$$

# Convergence result

$$\begin{aligned}y^{(k)} &= (1 - \rho_k)y^{(k-1)} + \rho_k F(y^{(k-1)}) \\ &= y^{(k-1)} - \rho_k G(y^{(k-1)})\end{aligned}$$

## Assumptions

- $F$  has fixed points (points  $x$  that satisfy  $0 \in \partial f(x) + \partial g(x)$ )
- $\rho_k \in [\rho_{\min}, \rho_{\max}]$  with  $0 < \rho_{\min} < \rho_{\max} < 2$

## Result

- $y^{(k)}$  converges to a fixed point  $y^*$  of  $F$
- $x^{(k)} = \text{prox}_f(y^{(k-1)})$  converges to a solution  $x^* = \text{prox}_f(y^*)$   
(follows from continuity of  $\text{prox}_f$ )

*Proof:* let  $y^*$  be any fixed point of  $F(y)$  (zero of  $G(y)$ )

consider iteration  $k$  (with  $y = y^{(k-1)}$ ,  $\rho = \rho_k$ ,  $y^+ = y^{(k)}$ ):

$$\begin{aligned}
 \|y^+ - y^*\|_2^2 - \|y - y^*\|_2^2 &= 2(y^+ - y)^T(y - y^*) + \|y^+ - y\|_2^2 \\
 &= -2\rho G(y)^T(y - y^*) + \rho^2 \|G(y)\|_2^2 \\
 &\leq -\rho(2 - \rho) \|G(y)\|_2^2 \\
 &\leq -M \|G(y)\|_2^2
 \end{aligned} \tag{1}$$

where  $M = \rho_{\min}(2 - \rho_{\max})$  (on line 3 we use firm nonexpansiveness of  $G$ )

- (1) implies that

$$M \sum_{k=0}^{\infty} \|G(y^{(k)})\|_2^2 \leq \|y^{(0)} - y^*\|_2^2, \quad \|G(y^{(k)})\|_2 \rightarrow 0$$

- (1) implies that  $\|y^{(k)} - y^*\|_2$  is nonincreasing; hence  $y^{(k)}$  is bounded
- since  $\|y^{(k)} - y^*\|_2$  is nonincreasing, the limit  $\lim_{k \rightarrow \infty} \|y^{(k)} - y^*\|_2$  exists



*Proof (continued)*

- since the sequence  $y^{(k)}$  is bounded, it has a convergent subsequence
- let  $\bar{y}_k$  be a convergent subsequence with limit  $\bar{y}$ ; by continuity of  $G$ ,

$$0 = \lim_{k \rightarrow \infty} G(\bar{y}_k) = G(\bar{y})$$

hence,  $\bar{y}$  is a zero of  $G$  and the limit  $\lim_{k \rightarrow \infty} \|y^{(k)} - \bar{y}\|_2$  exists

- let  $\bar{y}_1$  and  $\bar{y}_2$  be two limit points; the limits

$$\lim_{k \rightarrow \infty} \|y^{(k)} - \bar{y}_1\|_2, \quad \lim_{k \rightarrow \infty} \|y^{(k)} - \bar{y}_2\|_2$$

exist, and subsequences of  $y^{(k)}$  converge to  $\bar{y}_1$ , resp.  $\bar{y}_2$ ; therefore

$$\|\bar{y}_2 - \bar{y}_1\|_2 = \lim_{k \rightarrow \infty} \|y^{(k)} - \bar{y}_1\|_2 = \lim_{k \rightarrow \infty} \|y^{(k)} - \bar{y}_2\|_2 = 0$$

# References

- P. L. Lions and B. Mercier, *Splitting algorithms for the sum of two nonlinear operators*, SIAM Journal on Numerical Analysis (1979).
- D. Gabay, *Applications of the method of multipliers to variational inequalities*, in: Studies in Mathematics and Its Applications (1983).
- J. E. Spingarn, *Applications of the method of partial inverses to convex programming: decomposition*, Mathematical Programming (1985).
- J. Eckstein and D. Bertsekas, *On the Douglas-Rachford splitting method and the proximal algorithm for maximal monotone operators*, Mathematical Programming (1992).
- P.L. Combettes and J.-C. Pesquet, *A Douglas-Rachford splitting approach to nonsmooth convex variational signal recovery*, IEEE Journal of Selected Topics in Signal Processing (2007).
- S. Boyd, N. Parikh, E. Chu, B. Peleato, J. Eckstein, *Distributed optimization and statistical learning via the alternating direction method of multipliers*, Foundations and Trends in Machine learning (2010).
- D. O'Connor and L. Vandenberghe, *Primal-dual decomposition by operator splitting and applications to image deblurring*, SIAM J. Imaging Sciences (2014).

The image deblurring example is taken from this paper.