



ACCESS-CM2 - AMIP realisations


24 May 2019

Roger Bodman | University of Melbourne & CSIRO

AMIP Surface air temperature


Global annual-mean surface air temperature anomalies

HadCRUT4 anomalies, ACCESS-CM2 AMIP realisations: CABLE & JULES



Land-only 65N to 65S global annual-mean surface air temperature





CRUTEM anomalies, ACCESS-CM2 AMIP realisations: CABLE & JULES



Setup for CMIP6

CABLE and JULES versions

Global-mean temperature seasonal bias, AMIP (CABLE) – ERAI, 1979-2014



Annual RMSE = 1.43

DJF RMSE = 1.49

JJA RMSE = 2.17





Annual AWRMSE = 1.04

DJF AWRMSE = 1.29

JJA AWRMSE = 1.34

CABLE version (bg209). ERAI = ERA-Interim reanalysis.

Global-mean rainfall seasonal bias, AMIP (CABLE) – GPCP, 1979-2014



Annual RMSE = 1.11

DJF RMSE = 1.26

JJA RMSE = 1.57

Annual AWRMSE = 1.34

DJF AWRMSE = 1.53

JJA AWRMSE = 1.88

CABLE version (bg209). GPCP observations.


Temp and Precip: global scale biases

AMIP AWRMSE bias			
	A1.3	CABLE	JULES ensemble*
Temperature (model – ERAI)			
annual	1.21	1.04	1.06
djf	1.50	1.29	1.44
jja	1.44	1.34	1.33
Precipitation (model – GPCP)			
annual	1.26	1.34	1.23
djf	1.57	1.53	1.34
jja	2.02	1.88	1.72

A1.3, ACCESS1.3
AMIP from CMIP5

* 3 realisations

Australian land-only temperature anomalies & rainfall amounts



Bars: AWAP observations. Correlations between AWAP and model annual time series

A large, dark, irregular ink blot with splatters on a white background. The blot is roughly circular but has jagged, uneven edges. It is surrounded by numerous small, dark ink droplets and splatters of varying sizes, creating a sense of motion or a recent splash. The overall effect is dramatic and artistic.

The End