UTROGESTAN®
micronised progesterone

#1 GLOBAL LEADER in PROGESTERONE

Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

BESINS HEALTHCARE
Innovating for Well-being

Œstrogel®
ESTRADIOL-17ß

#1 GLOBAL LEADER in ESTRADIOL GEL for Menopausal Hormone Therapy

Source: IQVIA-IMS Health 2021, Q4 MAT Actual in Value

Women takes #NoPause

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN®
micronised progesterone

#1 GLOBAL LEADER in PROGESTERONE

Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Fertility

ORAL

CHRONIC AUB-O[1,2]

Dose	No. of Days	Cycle Days
200 mg OD (1 capsule)	**12 days**	**D15 - 26**

CHRONIC AUB-E [3]

Dose	No. of Days	Cycle Days
200 mg OD (1 capsule)	**22 days**	**D5 - 26**

SECONDARY AMENORRHEA [2,4]

Dose	No. of Days	Cycle Days
400 mg OD (2 capsules)	**10 days**	**at bedtime**

VAGINAL

LUTEAL PHASE SUPPORT (LPS) AFTER IVF/ICSI [5]

200 mg (1 CAPSULE) THREE TIMES DAILY

Should be in the window* **between the evening of the day of oocyte retrieval and Day 3 post oocyte retrieval.

Should be administered* **at least until the day of the pregnancy test.

REFERENCES:
1. https://emedicine.medscape.com/article/257007-treatment#d8
2. https://reference.medscape.com/drug/prometrium-progesterone-micronized-342792
3. Schindler AD, 2003
4. https://www.mayoclinic.org/drugs-supplements/progestin-oral-route-parenteral-route-vaginal-route/proper-use/drg-20069443
5. The ESHRE Guideline Group on Ovarian Stimulation, Bosch E, Broer S, Griesinger G, et al. ESHRE guideline: ovarian stimulation for IVF/ICSI. Hum Reprod Open, 2020(2), hoaa009. https://doi.org/10.1093/hropen/hoaa009

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Maternity

GIVEN UP TO
16
WEEKS
Threatened Miscarriage management

PRISM

PRogesterone in Sponteneous Miscarriage Trial

(UTROGESTAN is the brand used in the PRISM Trial)

NICE National Institute for Health and Care Excellence 2021

1.5 Management of miscarriage

GIVEN UP TO
36
WEEKS
Preterm Birth prevention

FIGO International Federation of Gynecology and Obstetrics
the Global Voice for Women's Health

FIGO good practice recommendations on progestogens for prevention of preterm delivery



Philippine Society of Maternal Fetal Medicine (PSMFM), Inc.

Clinical Practice Guidelines on Preterm Labor and Birth

REFERENCES:

1. https://www.nice.org.uk/terms-andconditions#notice-of-rights). Last updated 24 November 2021
2. Philippine Society of Maternal Fetal Medicine Clinical Practice Guidelines on Preterm Labor and Birth. Fourth Edition. November 2019

THREATENED MISCARRIAGE[1] | **PRETERM BIRTH[2]** | **PRETERM LABOR[2]**

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Maternity

GIVEN UP TO
16
WEEKS
Threatened Miscarriage management

GIVEN UP TO
36
WEEKS
Preterm Birth prevention

DUAL RISK

With early pregnancy bleeding and history of miscarriage

Dose:
400 mg (2 capsules) BID

Timing:
At the time of presentation until 16 completed weeks of gestation

VAGINAL

THREATENED MISCARRIAGE[1]

PRETERM BIRTH[2]

PRETERM LABOR[2]

REFERENCES:

1. https://www.nice.org.uk/terms-andconditions#notice-of-rights). Last updated 24 November 2021
2. Philippine Society of Maternal Fetal Medicine Clinical Practice Guidelines on Preterm Labor and Birth. Fourth Edition. November 2019

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Maternity

GIVEN UP TO
16
WEEKS
Threatened Miscarriage management

GIVEN UP TO
36
WEEKS
Preterm Birth prevention

HIGH RISK

With prior history of Preterm Birth

Dose:
200 mg (1 capsule) OD

Timing:
16 to 36 wks

VAGINAL | ORAL (ALTERNATIVE)

SHORT CERVIX

Asymptomatic but with cervical length ≤ 25 mm

Dose:
200 mg (1 capsule) OD

Timing:
From diagnosis to 36 6/7 weeks

VAGINAL | ORAL (ALTERNATIVE)

MULTIFETAL

Asymptomatic but with cervical length ≤ 25 mm

Timing:
From diagnosis to 36 weeks

Dose:
200 mg (1 capsule) BID

VAGINAL

THREATENED MISCARRIAGE[1]

PRETERM BIRTH[2]

PRETERM LABOR[2]

REFERENCES:

1. https://www.nice.org.uk/terms-andconditions#notice-of-rights). Last updated 24 November 2021
2. Philippine Society of Maternal Fetal Medicine Clinical Practice Guidelines on Preterm Labor and Birth. Fourth Edition. November 2019

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Maternity

GIVEN UP TO
16
WEEKS
Threatened Miscarriage management

GIVEN UP TO
36
WEEKS
Preterm Birth prevention

ACUTE PRETERM LABOR

Combining tocolytics with 400 mg (2 capsules) Q8H

MAINTENANCE TOCOLYSIS

200mg (1 capsule) BID or TID

VAGINAL

THREATENED MISCARRIAGE[1]

PRETERM BIRTH[2]

PRETERM LABOR[2]

REFERENCES:

1. https://www.nice.org.uk/terms-andconditions#notice-of-rights). Last updated 24 November 2021
2. Philippine Society of Maternal Fetal Medicine Clinical Practice Guidelines on Preterm Labor and Birth. Fourth Edition. November 2019

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Maternity

GIVEN UP TO **16** WEEKS
Threatened Miscarriage management

GIVEN UP TO **36** WEEKS
Preterm Birth prevention

DUAL RISK

With early pregnancy bleeding and history of miscarriage

Dose:
400 mg (2 capsules) BID

Timing:
At the time of presentation until 16 completed weeks of gestation

VAGINAL

HIGH RISK

With prior history of Preterm Birth

Dose:
200 mg (1 capsule) OD

Timing:
16 to 36 wks

VAGINAL | ORAL (ALTERNATIVE)

SHORT CERVIX

Asymptomatic but with cervical length ≤ 25 mm

Dose:
200 mg (1 capsule) OD

Timing:
From diagnosis to 36 6/7 weeks

VAGINAL | ORAL (ALTERNATIVE)

MULTIFETAL

Asymptomatic but with cervical length ≤ 25 mm

Timing:
From diagnosis to 36 weeks

Dose:
200 mg (1 capsule) BID

VAGINAL

ACUTE PRETERM LABOR

Combining tocolytics with 400 mg (2 capsules) Q8H

MAINTENANCE TOCOLYSIS

200mg (1 capsule) BID or TID

VAGINAL

THREATENED MISCARRIAGE[1]

PRETERM BIRTH[2]

PRETERM LABOR[2]

REFERENCES:

1. https://www.nice.org.uk/terms-andconditions#notice-of-rights). Last updated 24 November 2021
2. Philippine Society of Maternal Fetal Medicine Clinical Practice Guidelines on Preterm Labor and Birth. Fourth Edition. November 2019

BESINS HEALTHCARE
Innovating for Well-being

UTROGESTAN® micronised progesterone
#1 GLOBAL LEADER in PROGESTERONE
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Taking care of every woman.

Menopause

For women using Menopausal Hormone Therapy (MHT) and still **with an intact uterus**,
the recommended dose of oral micronized progesterone should be administered **for endometrial protection.**

Œstrogel®
ESTRADIOL-17ß

+

UTROGESTAN®
micronised progesterone

100mg daily
(continuous combined MHT regimen)

REFERENCES:

North American Menopause Society (NAMS),2017

BESINS HEALTHCARE
Innovating for Well-being

Œstrogel ESTRADIOL-17B

#1 GLOBAL LEADER in ESTRADIOL GEL for Menopausal Hormone Therapy

Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Let life continue to bloom!

Women takes #NoPause

Estradiol hemihydrate

Oestrogel

Not likely to produce thrombotic risk stroke, and coronary artery disease[1]

Only FDA-approved transdermal Estradiol gel available in the market[2]

Safe and effective in managing symptoms of meno **Pause**[3]

How to use Oestrogel

Endometrial Preparation

REFERENCES:

1. Cobin RH, Goodman NF on behalf of the AACE Committee. Endocrine Practice 2017; 23(7): 869-880
2. FDA Verification Portal. (2021). Food And Drug Administration Philippines. https://verification.fda.gov.ph/drug_productsview.php?showdetail=&registration_number=DR-XY42910
3. NAMS 2017 Position Statement. Menopause 2017; 24(7): 728-753

BESINS HEALTHCARE
Innovating for Well-being

Œstrogel ESTRADIOL-17β
#1 GLOBAL LEADER in ESTRADIOL GEL for Menopausal Hormone Therapy
Source: IQVIA MIDAS®, MAT Q2 2022, 75 countries, LEU MNF

Let life continue to bloom!

Women takes **#NoPause**

3 EASY STEPS TO USE

Estradiol hemihydrate

Oestrogel

1

Squeeze out the gel on the spatula evenly. The usual starting dose is a **single spatula measure (2.5g)once daily.**

2

Gently spread the gel to clean, dry,intact areas of the skin (e.g. **on the abdomen, arms, shoulders, and/or inner thighs**).The area of application should be as large as possible.

3

Allow the gel to dry for 2 minutes before covering the skin with clothing.

Gel does NOT need to be massaged into skin
Gel must NOT be applied on the breasts
or mucous surfaces such as the vulval region
*If the patient forgets to apply a dose, then she should apply
it as soon as possible and apply the next dose at the normal time.

#NoPause

Endometrial Preparation

BESINS HEALTHCARE
Innovating for Well-being



For endometrial preparation in embryo transfer (FET) cycles of women at high risk for deep vein thrombosis, hyperlipdemia, and clotting disturbance you can **use OESTROGEL!**

Estradiol hemihydrate

Oestrogel

(1 spatula measure of Oestrogel = 1.5mg estradiol)

6mg (4 spatula) per day of OESTROGEL

after GnRH agonist supression[1]

Start in the first day of menstruation cycle and continued until endometrial thickness reached 8 mm.[1]

- Clinically proven (**with lower misscarraige rate and higher ongoing pregnancy and live birth rates in comparison to estradiol valerate oral tablet**)[1]
- Avoids first pass hepatic metabolism
- Convenient (easy to administer) and affordable

REFERENCES:

Tehraninejad ES, Kabodmehri R, Rashidi BH, Jafarabadi M, Keikha F, Masomi M, Hagholahi F. International Journal of Reproductive Biomedicine. 2018 Jan;16(1):51.

#NoPause

How to use Oestrogel