

Mathematical Logic (VII)

Yijia Chen

1 Completeness

1.1 Henkin's Theorem

Recall that we fix a set Φ of S -formulas.

Definition 1.1. Let $t_1, t_2 \in T^S$. Then $t_1 \sim t_2$ if $\Phi \vdash t_1 \equiv t_2$. ⊢

Lemma 1.2. (i) \sim is an **equivalence** relation.

(ii) \sim is a **congruence** relation. That is:

- For every n -ary function symbol $f \in S$ and $2 \cdot n$ S -terms $t_1 \sim t'_1, \dots, t_n \sim t'_n$, we have

$$ft_1 \dots t_n \sim ft'_1 \dots t'_n.$$

- For every n -ary relation symbol $R \in S$ and $2 \cdot n$ S -terms $t_1 \sim t'_1, \dots, t_n \sim t'_n$, we have

$$\Phi \vdash Rt_1 \dots t_n \iff \Phi \vdash Rt'_1 \dots t'_n. \quad \vdash$$

Proof: By the equality rule and the substitution rule. □

Now for every $t \in T^S$ we define

$$\bar{t} := \{t' \in T^S \mid t' \sim t\},$$

i.e., the equivalence class of t .

Definition 1.3. The **term structure** for Φ , denoted by \mathcal{T}^Φ , is defined as below. → Mathematical $\{T\}$

(i) The universe is $T^\Phi := \{\bar{t} \mid t \in T^S\}$.

(ii) For every n -ary relation symbol $R \in S$, and $\bar{t}_1, \dots, \bar{t}_n \in T^\Phi$

$$(\bar{t}_1, \dots, \bar{t}_n) \in R^{\mathcal{T}^\Phi} \text{ if } \Phi \vdash Rt_1 \dots t_n.$$

(iii) For every n -ary function symbol $f \in S$, and $\bar{t}_1, \dots, \bar{t}_n \in T^\Phi$

$$f^{\mathcal{T}^\Phi}(\bar{t}_1, \dots, \bar{t}_n) := \overline{ft_1 \dots t_n}.$$

(iv) For every constant $c \in S$

$$c^{\mathcal{T}^\Phi} := \bar{c}.$$

This finishes the construction of \mathcal{T}^Φ . ⊢

Using Lemma 1.2, in particular (ii), it is easy to verify that:

Lemma 1.4. \mathcal{T}^Φ is well-defined. ⊢

To complete the definition of an S-interpretation, we still need to provide an assignment of the variables v_0, v_1, \dots in \mathcal{I}^Φ .

Definition 1.5. For every variable v_i we let

$$\beta^\Phi(v_i) := \bar{v}_i. \quad \dashv$$

Finally we let

$$\mathcal{I}^\Phi := (\mathcal{I}^\Phi, \beta^\Phi).$$

Lemma 1.6. (i) For any $t \in \mathcal{T}^S$

$$\mathcal{I}^\Phi(t) = \bar{t}.$$

(ii) For every **atomic** φ

$$\mathcal{I}^\Phi \models \varphi \iff \Phi \vdash \varphi. \quad \dashv$$

Proof: (i) We proceed by **structural** induction on t .

- $t = v_i$ is a variable. Then

$$\mathcal{I}^\Phi(v_i) = \beta^\Phi(v_i) = \bar{v}_i.$$

- $t = c$ is a constant. Then

$$\mathcal{I}^\Phi(c) = c^{\mathcal{I}^\Phi} = \bar{c}$$

- $t = ft_1 \dots t_n$. Then

$$\begin{aligned} \mathcal{I}^\Phi(ft_1 \dots t_n) &= f^{\mathcal{I}^\Phi}(\mathcal{I}^\Phi(t_1), \dots, \mathcal{I}^\Phi(t_n)) \\ &= f^{\mathcal{I}^\Phi}(\bar{t}_1, \dots, \bar{t}_n) && \text{(by induction hypothesis)} \\ &= \overline{ft_1 \dots t_n}. \end{aligned}$$

(ii) Recall that there are two types of atomic formulas. For the first type, let $\varphi = t_1 \equiv t_2$. Then

$$\begin{aligned} \mathcal{I}^\Phi \models t_1 \equiv t_2 &\stackrel{\text{def}}{\iff} \mathcal{I}^\Phi(t_1) = \mathcal{I}^\Phi(t_2) \\ &\iff \bar{t}_1 = \bar{t}_2 && \text{(by (i))} \\ &\iff t_1 \sim t_2 \\ &\stackrel{\text{def}}{\iff} \Phi \vdash t_1 \equiv t_2. \end{aligned}$$

Second, let $\varphi = Rt_1 \dots t_n$. We deduce

$$\begin{aligned} \mathcal{I}^\Phi \models Rt_1 \dots t_n &\iff (\mathcal{I}^\Phi(t_1), \dots, \mathcal{I}^\Phi(t_n)) \in R^{\mathcal{I}^\Phi} \\ &\iff (\bar{t}_1, \dots, \bar{t}_n) \in R^{\mathcal{I}^\Phi} && \text{(by (i))} \\ &\iff \Phi \vdash Rt_1 \dots t_n. \end{aligned}$$

□

Lemma 1.7. Let φ be an S-formula and x_1, \dots, x_n pairwise distinct variables. Then

(i) $\mathcal{I}^\Phi \models \exists x_1 \dots \exists x_n \varphi$ if and only if there are S-terms t_1, \dots, t_n such that

$$\mathcal{I}^\Phi \models \varphi \frac{t_1 \dots t_n}{x_1 \dots x_n}.$$

(ii) $\mathcal{I}^\Phi \models \forall x_1 \dots \forall x_n \varphi$ if and only if for all S -terms t_1, \dots, t_n we have

$$\mathcal{I}^\Phi \models \varphi \frac{t_1 \dots t_n}{x_1 \dots x_n}.$$

Proof: We prove (i), then (ii) follows immediately.

$$\mathcal{I}^\Phi \models \exists x_1 \dots \exists x_n \varphi$$

$$\stackrel{\text{def}}{\iff} \mathcal{I}^\Phi \frac{a_1 \dots a_n}{x_1 \dots x_n} \models \varphi \text{ for some } a_1, \dots, a_n \in T^\Phi,$$

$$\text{i.e., } \mathcal{I}^\Phi \frac{\bar{t}_1 \dots \bar{t}_n}{x_1 \dots x_n} \models \varphi \text{ for some } t_1, \dots, t_n \in T^S,$$

$$\iff \mathcal{I}^\Phi \frac{\mathcal{I}^\Phi(t_1) \dots \mathcal{I}^\Phi(t_n)}{x_1 \dots x_n} \models \varphi \text{ for some } t_1, \dots, t_n \in T^S, \quad (\text{by Lemma 1.6 (i)})$$

$$\iff \mathcal{I}^\Phi \models \varphi \frac{t_1 \dots t_n}{x_1 \dots x_n} \text{ for some } t_1, \dots, t_n \in T^S, \quad (\text{by the Substitution Lemma}).$$

$$\text{e.g. } \mathcal{I}^\Phi = \{\neg x \equiv x\} \quad \mathcal{I}^\Phi(\mathcal{I}) = \{\varphi \in L^S \mid \mathcal{I} \models \varphi\}$$

Definition 1.8. (i) Φ is **negation complete** if for every S -formula φ

$$\Phi \vdash \varphi \quad \text{or} \quad \Phi \vdash \neg \varphi.$$

\mathcal{I} is a S -interpretation

(ii) Φ contains witnesses if for every S -formula φ and every variable x there is a term $t \in T^S$ with

$$\Phi \vdash \left(\exists x \varphi \rightarrow \varphi \frac{t}{x} \right).$$

$\star \exists x$ will traverse universe

$\mathcal{I}(t)$ might not cover all universe.

Lemma 1.9. Assume that Φ is consistent, negation complete, and contains witnesses. Then for all S -formulas φ and ψ :

(i) $\Phi \vdash \varphi$ if and only if $\Phi \not\vdash \neg \varphi$.

(ii) $\Phi \vdash (\varphi \vee \psi)$ if and only if $\Phi \vdash \varphi$ or $\Phi \vdash \psi$.

(iii) $\Phi \vdash \exists x \varphi$ if and only if there is a term $t \in T^S$ such that $\Phi \vdash \varphi \frac{t}{x}$.

Proof: (i) Assume that $\Phi \vdash \varphi$. Since Φ is consistent, we conclude that $\Phi \not\vdash \neg \varphi$. Conversely, if $\Phi \not\vdash \neg \varphi$, then $\Phi \vdash \varphi$ by the negation completeness.

(ii) The direction from right to left is trivial by \vee -introduction in succedent. For the other direction, assume that $\Phi \vdash (\varphi \vee \psi)$ and $\Phi \not\vdash \varphi$. By the negation completeness, $\Phi \vdash \neg \varphi$. Then for some finite $\Gamma \subseteq \Phi$ we have the following sequent proof.

			\vdots	
m.		Γ_1	$(\varphi \vee \psi)$	
			\vdots	
n.		Γ_2	$\neg \varphi$	
(n+1).	Γ_1	Γ_2	φ	(antecedent by n)
(n+2).	Γ_1	Γ_2	φ	(assumption)
(n+3).	Γ_1	Γ_2	φ	(modified contradiction by n+1 and n+2)
(n+4).	Γ_1	Γ_2	ψ	(assumption)
(n+5).	Γ_1	Γ_2	$(\varphi \vee \psi)$	(V-introduction in antecedent)
(n+6).		Γ_1	ψ	(chain rule by m and n+5)

(iii) Let $\Phi \vdash \exists x\varphi$ and Φ contain witnesses. Thus there is a term $t \in T^S$ such that

$$\Phi \vdash \left(\exists x\varphi \rightarrow \varphi \frac{t}{x} \right).$$

By Modus ponens¹, we conclude $\Phi \vdash \varphi \frac{t}{x}$. The converse is by the rule of the \exists -introduction in succedent. \square

Theorem 1.10 (Henkin's Theorem). *Let $\Phi \subseteq L^S$ be consistent, negation complete, and contain witnesses. Then for every S -formula φ*

$$\mathcal{I}^\Phi \models \varphi \iff \Phi \vdash \varphi.$$

Proof: We proceed by induction on φ .

- φ is atomic. This is Lemma 1.6 (ii).
- $\varphi = \neg\psi$. Then

$$\begin{aligned} \mathcal{I}^\Phi \models \neg\psi &\iff \mathcal{I}^\Phi \not\models \psi \\ &\iff \Phi \not\vdash \psi && \text{(by induction hypothesis)} \\ &\iff \Phi \vdash \neg\psi && \text{(by Lemma 1.9 (i)).} \end{aligned}$$

- $\varphi = (\psi_1 \vee \psi_2)$. We deduce

$$\begin{aligned} \mathcal{I}^\Phi \models (\psi_1 \vee \psi_2) &\iff \mathcal{I}^\Phi \models \psi_1 \text{ or } \mathcal{I}^\Phi \models \psi_2 \\ &\iff \Phi \vdash \psi_1 \text{ or } \Phi \vdash \psi_2 && \text{(by induction hypothesis)} \\ &\iff \Phi \vdash (\psi_1 \vee \psi_2) && \text{(by Lemma 1.9 (ii)).} \end{aligned}$$

- $\varphi = \exists x\psi$.

$$\begin{aligned} \mathcal{I}^\Phi \models \exists x\psi &\iff \mathcal{I}^\Phi \models \psi \frac{t}{x} \text{ for some } t \in T^S && \text{(by Lemma 1.7)} \\ &\iff \Phi \vdash \psi \frac{t}{x} \text{ for some } t \in T^S && \text{(by induction hypothesis)} \\ &\iff \Phi \vdash \exists x\psi && \text{(by Lemma 1.9 (iii)).} \end{aligned}$$

Here, note that the length of $\psi \frac{t}{x}$ could be well larger than that $\exists x\psi$. Thus, our induction is on the so-called **connective rank** of ψ , denoted by $\text{rk}(\varphi)$, which is defined as follows:

$$\text{rk}(\varphi) := \begin{cases} 0 & \text{if } \varphi \text{ is atomic,} \\ 1 + \text{rk}(\psi) & \text{if } \varphi = \neg\psi, \\ 1 + \text{rk}(\psi_1) + \text{rk}(\psi_2) & \text{if } \varphi = (\psi_1 \vee \psi_2), \\ 1 + \text{rk}(\psi) & \text{if } \varphi = \exists x\psi. \end{cases}$$

\square

Corollary 1.11. *Let $\Phi \subseteq L^S$ be consistent, negation complete, and contain witnesses. Then*

$$\mathcal{I}^\Phi \models \Phi.$$

In particular, Φ is satisfiable.

¹That is, if $\Phi \vdash \varphi$ and $\Phi \vdash \varphi \rightarrow \psi$, then $\Phi \vdash \psi$.

2 Exercises

Exercise 2.1. Assume that Φ is inconsistent. Please describe the structure \mathcal{T}^Φ .

Exercise 2.2. Again let $S := \{R\}$ with unary relation symbol R , and

$$\Phi := \{Rx \vee Ry\}.$$

Prove that:

- Φ is consistent.
- $\Phi \not\models Rx$ and $\Phi \not\models Ry$.
- $\mathcal{T}^\Phi \not\models \Phi$.

Exercise 2.3. Let

$$\Phi := \{v_0 \equiv t \mid t \in T^S\} \cup \{\exists v_0 \exists v_1 \neg v_0 \equiv v_1\}.$$

Prove that Φ is consistent, but there is no consistent Ψ with $\Phi \subseteq \Psi \subseteq L^S$ which contains witnesses.
 \dashv

Exercise 2.4. Let \mathcal{A} and \mathcal{B} be two S -structures. A mapping $h : A \rightarrow B$ is a **homomorphism** from \mathcal{A} to \mathcal{B} if the following properties hold.

1. For every n -ary relation symbol $R \in S$ and $a_1, \dots, a_n \in A$ we have

$$(a_1, \dots, a_n) \in R^{\mathcal{A}} \text{ implies } (h(a_1), \dots, h(a_n)) \in R^{\mathcal{B}}.$$

2. For every n -ary function symbol $f \in S$ and $a_1, \dots, a_n \in A$ we have

$$h(f^{\mathcal{A}}(a_1, \dots, a_n)) = f^{\mathcal{B}}(h(a_1), \dots, h(a_n)).$$

3. For every constant $c \in S$

$$h(c^{\mathcal{A}}) = c^{\mathcal{B}}.$$

Now let $\Phi \subseteq L^S$ and \mathcal{A} be an S -structure with $\mathcal{A} \models \Phi$. Prove that there is a homomorphism from the term model \mathcal{T}^Φ to \mathcal{A} .
 \dashv