

# Mathematical Logic (XII)

Yijia Chen

## 1. Theories and Decidability

**Definition 1.1.** A set  $T \subseteq L_0^S$  of  $L$ -sentences is a *theory* if

- $T$  is satisfiable,
- and  $T$  is closed under consequences, i.e., for every  $\varphi \in L_0^S$ , if  $T \vdash \varphi$ , then  $\varphi \in T$ . +

*→ sequent calculus*

**Example 1.2.** Let  $\mathfrak{A}$  be an  $S$ -structure. Then

$$\text{Th}(\mathfrak{A}) := \{\varphi \in L_0^S \mid \mathfrak{A} \models \varphi\}$$

is a theory. +

**Definition 1.3.** Let  $\mathfrak{N} := (\mathbb{N}, +, \cdot, 0, 1)$ . Then  $\text{Th}(\mathfrak{N})$  is called (*elementary*) *arithmetic*. +

**Definition 1.4.** Let  $T \subseteq L_0^S$ . We define

$$T^F := \{\varphi \in L_0^S \mid T \models \varphi\}. \quad +$$

**Lemma 1.5.** All the following are equivalent.

- ① -  $T^F$  is a theory.
- ② -  $T$  is satisfiable.
- ③ -  $T^F \neq L_0^S$ .

*① ⇒ ②  $T^F$  is theory ⇒  $T^F$  satisfiable ⇒  $T \subseteq T^F$  is satisfiable*  
*② ⇒ ③ if  $T^F = L_0^S \Rightarrow T \models \exists x \neg x \equiv x \Rightarrow T$  not satisfiable.*  
*③ ⇒ ① Claim.  $T^F \models \varphi \Rightarrow T \models \varphi$ ;  $\forall A \in T, A \models T^F$*

*☆会考!*

**Definition 1.6.** The *Peano Arithmetic*  $\Phi_{PA}$  consists of the following  $S_{ar}$ -sentences, where  $S_{ar} = \{+, \cdot, 0, 1\}$ :

$$\begin{aligned} \forall x \neg x + 1 \equiv 0, & \quad \forall x \forall y (x + 1 \equiv y + 1 \rightarrow x \equiv y), \\ \forall x x + 0 \equiv x, & \quad \forall x \forall y x + (y + 1) \equiv (x + y) + 1, \\ \forall x x \cdot 0 \equiv 0, & \quad \forall x \forall y x \cdot (y + 1) \equiv x \cdot y + x, \end{aligned}$$

and for all  $n \in \mathbb{N}$ , all variables  $x_1, \dots, x_n, y$ , and all  $\varphi \in L^{S_{ar}}$  with

$$\text{free}(\varphi) \subseteq \{x_1, \dots, x_n, y\}$$

the sentence

$$\forall x_1 \dots \forall x_n \left( \left( \varphi \frac{0}{y} \wedge \forall y \left( \varphi \rightarrow \varphi \frac{y+1}{y} \right) \right) \rightarrow \forall y \varphi \right).$$

*∴  $A \models T^F$   
 ⇒  $T^F$  is closed under  $\vdash$   
 remains to show  
 $T^F$  is satisfiable.  
 proof by contradiction.  
 $\forall \varphi \in L_0^S, T^F \models \varphi$   
 ⇒  $\varphi \in T^F \Rightarrow T^F = L_0^S$   
 → 归谬 (用 FO 表示的命题)*

**Remark 1.7.** It is easy to see that  $\mathfrak{N} \models \Phi_{PA}$ , i.e.,  $\Phi_{PA}^F \subseteq \text{Th}(\mathfrak{N})$ . We will show that  $\Phi_{PA}^F \subsetneq \text{Th}(\mathfrak{N})$ . +

*↓  
 if  $\Phi_{PA} \models \varphi$   
 then  $\mathfrak{N} \models \varphi$*

**Definition 1.8.** Let  $T \subseteq L_0^S$  be a theory.



$$\begin{array}{ccc} \varphi \in \text{Th}(\mathcal{U}) & & \varphi \in \mathcal{D}^k \\ \uparrow & & \uparrow \\ \mathcal{U} \models \varphi \Leftrightarrow \mathcal{D} \models \varphi & & \end{array}$$

(i)  $T$  is *R-axiomatizable* if there exists an R-decidable  $\Phi \subseteq L_0^S$  with  $T = \Phi^\models$ . e.g.  $T = \text{Th}(\mathcal{U})$

(ii)  $T$  is *finitely axiomatizable* if there exists a finite  $\Phi \subseteq L_0^S$  with  $T = \Phi^\models$ .

Clearly any finitely axiomatizable  $T$  is R-axiomatizable.

e.g.  $T = \{\varphi \in L_0^{S_{gr}} \mid G \models \varphi \ \forall G \text{ is group}\}$

**Theorem 1.9.** Every R-axiomatizable theory is R-enumerable.

*Proof:* Let  $T = \Phi^\models$  where  $\Phi \subseteq L_0^S$  is R-decidable. We can effectively generate all derivable sequent proofs and check for each proof whether all the used assumptions belong to  $\Phi$  (by the R-decidability of  $\Phi$ ).

Recall  $\{\varphi \mid \vdash \varphi\}$  is enumerable  
 $\uparrow$   
 additionally check all assumptions

**Remark 1.10.** There are R-axiomatizable theories that are not R-decidable, e.g., for  $S = S_\infty$  and  $\Phi = \emptyset$

$$\Phi^\models = \{\varphi \in L_0^{S_\infty} \mid \vdash \varphi\}. \Rightarrow \text{enumerable but not decidable}$$

**Definition 1.11.** A theory  $T \subseteq L_0^S$  is *complete* if for any  $\varphi \in L_0^S$ , either  $\varphi \in T$  or  $\neg\varphi \in T$ .

**Remark 1.12.** Let  $\mathcal{A}$  be an S-structure. Then the theory  $\text{Th}(\mathcal{A})$  is complete

**Theorem 1.13.** (i) Every R-axiomatizable complete theory is R-decidable.

(ii) Every R-enumerable complete theory is R-decidable.

negation complete  
 $\neg$   
 运行此程序, 要么  $\varphi$  被 print, 要么  $\neg\varphi$ .  
 Remark  $S = \emptyset, \mathcal{D} = \emptyset, T = \mathcal{D}^\models$ .  
 $\exists x \exists y. \neg x = y \in T?$   
 $\Rightarrow$  not complete  
 R-axiomatizable

## 2. The Undecidability of Arithmetic

**Theorem 2.1.**  $\text{Th}(\mathcal{N})$  is not R-decidable.

Corollary  $\text{Th}(\mathcal{U})$  is not R-axiomatizable

Again, for the alphabet  $\mathcal{A} = \{\}$  we consider the halting problem

$$\Pi_{\text{halt}} := \{\omega_{\mathbb{P}} \mid \mathbb{P} \text{ a program over } \mathcal{A} \text{ and } \mathbb{P} : \square \rightarrow \text{halt}\}.$$

For any program  $\mathbb{P}$  over  $\mathcal{A}$  we will construct effectively an  $S_{\text{ar}}$ -sentence  $\varphi_{\mathbb{P}}$  (i.e.,  $\varphi_{\mathbb{P}}$  can be computed by a register machine) such that

$$\mathcal{N} \models \varphi_{\mathbb{P}} \iff \mathbb{P} : \square \rightarrow \text{halt}.$$

将停机问题转化为一个  $\mathcal{N}$  上的  $\varphi$

Assume that  $\mathbb{P}$  consists of instructions  $\alpha_0, \dots, \alpha_k$ . Let  $n$  be the maximum index  $i$  such that  $R_i$  is used by  $\mathbb{P}$ . Recall that a configuration of  $\mathbb{P}$  is an  $(n+2)$ -tuple

$$(L, m_0, \dots, m_n),$$

where  $L \leq k$  and  $m_0, \dots, m_n \in \mathbb{N}$ , meaning that  $\alpha_L$  is the instruction to be executed next and every register  $R_i$  contains  $m_i$ , i.e., the word  $\underbrace{|| \dots ||}_{m_i \text{ times}}$ .

**Lemma 2.2.** For every program  $\mathbb{P}$  over  $\mathcal{A}$  we can compute an  $S_{\text{ar}}$ -formula

$$\chi_{\mathbb{P}}(x_0, \dots, x_n, z, y_0, \dots, y_n)$$

such that for all  $\ell_0, \dots, \ell_n, L, m_0, \dots, m_n \in \mathbb{N}$

$$\mathcal{N} \models \chi_{\mathbb{P}}[\ell_0, \dots, \ell_n, L, m_0, \dots, m_n]$$

if and only if  $\mathbb{P}$ , beginning with the configuration  $(0, \ell_0, \dots, \ell_n)$ , after finitely many steps, reaches the configuration  $(L, m_0, \dots, m_n)$ .



Using the formula  $\chi_{\mathbb{P}}$  in Lemma 2.2, we define

$$\varphi_{\mathbb{P}} := \exists y_0 \cdots \exists y_n \chi_{\mathbb{P}}(0, \dots, 0, \bar{k}, y_0, \dots, y_n),$$

where  $\bar{k} := \underbrace{1 + \cdots + 1}_{k \text{ times}}$ . Then By Lemma 2.2, we conclude  $\mathfrak{N} \models \varphi_{\mathbb{P}}$  if and only if  $\mathbb{P}$ , beginning with the initial configuration  $(0, 0, \dots, 0)$ , after finitely many steps, reaches the configuration  $(k, m_0, \dots, m_n)$ , i.e.,  $\mathbb{P} : \square \rightarrow \text{halt}$ . This finishes our proof of Theorem 2.1.  $\square$

By Theorem 2.1, Theorem 1.13, and Remark 1.12:

**Corollary 2.3.**  $\text{Th}(\mathfrak{N})$  is neither  $R$ -axiomatizable nor  $R$ -enumerable. Thus

$$\Phi_{\text{PA}}^{\models} \subsetneq \text{Th}(\mathfrak{N}). \quad \dashv$$

**Proof of Lemma 2.2.** Recall that  $\chi_{\mathbb{P}}$  expresses in  $\mathfrak{N}$  that there is an  $s \in \mathbb{N}$  and a sequence of configurations  $C_0, \dots, C_s$  such that

- $C_0 = (0, x_0, \dots, x_n)$ ,
- $C_s = (z, y_0, \dots, y_n)$ ,
- for all  $i < s$  we have  $C_i \xrightarrow{\mathbb{P}} C_{i+1}$ , i.e., from the configuration  $C_i$  the program  $\mathbb{P}$  will reach  $C_{i+1}$  in one step.

We slightly rewrite the above formulation as that there is an  $s \in \mathbb{N}$  and a sequence of natural numbers

$$\underbrace{a_0, \dots, a_{n+1}}_{C_0} \underbrace{a_{n+2}, \dots, a_{(n+2)+(n+1)}}_{C_1} \cdots \underbrace{a_{s \cdot (n+2)}, \dots, a_{s \cdot (n+2) + (n+1)}}_{C_s} \quad (1)$$

such that

- $a_0 = 0, a_1 = x_0, \dots, a_{n+1} = x_n$ ,
- $a_{s \cdot (n+2)} = z, a_{s \cdot (n+2) + 1} = y_0, \dots, a_{s \cdot (n+2) + (n+1)} = y_n$ ,
- for all  $i < s$  we have

$$\left( a_{i \cdot (n+2)}, \dots, a_{i \cdot (n+2) + (n+1)} \right) \xrightarrow{\mathbb{P}} \left( a_{(i+1) \cdot (n+2)}, \dots, a_{(i+1) \cdot (n+2) + (n+1)} \right).$$

Observe that the length of the sequence (1) is unbounded, so we cannot quantify it directly in  $\mathfrak{N}$ . So we need the following beautiful (elementary) number-theoretic tool.

**Lemma 2.4** (Gödel's  $\beta$ -function). There is a function  $\beta : \mathbb{N}^3 \rightarrow \mathbb{N}$  with the following properties.

(i) For every  $r \in \mathbb{N}$  and every sequence  $(a_0, \dots, a_r)$  in  $\mathbb{N}$  there exist  $t, p \in \mathbb{N}$  such that for all  $i \leq r$

$$\beta(t, p, i) = a_i. \quad (\text{if no such } a \text{ exists, } \beta(t, p, i) := 0)$$

(ii)  $\beta$  is definable in  $L^{S_{\text{ar}}}$ . That is, there is an  $S_{\text{ar}}$ -formula  $\varphi_{\beta}(x, y, z, w)$  such that for all  $t, q, i, a \in \mathbb{N}$

$$\mathfrak{N} \models \varphi_{\beta}[t, q, i, a] \iff \beta(t, q, i) = a.$$

$\hookrightarrow$  孔的  $\beta$ (assignment) 满足  $\beta(x) = t \quad \beta(y) = q$   
 $\beta(z) = i \quad \beta(w) = a$

if Lemma 2.4 is correct:



$$x_p := \exists p \exists t \exists s \left( \varphi_\beta(t, p, 0, 0) \wedge \varphi_\beta(t, p, 1, x_0) \wedge \dots \wedge \varphi_\beta(t, p, \overline{n+1}, x_n) \right. \\
\left. \wedge \forall i (i < s \rightarrow \forall u \forall u_0 \dots \forall u_n \forall u' \forall u'_0 \dots \forall u'_n \right. \\
\left. (\varphi_\beta(t, p, i \cdot (u+2), u) \wedge \dots \wedge \varphi_\beta(t, p, i \cdot (u+2) + n+1, u_n) \right. \\
\left. \wedge \varphi_\beta(t, p, (i+1) \cdot (u+2), u') \wedge \dots \wedge \varphi_\beta(t, p, (i+1) \cdot (u+2) + n+1, u'_n) \right) \\
\rightarrow "(u, u_0, \dots, u_n) \xrightarrow{IP} (u', u'_0, \dots, u'_n))"$$

Proof: Let  $(a_0, \dots, a_r)$  be a sequence over  $\mathbb{N}$ . Choose a prime

$$p > \max\{a_0, \dots, a_r, r+1\},$$

and set

$$t := 1 \cdot p^0 + a_0 \cdot p^1 + 2 \cdot p^2 + a_1 \cdot p^3 + \dots + (i+1) \cdot p^{2i} + a_i \cdot p^{2i+1} \\
+ \dots + (r+1) \cdot p^{2r} + a_r \cdot p^{2r+1}. \quad (2)$$

In other words, the  $p$ -adic representation of  $t$  is precisely

$$a_r(r+1) \dots a_i(i+1) \dots a_1 2 a_0 1.$$

Claim. Let  $i \leq r$  and  $a \in \mathbb{N}$ . Then  $a = a_i$  if and only if there are  $b_0, b_1, b_2 \in \mathbb{N}$  such that:

$$(B1) \quad t = b_0 + b_1((i+1) + a \cdot p + b_2 \cdot p^2),$$

$$(B2) \quad a < p,$$

$$(B3) \quad b_0 < b_1,$$

$$(B4) \quad b_1 = p^{2m} \text{ for some } m \in \mathbb{N}.$$

Proof of the claim. Assume  $a = a_i$ . We set

$$b_0 := 1 \cdot p^0 + a_0 \cdot p^1 + 2 \cdot p^2 + a_1 \cdot p^3 + \dots + i \cdot p^{2i-2} + a_{i-1} \cdot p^{2i-1}$$

$$b_1 := p^{2i}$$

$$b_2 := (i+2) + a_{i+1} \cdot p + \dots + a_r \cdot p^{2(r-i)-1}.$$

By (2) it is routine to verify that all (B1)–(B4) hold.

Conversely,

$$t = (1 \cdot p^0 + a_0 \cdot p^1 + 2 \cdot p^2 + a_1 \cdot p^3 + \dots + i \cdot p^{2i-2} + a_{i-1} \cdot p^{2i-1})$$

$$+ (i+1) \cdot p^{2i} + a \cdot p^{2i+1}$$

$$+ ((i+2) + a_{i+1} \cdot p + \dots + a_r \cdot p^{2(r-i)-1}) \cdot p^{2i+2}$$

$$= b_0 + (i+1) \cdot p^{2m} + a \cdot p^{2m+1} + b_2 \cdot p^{2m+2};$$

用偶数位定位  $m$ , s.t.  $m \equiv i$

→  $p$ -adic 表示唯一

It is well known that the  $p$ -adic representation of any number is unique. Together with  $b_0 < p^{2m}$ , we conclude  $a = a_i$ .

Since  $p$  is chosen to be a prime, it is easy to verify that (B4) is equivalent to

(B4')  $b_1$  is a square, and for any  $d > 1$  if  $d \mid b_1$ , then  $p \mid d$ .

Finally for every  $t, q, i \in \mathbb{N}$  we define  $\beta(t, q, i)$  to be *smallest*  $a \in \mathbb{N}$  such that there are  $b_0, b_1, b_2 \in \mathbb{N}$  such that

$$- \quad t = b_0 + b_1((i+1) + a \cdot q + b_2 i^2),$$

$$- \quad a < q,$$

$$- \quad b_0 < b_1,$$

$$- \quad b_1 \text{ is a square, and for any } d > 1 \text{ if } d \mid b_1, \text{ then } q \mid d.$$

对相同序列, 用相同  $q$  去映射

再用  $t$  去寻址.



If no such  $a$  exists, then we let  $\beta(t, q, i) := 0$ .

By the above argument, (i) holds by choosing  $q$  to be a sufficiently large prime. To show (ii) we define

$$\varphi_\beta(x, y, z, w) := \left( \psi(x, y, z, w) \wedge \forall w' (\psi(x, y, z, w') \rightarrow (w' \equiv w \vee w < w'^1)) \right) \\ \vee \left( \neg \psi(x, y, z, w) \wedge w \equiv 0 \right).$$

Here  $\psi(x, y, z, w)$  expresses the properties (B1), (B2), (B3), and (B4'):

$$\psi(x, y, z, w) := \exists u_0 \exists u_1 \exists u_2 \left( x \equiv u_0 + u_1 \cdot ((z + 1) + w \cdot y + u_2 \cdot y \cdot y) \right. \\ \wedge w < y \wedge u_0 < u_1 \\ \left. \wedge \exists v u_1 \equiv v \cdot v \wedge \forall v (\exists v' u_1 \equiv v \cdot v' \rightarrow (v \equiv 1 \vee \exists v' v \equiv y \cdot v')) \right).$$

□

### 3. Exercises

**Exercise 3.1.** Prove that

$$\Phi_{\text{PA}} \models \forall x \forall y \ x + y \equiv y + x. \quad \dashv$$

**Exercise 3.2.** Let  $T$  be an  $R$ -enumerable theory. Show that  $T$  is  $R$ -axiomatizable. \dashv

**Exercise 3.3.** Construct an  $S_{\text{ar}}$ -formula  $\varphi_{\text{exp}}(x, y, z)$  such that for every  $a, b, c \in \mathbb{N}$

$$c = a^b \iff \mathfrak{N} \models \varphi_{\text{exp}}[a, b, c]. \quad \dashv$$

---

<sup>1</sup> $w < w'$  stands for the formula  $\exists v (\neg v \equiv 0 \wedge w + v \equiv w')$ .