

South Africa drought and wildlife survival

Andrea Corrado 20205529

Contents

1	Introduction	2
2	Technical Background	2
2.1	Generalized Lienar Model	2
2.2	Generalized Additive Model	2
2.3	Weighted Least Square	3
2.4	Local Regression	3
2.5	Ordinary Differential Equation	3
2.6	Levenberg-Marquardt algorithm	3
2.7	Sensitivity Analysis	3
3	Exploratory Data Analysis	3
3.1	Rainfall amount	3
4	Climate Modeling	4
5	Species count Modelling	7
6	Conclusion	7

Contents

Abstract

1 Introduction

During last decades, interest and awareness about the climate change has steeply increased. There have been several studies examining and reporting the potential consequences of these phenomenon. For instance, the temperature and rainfall change has been of particular interest. Since last century, we have witnessed a increasing frequency in the number and severity of droughts in particular territories such as South Africa *SA* [1]. The consequences here can be very dramatic. Due to the fact that area such as *SA* suffer from low economic power, a drought may drive towards dramatic consequences. It can affect either agriculture, for a period that lasts up to 6 months, either hydrological up to 24 months [1].

There have been several studies showing how to reduce the impact of drought on the agriculture field by growing restraint to higher temperature and computational model to predict droughts [2].

However, in this paper, we wish to focus our attention on the drought consequences on wildlife population evolution. For instance, different studies have shown as animals mortality grows during the highest temperature period within the driest areas [3]. We will expand this idea to more recent data about waterbirds [4]. This data set will provide the number of individual for different species in each year within the last 45 years. By doing so, we are able to track the evolution of the species during time. However, the data set will not provide the number of new death, new born and the relative causes. It will be our job to build a model which allows us to investigate the relation other factors. In particular the data we are going to analyse regard the global temperature and rain fall amount [5] from 1901 up to 2020 which would give a foundation to build on the model of interest.

Having access to this information, our goal is to build a model which relates the climate change to the number of individual for each species and predict how, following the current trend, these would evolve over time and the potential consequences these could drive towards.

In section 2 we give a brief introduction to the methodology used within this research project. In section 3 we present the data with an exploratory data analysis. In section 4 we model the climate data and provide prediction on a possible future scenario. In section 5 we model the species count in

relation to the climate data and provide prediction about the future number of individuals. In section 6 we discuss the finding of the project.

2 Technical Background

2.1 Generalized Lienar Model

In order to achieve the goal, we are going to employ different methodology which would allow us to model the data of interest. In particular, first of all we will related the temperature data to the rain fall amount. In order to do so, we will employ the Generalized Linear Model [6] which theory would allow to establish a linear relationship between the quantity of interest (rain) and a set of covariates (temperature) plus a stochastic term which models the uncertainty about the random variable realisation

$$g(\mathbf{Y}) = \mathbf{X}\beta + \epsilon$$

where we define a distribution \mathcal{D} of interest on the stochastic term ϵ

$$\epsilon \sim \mathcal{D}(\theta)$$

and a link function $g(\cdot)$ which is distribution dependent.

The model will be validate by a set of diagnostic on the residuals $\mathbf{Y} - \hat{\mathbf{Y}}$ which provide meaningful insight on the goodness of fitness.

2.2 Generalized Additive Model

We will further extend the *GLM* [6] theory to allow for non-linear relationship between the predictors \mathbf{X} and the response \mathbf{Y} exploiting the *Generalized Additive Model* theory [7] while maintaining the model linear in the parameters

$$g(\mathbf{Y}) = f(\mathbf{X}) + \epsilon$$

where we define a distribution \mathcal{D} of interest on the stochastic term ϵ

$$\epsilon \sim \mathcal{D}(\theta)$$

and a link function $g(\cdot)$ which is distribution dependent. Moreover, we will define a function $f(\cdot)$ which usually is a non linear function such as a *spline* [8].

2.3 Weighted Least Square

During the research project, we perform different model estimation and to remedy to inadequate diagnostics, such as those violating the homoskedasticity assumption, we will employ *Weighted Least Square* estimation theory [9] which allows us to iteratively estimate a weight matrix \mathbf{W} to be employed in the model estimation as a measure of the importance for each unit \mathbf{x}_i $i = \{1, \dots, N\}$ where N is the total number of observations.

2.4 Local Regression

2.5 Ordinary Differential Equation

To track the evolution of a phenomenon over time, it is natural to think about *Ordinary Differential Equation (ODE)* theory [10] which allows us to keep track of the change of a quantity in continuous time

$$\frac{d}{dt}x = f(x)$$

given initial condition $f(x, t = 0) = x_0$

2.6 Levenberg-Marquardt algorithm

Very often, the function defined within an ODE is characterized by a set of parameters β . In empirical studies, we often have access to data realization $\{(y_i, \mathbf{x}_i)\}_{i=1}^n$ we want to find the parameters β so that $y = f(\mathbf{x}, \beta)$ best fits the data of interest. The *Levenberg-Marquardt* [11] provide a useful and efficient solution to the problem by making smart combination of *Gauss-Newton* and *Gradient Descent* theory.

2.7 Sensitivity Analysis

Mathematical modeling is often subject to very strong assumption which may be difficult to evaluate. In order to provide a robust framework, it is necessary to validate the model under perturbations on the findings. This is why, we will exploit *Local Sensitivity Analysis* theory [12] to evaluate the robustness of the estimated models.

3 Exploratory Data Analysis

3.1 Rainfall amount

Within this exploratory data analysis context, we give a short introduction to the data by illustrating some characteristics. In the first place, we investigate the rainfall data set. We have access to the information between January 1901 up to December 2020. For simplicity, we decide to compute an annual average and inspect the resulting distribution.




Figure 1: Yearly rainfall distribution

From figure 1 we can observe that the data distribution is somewhat bell-shaped. However, we can easily notice that the distribution is right skewed. Clearly, the phenomenon is left bounded by $0 \leq x \in \mathbb{R}_+$ and we can rarely observe very extreme event such as very heavy precipitation. We are able to quantify the amount of skewness by computing the data sample moments ratio known as the *Skewness* index which result to be 0.42, agreeing with the histogram depicted.

Afterwards, we investigate the temperature data. As for the rain data set, we have availability of the information from 1901 up to 2020. In figure ?? we depict the annual temperature distribution.

We can clearly observe a fairly strong right skewness indicating more extreme events such as sweltering days than very cold days. The phenomenon now takes values in the real domain $x \in \mathbb{R}$ and present a skewness index of 0.43.

In order to give track the evolution of the phenomenon over time and to give an idea of the overall trend we make use of non parametric model, meaning that we do not assume anything but a relation between \mathbf{x} and \mathbf{y} governed by a function




Figure 2: Annual temperature distribution

$f(\cdot)$. The method employed is the locally weighted regression discussed in section 2.

Therefore, to track the evolution of the phenomenon we fit the model for different values of the span. In particular:

- 100% of the neighbor data to take into account the global context (violet)
- 50% of the neighbor data to allow for a more local structure (blue)

By investigating the results in ?? we can notice how the global fit to the rain data shows a constant trend up to the 1950 which evolves in a slightly linear decreasing trend onward. On the other hand, the local fit shows a constant trend up to the 1980s which evolves in a steeper decrease for the later year. Nevertheless, both the models suggest an evolution of the amount of rainfall which changes over time.





Figure 3: Annual rainfall

Afterwards, we fit a the same two models for the temperature data. Here the relation with time is much clearer. For instance, we can observe how the global fit shows a linearly increasing trend over time, with constant velocity. On the other hand, the local fit shows an increasing trend with different intensities in different period allowing for non linear relationship. Nevertheless, the overall trend does not change dramatically. Annual temperature has clearly been increasing over the last century.



	year	rain	temp
year	1.00	-0.06	0.82
rain	-0.06	1.00	-0.33
temp	0.82	-0.33	1.00




Table 1: Correlation table

In order to give a global idea of the data we are investigating we would like to present the pairs plot showing each pair combination plus the estimated correlation matrix \mathbf{R} in table ?. As mentioned earlier, in figure ?? we can notice the strong relationship between temperature and year $\hat{\rho}_{t,y} = 0.82$ and a weaker relationship between rain and year $\hat{\rho}_{r,y} = -0.06$. However, we can see a clear relationship between year and rain $\hat{\rho}_{r,t} = -0.33$ which may be worth further investigation.

4 Climate Modeling

We are now going to propose a very simple physical model to describe the temperature behavior. Ascertained that temperature heavily evolves over time, we present a very simple ordinary different equation given the temperature τ and time t

$$\frac{d}{dt} \tau = \gamma \tau$$



which posit a linear evolution of the phenomenon over time. Moreover, we add initial condition equal to the average temperature values of the first 5 year. The reason behind this choice is that, within this project, we do not have access to older data and during the exploratory data analysis in section ref{sec:eda} we have seen how the phenomenon is highly variable. Therefore, we choose the mean as initial condition

$$f(\tau, t = 0) = \tau_0 = 16.8$$

Then we need to tune the parameter γ and we are going to do so exploiting the Levenberg-Marquardt algorithm discussed in section 2.

The algorithm applied to the data set of interest returns a point estimate of $\hat{\gamma} = 7.6528 \times 10^{-4}$ and an estimated standard error of $\hat{SE}(\hat{\gamma}) = 2.43 \times 10^{-5}$, resulting in the 95% confidence interval

$$P(\gamma \in [0.00071764, 0.00081291]) = 0.95 \quad (1)$$

Overall, we can claim that, according to the model of interest, the global temperature increases over time by a factor within the region in (1). We graphically present the estimated model in figure 5. From the plot we can easily notice how the mean temperature increases over time. It also very noticeable how the model may underfit the data. It could be worth trying to fit a more complex model including higher order terms to address a certain amount of non-linearity. However, while these models may outperform the current in the data representation, they may be very unstable when employed in extrapolation task. The final goal here is to produce prediction about the future and a non-linear model may lead to very erratic conclusion. Therefore, we decide that this simple model fairly represents the phenomenon of interest.

We have now available a model which may lead us to future prediction. In particular, we are interested in what would happen in a 50 years time following the current trend. According to the estimated model and depicted in figure 6, the global temperature may continue to rise up to reaching 19 Celcius average degree in 50 years time, with a reasonably tight confidence interval.

A fundamental requisite for a mathematical model is the robustness to its input. Here, the data is empirically observed and the parameter γ is estimated from it. The question we want to address is whether we can rely on this parameter and whether the model output would change if the parameters in input changed. In

Figure 4: Year, Rain, Temperature scatterplot

order to tackle this question, a common and reliable choice is local sensitivity analysis discussed in section 2. Given the model

$$f(\tau) = \gamma \tau$$

the method allow us to derive the sensitivity equation as

$$\begin{aligned} \frac{d}{dt}s &= \frac{d}{d\gamma}f s + \frac{d}{d\gamma}f \\ &= \gamma s + \tau \end{aligned} \quad (2)$$

Therefore, we will need to deal with the system of ordinary differential equation

$$\begin{cases} \frac{d}{dt}\tau = \gamma\tau \\ \frac{d}{dt}s = \gamma s + \tau \end{cases} \quad (3)$$




Figure 5: Annual temperature modeled by Ordinary Differential Equation




Figure 6: 50 out of sample years temperature prediction





Figure 7: a) evolution of temperature in time with input parameter $\pm 10 SE$; b) evolution of γ parameter in time $\pm 10 SE$

5 Species count Modelling

6 Conclusion

References

- [1] M. R. . Y. Richard, “Intensity and spatial extension of drought in South Africa at different time scales,” *African Journals Online*, vol. 29, no. 4, 2003.
- [2] M. Grayson, “Agriculture and drought,” *Nature Outlook*, vol. 501, no. 7468, 2013.
- [3] J. C. H. et. al, “Mortality of wildlife in Nairobi National Park, during the drought of 1973-1974,” *African Journal of Ecology*, vol. 15, no. 1-18, 1977.
- [4] G. G. B. I. Facility, “Free and open access to biodiversity data.” www.gbif.org/occurrence/.
- [5] W. B. Group, “Climate change knowledge portal.” <https://climateknowledgeportal.worldbank.org/download-data>.
- [6] P. McCullagh and J. Nelder, *Generalized Linear Models*. International series of monographs on physics, Chapman and Hall, 1989.
- [7] R. T. Trevor Hastie, *Generalized Additive Models*. 1990.
- [8] M. W. M. Aertsa, G. Claeskensb, “Some theory for penalized spline generalized additive models,” *Journal of Statistical Planning and Inference*, vol. 103, no. 455-470, 2002.
- [9] P. M. Hooper, “Iterative Weighted Least Squares Estimation in Heteroscedastic Linear Models,” *Journal of the American Statistical Association*, vol. 103, no. 88:421, 2012.
- [10] G. Gbur, *Mathematical methods for optical physics and engineering*. 2011.
- [11] M. J.J., *The Levenberg-Marquardt algorithm: Implementation and theory*. Lecture Notes in Mathematics, Springer, 1978.
- [12] J. Morio, “Global and local sensitivity analysis methods for a physical system,” *European Journal of Physics*, vol. 32, no. 6, 2011.