Definition (Traced monoidal category). A symmetric monoidal category $\langle \mathbf{C}, \boldsymbol{\otimes}_{\mathbf{C}}, \mathbf{1}_{\mathbf{C}}, \mathbf{br} \rangle$ is said to be *traced* if it is equipped with a family of functions

$$\operatorname{Tr}_{X,Y}^{Z}$$
: $\operatorname{Hom}_{\mathbb{C}}(X \otimes_{\mathbb{C}} Z; Y \otimes_{\mathbb{C}} Z) \to \operatorname{Hom}_{\mathbb{C}}(X; Y)$,

satisfying the following axioms:

1. Vanishing: For all morphisms $f: X \to Y$ in \mathbb{C} ,

$$\operatorname{Tr}^1_{X,Y}(f) = f.$$

Furthermore, for all morphisms $f: X \otimes_{\mathbb{C}} Z \otimes_{\mathbb{C}} U \to Y \otimes_{\mathbb{C}} Z \otimes_{\mathbb{C}} U$ in \mathbb{C} :

$$\operatorname{Tr}_{X,Y}^{Z \otimes_{\mathbf{C}} U}(f) = \operatorname{Tr}_{X,Y}^{Z} \left(\operatorname{Tr}_{X \otimes_{\mathbf{C}} Z, Y \otimes_{\mathbf{C}} Z}^{U}(f) \right).$$

2. Superposing: For all morphisms $f: X \otimes_{\mathbb{C}} Z \to Y \otimes_{\mathbb{C}} Z$ in \mathbb{C} :

$$\operatorname{Tr}^{Z}_{V \otimes_{\mathbf{C}} X, V \otimes_{\mathbf{C}} Y}(\operatorname{Id}_{V} \otimes_{\mathbf{C}} f) = \operatorname{Id}_{V} \otimes_{\mathbf{C}} \operatorname{Tr}^{Z}_{X, Y}(f).$$

3. Yanking:

$$\operatorname{Tr}_{Z,Z}^{Z}(\operatorname{br}_{Z,Z}) = \operatorname{Id}_{Z}.$$