

Gap Crossing Problem with CasADi

Francesco Roscia, Matteo Sodano

Consider a robot with $n_c = 4$ contact points with the ground.

The quadruped is modeled as two compass bipeds linked by a prismatic joint. We denote with $\mathbf{r}_{CoM,j}$ the position of the CoM of the j -th biped, where $j \in \{1(B), 2(F)\}$.

State: $\mathbf{x} = [\mathbf{r}_{CoM,j}^{[1]} \dots \mathbf{r}_{CoM,j}^{[n_{step}]} \mathbf{r}_{C,i}^{[1]} \dots \mathbf{r}_{C,i}^{[n_{step}]} \mathbf{F}_{C,i}^{[1]} \dots \mathbf{F}_{C,i}^{[n_{step}]}]$,

with $i \in \{1(BR), 2(BL), 3(FR), 4(FL)\}$. The superscript $[k]$ denotes the value of the variable at the k -th step. B: back, F: front, R: right, L: left.

We want to find the sequence \mathbf{x}^* that moves the robot from $\mathbf{r}_{CoM,j}^{start}$ to $\mathbf{r}_{CoM,j}^{goal}$ solving the following NLP problem:

$$\min_{\mathbf{x}} w_{CoM} \sum_{k=1}^{n_{step}} \sum_{j=1}^2 \|\mathbf{r}_{CoM,j}^{[k]} - \mathbf{r}_{CoM,j}^{goal}\|^2 + w_F \sum_{k=1}^{n_{step}} \|\mathbf{F}_C^{[k]}\|^2 \quad (1)$$

$$m\mathbf{g} + \mathbf{G}_{CD}\mathbf{F}_c^{[k]} = \mathbf{0} \quad \text{centroidal dynamics} \quad (2)$$

$$\mathbf{f}_{env}(\mathbf{r}_{C,i}) = \mathbf{0} \quad \text{feet touch the floor} \quad (3)$$

$$\begin{cases} \mathbf{F}_{C,i} \cdot \mathbf{n}_{C,i} > F_{thr} \\ \|\mathbf{F}_{C,i}^t\| \leq \mu_i(\mathbf{F}_{C,i} \cdot \mathbf{n}_{C,i}) \end{cases} \quad \text{friction cones} \quad (4)$$

$$\|\mathbf{r}_{C,i}^{[k]} - \mathbf{r}_{C,i}^{[k+1]}\|^2 \leq 0.09 \quad \text{the step is inside a circle of radius 0.3, centered in } \mathbf{r}_{C,i}^{[k]} \quad (5)$$

$$1 \leq \|\mathbf{r}_{C,BR}^{[k]} - \mathbf{r}_{C,FR}^{[k]}\|^2 \leq 9 \quad \text{distance between feet is bounded} \quad (6)$$

$$1 \leq \|\mathbf{r}_{C,BL}^{[k]} - \mathbf{r}_{C,FL}^{[k]}\|^2 \leq 9 \quad " \quad (7)$$

$$0.25 \leq \|\mathbf{r}_{C,BR}^{[k]} - \mathbf{r}_{C,BL}^{[k]}\|^2 \leq 2.25 \quad " \quad (8)$$

$$0.25 \leq \|\mathbf{r}_{C,FR}^{[k]} - \mathbf{r}_{C,FL}^{[k]}\|^2 \leq 2.25 \quad " \quad (9)$$

$$\|\mathbf{r}_{C,i}^{[k]} - \mathbf{r}_{C,i}^{[k+1]}\|^2 \|\mathbf{r}_{C,j}^{[k]} - \mathbf{r}_{C,j}^{[k+1]}\|^2 = 0, \quad i \neq j \quad \text{only one foot can be moved} \quad (10)$$

$$\|\mathbf{r}_{CoM,j}^{[k]} - \mathbf{r}_{C,i}^{[k]}\|^2 = 0.5, \quad i = 1, 2 \text{ if } j = 1 \text{ or } i = 3, 4 \text{ if } j = 2 \quad (11)$$

the distance between CoMs and feet is bounded

where:

$$\mathbf{F}_{C,i}^n = (\mathbf{F}_{C,i} \cdot \mathbf{n}_{C,i})\mathbf{n}_{C,i}$$

$$\mathbf{F}_{C,i}^t = \mathbf{F}_{C,i} - \mathbf{F}_{C,i}^n$$

$$\mathbf{n}_{C,i} = -\frac{\nabla \mathbf{f}_{env}(\mathbf{r}_{C,i})}{\|\nabla \mathbf{f}_{env}(\mathbf{r}_{C,i})\|}.$$