

# 1 Probability

*Chapter abstract:* Lorem ipsum dolor sit amet, consectetur adipiscing elit. Suspendisse augue est, porttitor commodo velit a, tristique pharetra ante. Mauris pretium ante at lorem suscipit porttitor. Sed neque tortor, lacinia a aliquam quis, molestie tincidunt nisi. Vivamus congue cursus iaculis. Aenean id massa convallis, sodales metus a, imperdiet velit. In metus erat, suscipit vel mollis sed, tincidunt at ante. In hac habitasse platea dictumst. Cras malesuada mollis odio, eget mattis mauris tincidunt a.

## 1.1 ELEMENTARY PROBABILITY

### Outline

1.1	Elementary Probability . .	1
1.2	Conditional Probability . .	2
1.3	Independent Events . . .	2
1.4	Random Variables . . . .	3

### Permutation

#### Definition 1: Permutation

any alignment in  $n$  places

$$P_n = n! \quad (1.1)$$

### Disposition

#### Definition 2: Disposition

Any alignment in  $k$  places

##### 1. Simple:

$$D_{n,k} = n \cdot (n-1) \cdot \dots \cdot (n-k+1) = \frac{n!}{(n-k)!} \quad (1.2)$$

##### 2. With repetition:

$$D_{n,k}^* = n \cdot n \cdot \dots \cdot n = n^k \quad (1.3)$$

### Combination

#### Definition 3: Combination

any alignment of  $k$  objects from  $n$

$$C_{n,k} = \frac{n!}{k!(n-k)!} = \binom{n}{k} \quad (1.4)$$

## 1.2 CONDITIONAL PROBABILITY

In the probability space  $(\Omega, A, \mathbb{P})$ , conditional probability can be written as

$$\mathbb{P}(E|F) = \frac{\mathbb{P}(E \cap F)}{\mathbb{P}(F)} \quad (1.5)$$

with  $\mathbb{P}(F) > 0$

### Conditional Probability as Intersection of Events

$$\begin{aligned} \mathbb{P}(E \cap F) &= \mathbb{P}(F) \cdot \mathbb{P}(E|F) \\ \mathbb{P}(E_1 \cap E_2 \cap \dots \cap E_n) &= \mathbb{P}(E_1) \cdot \mathbb{P}(E_2|E_1) \cdot \mathbb{P}(E_3|E_2 \cap E_1) \dots \mathbb{P}(E_n|E_1 \cap E_2 \cap \dots \cap E_{n-1}) \end{aligned} \quad (1.6)$$

### Law of Total Probability

Given  $E_1, E_2, \dots, E_n$  partitions of  $\Omega$  and the conditions

1.  $E_i \cap E_j = \emptyset \forall i \neq j$
2.  $E_1 \cup E_2 \cup \dots \cup E_n = \Omega$

We can state that:

$$\mathbb{P}(F) = \sum_{i=1}^n \mathbb{P}(F|E_i) \mathbb{P}(E_i) \quad (1.7)$$

### Bayes Formula

Given

$$\mathbb{P}(E_n|F) = \frac{\mathbb{P}(F|E_n) \mathbb{P}(E_n)}{\sum_{i=1}^n \mathbb{P}(F|E_i) \mathbb{P}(E_i)} \rightarrow \mathbb{P}(A|P) = \frac{\mathbb{P}(B|A) \mathbb{P}(A)}{\mathbb{P}(B)} \quad (1.8)$$

## 1.3 INDEPENDENT EVENTS

### Stochastic Independence

#### Definition 4: Stochastic Independence

Given the probability space  $(\Omega, A, \mathbb{P})$ , two events  $E, F \in A$  are said to be **stochastically independent** if  $\mathbb{P}(E \cap F) = \mathbb{P}(E) \cdot \mathbb{P}(F)$

$$\mathbb{P}(E|F) = \mathbb{P}(E) \iff \mathbb{P}(F|E) = \mathbb{P}(F) \quad (1.9)$$

### Conditional Stochastic Independence

#### Definition 5: Conditional Stochastic Independence

Given the probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  and the events  $A, B, F$ , **Conditional Stochastic Independence** can be characterised as:

$$\mathbb{P}(A \cap B | F) = \mathbb{P}(A | F) \cdot \mathbb{P}(B | F) \quad (1.10)$$

**Warning :** Stochastic Independence **does not imply** Conditional Stochastic Independence, and vice versa.

## 1.4 RANDOM VARIABLES

Given the sample space of tossing a dice  $\Omega = \{1, 2, 3, 4, 5, 6\}$ , we can define:

$$X : \Omega \rightarrow \mathbb{R}$$

#### Remark 1.

In the equation above:

- if  $\Omega$  is finite  $\rightarrow$  it is a random variable
- if  $\Omega$  is discrete  $\rightarrow$  it is **not** a random variable

Properties to be considered a random variable:

- $\forall t \in \mathbb{R}$

$$\{\omega \in \Omega : X(\omega) \leq t\} = E_t \in \mathcal{A}$$

With  $X$  being **Borel measurable**

- $\mathbb{P}(E_t) = \mathbb{P}(\omega \in \Omega : X(\omega) \leq t) = \mathbb{P}(X(\omega) \leq t) = \mathbb{P}(X \leq t) = F_x(t)$
- $F_x : \mathbb{R} \rightarrow \mathbb{R}$ , with  $F_x(t)$  being the distribution function of the variable.

$$\lim_{t \rightarrow -\infty} F(t) = 0$$

$$\lim_{t \rightarrow \infty} F(t) = 1$$

- If  $X$  is discrete, the graph will possess the following properties:

1. Stepwise
2. Non-decreasing
3.  $F_x$  continuous from the right

- If  $X$  is **absolutely** continuous

1.  $F$  is continuous




Figure 1.1: Visual representation of an example where  $X$  is a **discrete** random variable




Figure 1.2: Visual representation of an example where  $X$  is an **absolutely continuous** random variable

# 2

## Discrete Probability Distributions

*Chapter abstract:* This chapter provides an overview of the most common discrete probability distributions. We start by introducing the Bernoulli distribution, which models a single coin toss. We then move on to the binomial distribution, which models the number of successes in a fixed number of Bernoulli trials. We also introduce the geometric and negative binomial distributions, for the number of trials needed to get the first success and a fixed number of successes, respectively. Finally, we introduce the hypergeometric distribution, which models the number of successes in a sample drawn without replacement from a finite population.

### 2.1 BERNOULLI DISTRIBUTION

Bernoulli Scheme can describe a series of coin tossing experiments, or the extraction of a ball from an urn with two colors, if there is replacement.

#### Outline

2.1	Bernoulli Distribution . . .	5
2.2	Binomial Distribution . . .	6
2.3	Geometric Distribution . .	7
2.4	Negative Binomial . . . .	8
2.5	Hypergeometric Distribution	8

#### Definition 6: Bernoulli Scheme

A Bernoulli random variable is a random variable that takes the value 1 with probability  $p$  and the value 0 with probability  $1 - p$ . We denote a Bernoulli random variable as  $X \sim \text{Bern}(p)$ .

We observe a Bernoulli Scheme when we have:

- A sequence of  $n$  independent trials.
- Each trial has two possible outcomes: success or failure.
- The probability of success of a single trial is constant ( $p$ ).

$$x = \begin{cases} 0 & 1 - p \\ 1 & p \end{cases}$$

$$X \sim \text{Bern}(p)$$

What is the expected value, and the variance?

$$\begin{aligned} E[X] &= 0q + 1p = p \\ E[X^2] &= 0^2q + 1^2p = p \\ \text{Var}[X] &= E[X^2] - E[X]^2 = p - p^2 = p(1 - p) = pq \end{aligned} \tag{2.1}$$

If, in any probability space there is an event,  $A$  linked to an indicator function  $\mathbb{1}$ , then the indicator function of  $A$  is a Bernoulli random variable with parameter  $p$ .

$$\mathbb{1}_A \sim \text{Bern}(p)$$

## 2.2 BINOMIAL DISTRIBUTION

### Definition 7: Binomial Distribution

The binomial distribution is a discrete probability distribution that models the number of successes in a fixed number of Bernoulli trials.

The binomial distribution is characterized by two parameters:

- $n$  - the number of trials.
- $p$  - the probability of success in each trial.

We can define the random variable  $X$  as the number of successes in  $n$  Bernoulli trials, with values  $0 \leq k \leq n$ . The probability of getting  $k$  successes in  $n$  trials is given by the binomial distribution.

$$X \sim \text{Bin}(n, p)$$

<sup>1</sup> The binomial coefficient  $\binom{n}{k}$  is the number of ways to choose  $k$  successes in  $n$  trials. It is equivalent to writing:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

The probability distribution function of  $X$  is given by<sup>1 2</sup>:

$$P(X = k) = \binom{n}{k} p^k q^{n-k} \quad (2.2)$$

### What is the link between the binomial and the Bernoulli distribution?

Defining  $Y_i$  as the  $i^{\text{th}}$  Bernoulli random variable, we can say that the sum of  $n$  independent Bernoulli random variables is a binomial random variable.

$$X = \sum_{i=1}^n Y_i \sim \text{Bin}(n, p) \quad (2.3)$$

### Example 1.

Adam and Barbara are playing table tennis. In a single game, Barbara wins with probability of 0.6. If they play a series of 7 games, what is the probability that Barbara wins exactly 5 games? Assume that the outcome of each game is independent.

We can model the number of games Barbara wins as a binomial random variable  $X \sim \text{Bin}(7, 0.6)$ . The probability of Barbara winning exactly 5 games is given by:


$$P(X = 5) = \binom{7}{5} 0.6^5 0.4^2$$

### Example 2.


In the previous setting, what is the probability that Barbara wins at least 5 games?

The probability of Barbara winning at least 5 games is given by:

$$P(X \geq 5) = P(X = 5) + P(X = 6) + P(X = 7)$$



**Figure 2.1:** Probability distribution function of a binomial random variable with  $p = 0.2$  and  $n = 20$ .



**Figure 2.2:** Cumulative distribution function.

**Example 3.**

In the previous setting, what is the probability of Barbara winning the last 2 games, given that Barbara won 4 and lost 3 games?

We need to find the probability that the last two games were wins given that Barbara has 4 wins and 3 losses in total.

The possible scenarios are:

- 2 wins, 3 losses in the first 5 games, then 2 wins in the last 2 games.
- 3 wins, 2 losses in the first 5 games, then 1 win and 1 loss in the last 2 games.
- 4 wins, 1 loss in the first 5 games, then 0 wins in the last 2 games.

The only favorable scenario is the first one.

Number of favorable outcomes (2 wins in the last three games):  $\binom{5}{2} = 10$

Number of possible outcomes:  $\binom{7}{4} = 35$

Therefore, the probability is:

$$P(2 \text{ wins in the last 2 games} \mid 4 \text{ wins and 3 losses}) = \frac{10}{35}$$

## 2.3 GEOMETRIC DISTRIBUTION

If  $X$  is the number of trials needed to get the first success in a sequence of Bernoulli trials.

$$P(X = k)$$

1. For the first success to occur on the  $k$ -th trial, the first  $k - 1$  trials must all result in failures. The probability of failure on each trial is  $1 - p$ . Therefore, the probability that the first  $k - 1$  trials all fail is  $(1 - p)^{k-1}$ .

2. The  $k$ -th trial must be a success. The probability of a success on this trial is  $p$ .

Hence, the probability that the first success occurs on the  $k$ -th trial is given by:

$$P(X = k) = (1 - p)^{k-1}p$$

**Definition 8: Geometric Distribution**

The geometric distribution is a probability distribution that models the number of Bernoulli trials needed to get the first success.

The geometric distribution is characterized by a single parameter:

- $p$  - the probability of success in each trial.

$$E[X] = \sum_{k=1}^{+\infty} k(q)^{k-1}p = \frac{1}{p} \quad \text{Var}[X] = \frac{q}{p^2}$$

## 2.4 NEGATIVE BINOMIAL

If we fix the number of successes desired, and we model the number of trials needed to get the desired number of successes, we get the negative binomial distribution.

### Definition 9: Negative Binomial Distribution

The negative binomial distribution is a probability distribution that models the number of Bernoulli trials needed to get a fixed number of successes.

The negative binomial distribution is characterized by two parameters:

- $n$  - the number of successes desired.
- $p$  - the probability of success in each trial.

$$X \sim \text{NegBin}(r, p)$$

We have  $n$  successes in  $k$  trials, the number of failures is therefore  $k - n$ . Since the last trial must be a success, the number of possible arrangements of outcomes is  $\binom{k-1}{n-1}$ . We can then write the probability as:

$$P(X = k) = \binom{k-1}{n-1} p^n q^{k-n}$$

## 2.5 HYPERGEOMETRIC DISTRIBUTION

### Definition 10: Hypergeometric Distribution

The hypergeometric distribution is a probability distribution that models the number of successes in a sample of size  $n$  drawn without replacement from a finite population of size  $N$  that contains  $K$  successes. It is characterized by three parameters:

- $N$  - the population size.
- $K$  - the number of successes in the population.
- $n$  - the sample size.

$$X \sim \text{Hypergeom}(N, K, n)$$

The probability distribution function of  $X$  is given by:

$$P(X = k) = \frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}}$$

Its expected value is:

$$E[X] = n \frac{K}{N}$$



# 3 Continuous Probability Distributions

*Chapter abstract:* This chapter is dedicated to continuous probability distributions. We start by introducing the Poisson distribution, which models the number of events occurring in a fixed interval of time or space. We then move on to the exponential distribution, which models the time until the first event occurs in a Poisson process. We also introduce the gamma distribution, which models the sum of  $n$  exponential random variables. Finally, we introduce the uniform distribution, which models the probability of all outcomes in an interval being equally likely.

## 3.1 POISSON DISTRIBUTION

This is the continuous time equivalent of a bernoulli random variable. It is used to model the number of events occurring in an interval of time or space.

An example is the number of phone calls received by a person. At any given time, the number of calls received is either zero or one, modeled by a Poisson distribution.

Fixing the time interval  $[0, T]$ ,  $X = \text{number of events in } [0, T]$  is a Poisson random variable<sup>3</sup>.

### Definition 11: Poisson Distribution

The Poisson distribution is a probability distribution that models the number of events occurring in a fixed interval of time or space.

The Poisson distribution is characterized by a single parameter:

- $\lambda$  - the average rate of events occurring in the interval

$$\lambda = \frac{\text{number of arrivals}}{\text{time interval}}$$

The probability of having 1 arrival in a time interval is given by:

$$P(X = 1) \approx \lambda \Delta t$$

$$P(X = 1) = \lambda \Delta t + o(\Delta t)$$

The probability distribution function of a Poisson random variable is given by:


$$P(X = k) = \frac{e^{-\lambda T} (\lambda T)^k}{k!}$$

## 3.2 EXPONENTIAL DISTRIBUTION


### Outline

3.1	Poisson Distribution . . .	9
3.2	Exponential Distribution .	10
3.3	Gamma Distribution . . .	10
3.4	Uniform Distribution . . .	12

3



**Figure 3.1:** Probability distribution function of a Poisson random variable with  $\lambda = 5$ .



**Figure 3.2:** Cumulative distribution function.

If  $X$  is the waiting time until the first event occurs in a Poisson process, then  $X$  is an exponential random variable


#### Definition 12: Exponential Distribution

The exponential distribution is a probability distribution that models the time until the first event occurs in a Poisson process.


The exponential distribution is characterized by a single parameter:

- $\lambda$  - the rate of events occurring in the Poisson process.

4



**Figure 3.3:** Probability distribution function of an exponential random variable with  $\lambda = 0.5$ .



**Figure 3.4:** Cumulative distribution function.

The probability distribution function of an exponential random variable is given by <sup>4</sup>:

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

The expected value and variance of an exponential random variable are given by:

$$E[T] = \int_0^{+\infty} x f(x) dx = \frac{1}{\lambda} \quad \text{Var}[T] = \frac{1}{\lambda^2}$$

The exponential distribution is memoryless, meaning that the probability of an event occurring in the next interval (waiting time) is independent of the time that has already passed.

### 3.3 GAMMA DISTRIBUTION

#### Definition 13: Gamma Function

The gamma function is defined as:

$$\Gamma(z) = \int_0^{+\infty} x^{z-1} e^{-x} dx$$

It is defined as long as the exponent  $z$  is positive.

<sup>5</sup> Integration by part:

$$\int u dv = uv - \int v du$$

Properties of the gamma function:

- $\Gamma(z+1) = z\Gamma(z)$
- $\Gamma(1) = 1$
- $\Gamma(n+1) = n!$

The most important property of the gamma function is its recursive definition<sup>5</sup>:

**Proposition 1.**

$$\Gamma(z+1) = z\Gamma(z)$$

**Proof.**

$$\begin{aligned}\Gamma(z+1) &= \int_0^{+\infty} x^z e^{-x} dx \\ &= [-x^z e^{-x}]_0^{+\infty} + z \int_0^{+\infty} x^{z-1} e^{-x} dx = z\Gamma(z)\end{aligned}\quad (3.1)$$

From this property, we can see that the gamma function is a generalization of the factorial function. For any positive integer  $n$ , we have  $\Gamma(n+1) = n!$ .

Starting from the gamma function, we can define the gamma distribution<sup>6</sup>.

**Definition 14: Gamma Distribution**

The gamma distribution is a probability distribution that models the sum of  $n$  exponential random variables.

The gamma distribution is characterized by two parameters:

- $n$  - the number of exponential random variables.
- $\lambda$  - the rate of events occurring in the Poisson process.

$$X \sim \text{Ga}(\alpha, \lambda)$$

$$f(x) = \begin{cases} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

**Proposition 2.**

The expected value of  $X \sim \text{Ga}(\alpha, \lambda)$  is given by:

$$E[X] = \frac{\alpha}{\lambda}$$


**Proof.**

$$\begin{aligned}E[X] &= \int_0^{+\infty} x f(x) dx = \int_0^{+\infty} x \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx \\ &= \int_0^{+\infty} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^\alpha e^{-\lambda x} dx\end{aligned}\quad (3.2)$$


We can see that this is similar to the integral of the gamma function with  $\alpha = \alpha + 1$ , to make it equal, we need to multiply by  $\frac{\lambda}{\lambda}$ .

$$\begin{aligned}E[X] &= \frac{1}{\lambda} \frac{\overbrace{\Gamma(\alpha+1)}^{=\alpha\Gamma(\alpha)}}{\Gamma(\alpha)} \underbrace{\int_0^{+\infty} \frac{\lambda^{\alpha+1}}{\Gamma(\alpha+1)} x^\alpha e^{-\lambda x} dx}_{=1} \\ &= \frac{\alpha}{\lambda}\end{aligned}\quad (3.3)$$

6



**Figure 3.5:** Probability distribution function of a gamma random variable with  $\alpha = 2$  and  $\lambda = 0.5$ .



**Figure 3.6:** Cumulative distribution function.

### 3.4 UNIFORM DISTRIBUTION

---

If all outcomes in an interval are equally likely, we have a uniform distribution. Its probability distribution function is given by:

$$f(x) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

The cumulative distribution function is:

$$F(t) = \begin{cases} 0 & x < a \\ \frac{t-a}{b-a} & a \leq x \leq b \\ 1 & x > b \end{cases}$$

The expected value and variance of a uniform random variable are given by:

$$E[X] = \frac{a+b}{2} \quad \text{Var}[X] = \frac{(b-a)^2}{12}$$

# 4

## Conditional Probability and Sequences of R.V.s

*Chapter abstract:* In day 6, we discuss the concept of conditional probability distributions in the context of discrete and continuous random variables. We then introduce the concept of sample mean and variance of a set of random variables. Finally, we discuss the concept sequences of random variables and their convergence.

### 4.1 EXERCISE

#### Outline

4.1	Exercise . . . . .	13
4.2	Conditional Distributions . . . . .	14
4.3	Sample Mean and Variance . . . . .	18
4.4	Sequence of R.V.s . . . . .	19
4.5	Central Limit Theorem . . . . .	22

$$X \sim \text{Gamma}(\alpha, \lambda) \quad Y \sim \text{Gamma}(\beta, \lambda)$$

$$\begin{cases} V = \frac{x}{y} \\ W = X + Y \end{cases}$$

$$\begin{cases} V = \frac{x}{y} \\ W = Y(1 + V) \end{cases}$$

$$\begin{cases} X = \frac{VW}{1+V} \\ Y = \frac{W}{1+V} \end{cases}$$

$$J = \det \begin{vmatrix} \frac{W(1+V)-vw}{(1+V)^2} & \frac{v}{1+v} \\ -\frac{w}{(1+v)^2} & \frac{1}{1+v} \end{vmatrix} = \frac{w}{(1+v)^2}$$

The density of  $X$  and  $Y$  is given by:

$$f_{X,Y}(x, y) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} \frac{\lambda^\beta}{\Gamma(\beta)} y^{\beta-1} e^{-\lambda y}$$

$X$  and  $Y$  are two independent random variables, therefore the joint distribution of  $V$  and  $W$  is given by:

$$f_{V,W}(v, w) = f_{X,Y}(x, y) |J| = f_X\left(\frac{vw}{1+v}\right) f_Y\left(\frac{w}{1+v}\right) \frac{w}{(1+v)^2}$$

$$f_{V,W}(v, w) = \frac{\lambda^{\alpha+\beta}}{\Gamma(\alpha)\Gamma(\beta)} \left(\frac{vw}{1+v}\right)^{\alpha-1} e^{-\lambda \frac{vw}{1+v}} \left(\frac{w}{1+v}\right)^{\beta-1} e^{-\lambda \frac{w}{1+v}} \frac{w}{(1+v)^2}$$

$$f_{V,W}(v, w) = \frac{\lambda^{\alpha+\beta}}{\Gamma(\alpha)\Gamma(\beta)} \frac{v^{\alpha-1}}{(1+v)^{\alpha+\beta}} w^{\alpha+\beta-1} e^{-\lambda w} \mathbb{1}_{(0,+\infty)}(v) \mathbb{1}_{(0,+\infty)}(w)$$

If the joint density is the product of two functions, then the two random variables are independent.

The joint density is therefore

$$f_{V,W}(v,w) = \frac{\gamma^{\alpha+\beta}}{\Gamma(\alpha)\Gamma(\beta)} \frac{v^{\alpha-1}}{(1+v)^{\alpha+\beta}} \frac{\lambda^{\alpha+\beta}}{(1+v)^{(\alpha+\beta)}} e^{-\lambda w} \mathbb{1}_{(0,+\infty)}(v) \mathbb{1}_{(0,+\infty)}(w)$$

## 4.2 CONDITIONAL DISTRIBUTIONS

### DISCRETE CASE

Consider two random variables  $X$  and  $Y$  with the following joint distribution:

	1	2	3
0	0.1	0.2	0.1
1	0.2	0.1	0.3

<sup>7</sup> **Conditional Distribution:** the probability distribution of a random variable, calculated according to the rules of conditional probability after observing the realization of another random variable.

We can calculate the conditional distribution<sup>7</sup> of  $Y$  given  $X$ :

$$P(Y=1|X=0) = \frac{P(Y=1, X=0)}{P(X=0)} = \frac{0.1}{0.4}$$

More in general:

$$P_{Y|X}(y_i|x_i) = \frac{P_{XY}(y_i, x_i)}{P_X(x_i)}$$

Expected values can be evaluated in the same way:

$$E(Y|X=0) = 1 \cdot 0.25 + 2 \cdot 0.5 + 3 \cdot 0.25 = 2$$

The same can be done for the variance:

$$\text{Var}(Y|X=0) = E(Y^2|X=0) - E(Y|X=0)^2$$

$$E(Y^2|X=0) = 1^2 \cdot 0.25 + 2^2 \cdot 0.5 + 3^2 \cdot 0.25 = 4.5$$

$$\text{Var}(Y|X=0) = 4.5 - 2^2 = 0.5$$

### CONTINUOUS CASE

$$f_{XY}(x,y) = \begin{cases} \frac{15}{8}xy^2 & (x,y) \in T \\ 0 & \text{otherwise} \end{cases} = \frac{15}{8} \mathbb{1}_{(0,1)}(x) \mathbb{1}_{(0,2x)}(y)$$

Can we construct the conditional distribution of  $X$  and  $Y$ ? In this case, we cannot use the formula  $P_{Y|X}(y_i|x_i) = \frac{P_{XY}(y_i, x_i)}{P_X(x_i)}$  because the probability of  $X$  is zero. We can, however, use the formula for the continuous case:

$$f_{Y|X}(y|x) = \frac{\frac{15}{8}xy^2 \mathbb{1}_{(0,1)}(x) \mathbb{1}_{(0,2x)}(y)}{5x^4 \mathbb{1}_{(0,1)}(x)} = \frac{3}{8} \frac{y^2}{x^3} \mathbb{1}_{(0,2x)}(y)$$

We can now calculate the expected value of  $Y$  given  $X$ , as the integral of  $y$  times the conditional density of  $Y$  given  $X$ :

$$E(Y|X=x) = \int_{-\infty}^{+\infty} y f_{Y|X}(y|x) dy = \int_{-\infty}^{+\infty} y \frac{3}{8} \frac{y^2}{x^3} \mathbb{1}_{(0,2x)}(y) dy$$

$$E(Y|X=x) = \int_0^{2x} \frac{3}{8} y^3 x^{-3} dy = \frac{3}{8} x^{-3} \frac{y^4}{4} \Big|_0^{2x} = \frac{3}{8} x^{-3} \frac{16x^4}{4} = \frac{3}{2} x$$

The variance can be calculated in the same way:

$$\begin{aligned} \text{Var}(Y|X=x) &= E(Y^2|X=x) - E(Y|X=x)^2 \\ E(Y^2|X=x) &= \int_{-\infty}^{+\infty} y^2 f_{Y|X}(y|x) dy = \int_0^{2x} \frac{3}{8} y^4 x^{-3} dy \\ &= \frac{3}{8} x^{-3} \frac{y^5}{5} \Big|_0^{2x} = \frac{12}{5} x^2 \end{aligned}$$

So the variance is:

$$\text{Var}(Y|X=x) = \frac{12}{5} x^2 - \left(\frac{3}{2} x\right)^2 = \frac{3}{20} x^2$$

#### Remark 2.

If, in the discrete or continuous case, you construct the conditional distribution of  $Y$  given  $X$ , the expected value and variance of  $Y$  given  $X$  are functions of  $X$ . This holds true unless  $X$  and  $Y$  are independent.

#### Example 4.

Let's go back to the previous example:

	1	2	3
0	0.1	0.2	0.1
1	0.2	0.1	0.3

We know that  $E(Y|X=0) = 2$ , what is  $E(Y|X=1)$ ?

$$E(Y|X=1) = 1 \cdot \frac{0.2}{0.6} + 2 \cdot \frac{0.1}{0.6} + 3 \cdot \frac{0.3}{0.6} = \frac{13}{6}$$

As in the continuous case, the expected value of  $Y$  given  $X$  is a function of  $X$ .

$$E(X|Y=x) = h(x) = \begin{cases} 2 & x=0 \\ \frac{13}{6} & x=1 \end{cases}$$

## GENERAL CASE

$$E(Y|X) = h(X) \leftarrow \text{random variable}$$

In this case, there is no  $X=x$  in the conditional expectation, so we need to calculate the expected value of  $Y$  given  $X$  as a function of  $X$ , not of  $x$ .

So, in the example above:

$$E(Y|X) = \frac{3}{2} X$$

**Definition 15: Conditional Expectation**

The conditional expectation of  $Y$  given  $X$  is a random variable  $h(X)$ . (a function of  $X$ )

$$E(Y|X) = h(X)$$

**Properties:**

- $E(E(Y|X)) = E(Y)$  (Tower Property)
- $E(Yg(X)|X) = g(X)E(Y|X)$
- $Var(Y|X)$  is a r.v.
- $Var(Y) = Var(E(Y|X)) + E(Var(Y|X))$

**Example 5.**

Suppose that  $Y$  is a random variable “duration of battery”, and  $X$  is the r.v. “percentage of an element”

$$X \sim \text{Uniform}(1, 3) \quad (Y|X = x) \sim \text{Exp}(\lambda = x)$$

What is the average duration of the battery? i.e.  $E(Y)$

$$E(Y|X = x) = \frac{1}{x} \quad E(Y|X) = \frac{1}{X}$$

Therefore, we can use the Tower Property:

$$E(Y) = E(E(Y|X)) = E\left[\frac{1}{X}\right] = \int_1^3 \frac{1}{x} \frac{1}{2} dx = \frac{1}{2} \int_1^3 \frac{1}{x} dx = \frac{1}{2} \ln(3)$$

**Example 6.**

The duration of a call is:  $T_1 \sim \text{Exp}(\lambda = \frac{1}{2})$

The number of calls is:  $N \sim \text{Poisson}(\lambda = 60)$

The total time spent on calls is therefore:

$$Y = \sum_{i=1}^N T_i$$

What is the expected value and variance of the total time spent on calls?

$$(Y|N = n) = \sum_{i=1}^n T_i \sim \text{Gamma}(n, \frac{1}{2})$$

$$E(Y|N = n) = E(Ga(n, \frac{1}{2})) = \frac{n}{\lambda} = 2n$$

$$E(Y) = E(E(Y|N)) = E(2N) = 2E(N) = 2 \cdot 60 = 120$$

Calculating the variance:

$$Var(Y|N = n) = n \cdot \frac{1}{\lambda^2} = 4n$$



$$\text{Var}(Y) = \text{Var}(E(Y|N)) + E(\text{Var}(Y|N))$$

$$\text{Var}(Y) = \text{Var}(2N) + E(4N) = 4\text{Var}(N) + 4E(N) = 4 \cdot 60 + 4 \cdot 60 = 480$$

**Example 7.**

Suppose  $X_1, \dots, X_n$  are independent and identically distributed random variables with  $X_n \sim \text{Bern}(p)$ .

Let  $Y = \sum_{i=1}^n X_i$ .

What is  $E(X_1|Y)$ ?

The distribution of  $Y$  is a binomial:  $Y \sim \text{Bin}(n, p)$ .

$$E(X_1|Y = k) = 0 \cdot P(X_1 = 0|Y = k) + 1 \cdot P(X_1 = 1|Y = k)$$

$$E(X_1|Y = k) = P(X_1 = 1|Y = k) = \frac{P(X_1 = 1, Y = k)}{P(Y = k)}$$

The possible values of  $k$  are  $0, 1, \dots, n$ , so:

$$E(X_1|Y = k) = \begin{cases} 0 & k = 0 \\ ? & k = 1, \dots, n \end{cases}$$

We can rewrite the conditional expectation as:

$$E\left(X_1 \mid \sum_{i=1}^n X_i = k\right) = \frac{P(X_1 = 1, \sum_{i=1}^n X_i = k)}{P(\sum_{i=1}^n X_i = k)}$$

The two events in the numerator are not independent, so we have to rewrite it to solve the problem. The probability of  $X_1 = 1$  and  $\sum_{i=1}^n X_i = k$  is the same as the probability of  $X_1 = 1$  and  $X_2 + \dots + X_n = k - 1$ :

$$\begin{aligned} P(X_1 = 1, \sum_{i=1}^n X_i = k) &= \frac{P(X_1 = 1, \sum_{i=2}^n X_i = k-1)}{P(\sum_{i=1}^n X_i = k)} \\ &= \frac{P(X_1 = 1)P(\sum_{i=2}^n X_i = k-1)}{P(\sum_{i=1}^n X_i = k)} = \frac{p \cdot \binom{n-1}{k-1} p^{k-1} q^{n-k}}{\binom{n}{k} p^k q^{n-k}} \\ &= \frac{\binom{n-1}{k-1}}{\binom{n}{k}} = \frac{\frac{(n-1)!}{(k-1)!(n-k)!}}{\frac{n!}{k!(n-k)!}} = \frac{k}{n} \end{aligned}$$

The final result is therefore:

$$E(X_1|Y = k) = \begin{cases} 0 & k = 0 \\ \frac{k}{n} & k = 1, \dots, n \end{cases} = \frac{k}{n}$$

Therefore:

$$E(X_1|Y) = \frac{Y}{n}$$

### 4.3 SAMPLE MEAN AND VARIANCE

Take  $X_1, \dots, X_n$  i.i.d. We can define the sample mean  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ .

Defining  $m = E(X_1)$ , we can calculate the expected value of the sample mean:

$$E(\bar{X}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot n \cdot m = m$$

With the variance of  $X_1$  defined as  $v$ , we can calculate the variance of the sample mean:

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n Var(X_i) = \frac{1}{n^2} \cdot n \cdot v = \frac{v}{n}$$

This holds true for any sampling distribution.

### SAMPLE VARIANCE

The sample variance is defined differently, depending on whether  $m$  is known or not.

If  $m$  is known, the sample variance is:

$$S_0^2 = \frac{1}{n} \sum_{i=1}^n (X_i - m)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - 2m \sum_{i=1}^n X_i + nm^2$$

The expected value of the sample variance is  $E(S_0^2) = v$ .

If  $m$  is unknown, the sample variance is:

$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

This can be rewritten as:

$$S_n^2 = \frac{1}{n-1} \left[ \sum_{i=1}^n X_i^2 - \underbrace{2\bar{X}_n \sum_{i=1}^n X_i}_{=2\bar{X}_n^2} + n\bar{X}_n^2 \right]$$

$$S_n^2 = \frac{1}{n-1} \left[ \sum_{i=1}^n X_i^2 - n\bar{X}_n^2 \right]$$

### CASE OF A NORMAL DISTRIBUTION

Take  $X_1, \dots, X_n$  i.i.d. with  $X_i \sim N(\mu, \sigma^2)$ . Then the sample mean  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  is normally distributed with:  $\bar{X}_n \sim N(\mu, \frac{\sigma^2}{n})$ .

The sample variance in the case where  $m$  is known is distributed as:

$$\frac{nS_0^2}{\sigma^2} \sim \chi^2(n)$$

In the case where  $m$  is unknown, the sample variance is distributed as:

$$\frac{(n-1)S_n^2}{\sigma^2} \sim \chi^2(n-1)$$

## 4.4 SEQUENCE OF R.V.S

### Definition 16: Limit of a Sequence

In a sequence of real numbers  $a_1, a_2, \dots, a_n$ , the limit is defined as:

$$\lim_{n \rightarrow \infty} a_n = L \in \mathbb{R} \quad \forall \epsilon > 0 \quad \exists n_\epsilon \in \mathbb{N} \quad | \quad \forall n \geq n_\epsilon \implies |a_n - L| < \epsilon$$

Intuitively, the limit exists if there exists an  $n$  large enough so that after that  $n$  all the terms are within  $\epsilon$  distance of the limit. ( $\text{dist}(a_n, L) < \epsilon$ )

Limits can be defined in spaces other than  $\mathbb{R}$ , as long as there is a way to define the distance between two elements.

### Sequence of Random Variables

Take a sequence of random variables  $X_1, X_2, \dots, X_n, \dots$ . Suppose that all the random variables are defined on the same probability space  $(\Omega, \mathcal{A}, P)$ .

### Definition 17: Sure Convergence

We say that  $X_n \rightarrow Y$  surely if:

$$\forall \omega \in \Omega \quad X_n(\omega) \rightarrow Y(\omega)$$

In other words, for every sequence of outcomes  $X_n(\omega)$ , the limit of the sequence is  $Y(\omega)$ .

This definition is very strong, and is not very useful in practice.

### Definition 18: Almost Sure Convergence

We say that  $X_n \rightarrow Y$  almost surely if:

$$P(\{\omega \in \Omega | X_n(\omega) \rightarrow Y(\omega)\}) = 1$$

In other words, the set of outcomes for which the sequence of random variables converges to the limit has probability 1.

Property:

$$g : \mathbb{R} \rightarrow \mathbb{R} \quad \text{continuous} \quad \implies \quad g(X_n) \rightarrow g(Y) \quad \text{almost surely}$$

### Definition 19: Convergence in Probability

We say that  $X_n \rightarrow Y$  in probability if:

$$\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|X_n - Y| < \epsilon) = 1$$

In other words, the probability that the distance between  $X_n$  and  $Y$  is less than  $\epsilon$  converges to 1 as  $n$  goes to infinity.

**Definition 20: Convergence in Mean of Order  $k$**

We say that  $X_n \rightarrow Y$  in mean of order  $k \geq 1$  if:

$$E(|X_n - Y|^k) \rightarrow 0$$

**Definition 21: Convergence in Quadratic Mean**

We say that  $X_n \rightarrow Y$  in quadratic mean if:

$$E(|X_n - Y|^2) \rightarrow 0$$

From a sequence of random variables, we can define the sequence of sample means:

$$\bar{X}_1 = X_1 \quad \bar{X}_2 = \frac{X_1 + X_2}{2} \quad \bar{X}_3 = \frac{X_1 + X_2 + X_3}{3} \quad \dots$$

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

We can demonstrate that the sample mean converges to the expected value of the random variable:

$$Var(\bar{X}_n) = \frac{1}{n^2} \sum_{i=1}^n Var(X_i) = \frac{1}{n^2} \cdot n \cdot v = \frac{v}{n}$$

$$E(|\bar{X}_n - m|^2) = Var(\bar{X}_n) = \frac{v}{n} \rightarrow 0$$

**Definition 22: Strong Law of Large Numbers**

If  $X_1, X_2, \dots, X_n$  are i.i.d. with  $E(X_i) = m$  and  $Var(X_i) = v$ , then:

$$\bar{X}_n \rightarrow m \quad \text{almost surely}$$

If  $X_n \rightarrow Y$  almost surely, then  $X_n \rightarrow Y$  in probability. The converse is not true.

$$X_n \rightarrow Y \text{ almost surely} \quad \Rightarrow \quad X_n \rightarrow Y \text{ in probability}$$

Moreover, if  $X_n \rightarrow Y$  in order  $k$ , then  $X_n \rightarrow Y$  in probability. The converse is again not true.

$$X_n \rightarrow Y \text{ in order } k \quad \Rightarrow \quad X_n \rightarrow Y \text{ in probability}$$

## CONVERGENCE IN DISTRIBUTION

### Definition 23: Convergence in Distribution

We say that  $X_n \rightarrow Y$  in distribution if:

$$F_{X_n}(t) \rightarrow F_Y(t) \quad \forall t \quad \text{where } F_Y \text{ is continuous}$$

Convergence in distribution is a weaker form of convergence than other forms of convergence. Link between convergence in distribution and convergence in probability:

$$X_n \rightarrow Y \text{ in probability} \Rightarrow X_n \rightarrow Y \text{ in distribution}$$

The converse is generally not true. However, if  $Y$  is a constant, then the two forms of convergence are equivalent.

$$X_n \rightarrow y \in \mathbb{R} \text{ in distribution} \Leftrightarrow X_n \rightarrow y \in \mathbb{R} \text{ in probability}$$

### Example 8.

Take a sequence of i.i.d. random variables  $X_1, X_2, \dots, X_n, \dots$  with:

$$X_i \sim \text{Unif}(0, 1)$$

We take the sequence  $Y_n = \min(X_1, \dots, X_n)$ . What is the limit of  $Y_n$ ?

Recall that if  $V_n = \min(X_1, \dots, X_n)$ , then  $F_{V_n}(t) = 1 - [1 - F_X(t)]^n$ .

Since the distribution function of  $X$  is:

$$F_X(t) = \begin{cases} 0 & t < 0 \\ t & 0 \leq t \leq 1 \\ 1 & t > 1 \end{cases}$$

The distribution function of  $V_n$  is:

$$F_{V_n}(t) = \begin{cases} 0 & t < 0 \\ 1 - (1 - t)^n & 0 \leq t \leq 1 \\ 1 & t > 1 \end{cases}$$

For the convergence in distribution, we need to calculate the limit of  $F_{V_n}(t)$  as  $n$  goes to infinity:

$$\lim_{n \rightarrow \infty} F_{V_n}(t) = \begin{cases} 0 & t \leq 0 \\ 1 & t > 0 \end{cases}$$

This is not the distribution function of any random variable, because it has a jump at 0, and in that jump the value is continuous from the left but not from the right.

This is the distribution of a discrete random variable, as it is not continuous but piece-wise constant. From the definition of the limit of a sequence of random variables, we can define the limit of  $F_{V_n}(t)$  as:

$$F_Y(t) = \begin{cases} 0 & t < 0 \\ 1 & t \geq 0 \end{cases}$$

Which is the distribution function of a random variable  $Y$  that is equal to 0 with probability 1.

## 4.5 CENTRAL LIMIT THEOREM

### Definition 24: Central Limit Theorem

Let  $X_1, \dots, X_n$  be i.i.d random variables with  $m = E(X_i)$  and  $v = Var(X_i)$ .

$$(P) \left( \frac{\bar{X}_n - m}{\sqrt{\frac{v}{n}}} \leq t \right) \rightarrow \Phi(t) \quad \text{as } n \rightarrow +\infty$$

or

$$\frac{\bar{X}_n - m}{\sqrt{\frac{v}{n}}} \rightarrow N(0, 1) \quad \text{in distribution}$$

For this to hold, we have to assume that the random variables have finite mean and variance. In other words,  $E(X_i^2) < +\infty$ .

In other words, if  $n$  is “large enough”, then the distribution of the sample mean is approximately normal with  $\bar{X}_n \approx N(m, \frac{v}{n})$ .

If we multiply by  $n$ , we get:

$$\sum_{i=1}^n X_i \approx N(nm, nv)$$

<sup>8</sup> The Poisson distribution with parameter  $\lambda$  is the sum of  $\lambda$  Poisson random variables with parameter 1.

<sup>9</sup> The Gamma distribution with parameters  $\alpha$  and  $\lambda$  is the sum of  $\alpha$  exponential random variables with parameter  $\lambda$ .

<sup>10</sup> The Chi-squared distribution with  $n$  degrees of freedom is the sum of  $n$  standard normal random variables squared ( $\chi^2$  with 1 d.f.).

Take  $Y \sim \text{Bin}(n, p)$ , then if  $n$  is large enough and  $p$  is not too close to 0 or 1, then  $Y \approx N(np, npq)$ .

With a binomial distribution  $Y \sim \text{Bin}(n, p)$ , if  $n \rightarrow +\infty$  and  $p \rightarrow 0$  such that  $np \rightarrow \lambda$ , then  $Y \approx \text{Po}(\lambda)$ . In other words,  $X_n \rightarrow P(\lambda)$  in distribution.

Poisson random variables are also approximately normal if  $\lambda$  is large enough.<sup>8</sup>

In the case of Gamma distributions  $Y \sim \text{Ga}(\alpha, \lambda)$ , if  $\alpha$  is large enough, then  $Y \approx N(\alpha/\lambda, \alpha/\lambda^2)$ .<sup>9</sup>

For a Chi-squared distribution with  $n$  degrees of freedom, if  $n$  is large enough, then  $X \approx N(n, 2n)$ .<sup>10</sup>

**Definition 25: Slutsky's Theorem**

Let  $X_n \rightarrow X$  in distribution. If  $Y_n \rightarrow Y$  in probability, then:

$$X_n + Y_n \rightarrow X + Y \quad \text{in distribution}$$

$$X_n \cdot Y_n \rightarrow X \cdot Y \quad \text{in distribution}$$

$$\forall n \quad P(Y_n = 0) = 0 \text{ and } y \neq 0.$$

**Sampling from a Normal Distribution**

Take  $X_1, X_2, \dots, X_n$  i.i.d. with  $X_i \sim N(\mu, \sigma^2)$ .

$$\frac{\bar{X}_n - \mu}{\sqrt{\frac{S_n^2}{n}}} = \frac{\bar{X}_n - \mu}{\underbrace{\sqrt{\frac{\sigma^2}{n}}}_{\sim N(0,1)}} \underbrace{\sqrt{\frac{\sigma^2}{S_n^2}}}_{\rightarrow 1} \rightarrow Z \sim N(0,1) \text{ in distribution}$$

Therefore, the distribution is approximately  $N(0,1)$ .

**Sampling not from a Normal Distribution** If  $m = E(X_i)$  and  $v = Var(X_i)$ , then:

$$\frac{\bar{X}_n - \mu}{\sqrt{\frac{S_n^2}{n}}} = \frac{\bar{X}_n - \mu}{\underbrace{\sqrt{\frac{v}{n}}}_{\sim Z}} \underbrace{\sqrt{\frac{v}{S_n^2}}}_{\rightarrow 1} \rightarrow Z \sim N(0,1) \text{ in distribution}$$