

# Lecture 8

## Continuous time dynamic models

---

Ivan Rudik  
AEM 7130

# Roadmap

1. The theory behind continuous time models
2. Numerical methods for solving continuous time model

# Model setup

Consider a problem where each period an agent obtains flow utility  $J(x(t), u(t))$ , where  $x$  is our **state** and  $u$  is our **control**

# Model setup

Consider a problem where each period an agent obtains flow utility  $J(x(t), u(t))$ , where  $x$  is our **state** and  $u$  is our **control**

Suppose there is a finite horizon with a **terminal time**  $T$

# Model setup

The agent's objective is to maximize the total payoff, subject to the transitions of the states

$$\max_{u, x_T} \int_0^T J(x(t), u(t)) dt$$

$$\text{subject to: } \dot{x}(t) = g(x(t), u(t)), x(0) = x_0, x(T) = x_T$$

This is an open-loop solution so we optimize our entire policy trajectory from time  $t = 0$

We will not be solving for functions of states, but functions of time:  $u(t), x(t)$

# Hamiltonians

In a dynamic optimization problem, we will have an auxiliary function that yields the first-order conditions

# Hamiltonians

In a dynamic optimization problem, we will have an auxiliary function that yields the first-order conditions

This function is called the *Hamiltonian*:

$$H(x(t), u(t), \lambda(t)) \equiv J(x(t), u(t)) + \lambda(t)g(x(t), u(t))$$

It is a function that treats the transitions as quasi-constraints so it appears similar to the Lagrangian you know

# Hamiltonians

Pontryagin's Maximum Principle states that the following conditions are necessary for an optimal solution:

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial u} = 0 \quad \forall t \in [0, T] \quad (\text{Maximality})$$

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial x} = -\dot{\lambda}(t) \quad (\text{Co-state})$$

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial \lambda} = \dot{x}(t) \quad (\text{State transitions})$$

$$x(0) = x_0 \quad (\text{Initial condition})$$

$$\lambda(T) = 0 \quad (\text{Transversality})$$

What do these conditions mean?



# Necessary conditions

First, what *is* the Hamiltonian?

# Necessary conditions

First, what *is* the Hamiltonian?

The Hamiltonian tells us the contribution of that instant  $t$  to overall utility via the change in flow utility and the change in the state (which affects future flow utilities)

# Necessary conditions

First, what *is* the Hamiltonian?

The Hamiltonian tells us the contribution of that instant  $t$  to overall utility via the change in flow utility and the change in the state (which affects future flow utilities)

The decisionmaker can use her control to increase the contemporaneous flow utility and reap immediate rewards, or to alter the state variable to increase future rewards

$$H(x(t), u(t), \lambda(t)) \equiv \underbrace{J(x(t), u(t))}_{\text{current flow}} + \underbrace{\lambda(t)g(x(t), u(t))}_{\text{change in future value}}$$

# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial u} = 0 \quad \forall t \in [0, T]$$

The **maximality condition**: in every instant, we select the control so that we can no longer increase our total payoff

# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial u} = 0 \quad \forall t \in [0, T]$$

The **maximality condition**: in every instant, we select the control so that we can no longer increase our total payoff

It effectively sets the net marginal benefits of the control to zero

# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial \lambda} = \dot{x}(t)$$

The **state transition** is just a definition

# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial \lambda} = \dot{x}(t)$$

The **state transition** is just a definition

Taking the derivative of the Hamiltonian with respect to the shadow value, just like a Lagrangian, yields this constraint back

# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial x} = -\dot{\lambda}(t)$$

The **co-state condition** defines how the shadow value of our state transition, called the **co-state variable**, evolves over time



# Necessary conditions

$$\frac{\partial H(x(t), u(t), \lambda(t))}{\partial x} = -\dot{\lambda}(t)$$

The **co-state condition** defines how the shadow value of our state transition, called the **co-state variable**, evolves over time

What is the co-state?

# Necessary conditions

What is the co-state?

# Necessary conditions

What is the co-state?

The additional future value of having one more unit of our state variable

# Necessary conditions

What is the co-state?

The additional future value of having one more unit of our state variable

Suppose we increase today's stock of  $x$  by one unit and this increases the instantaneous change in our value ( $H$ )

# Necessary conditions

What is the co-state?

The additional future value of having one more unit of our state variable

Suppose we increase today's stock of  $x$  by one unit and this increases the instantaneous change in our value ( $H$ )

Then the shadow value of that stock ( $\lambda$ ) must decrease along an optimal trajectory

# Necessary conditions

What is the co-state?

The additional future value of having one more unit of our state variable

Suppose we increase today's stock of  $x$  by one unit and this increases the instantaneous change in our value ( $H$ )

Then the shadow value of that stock ( $\lambda$ ) must decrease along an optimal trajectory

Why?

# Necessary conditions

If it didn't, we could increase value by accumulating more of the stock variable  
→ there is a profitable deviation and what we were doing cannot be optimal

We can re-write the co-state equation as

$$\frac{\partial J}{\partial x} + \lambda(t) \frac{\partial g}{\partial x} + \dot{\lambda}(t) = 0$$

# Necessary conditions

$$\frac{\partial J}{\partial x} + \lambda(t) \frac{\partial g}{\partial x} + \dot{\lambda}(t) = 0$$

We must have that the a unit of the stock's value must change (third term), so that is exactly offsets the change in value from increasing the stock in the immediate instant of time



# Necessary conditions

$$\frac{\partial J}{\partial x} + \lambda(t) \frac{\partial g}{\partial x} + \dot{\lambda}(t) = 0$$

We must have that the a unit of the stock's value must change (third term), so that is exactly offsets the change in value from increasing the stock in the immediate instant of time

The immediate value is made up of the actual utility payoff (first term), and the future utility payoff payoff from how increasing the stock today affects the stock in the future (second term)

# Necessary conditions

These necessary conditions give us the shape of the optimal path but they do not tell us what the optimal path actually is

# Necessary conditions

These necessary conditions give us the shape of the optimal path but they do not tell us what the optimal path actually is

Many different paths are consistent with these differential equations, depends on the constant of integration

# Necessary conditions

These necessary conditions give us the shape of the optimal path but they do not tell us what the optimal path actually is

Many different paths are consistent with these differential equations, depends on the constant of integration

We need additional optimality conditions to use as constraints to impose the optimal path

# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

We need two constraints

# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

We need two constraints

**Constraint 1:** Initial condition on the state

# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

We need two constraints

**Constraint 1:** Initial condition on the state

This directly pins down where the state path starts



# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

We need two constraints

**Constraint 1:** Initial condition on the state

This directly pins down where the state path starts

We need to pin down the co-state path

# Pinning down the optimal path

We have effectively two ODEs, one for  $\dot{\lambda}(t)$  and one for  $\dot{x}(t)$

We need two constraints

**Constraint 1:** Initial condition on the state

This directly pins down where the state path starts

We need to pin down the co-state path

We do this using **transversality conditions**

# Pinning down the optimal path

In general there are four types of transversality conditions

# Pinning down the optimal path

In general there are four types of transversality conditions

The first two are for free initial or terminal time problems: these are problems where the agent can select when the problem starts or ends

# Pinning down the optimal path

In general there are four types of transversality conditions

The first two are for free initial or terminal time problems: these are problems where the agent can select when the problem starts or ends

The second two are for pinning down the initial and terminal state variables if they're free

# Pinning down the optimal path

In general there are four types of transversality conditions

The first two are for free initial or terminal time problems: these are problems where the agent can select when the problem starts or ends

The second two are for pinning down the initial and terminal state variables if they're free

Usually terminal conditions are free and initial conditions are not

# Pinning down the optimal path example

If the terminal state is free, the transversality condition is that its shadow value must be zero

# Pinning down the optimal path example

If the terminal state is free, the transversality condition is that its shadow value must be zero

Why?



# Pinning down the optimal path example

If the terminal state is free, the transversality condition is that its shadow value must be zero

Why?

If it were positive the policymaker could profitably deviate by altering the level of the stock. Finally, these are all necessary conditions of the problem

# Discounting

We discount the future using exponential discounting → now time can directly affect value

# Discounting

We discount the future using exponential discounting  $\rightarrow$  now time can directly affect value

Assume that time does not directly affect instantaneous payoffs or the transitions equations

# Discounting

We discount the future using exponential discounting → now time can directly affect value

Assume that time does not directly affect instantaneous payoffs or the transitions equations

Then our value is  $J(x(t), u(t), t) = e^{-rt} V(x(t), u(t))$

# Discounting

We discount the future using exponential discounting → now time can directly affect value

Assume that time does not directly affect instantaneous payoffs or the transitions equations

Then our value is  $J(x(t), u(t), t) = e^{-rt} V(x(t), u(t))$

$J$  yields the present, time 0 value of the change in value at time  $t$

# Discounting

We discount the future using exponential discounting → now time can directly affect value

Assume that time does not directly affect instantaneous payoffs or the transitions equations

Then our value is  $J(x(t), u(t), t) = e^{-rt} V(x(t), u(t))$

$J$  yields the present, time 0 value of the change in value at time  $t$

$J$  is the *present value* and  $V$  is the *current value*

# Discounting

We discount the future using exponential discounting → now time can directly affect value

Assume that time does not directly affect instantaneous payoffs or the transitions equations

Then our value is  $J(x(t), u(t), t) = e^{-rt} V(x(t), u(t))$

$J$  yields the present, time 0 value of the change in value at time  $t$

$J$  is the *present value* and  $V$  is the *current value*

Present value refers to the value with respect to a specific period that we call the present

# Current value terms

Our previous necessary conditions apply to present value Hamiltonians, but let us analyze a current value Hamiltonian to avoid including time terms,

$$\begin{aligned} H^{cv}(x(t), u(t), \mu(t)) &\equiv e^{rt} H(x(t), u(t), \lambda(t), t) \\ &= e^{rt} J(x(t), u(t), t) + e^{rt} \lambda(t) g(x(t), u(t)) \end{aligned}$$

$\mu(t)$  is the shadow value  $\lambda$  brought into current value terms:  $\mu(t) = e^{rt} \lambda(t)$



# Current value terms

We can then re-write our necessary conditions in current value by substituting in for:

- the shadow value (which implies that  $\dot{\lambda}(t) = -re^{-rt}\mu(t) + e^{-rt}\dot{\mu}(t)$ )
- $\partial H / \partial x = e^{-rt} \partial H^{cv} / \partial x$  into our co-state condition:

$$e^{-rt} \frac{\partial H^{cv}(x(t), u(t), \mu(t))}{\partial x} = e^{-rt} [r\mu(t) - \dot{\mu}(t)]$$

# Current value

$$e^{-rt} \frac{\partial H^{cv}(x(t), u(t), \mu(t))}{\partial x} = e^{-rt} [r\mu(t) - \dot{\mu}(t)]$$

# Current value

$$e^{-rt} \frac{\partial H^{cv}(x(t), u(t), \mu(t))}{\partial x} = e^{-rt} [r\mu(t) - \dot{\mu}(t)]$$

Before, the present value form of the co-state condition required the change in the present shadow value precisely equal the effect of the state variable on instantaneous value

# Current value

$$e^{-rt} \frac{\partial H^{cv}(x(t), u(t), \mu(t))}{\partial x} = e^{-rt} [r\mu(t) - \dot{\mu}(t)]$$

In current value form, the co-state condition recognizes that the change in the present shadow value is comprised of two parts:

1. The change in the current shadow value
2. The reduction in present value purely from the passage of time

# Current value

$$e^{-rt} \frac{\partial H^{cv}(x(t), u(t), \mu(t))}{\partial x} = e^{-rt} [r\mu(t) - \dot{\mu}(t)]$$

In current value form, the co-state condition recognizes that the change in the present shadow value is comprised of two parts:

1. The change in the current shadow value
2. The reduction in present value purely from the passage of time

If discounting is high (large  $r$ ), then the current shadow value must change quicker in order to compensate the policymaker for leaving stock for the future

# Numerical methods for continuous time models

Continuous time models are systems of ODEs

# Numerical methods for continuous time models

Continuous time models are systems of ODEs

A first-order ordinary differential equation has the form

$$dy/dt = f(y, t)$$

# Numerical methods for continuous time models

Continuous time models are systems of ODEs

A first-order ordinary differential equation has the form

$$dy/dt = f(y, t)$$

Where  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$



# Numerical methods for continuous time models

Continuous time models are systems of ODEs

A first-order ordinary differential equation has the form

$$dy/dt = f(y, t)$$

Where  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$

The unknown is the function  $y(t) : [t_0, T] \rightarrow \mathbb{R}^n$

# Numerical methods for continuous time models

Continuous time models are systems of ODEs

A first-order ordinary differential equation has the form

$$dy/dt = f(y, t)$$

Where  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$

The unknown is the function  $y(t) : [t_0, T] \rightarrow \mathbb{R}^n$

$n$  determines the number of differential equations that we have

# Numerical methods for continuous time models

The differential equation will give us the shape of the path that is the solution, but not where that path lies in our state space

# Numerical methods for continuous time models

The differential equation will give us the shape of the path that is the solution, but not where that path lies in our state space

We need additional conditions to pin down  $y(t)$ , how we pin down  $y(t)$  is what defines the different types of problems we have

# Numerical methods for continuous time models

The differential equation will give us the shape of the path that is the solution, but not where that path lies in our state space

We need additional conditions to pin down  $y(t)$ , how we pin down  $y(t)$  is what defines the different types of problems we have

If we pin down  $y(t_0) = y_0$  or  $y(T) = y_T$  we have an initial value problem

# Numerical methods for continuous time models

The differential equation will give us the shape of the path that is the solution, but not where that path lies in our state space

We need additional conditions to pin down  $y(t)$ , how we pin down  $y(t)$  is what defines the different types of problems we have

If we pin down  $y(t_0) = y_0$  or  $y(T) = y_T$  we have an initial value problem

IVPs are defined by the function being pinned down at one end or the other

# Numerical methods for continuous time models

In one dimension we must pin down the function with either an initial or terminal condition so they are all IVPs by default

# Numerical methods for continuous time models

In one dimension we must pin down the function with either an initial or terminal condition so they are all IVPs by default

If  $n > 1$  then we can have a *boundary value problem* where we impose  $n$  conditions on  $y$

$$\begin{aligned} g_i(y(t_0)) &= 0, & i &= 1, \dots, n', \\ g_i(y(T)) &= 0, & i &= n' + 1, \dots, n \end{aligned}$$

where  $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$



# Numerical methods for continuous time models

In general we have that

$$g_i(y(t_i)) = 0$$

for some set of points  $t_i, t_0 \leq t_i \leq T, 1 \leq i \leq n$

# Numerical methods for continuous time models

In general we have that

$$g_i(y(t_i)) = 0$$

for some set of points  $t_i, t_0 \leq t_i \leq T, 1 \leq i \leq n$

Often we set  $T = \infty$  so we will need some condition in the limit:  $\lim_{t \rightarrow \infty} y(t)$

# Numerical methods for continuous time models

Note two more things:

1. We are implicitly assuming that these  $n$  conditions are independent, otherwise we will not have a unique solution
2. IVPs and BVP are fundamentally different: IVPs are problems where the auxiliary conditions that pin down the solution are all at one point, in BVPs they can be at different points, this has significant implications for how we can solve the problems

# Numerical methods for continuous time models

If we have higher-order ODEs we can use a simple change of variables

$$d^2y/dx^2 = g(dy/dx, y, x)$$

for  $x, y \in \mathbb{R}$

# Numerical methods for continuous time models

If we have higher-order ODEs we can use a simple change of variables

$$d^2y/dx^2 = g(dy/dx, y, x)$$

for  $x, y \in \mathbb{R}$

Define  $z = dy/dx$  and then we can study the alternative system

$$dy/dx = z \quad dz/dx = g(z, y, x)$$

of two first-order ODEs

# Numerical methods for continuous time models

If we have higher-order ODEs we can use a simple change of variables

$$d^2y/dx^2 = g(dy/dx, y, x)$$

for  $x, y \in \mathbb{R}$

Define  $z = dy/dx$  and then we can study the alternative system

$$dy/dx = z \quad dz/dx = g(z, y, x)$$

of two first-order ODEs

In general you can always transform a  $n$ th-order ODE into  $n$  first-order ODEs

# Finite difference methods for IVPs

We solve IVPs using finite-difference methods

# Finite difference methods for IVPs

We solve IVPs using finite-difference methods

Consider the following IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$



# Finite difference methods for IVPs

We solve IVPs using finite-difference methods

Consider the following IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

A finite-difference method solves this IVP by first specifying a grid/mesh over  $t$ :  $t_0 < t_1 < \dots < t_i < \dots$

# Finite difference methods for IVPs

We solve IVPs using finite-difference methods

Consider the following IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

A finite-difference method solves this IVP by first specifying a grid/mesh over  $t$ :  $t_0 < t_1 < \dots < t_i < \dots$

Assume the grid is uniformly spaced:  $t_i = t_0 + ih$ ,  $i = 0, 1, \dots, N$  where  $h$  is the mesh size

# Finite difference methods for IVPs

We solve IVPs using finite-difference methods

Consider the following IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

A finite-difference method solves this IVP by first specifying a grid/mesh over  $t$ :  $t_0 < t_1 < \dots < t_i < \dots$

Assume the grid is uniformly spaced:  $t_i = t_0 + ih$ ,  $i = 0, 1, \dots, N$  where  $h$  is the mesh size

Our goal is to find for each  $t_i$ , a value  $Y_i$  that closely approximates  $y(t_i)$

# Finite difference methods for IVPs

To do this, we replace our differential equation with a difference system on the grid

# Finite difference methods for IVPs

To do this, we replace our differential equation with a difference system on the grid

For example we might have  $Y_{i+1} = F(Y_i, Y_{i-1}, \dots, t_{i+1}, t_i, \dots)$

# Finite difference methods for IVPs

To do this, we replace our differential equation with a difference system on the grid

For example we might have  $Y_{i+1} = F(Y_i, Y_{i-1}, \dots, t_{i+1}, t_i, \dots)$

We then solve the difference equation for the  $Y$ 's in sequence where the initial  $Y_0$  is fixed by the initial condition  $Y_0 = y_0$

# Finite difference methods for IVPs

To do this, we replace our differential equation with a difference system on the grid

For example we might have  $Y_{i+1} = F(Y_i, Y_{i-1}, \dots, t_{i+1}, t_i, \dots)$

We then solve the difference equation for the  $Y$ 's in sequence where the initial  $Y_0$  is fixed by the initial condition  $Y_0 = y_0$

This approximates the solution only at the grid points, but then we can interpolate using standard procedures to get the approximate solution off the grid points

# Euler's method

The workhorse finite-difference method is Euler's method



# Euler's method


The workhorse finite-difference method is Euler's method

Euler's method is the difference equation

$$Y_{i+1} = Y_i + hf(t_i, Y_i)$$


where  $Y_0$  is given by the initial condition

# Euler's method



Suppose the current iterate is  $P = (t_i, Y_i)$  and  $y(t)$  is the true solution


# Euler's method



Suppose the current iterate is  $P = (t_i, Y_i)$  and  $y(t)$  is the true solution

At  $P$ ,  $y'(t_i)$  is the tangent vector  $\vec{PQ}$

# Euler's method




Suppose the current iterate is  $P = (t_i, Y_i)$  and  $y(t)$  is the true solution

At  $P$ ,  $y'(t_i)$  is the tangent vector  $\vec{PQ}$

Euler's method follows that direction until  $t = t_{i+1}$  at  $Q$

# Euler's method



Suppose the current iterate is  $P = (t_i, Y_i)$  and  $y(t)$  is the true solution

At  $P$ ,  $y'(t_i)$  is the tangent vector  $\vec{PQ}$

Euler's method follows that direction until  $t = t_{i+1}$  at  $Q$

The Euler estimate of  $y(t_{i+1})$  is then  $Y_{i+1}^E$

# Euler's method

This sounds very similar to Newton's method, because it is

# Euler's method

This sounds very similar to Newton's method, because it is

Euler's method can be motivated by a similar Taylor approximation argument

# Euler's method

This sounds very similar to Newton's method, because it is

Euler's method can be motivated by a similar Taylor approximation argument

If  $y(t)$  is the true solution, the second order Taylor expansion around  $t_i$  is

$$y(t_{i+1}) = y(t_i) + hy'(t_i) + \frac{h^2}{2}y''(\xi)$$

for some  $\xi \in [t_i, t_{i+1}]$



# Euler's method

$$y(t_{i+1}) = y(t_i) + hy'(t_i) + \frac{h^2}{2}y''(\xi)$$

If we drop the second order term and assume  $f(t_i, Y_i) = y'(t_i)$  and  $Y_i = y(t_i)$  we have exactly Euler's formula

# Euler's method

$$y(t_{i+1}) = y(t_i) + hy'(t_i) + \frac{h^2}{2}y''(\xi)$$

If we drop the second order term and assume  $f(t_i, Y_i) = y'(t_i)$  and  $Y_i = y(t_i)$  we have exactly Euler's formula

For small  $h$ ,  $y(x)$  should be a close approximation to the solution of the truncated Taylor expansion, so  $Y_i$  should be a good approximation to  $y(t_i)$

# Euler's method

This approach approximated  $y(t)$  with a linear function with slope  $f(t_i, Y_i)$  on the interval  $[t_i, t_{i+1}]$

# Euler's method

This approach approximated  $y(t)$  with a linear function with slope  $f(t_i, Y_i)$  on the interval  $[t_i, t_{i+1}]$

We can motivate Euler's method with an integration argument instead of a Taylor expansion argument

# Euler's method

This approach approximated  $y(t)$  with a linear function with slope  $f(t_i, Y_i)$  on the interval  $[t_i, t_{i+1}]$

We can motivate Euler's method with an integration argument instead of a Taylor expansion argument

The fundamental theorem of calculus tells us that

$$y(t_{i+1}) = y(t_i) + \int_{t_i}^{t_{i+1}} f(s, y(s)) ds$$

# Euler's method

$$y(t_{i+1}) = y(t_i) + \int_{t_i}^{t_{i+1}} f(s, y(s)) ds$$

If we approximate the integral with  $hf(t_i, y(t_i))$ , a box of width  $h$  and height  $f(t_i, y(t_i))$ , then  $y(t_{i+1}) = y(t_i) + hf(t_i, y(t_i))$  which implies the Euler method difference equation above if  $Y_i = y(t_i)$

# Euler's method

$$y(t_{i+1}) = y(t_i) + \int_{t_i}^{t_{i+1}} f(s, y(s)) ds$$

If we approximate the integral with  $hf(t_i, y(t_i))$ , a box of width  $h$  and height  $f(t_i, y(t_i))$ , then  $y(t_{i+1}) = y(t_i) + hf(t_i, y(t_i))$  which implies the Euler method difference equation above if  $Y_i = y(t_i)$

Thus this also approximate  $y(t)$  with a linear function over each subinterval with slope  $f(t_i, Y_i)$

# Euler's method

$$y(t_{i+1}) = y(t_i) + \int_{t_i}^{t_{i+1}} f(s, y(s)) ds$$

If we approximate the integral with  $hf(t_i, y(t_i))$ , a box of width  $h$  and height  $f(t_i, y(t_i))$ , then  $y(t_{i+1}) = y(t_i) + hf(t_i, y(t_i))$  which implies the Euler method difference equation above if  $Y_i = y(t_i)$

Thus this also approximate  $y(t)$  with a linear function over each subinterval with slope  $f(t_i, Y_i)$

As  $h$  decreases, we would expect the solutions to become more accurate



# Euler's method

$$y(t_{i+1}) = y(t_i) + \int_{t_i}^{t_{i+1}} f(s, y(s)) ds$$

If we approximate the integral with  $hf(t_i, y(t_i))$ , a box of width  $h$  and height  $f(t_i, y(t_i))$ , then  $y(t_{i+1}) = y(t_i) + hf(t_i, y(t_i))$  which implies the Euler method difference equation above if  $Y_i = y(t_i)$

Thus this also approximate  $y(t)$  with a linear function over each subinterval with slope  $f(t_i, Y_i)$

As  $h$  decreases, we would expect the solutions to become more accurate

As  $h \rightarrow 0$ , we are back in the ODE world

# Euler's method errors

Consider a system,  $y'(t) = y(t)$ ,  $y(0) = 1$

# Euler's method errors

Consider a system,  $y'(t) = y(t)$ ,  $y(0) = 1$

The solution to this is simply  $y(t) = e^t$

# Euler's method errors

Consider a system,  $y'(t) = y(t)$ ,  $y(0) = 1$

The solution to this is simply  $y(t) = e^t$

The Euler method gives us a difference equation of

$$Y_{i+1} = Y_i + hY_i = (1 + h)Y_i$$

# Euler's method errors

Consider a system,  $y'(t) = y(t)$ ,  $y(0) = 1$

The solution to this is simply  $y(t) = e^t$

The Euler method gives us a difference equation of

$$Y_{i+1} = Y_i + hY_i = (1 + h)Y_i$$

This difference equation has solution  $Y_i = (1 + h)^i$  and implies the approximation is  $Y(t) = (1 + h)^{t/h}$

# Euler's method errors

Thus the relative error between the two is

$$\log(|Y(t)/y(t)|) = \frac{t}{h} \log(1 + h) - t = \frac{t}{h} (h - h^2 + \dots) - t = -th + \dots$$

where excluded terms have order higher than  $h$

# Euler's method errors

Thus the relative error between the two is

$$\log(|Y(t)/y(t)|) = \frac{t}{h} \log(1 + h) - t = \frac{t}{h} (h - h^2 + \dots) - t = -th + \dots$$

where excluded terms have order higher than  $h$

Thus the relative error in the Euler approximation has order  $h$  and as  $h$  goes to zero so does the approximation error

# Euler's method errors

In general we can show that Euler's method has linear convergence

*Suppose the solution to  $y'(t) = f(t, y(t))$ ,  $y(t_0) = y_0$  is  $C^3$  on  $[t_0, T]$ , that  $f$  is  $C^2$ , and that  $f_y$  and  $f_{yy}$  are bounded for all  $y$  and  $t_0 \leq t \leq T$ . Then the error of the Euler scheme with step size  $h$  is  $O(h)$*



# Euler's method code

# Euler's method code

```
function euler_ode(df, t0, y0, h, n)

    t = zeros(n+1)
    y = zeros(n+1)

    # set the initial values
    t[1] = t0
    y[1] = y0

    # use Euler's method to approximate the solution at each step
    for i in 1:n
        t[i+1] = t[i] + h
        y[i+1] = y[i] + h * df(t[i], y[i])
    end

    return (t, y)

end
```


## euler\_ode (generic function with 1 method)

# Euler's method code

```
#  $dy/dt = y \rightarrow y = C_0 * \exp(t)$   
df(t, y) = y  
t1, y1 = euler_ode(df, 0., 1., .1, 10)
```

Define  $df/dt$  and send it to the `euler_ode` function

# Euler's method code



# Implicit Euler method

We expanded  $y$  around  $t_i$ , but we could always expand around  $t_{i+1}$  so that we have

$$y(t_i) = y(t_{i+1}) - hy'(t_{i+1}) = y(t_{i+1}) - hf(t_{i+1}, y(t_{i+1}))$$

# Implicit Euler method

We expanded  $y$  around  $t_i$ , but we could always expand around  $t_{i+1}$  so that we have

$$y(t_i) = y(t_{i+1}) - hy'(t_{i+1}) = y(t_{i+1}) - hf(t_{i+1}, y(t_{i+1}))$$

This yields the implicit Euler method

$$Y_{i+1} = Y_i + hf(t_{i+1}, Y_{i+1})$$

# Implicit Euler method

We expanded  $y$  around  $t_i$ , but we could always expand around  $t_{i+1}$  so that we have

$$y(t_i) = y(t_{i+1}) - hy'(t_{i+1}) = y(t_{i+1}) - hf(t_{i+1}, y(t_{i+1}))$$

This yields the implicit Euler method

$$Y_{i+1} = Y_i + hf(t_{i+1}, Y_{i+1})$$

Notice that now  $Y_{i+1}$  is only implicitly defined in terms of  $t_i$  and  $Y_i$  so we will need to solve a non-linear equation in  $Y_{i+1}$

# Implicit Euler method

This seems not great, before we simply computed  $Y_{i+1}$  from values known at  $i$  but now we have to perform a rootfinding problem



# Implicit Euler method

This seems not great, before we simply computed  $Y_{i+1}$  from values known at  $i$  but now we have to perform a rootfinding problem

However,  $Y_{i+1}$  does not simply depend on only  $Y_i$  and  $t_{i+1}$  but also  $f$  at  $t_{i+1}$

# Implicit Euler method

This seems not great, before we simply computed  $Y_{i+1}$  from values known at  $i$  but now we have to perform a rootfinding problem

However,  $Y_{i+1}$  does not simply depend on only  $Y_i$  and  $t_{i+1}$  but also  $f$  at  $t_{i+1}$

Thus the implicit Euler method will get us better approximation properties, often times much better

# Implicit Euler method

This seems not great, before we simply computed  $Y_{i+1}$  from values known at  $i$  but now we have to perform a rootfinding problem

However,  $Y_{i+1}$  does not simply depend on only  $Y_i$  and  $t_{i+1}$  but also  $f$  at  $t_{i+1}$

Thus the implicit Euler method will get us better approximation properties, often times much better

Because of this we can typically use larger  $h$ 's with the implicit Euler method

# Implicit Euler's method code

```
# rootfinding portion of implicit Euler
function find_euler_root(df, y, t, h, y0, tol)

    y_new = y0
    y_old = y0
    error = Inf

    while error > tol
        y_new = y + h * df(t, y_new)
        error = abs((y_new - y_old)/y_old)
        y_old = deepcopy(y_new)
    end

    return y_new

end
```

```
## find_euler_root (generic function with 1 method)
```

# Implicit Euler's method code

```
function euler_implicit_ode(df, t0, y0, h, n, tol = 1e-6)

    t = zeros(n+1)
    y = zeros(n+1)
    t[1] = t0
    y[1] = y0

    for i in 1:n
        t[i+1] = t[i] + h
        y[i+1] = find_euler_root(df, y[i], t[i+1], h, y[i], tol)
    end

    return (t, y)

end
```


```
## euler_implicit_ode (generic function with 2 methods)
```

# Implicit Euler's method code

```
df(t, y) = y  
t2, y2 = euler_implicit_ode(df, 0., 1., .1, 10, 1e-6)
```

Define  $df/dt$  and send it to the `euler_implicit_ode` function

# Implicit Euler's method code




# Comparison

```
df(t, y) = y  
t1, y1 = euler_ode(df, 0., 1., .1, 10)  
t2, y2 = euler_implicit_ode(df, 0., 1., .1, 10, 1e-6)  
y_real = exp.(t1)
```



# Comparison



# Runge-Kutta methods

Runge-Kutta methods take Euler methods but adapts  $f$

# Runge-Kutta methods

Runge-Kutta methods take Euler methods but adapts  $f$

In the standard Euler approach we implicitly assume that the slope at  $(t_{i+1}, Y_{i+1}^E)$  is the same as the slope at  $(t_i, Y_i^E)$  which is a bad assumption unless  $y$  is linear

# Runge-Kutta methods

Runge-Kutta methods take Euler methods but adapts  $f$

In the standard Euler approach we implicitly assume that the slope at  $(t_{i+1}, Y_{i+1}^E)$  is the same as the slope at  $(t_i, Y_i^E)$  which is a bad assumption unless  $y$  is linear

For example if  $y$  is concave we will overshoot the true value

# Runge-Kutta methods

Runge-Kutta methods take Euler methods but adapts  $f$

In the standard Euler approach we implicitly assume that the slope at  $(t_{i+1}, Y_{i+1}^E)$  is the same as the slope at  $(t_i, Y_i^E)$  which is a bad assumption unless  $y$  is linear

For example if  $y$  is concave we will overshoot the true value

We could instead use the slope at  $(t_{i+1}, Y_{i+1}^E)$  but this will give the same problem but in the opposite direction, we will undershoot

# Runge-Kutta methods

Runge-Kutta methods recognizes these two facts

# Runge-Kutta methods

Runge-Kutta methods recognizes these two facts

A first-order Runge-Kutta method will take the average of these two slopes to arrive at the formula

$$Y_{i+1} = Y_i + \frac{h}{2} [f(t_i, Y_i) + f(t_{i+1}, Y_{i+1})]$$

# Runge-Kutta methods

Runge-Kutta methods recognizes these two facts

A first-order Runge-Kutta method will take the average of these two slopes to arrive at the formula

$$Y_{i+1} = Y_i + \frac{h}{2} [f(t_i, Y_i) + f(t_{i+1}, Y_{i+1})]$$

This converges quadratically to the true solution in  $h$ , but now uses two evaluations per step



# Runge-Kutta methods

Runge-Kutta methods recognizes these two facts

A first-order Runge-Kutta method will take the average of these two slopes to arrive at the formula

$$Y_{i+1} = Y_i + \frac{h}{2} [f(t_i, Y_i) + f(t_{i+1}, Y_{i+1})]$$

This converges quadratically to the true solution in  $h$ , but now uses two evaluations per step

This has the same flavor as forward vs central finite differences

# Runge-Kutta methods

Runge-Kutta methods recognizes these two facts

A first-order Runge-Kutta method will take the average of these two slopes to arrive at the formula

$$Y_{i+1} = Y_i + \frac{h}{2} [f(t_i, Y_i) + f(t_{i+1}, Y_{i+1})]$$

This converges quadratically to the true solution in  $h$ , but now uses two evaluations per step

This has the same flavor as forward vs central finite differences

There are higher order Runge-Kutta rules that have even more desirable

# Runge-Kutta code

```
function find_euler_root_rk(df, y, t, tp, h, y0, tol)

    y_new = y0
    y_old = y0
    error = Inf

    while error > tol
        y_new = y + h/2 * (df(t, y) + df(tp, y_new))
        error = abs((y_new - y_old)/y_old)
        y_old = deepcopy(y_new)
    end

    return y_new

end
```

```
## find_euler_root_rk (generic function with 1 method)
```

# Runge-Kutta code

```
function euler_rk_ode(df, t0, y0, h, n, tol = 1e-6)

    t = zeros(n+1)
    y = zeros(n+1)
    t[1] = t0
    y[1] = y0

    for i in 1:n
        t[i+1] = t[i] + h
        y[i+1] = find_euler_root_rk(df, y[i], t[i], t[i+1], h, y0, tol)
    end

    return (t, y)

end
```


```
## euler_rk_ode (generic function with 2 methods)
```

# Comparison


```
df(t, y) = y
t1, y1 = euler_ode(df, 0., 1., .1, 100)
t2, y2 = euler_implicit_ode(df, 0., 1., .1, 100, 1e-7)
t2, y3 = euler_rk_ode(df, 0., 1., .1, 100, 1e-7)
y_real = exp.(t1)
```

Check the time/memory with `@btime` in `BenchmarkTools`

# Comparison: RK has minimal error



# Comparison: RK has minimal error



# Boundary Value Problems

IVPs are easy to solve because the solution depends only on local conditions so we can use local solution algorithms which are convenient



# Boundary Value Problems

IVPs are easy to solve because the solution depends only on local conditions so we can use local solution algorithms which are convenient

BVPs have auxiliary conditions that are imposed at different points in time so we lose the local nature of the problem and our solutions must now be global in nature

# Boundary Value Problems

Consider the following BVP

$$\dot{x} = f(x, y, t)$$

$$\dot{y} = g(x, y, t)$$

$$x(t_0) = x_0, \quad y(T) = y_T$$

where  $x \in \mathbb{R}^n, y \in \mathbb{R}^m$

# Boundary Value Problems

Consider the following BVP

$$\begin{aligned}\dot{x} &= f(x, y, t) \\ \dot{y} &= g(x, y, t) \\ x(t_0) &= x_0, \quad y(T) = y_T\end{aligned}$$

where  $x \in \mathbb{R}^n, y \in \mathbb{R}^m$

We cannot use standard IVP approaches because at  $t_0$  or  $T$  we only know the value of either  $x$  or  $y$  but not both

# Boundary Value Problems

Consider the following BVP

$$\begin{aligned}\dot{x} &= f(x, y, t) \\ \dot{y} &= g(x, y, t) \\ x(t_0) &= x_0, \quad y(T) = y_T\end{aligned}$$

where  $x \in \mathbb{R}^n, y \in \mathbb{R}^m$

We cannot use standard IVP approaches because at  $t_0$  or  $T$  we only know the value of either  $x$  or  $y$  but not both

Thus we cannot find the next value of both of them using only local information: we need alternative approaches

# Boundary Value Problems: Shooting

The core method for solving BVPs is called **shooting**

# Boundary Value Problems: Shooting

The core method for solving BVPs is called **shooting**

The idea behind shooting is that we guess the value of  $y(t_0)$ , and use an IVP method to see what that means about  $y(T)$

# Boundary Value Problems: Shooting

The core method for solving BVPs is called **shooting**

The idea behind shooting is that we guess the value of  $y(t_0)$ , and use an IVP method to see what that means about  $y(T)$

For any given guess, we generally won't hit the terminal condition exactly and might not even be that close

# Boundary Value Problems: Shooting

The core method for solving BVPs is called **shooting**

The idea behind shooting is that we guess the value of  $y(t_0)$ , and use an IVP method to see what that means about  $y(T)$

For any given guess, we generally won't hit the terminal condition exactly and might not even be that close

But we do get some information from where we end up at  $y(T)$  and can use that information to update our guesses for  $y(t_0)$  until we are sufficiently close to  $y_T$



# Boundary Value Problems: Shooting

The core method for solving BVPs is called **shooting**

The idea behind shooting is that we guess the value of  $y(t_0)$ , and use an IVP method to see what that means about  $y(T)$

For any given guess, we generally won't hit the terminal condition exactly and might not even be that close

But we do get some information from where we end up at  $y(T)$  and can use that information to update our guesses for  $y(t_0)$  until we are sufficiently close to  $y_T$

There are two components to a shooting method

# Shooting

First we guess some  $y(0) = y_0$  and then solve the IVP problem with methods we've already used

$$\dot{x} = f(x, y, t)$$

$$\dot{y} = g(x, y, t)$$

$$x(t_0) = x_0, \quad y(0) = y_0$$

to find some  $y(T)$  which we call  $Y(T, y_0)$  since it depends on our initial guess  $y_0$

# Shooting

Second we need to find the *right*  $y_0$

# Shooting

Second we need to find the *right*  $y_0$

We want to find a  $y_0$  such that  $y_T = Y(T, y_0)$

# Shooting

Second we need to find the *right*  $y_0$

We want to find a  $y_0$  such that  $y_T = Y(T, y_0)$

This is a nonlinear equation in  $y_0$  so we need to solve nonlinear equations

# Shooting

Second we need to find the *right*  $y_0$

We want to find a  $y_0$  such that  $y_T = Y(T, y_0)$

This is a nonlinear equation in  $y_0$  so we need to solve nonlinear equations

We can write the algorithm as

1. Initialize: Guess  $y_0^i$ . Choose a stopping criterion  $\epsilon > 0$
2. Solve the IVP for  $x(T), y(T)$  given the initial condition  $y_0 = y_0^i$
3. If  $\|y(T) - y_T\| < \epsilon$ , STOP. Else choose  $y_0^{i+1}$  based on the previous values of  $y$  and go back to step 1

# Shooting

This is an example of a two layer algorithm

# Shooting

This is an example of a two layer algorithm

The inner layer (step 1) uses an IVP method that solves  $Y(T, y_0)$  for any  $y_0$



# Shooting

This is an example of a two layer algorithm

The inner layer (step 1) uses an IVP method that solves  $Y(T, y_0)$  for any  $y_0$

This can be Euler, Runge-Kutta or anything else

# Shooting

This is an example of a two layer algorithm

The inner layer (step 1) uses an IVP method that solves  $Y(T, y_0)$  for any  $y_0$

This can be Euler, Runge-Kutta or anything else

In the outer layer (step 2) we solve the nonlinear equation  $Y(T, y_0) = y_T$

# Shooting

This is an example of a two layer algorithm

The inner layer (step 1) uses an IVP method that solves  $Y(T, y_0)$  for any  $y_0$

This can be Euler, Runge-Kutta or anything else

In the outer layer (step 2) we solve the nonlinear equation  $Y(T, y_0) = y_T$

We can use any nonlinear solver here, typically we do this by defining a subroutine that computes  $Y(T, y_0) - y_T$  as a function of  $y_0$  and then sends that subroutine to a rootfinding program

# Example: Lifecycle model

A simple lifecycle model is given by

$$\begin{aligned} \max_{c(t)} \quad & \int_0^T e^{-rt} u(c(t)) dt \\ \text{s. t.} \quad & \dot{A}(t) = f(A(t)) + w(t) - c(t) \\ & A(0) = A(T) = 0. \end{aligned}$$

$u(c(t))$  is utility from consumption,  $w(t)$  is the wage rate,  $A(t)$  are assets and  $f(A(t))$  is the return on invested assets

# Example: Lifecycle model

A simple lifecycle model is given by

$$\begin{aligned} \max_{c(t)} \quad & \int_0^T e^{-rt} u(c(t)) dt \\ \text{s. t.} \quad & \dot{A}(t) = f(A(t)) + w(t) - c(t) \\ & A(0) = A(T) = 0. \end{aligned}$$

$u(c(t))$  is utility from consumption,  $w(t)$  is the wage rate,  $A(t)$  are assets and  $f(A(t))$  is the return on invested assets

We assume that assets are initially and terminally zero where the latter would come about naturally from a transversality condition

# Example: Lifecycle model

The Hamiltonian is

$$H = u(c(t)) + \lambda(t) [f(A(t)) + w(t) - c(t)]$$

# Example: Lifecycle model

The Hamiltonian is

$$H = u(c(t)) + \lambda(t) [f(A(t)) + w(t) - c(t)]$$

and the co-state condition is given by

$$\dot{\lambda}(t) = r\lambda(t) - \lambda(t)f'(A(t))$$

# Example: Lifecycle model

The Hamiltonian is

$$H = u(c(t)) + \lambda(t) [f(A(t)) + w(t) - c(t)]$$

and the co-state condition is given by

$$\dot{\lambda}(t) = r\lambda(t) - \lambda(t)f'(A(t))$$

The maximum principle implies that  $u'(c(t)) = \lambda(t)$



# Example: Lifecycle model

This gives us a two equation system of differential equations (1 for the  $A$  transition, 1 for the costate condition) and the boundary conditions on  $A$  are what pin down the problem

# Example: Lifecycle model

This gives us a two equation system of differential equations (1 for the  $A$  transition, 1 for the costate condition) and the boundary conditions on  $A$  are what pin down the problem

The issue here is that we never know  $A$  and  $\lambda$  at either  $t = 0$  or  $t = T$

# Example: Lifecycle model

This gives us a two equation system of differential equations (1 for the  $A$  transition, 1 for the costate condition) and the boundary conditions on  $A$  are what pin down the problem


The issue here is that we never know  $A$  and  $\lambda$  at either  $t = 0$  or  $t = T$

We can use the maximum principle to convert the costate condition into a condition on consumption

$$\dot{c}(t) = -\frac{u'(c(t))}{u''(c(t))} [f'(A(t)) - r]$$

# Example: Lifecycle model

The Figure shows the phase diagram assuming that  $f'(A) > r$  for all  $A$



If  $A(T) < 0$  when we guess  $c(0) = c_H$ , but  $A(T) > 0$  when we guess  $c(0) = c_L$ , we know the correct guess lies in between and we can solve for it using the

# Example: Lifecycle model

Let's code it up

$$\dot{A}(t) = f(A(t)) + w(t) - c(t) \qquad \dot{c}(t) = -\frac{u'(c(t))}{u''(c(t))} [f'(A(t)) - r]$$

- $f(A(t)) = 1.05A(t)$
- $u(c(t)) = \log(c(t))$
- $w(t) = 5$
- $r = .02$

# Example: Lifecycle model

Let's code it up

$$\dot{A}(t) = f(A(t)) + w(t) - c(t) \quad \dot{c}(t) = -\frac{u'(c(t))}{u''(c(t))} [f'(A(t)) - r]$$

- $f(A(t)) = 1.05A(t)$
- $u(c(t)) = \log(c(t))$
- $w(t) = 5$
- $r = .02$

```
df(t,a,c) = (1.05*a + 5 - c, -1 * (1/c) / (-1/c^2) * (1.05 - .02))
```

```
## df (generic function with 2 methods)
```

# Example: Lifecycle model

We need a 2 variable ODE solver next

# Example: Lifecycle model

We need a 2 variable ODE solver next

```
function euler_ode(df, t0, a0, c0, h, n)

    t = zeros(n+1)
    a = zeros(n+1)
    c = zeros(n+1)

    t[1] = t0
    a[1] = a0
    c[1] = c0

    for i in 1:n
        t[i+1] = t[i] + h
        a[i+1] = a[i] + h * df(t[i], a[i], c[i])[1]
        c[i+1] = c[i] + h * df(t[i], a[i], c[i])[2]
    end

    return t, a, c

end
```



# Example: Lifecycle model

Last, wrap it in bisection method

```
function solve_bvp(df, t0, a0, aend, c0low, c0high, h, n, tol = 1e-6)

    t = zeros(n+1)
    a = zeros(n+1)
    c = zeros(n+1)

    while abs.(c0low - c0high) > tol

        c0guess = (c0low + c0high)/2
        t, a, c = euler_ode(df, t0, a0, c0guess, h, n)
        anew = a[end]

        if sign(anew) > 0
            c0low = c0guess
        else
            c0high = c0guess
        end

    end
```

# Example: Lifecycle model

Now we have to find the initial bounds, one where  $A(T) > 0$ , one where  $A(T) < 0$

```
aend = 0.  
a0 = 0.  
t0 = 0.  
h = .01  
n = 100  
c0low = 1      # low c0 guess  
c0high = 10    # high c0 guess
```

# Example: Lifecycle model

```
a0high = euler_ode(df, t0, a0, c0high, h, n)[2][end]
```

```
## -19.07974813179398
```

```
a0low = euler_ode(df, t0, a0, c0low, h, n)[2][end]
```


```
## 5.986527176940143
```

As expected, too high consumption  $C(0)$  yields negative assets  $A(T)$ , too low consumption  $C(0)$  yields positive assets  $A(T)$

The  $C(0)$  that solves the problem will fall somewhere in between

# Example: Lifecycle model

```
t, a, c = solve_bvp(df, t0, a0, aend, c0low, c0high, h, n)
```



# Reverse shooting for $\infty$ horizon problems

The standard infinite horizon optimal control problem is

$$\begin{aligned} \max_{u(t)} \quad & \int_0^{\infty} e^{-rt} \pi(x(t), u(t)) dt \\ \text{s. t.} \quad & \dot{x}(t) = f(x(t), u(t)) \\ & x(0) = x_0. \end{aligned}$$

# Reverse shooting for $\infty$ horizon problems

The standard infinite horizon optimal control problem is

$$\begin{aligned} \max_{u(t)} \quad & \int_0^{\infty} e^{-rt} \pi(x(t), u(t)) dt \\ \text{s. t.} \quad & \dot{x}(t) = f(x(t), u(t)) \\ & x(0) = x_0. \end{aligned}$$

We still have  $x(0) = x_0$  as before, but we no longer have the terminal condition

# Reverse shooting for $\infty$ horizon problems

The standard infinite horizon optimal control problem is

$$\begin{aligned} \max_{u(t)} \quad & \int_0^{\infty} e^{-rt} \pi(x(t), u(t)) dt \\ \text{s. t.} \quad & \dot{x}(t) = f(x(t), u(t)) \\ & x(0) = x_0. \end{aligned}$$

We still have  $x(0) = x_0$  as before, but we no longer have the terminal condition

We replace it with a transversality condition that  $\lim_{t \rightarrow \infty} e^{-rt} |\lambda(t)^T x(t)| \leq \infty$

# Reverse shooting for $\infty$ horizon problems

Shooting methods do not really work for infinite horizon problems since we need to integrate the problem over a very long time horizon and so  $x(T)$  will be particularly sensitive to  $\lambda(0)$  when  $T$  is large



# Reverse shooting for $\infty$ horizon problems

Shooting methods do not really work for infinite horizon problems since we need to integrate the problem over a very long time horizon and so  $x(T)$  will be particularly sensitive to  $\lambda(0)$  when  $T$  is large

The primary issue is that the terminal state, because of the long time horizon, is very sensitive to the initial guess

# Reverse shooting for $\infty$ horizon problems

Shooting methods do not really work for infinite horizon problems since we need to integrate the problem over a very long time horizon and so  $x(T)$  will be particularly sensitive to  $\lambda(0)$  when  $T$  is large

The primary issue is that the terminal state, because of the long time horizon, is very sensitive to the initial guess

But this implies something very convenient: that the initial state corresponding to any terminal state is not very sensitive to the value of the terminal state

# Reverse shooting for $\infty$ horizon problems

Shooting methods do not really work for infinite horizon problems since we need to integrate the problem over a very long time horizon and so  $x(T)$  will be particularly sensitive to  $\lambda(0)$  when  $T$  is large

The primary issue is that the terminal state, because of the long time horizon, is very sensitive to the initial guess

But this implies something very convenient: that the initial state corresponding to any terminal state is not very sensitive to the value of the terminal state

Thus what we will do is not guess the value of the initial condition and integrate forward, we will guess the terminal condition and integrate

# Example: Reverse shooting for $\infty$ horizon problems

Consider the simplest growth model

$$\begin{aligned} \max_{c(t)} \quad & \int_0^{\infty} e^{-rt} u(c(t)) dt \\ \dot{k}(t) = & f(k(t)) - c(t) \\ \text{s.t.} \quad & k(0) = k_0, \end{aligned}$$

where  $c$  is consumption,  $k$  is the capital stock, and  $f$  is production

# Example: Reverse shooting for $\infty$ horizon problems

We can use Pontryagin's necessary conditions to get that consumption and capital are governed by the following differential equations


$$\begin{aligned}\dot{c}(t) &= -\frac{u'(c(t))}{u''(c(t))} (f'(k) - r) \\ \dot{k}(t) &= f(k(t)) - c(t),\end{aligned}$$

with boundary conditions,

$$k(0) = k_0, \quad 0 < \lim_{t \rightarrow \infty} |k(t)| \leq \infty$$

# Example: Reverse shooting for $\infty$ horizon problems

Assume  $u$  and  $f$  are concave, the Figure shows the phase diagram for the problem



We have a steady state at  $k = k^*$ , this occurs when  $f'(k(t)) = r$  and

# Example: Reverse shooting for $\infty$ horizon problems

For this problem there exists a stable manifold  $M_S$  and an unstable manifold  $M_U$

so that the steady state is **saddle point stable**

# Example: Reverse shooting for $\infty$ horizon problems

For this problem there exists a stable manifold  $M_S$  and an unstable manifold  $M_U$

so that the steady state is **saddle point stable**

Both are invariant manifolds because any system that starts on either of these manifolds will continue to move along the manifold



# Example: Reverse shooting for $\infty$ horizon problems

For this problem there exists a stable manifold  $M_S$  and an unstable manifold  $M_U$

so that the steady state is **saddle point stable**

Both are invariant manifolds because any system that starts on either of these manifolds will continue to move along the manifold

However  $M_S$  is stable because it will converge to the steady state while  $M_U$  diverges away from the steady state

# Example: Reverse shooting for $\infty$ horizon problems

Lets first use standard shooting to try to compute the stable manifold

# Example: Reverse shooting for $\infty$ horizon problems

Lets first use standard shooting to try to compute the stable manifold

We want  $k$  and  $c$  to equal their steady state values at  $t = \infty$ , but we can't quite do that so we search for a  $c(0)$  so that  $(c(t), k(t))$  has a path that is close to the steady state

# Example: Reverse shooting for $\infty$ horizon problems

Lets first use standard shooting to try to compute the stable manifold

We want  $k$  and  $c$  to equal their steady state values at  $t = \infty$ , but we can't quite do that so we search for a  $c(0)$  so that  $(c(t), k(t))$  has a path that is close to the steady state

Suppose we start with  $k_0 < k^*$ , if we guess  $c(0)$  too big we will cross the  $k$  isoquant and have a falling capital stock, but if we guess  $c(0)$  too small we will get a path that crosses the  $c$  isoquant and results in a falling consumption level

# Example: Reverse shooting for $\infty$ horizon problems

This gives us our algorithm

1. Initialize: set  $c_H = f(k_0)$  and set  $c_L = 0$ , choose a stopping criterion  $\epsilon > 0$
2. Set  $c_0 = \frac{1}{2}(c_L + c_H)$
3. Solve the IVP with initial conditions  $c(0) = c_0, k(0) = k_0$ . Stop the IVP at the first  $t$  when  $\dot{c}(t) < 0$  or  $\dot{k}(t) < 0$ , denote this  $T$
4. If  $|c(T) - c^*| < \epsilon$ , STOP. If  $\dot{c}(t) < 0$ , set  $c_L = c_0$ , else set  $c_H = c_0$ . Go to step 2.

# Example: Reverse shooting for $\infty$ horizon problems

This algorithm makes sense but the phase diagram shows why it will have trouble finding the stable manifold

# Example: Reverse shooting for $\infty$ horizon problems

This algorithm makes sense but the phase diagram shows why it will have trouble finding the stable manifold

Any small deviation from  $M_S$  is magnified and results in a path that increasingly gets far away from  $M_S$

# Example: Reverse shooting for $\infty$ horizon problems

This algorithm makes sense but the phase diagram shows why it will have trouble finding the stable manifold

Any small deviation from  $M_S$  is magnified and results in a path that increasingly gets far away from  $M_S$

Unless we happen to pick a point precisely on the stable manifold we will move away from it, so it is hard to search for the solution since changes in our guesses will lead to wild changes in terminal values



# Example: Reverse shooting for $\infty$ horizon problems

Now suppose we wanted to find a path on  $M_U$ , notice that the flow pushes points *toward*  $M_U$  so the deviations are smushed together

# Example: Reverse shooting for $\infty$ horizon problems

Now suppose we wanted to find a path on  $M_U$ , notice that the flow pushes points *toward*  $M_U$  so the deviations are smushed together

If we wanted to compute a path that lies near the unstable manifold, we could simply pick a point near the steady state as the initial condition and integrate the system

# Example: Reverse shooting for $\infty$ horizon problems

Now suppose we wanted to find a path on  $M_U$ , notice that the flow pushes points *toward*  $M_U$  so the deviations are smushed together

If we wanted to compute a path that lies near the unstable manifold, we could simply pick a point near the steady state as the initial condition and integrate the system

We don't actually want to solve for a path on  $M_U$  but this gives us some insight

# Example: Reverse shooting for $\infty$ horizon problems

We want to change the system so that the stable manifold becomes the unstable manifold

# Example: Reverse shooting for $\infty$ horizon problems

We want to change the system so that the stable manifold becomes the unstable manifold

We can do this easily by reversing time

$$\begin{aligned}\dot{c}(t) &= \frac{u'(c(t))}{u''(c(t))} (f'(k) - r) \\ \dot{k}(t) &= -(f(k(t)) - c(t))\end{aligned}$$

# Example: Reverse shooting for $\infty$ horizon problems

We want to change the system so that the stable manifold becomes the unstable manifold

We can do this easily by reversing time

$$\begin{aligned}\dot{c}(t) &= \frac{u'(c(t))}{u''(c(t))} (f'(k) - r) \\ \dot{k}(t) &= -(f(k(t)) - c(t))\end{aligned}$$

This yields the same phase diagram but with the arrows turned  $180^\circ$  so the stable manifold forward in time is now the unstable manifold reverse in time

# Example: Reverse shooting for $\infty$ horizon problems

We want to change the system so that the stable manifold becomes the unstable manifold


We can do this easily by reversing time

$$\begin{aligned}\dot{c}(t) &= \frac{u'(c(t))}{u''(c(t))} (f'(k) - r) \\ \dot{k}(t) &= -(f(k(t)) - c(t))\end{aligned}$$

This yields the same phase diagram but with the arrows turned  $180^\circ$  so the stable manifold forward in time is now the unstable manifold reverse in time

This allows us to exploit how paths tend to converge toward the stable

# Reverse shooting for $\infty$ horizon problems





# Reverse shooting example

Let's code it up

- $f(k) = \sqrt{k}$
- $r = .02$
- $u(c) = \log(c)$

# Reverse shooting example

Let's code it up

- $f(k) = \sqrt{k}$
- $r = .02$
- $u(c) = \log(c)$

We know the steady state is where  $f'(k) = r$  and  $f(k) = c$

# Reverse shooting example

Let's code it up

- $f(k) = \sqrt{k}$
- $r = .02$
- $u(c) = \log(c)$

We know the steady state is where  $f'(k) = r$  and  $f(k) = c$

This is  $k = 625, c = 25$

# Reverse shooting example

Let's code it up

- $f(k) = \sqrt{k}$
- $r = .02$
- $u(c) = \log(c)$

We know the steady state is where  $f'(k) = r$  and  $f(k) = c$

This is  $k = 625, c = 25$

Pretend we only knew  $f(k) = c$  (from  $\dot{K} = 0$ ) and solve by searching over terminal capital

# Example: Lifecycle model

```
df(t,k,c) = (  
    -(sqrt(k) - c),  
    -(-1 * (1/c) / (-1/c^2) * (0.5*k^(-0.5) - .02))  
)
```

```
## df (generic function with 2 methods)
```

# Reverse shooting example

We need a 2 variable ODE solver next

# Reverse shooting example

We need a 2 variable ODE solver next

```
function euler_ode(df, t0, k0, c0, h, n)

    t = zeros(n+1)
    k = zeros(n+1)
    c = zeros(n+1)

    t[1] = t0
    k[1] = k0
    c[1] = c0

    for i in 1:n
        t[i+1] = t[i] + h
        k[i+1] = max(1e-6, k[i] + h * df(t[i], k[i], c[i])[1])
        c[i+1] = max(1e-6, c[i] + h * df(t[i], k[i], c[i])[2])
    end

    return t, k, c

end
```

# Reverse shooting example

Last, wrap it in bisection method

```
function solve_bvp_rev(df, t0, k0, klow, khigh, h, n, tol = 1e-6)

    t = zeros(n+1)
    k = zeros(n+1)
    c = zeros(n+1)

    while abs.(klow - khigh) > tol
        kguess = (klow + khigh)/2
        t, k, c = euler_ode(df, t0, kguess, sqrt(kguess), h, n)
        anew = k[end]
        if anew < k0
            klow = kguess
        else
            khigh = kguess
        end
    end

    return t, k, c
```



# Reverse shooting example

Now we have to find the initial bounds, one where  $A(T) > 0$ , one where  $A(T) < 0$

```
k0 = 10 # initial condition to hit
t0 = 0. # time starts at 0


klow = 100 # below closed-form solution of k = 625
khigh = 1000 # above closed-form solution of k = 625

aend = (.5 / .02)^2 # closed-form solution
cend = sqrt(aend) # closed-form solution

h = .1
n = 10000 # make the horizon long to approx infinite-horizon
```

# Reverse shooting example

```
t, k, c = solve_bvp_rev(df, t0, k0, klow, khigh, h, n)
```



# Reverse shooting example

```
k0 = 800
```

```
t, k, c = solve_bvp_rev(df, t0, k0, klow, khigh, h, n)
```

