



## Supplementary Figures




**Supplementary Fig. 1 | Experimental tomographic tilt series of the multi-component glass-forming nanoparticle (particle 1) after denoising.** 55 ADF-STEM images of the nanoparticle with a tilt range from  $-69.4^{\circ}$  to  $+72.6^{\circ}$ . Scale bar, 2 nm.




**Supplementary Fig. 2 | 55 multislice STEM images of the multi-component glass-forming nanoparticle (particle 1).** The multislice simulation was calculated from the experimental 3D atomic model (Methods). To account for the source size and incoherent effects, each multislice image was convolved with a Gaussian function. Scale bar, 2 nm.





**Supplementary Fig. 3 | 53 raw experimental images of a polycrystalline nanoparticle (particle 5).** The insets show the 2D power spectra of the experimental images, where the Bragg peaks are visible. Scale bar, 2 nm.



**Supplementary Fig. 4 | Determination of the 3D atomic structure of the polycrystalline nanoparticle (particle 5).** **a**, Representative raw experimental image at  $0^\circ$ . Scale bar, 2 nm. **b**, Average 2D power spectrum of the 53 experimental images (Supplementary Fig. 3), in which the Bragg peaks are visible. **c**, **d**, Two representative 2.4-Å-thick internal slices of the 3D reconstruction in the  $xy$  and  $xz$  planes, respectively. **e**, Experimental 3D atomic model of the polycrystalline nanoparticle, where green, blue and red balls represent type 1, 2 and 3 atoms, respectively. **f**, Distribution of ten crystalline grains (in different colors) inside the nanoparticle. **g**, The local BOO parameters of all the atoms in the polycrystalline nanoparticle. Based on the criterion of the normalized BOO parameter  $\geq 0.5$  (dashed red curve), there are 94.25% crystalline atoms. **h**, The PDF of all the atoms in the nanoparticle, where the peaks of different crystal planes are visible. **i**, Ten most abundant Voronoi polyhedra in the polycrystalline nanoparticle.