

a
Design effect by cell type and tissue, GSE325587 (cell types with median proportion ≥0.5%)
Distinct proximal tubule 1 (23.6%)
Epithelial progenitor cell (15.2%)
Proliferating Proximal Tubule (11.1%)
Connecting tubule (10.9%)
MNP-c/dendritic cell (8.7%)
Fibroblast (7.1%)
Ascending vasa recta endothelium (4.7%)
Myofibroblast (3.0%)
Podocyte (2.1%)
B cell (2.1%)
Principal cell (1.4%)
CD4 T cell (1.2%)
Distinct proximal tubule 2 (0.7%)
Thick ascending limb of Loop of Henle (0.6%)
Ctrl_2
Ctrl_1
DKD_4-4
DKD_3-3
DKD_1-2
DKD_4-2
DKD_3-2
DKD_2-2
DKD_4-5
DKD_1-3
DKD_2-3
DKD_4-1
DKD_4-3
DKD_1-1
DKD_2-1
DKD_3-1
Ctrl_3
Design effect D (log scale)
b
Abundant types are most penalized
DKD (GSE325587)
Lupus nephritis
GSE211785
Design effect D
D = 1: cells independent
Cell-type proportion in tissue
c
Lineages (colours as in b); bars = medians
Other epithelial
Immune
Stromal
Endothelial
Glomerular epithelial
Design effect D (lineage)