

Differential Equations - Notes

Arun Felix

June 15, 2025

Contents

I	Basic Concepts	3
1	Definitions	3
1.1	What is a differential equation?	3
1.2	What is Order?	3
1.3	Ordinary vs. Partial Differential Equation	3
1.4	Linear Differential Equation	3
1.5	Initial Conditions	3
1.6	Implicit vs. Explicit Solutions	4
II	First Order Differential Equations	4
2	Linear Differential Equations	4
2.1	Introduction	4
2.2	Example 1	4
3	Separable Equations	5
3.1	Introduction	5
3.2	Example	6
4	Exact Equations	6
4.1	Introduction	6
4.2	What is a exact equation	7
4.3	Example	7
5	Bernoulli Differential Equation	8
5.1	Introduction	8
5.2	Example	8

6	Substitutions	9
6.1	Introduction	9
6.1.1	Homogenous equations	9
6.1.2	Other equation	10
7	Intervals of Validity	10
7.1	Theorem one	10
7.1.1	Example	10
7.2	Theorem 2	11
7.2.1	Example	11
III	Second Order Differential Equations	12
8	Basic Concepts	12
8.1	Second order linear differential equation	12
8.2	Homogenous vs. nonhomogenous differential equations	12
8.2.1	Example	12
8.3	Principle of Superposition	13
8.3.1	Example	13
8.4	Characteristic equations	13
8.4.1	Example	13
8.5	Real & distinct roots	14
8.5.1	Example	14
8.6	Complex Roots	15
8.6.1	Introduction	15
8.6.2	Example	16
8.7	Repeated Roots	16
8.7.1	Introduction	16
8.7.2	Example	17

Part I

Basic Concepts

1 Definitions

1.1 What is a differential equation?

- A differential equation is an equation that contains derivatives, either ordinary or partial derivatives
- A popular example is newtons second law of motion, $F = ma$
- This is a differential equation since $a = \frac{dv}{dt}$
- Given this, we can actually write Newton's second law as a differential equation: $m \frac{dv}{dt} = F(t, v)$

1.2 What is Order?

- The order is the largest derivative present in the differential equation.
- For example, $\frac{\partial^3 u}{\partial x^2 \partial t} = 1 + \frac{\partial u}{\partial y}$ is a third order differential equation since it's greatest derivative is 3.

1.3 Ordinary vs. Partial Differential Equation

- An ordinary differential equation only has original differential equations, ode.
- A partial differential equation will have partial derivatives in it.

1.4 Linear Differential Equation

A linear differential equation is a differential equation following the form:

$$a_n(t)y^{(n)}(t) + a_{n-1}(t)y^{(n-1)} + \dots + a_1(t)y'(t) + a_0(t)y(t) = g(t)$$

1.5 Initial Conditions

An initial condition allows us to determine which solution we're after. For example, $y(t_0) = y_0$ or $y^{(k)}(t_0) = y_k$.

1.6 Implicit vs. Explicit Solutions

An implicit solution is a solution given in the form of $y = y(t)$

An explicit solution is a solution that isn't implicit.

Part II

First Order Differential Equations

2 Linear Differential Equations

2.1 Introduction

A linear first order differential equation is an equation in the form:

$$\frac{dy}{dt} + p(t)y = g(t)$$

- Here, we can assume $p(t)$ and $g(t)$ are continuous.

Now, let's assume the existence of a variable, $\mu(t)$, where we multiple all factors by it

$$\mu(t)\frac{dy}{dt} + \mu(t)p(t)y = \mu(t)g(t)$$

To solve problems using this:

- Put the differential equation in the correct form above.
- Find the integrating factor, $\mu(t)$, where $\mu(t) = e^{\int p(t)dt}$
- Multiply everything in the differential equation with $\mu(t)$, and verify the leftside becomes the product rule, $(\mu(t)y(t))'$.
- Integrate both sides.
- Solve for $y(t)$.

2.2 Example 1

Let's try to find the solution to the following differential equation:

$$\frac{dv}{dt} = 9.8 - 0.196v$$

We can see that we need to put the equation into linear form.

$$\frac{dv}{dt} + 0.196v = 9.8$$

Let's solve for $\mu(t)$

$$\mu(t) = e^{\int 0.196 dt} = e^{0.196t}$$

Next, we can plug in $\mu(t)$ in all variables

$$e^{0.196t} \frac{dv}{dt} + e^{0.196t} 0.196v = 9.8e^{0.196t}$$

We can rearrange some numbers:

$$e^{0.196t} \frac{dv}{dt} + 0.196e^{0.196t}v = 9.8e^{0.196t}$$

this resembles the product rule, hence we can say:

$$(e^{0.196t}v)' = 9.8e^{0.196t}$$

Now, we can integrate:

$$\int (e^{0.196t}v)' dt = \int 9.8e^{0.196t} dt$$

$$e^{0.196t}v = 9.8 \int e^{0.196t} dt$$

for the right equation, we can use u sub

$$e^u dt, \text{ where } u = 0.196t, du = 0.196dt, \text{ and } dt = \frac{du}{0.196}$$

We can plug this into our equation

$$\begin{aligned} e^{0.196t}v + k &= 9.8 \cdot \frac{1}{0.196} \int e^u du \\ e^{0.196t}v + k &= 50 \int e^u du \\ e^{0.196t}v + k &= 50e^{0.196t} + c \end{aligned}$$

Finally, we can solve for y, or v in this context:

$$v = \frac{50e^{0.196t} + c}{e^{0.196t}}$$

3 Separable Equations

3.1 Introduction

A separable differential equation is an equation where you can separate the variables:

$$N(y) \frac{dy}{dx} = M(x)$$

To put it simply, all the ys need to be on one side, xs on the other. To solve such a differential equation, you'd integrate both sides.

$$\int N(y) \frac{dy}{dx} dx = \int M(x) dx$$

We can use substitution to get this instead:

$$\int N(u) du = \int M(x) dx$$

3.2 Example

Solve the following differential equation and determine the interval of validity for the solution.

$$\frac{dy}{dx} = 6y^2x, \text{ and } y(1) = \frac{1}{25}$$

The first step is separating the Xs and Ys

$$\begin{aligned} dx \cdot \frac{dy}{dx} &= 6y^2x \cdot dx \\ \frac{dy}{y^2} &= \frac{6y^2x \cdot dx}{y^2} \\ \frac{1}{y^2} dy &= 6x dx \end{aligned}$$

From here, we can do integration:

$$\begin{aligned} \int \frac{1}{y^2} dy &= \int 6x dx \\ \int y^{-2} dy &= 3x^2 + C \\ -y^{-1} &= 3x^2 + C \\ -\frac{1}{y} &= 3x^2 + C \end{aligned}$$

now, we can solve for y:

$$y = \frac{1}{3x^2 + C}$$

Finally, we can solve for C to get our differential equation. This is trivial, as we have $y(1) = \frac{1}{25}$

$$\frac{1}{25} = \frac{1}{3+C}$$

We can easily figure out that $C = 22$.

Hence our final differential equation is:

$$y = \frac{1}{3x^2 + 22}$$

4 Exact Equations

4.1 Introduction

An exact equation follows the form: $M(x, y) + N(x, y) \frac{dy}{dx} = 0$

Let's create two variables:

- $\psi_x = M(x, y)$
- $\psi_y = N(x, y)$

Plugging it into our equation, we get:

$$\psi_x + \phi_y \frac{dy}{dx} = 0$$

We can then use the chain rule to find the derivative of the differential equation:

$$\frac{d}{dx}(\psi(x, y(x))) = 0$$

this gives us an implicit solution: $\psi(x, y) = c$.

4.2 What is a exact equation

An exact equation is an equation, given M_y and N_x , if their partial derivatives are equal to eachother.

4.3 Example

Solve the following IVP and find the interval of validity for the solution

$$2xy - 9x^2 + (2y + x^2 + 1)\frac{dy}{dx} = 0, \text{ where } y(0) = -3.$$

- We can set two variables.
- $M(x, y) = 2xy - 9x^2$
- $N(x, y) = 2y + x^2 + 1$

The first thing we can do is check if this is an exact equation:

$$\begin{aligned}\frac{\partial M}{\partial y} &= 2x \\ \frac{\partial N}{\partial x} &= 2x\end{aligned}$$

We can see the partial derivatives are identical, hence it's an exact equation.
Next, we need to integrate one of the equations:

$$\int M(x, y) = \int 2xy - 9x^2 dx = x^2y - 3x^3 + h(y)$$

Notice how, instead of a constant, we use $h(y)$.

We actually have $h'(y)$, it's $N = 2y + x^2 + 1$.

Now, we find $\frac{\partial M}{\partial y}$ the integral of M, giving us $x^2 + h'(y)$

We can set $\frac{\partial M}{\partial y} = M$, or $x^2 + h'(y) = 2y + x^2 + 1$.

$$\begin{aligned}x^2 + h'(y) - x^2 &= 2y + x^2 + 1 - x^2 \\ h'(y) &= 2y + 1\end{aligned}$$

We can integrate $h'(y)$ to get $h(y)$.

$$\int 2y + 1 dy = y^2 + y + k$$

We can plug this into our equation:

$$\psi(x, y) = x^2y - 3x^3 + y^2 + y + k$$

Now, we need to solve for k, or c

$$\begin{aligned}\psi(x, y) &= x^2y - 3x^3 + y^2 + y + k = c \\ \psi(x, y) &= x^2y - 3x^3 + y^2 + y + k - k = c - k \\ \psi(x, y) &= x^2y - 3x^3 + y^2 + y = c \\ \psi(x, y) &= x^2y - 3x^3 + y^2 + y = c \text{ Here, we use } y(0) = -3 \\ \psi(x, y) &= (0)^2(-3) - 3(0)^3 + (-3)^2 + (-3) = c \text{ Here, we use } y(0) = -3 \\ \psi(x, y) &= 9 - 3 = c, \text{ where } c = 6 \\ \text{hence our solution is } \psi(x, y) &= x^2y - 3x^3 + y^2 + y - 6 = 0\end{aligned}$$

5 Bernoulli Differential Equation

5.1 Introduction

Bernoulli Equations are equations that follow the form:

$$y' + p(x)y = q(x)y^n$$

Here, we assume $p(x)$ and $q(x)$ are continuous functions, and n is a number. if $n = 0$ or 1 , we'll have a linear equation, which is easy to solve.

if n isn't 0 or 1 , we need to divide both sides by y^n , substitute $v = y^{1-n}$. Once we do this, we'll get a linear equation we can solve.

5.2 Example

Solve the following IVP and find the interval of validity for the solution:

$$y' + \frac{4}{x}y = x^3y^2, \text{ where } y(2) = -1 \text{ and } x > 0$$

We can see our differential equation resembles the bernoulli equation, where:

- $p(x) = \frac{4}{x}$
- $q(x) = x^3$
- $n = 2$

now, we need to divide both sides by y^n

$$\begin{aligned}\frac{y' + \frac{4}{x}y}{y^2} &= \frac{x^3y^2}{y^2} \\ \frac{y' + \frac{4}{x}y}{y^2} &= \frac{x^3y^2}{y^2} \\ y^{-2}y' + \frac{4}{x}y^{-1} &= x^3\end{aligned}$$

From here, we can substitute, using $v = y^{1-n} = y^{1-2} = y^{-1}$ and $v' = -y^{-2}y'$, giving us:

$$-v' + \frac{4}{x}v = x^3$$

Because $\frac{dv}{dt}$ needs to be positive, we need to multiply both sides by -1 to get:

$$v' - \frac{4}{x}v = -x^3$$

Next, we need to multiply all variables by $\mu(x)$

$$\mu(x)v' - \mu(x)\frac{4}{x}v = -\mu(x)x^3$$

next, we need to find $\mu(x)$

$$\mu(x) = e^{\int -\frac{4}{x} = -4 \int \frac{1}{x} = -4 \ln|x| \rightarrow \ln|x|^{-4}} = x^{-4}$$

We can plug this in to get

$$x^{-4}v' - x^{-4}\frac{4}{x}v = -x^{-4}x^3 = -x^4v' - 4x^{-5}v = -x^{-1}$$

You can see that this resembles the product rule, hence we can simplify this equation to:

$$-x^{-4}x^3 = -x^4v' - 4x^{-5}v = (x^{-4}v)'$$

We can then integrate:

$$\begin{aligned} \int (x^{-4}v)' &= \int -x^{-1}dx \\ x^{-4}v &= -\ln|x| + c \\ v(x) &= cx^4 - x^4\ln x, \text{ or } v = cx^4 - x^4\ln x \end{aligned}$$

We can then solve for c by using the condition $y(2) = -1$. We can also plug y^{-1} back in

$$\begin{aligned} y^{-1} &= cx^4 - x^4\ln x \\ -1^{-1} &= c2^4 - 2^4\ln 2 \\ -1 + 16\ln 2 &= 16c - 16\ln 2 + 16\ln 2 \\ \frac{-1+16\ln 2}{16} &= \frac{16c}{16} \quad c = \frac{-1+16\ln 2}{16} \end{aligned}$$

We can solve for y, bla bla bla.

6 Substitutions

6.1 Introduction

In addition to Bernoulli's Differential equation, there's also another one known as homogenous differential equations.

6.1.1 Homogenous equations

Homogenous equations are equations that take the form:

$$y' = F\left(\frac{y}{x}\right)$$

If this is the case, we can use the following substitutions:

$$\begin{aligned} v &= \frac{y}{x}, \text{ which can be written as } y = xv \\ y' &= v + xv' \end{aligned}$$

using this will give us a separable differential equation, which we can easily solve.

Example Solve the following IVP and find the interval of validity for the solution:

$$xyy' + 4x^2 + y^2 = 0, \text{ where } y(2) = -7 \text{ and } x > 0$$

We can move $4x^2 + y^2$ to the other side of the equation, giving us:

$$xyy' = -4x^2 - y^2$$

we can then divide both sides by two:

$$\frac{y}{x}y' = -4 - \frac{y^2}{x^2} = -4 - \left(\frac{y}{x}\right)^2$$

Hence, we proved our equation here is homogenous. Next, we can apply the substitution here.

$$v(v + xv') = -4 - v^2$$

We can then separate variables, then use integration to solve our differential equation.

$$\int \frac{v}{2v^2} dv = \int -\frac{1}{x} dx$$

6.1.2 Other equation

If you have a differential equation with the form: $y' = G(ax + by)$, then you can apply the following substitution:

$$\begin{aligned} v &= ax + by \\ v' &= a + by' \end{aligned}$$

7 Intervals of Validity

7.1 Theorem one

When it comes to intervals of validity, there's an important idea:

$$y' + p(t)y = g(t), \text{ where } y(t_0) = y_0$$

If $p(t)$ and $g(t)$ are continuous functions on an open interval $\alpha < t < \beta$, containing an interval t_0 , then there's a unique solution to the IVP on the interval.

7.1.1 Example

Let's try to determine the intervals of validity for the following initial value problem.

$$(t^2 - 9)y' + 2y = \ln|20 - 4t|, \text{ where } y(4) = -3$$

Let's start off by dividing both sides by $(t^2 - 9)$, giving us:

$$y' + \frac{2}{t^2-9}y = \frac{\ln|20-4t|}{t^2-9}$$

In order to find non continuous intervals, we need to find points that aren't defined. In this example, dividing by zero, or if you get $\ln|0|$.

$$20 - 4t = 0 \rightarrow t = 5$$

$$t^2 - 9 = 0 \rightarrow (t - 3)(t + 3) = 0 \rightarrow t = \pm 3$$

hence, our Intervals of Validity include:

$$-\infty < t < -3, -3 < t < 3, 3 < t < 5, 5 < t < \infty$$

7.2 Theorem 2

Let's consider the following IVP:

$$y' = f(t, y), \text{ where } y(t_0) = y_0$$

if $f(t, y)$ and $\frac{\partial f}{\partial y}$ are continuous functions in some rectangle $\alpha < t < \beta$, and $\gamma < y < \delta$, containing the point (t_0, y_0) , then there's a unique solution to the IVP in some interval $t_0 - h < t < t_0 + h$

7.2.1 Example

Determine the interval of validity for the initial value problem below and give its dependence on the value of y_0

$$y' = y^2, \text{ where } y(0) = y_0$$

Here, we can define the variables:

- $f(t, y) = y^2$
- $t_0 = 0$
- y_0 is the initial value.

We first need to check for continuity for $f(t, y)$ and $\frac{\partial f}{\partial y}$

- $f(t, y) = y^2$ is continuous
- $\frac{\partial f}{\partial y} = 2y$, which is also continuous.

Our equation also contains the point (t_0, y_0) , so we're good. we can now now, we can work on this:

$$\begin{aligned} \frac{dy}{dt} &= y^2 \\ dy &= y^2 dt \\ \frac{1}{y^2} dy &= dt \\ \int \frac{1}{y^2} dy &= \int dt \\ -\frac{1}{y} &= t + c \end{aligned}$$

We're given $y(0) = y_0$, so we can figure out c with it.

$$-\frac{1}{y_0} = c$$

therefore, our solution is $-\frac{1}{y} = t - \frac{1}{y_0}$.

We can rearrange the variables to get $y = \frac{y_0}{1 - y_0 t}$

We can then find out that y is undefined when $t = \frac{1}{y_0}$.

Part III

Second Order Differential Equations

8 Basic Concepts

8.1 Second order linear differential equation

The most general linear order differential equation follows the form:

$$P(t)y'' + q(t)y' + r(t)y = g(t)$$

If we were to assume constant coefficient, we can use the following:

$$ay'' + by' + cy = g(t)$$

8.2 Homogenous vs. nonhomogenous differential equations

Given the form, $ay'' + by' + cy = g(t)$:

- we say a differential equation is homogenous if $g(t) = 0$
- Otherwise, if $g(t) \neq 0$, it's nonhomogenous.

8.2.1 Example

Determine some solutions to $y'' - 9y = 0$

We can bring $9y$ to the other side:

$$y'' = 9y$$

- This tells us y'' needs to be the original function times 9.
- Right off the bat, we can think of using e .
- more specifically, $y = e^{\pm 3t}$

We can test this:

$$\begin{aligned}y &= e^{3t} \\ y' &= 3e^{3t} \\ y'' &= 9e^{3t}\end{aligned}$$

since $y = e^{3t}$, we can say $y'' = 9y$

8.3 Principle of Superposition

If $y_1(t)$ and $y_2(t)$ are solutions to a linear, homogenous differential equation, we can also say $y(t) = c_1y_1(t) + c_2y_2(t)$ is a solution to our differential equation. **This is true regardless of order**, applies to any homogenous linear differential equation.

Another important question to ask is, how do we find the constants, c_1 and c_2 ? We simply just specify values of the solutions at two distinct points.

8.3.1 Example

Solve the following IVP

$$y'' - 9y = 0, \text{ where } y(0) = 2 \text{ and } y'(0) = -1$$

- We solved y in the first part, it's $e^{\pm 3t}$
- Let's plug it in: $y(t) = c_1e^{3t} + c_2e^{-3t}$
- next, we can figure out the derivative of $y(t)$
 - $y'(t) = 3c_1e^{3t} - 3c_2e^{-3t}$
- We can plug in our initial conditions:
 - $2 = c_1 + c_2$, which is $y(0)$
 - $-1 = 3c_1 - 3c_2$, which is $y'(0)$
- We can use gaussian elimination to find the $c_1 = \frac{5}{6}$ and $c_2 = \frac{7}{6}$
- Plugging everything in, we can find that our differential equation is $y(t) = \frac{5}{6}e^{3t} + \frac{7}{6}e^{-3t}$

8.4 Characteristic equations

- We can notice that the equations above resembled $y(t) = e^{rt}$.
- How can we solve without guessing?

8.4.1 Example

Let's do the $y'' - 9y = 0$

Let's first calculate y , which we'll say is $y(t) = e^{rt}$

- $y(t) = e^{rt}$
- $y'(t) = re^{rt}$
- $y''(t) = r^2e^{rt}$

We can plug these into our equation:

$$r^2 e^{rt} - 9e^{rt} = 0$$

from here, we can bring the e^{rt} out the equation:

$$e^{rt}(r^2 - 9) = 0$$

- $r^2 - 9$ is known as a **characteristic equation**.
- Based on this, we can deduce that, in order for the equation to be equal to zero, $r^2 - 9 = 0$. We can use easy algebra to find that $r = 3$, and $r = -3$.

8.5 Real & distinct roots

Let's think about our characteristic equation:

$$ar^2 + br + c = 0$$

and the two solutions: $y_1(t) = e^{r_1 t}$ and $y_2(t) = e^{r_2 t}$

- We say that the roots, r_1 and r_2 are real and distinct, if the two are real numbers and $r_1 \neq r_2$.
- If this is true, then the two solutions are **nice enough** to form the solution:
 $y(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}$

8.5.1 Example

Solve the following IVP:

$$y'' + 11y' + 24y = 0, \text{ given } y(0) = 0, \text{ and } y'(0) = -7.$$

Let's use $y = e^{rt}$

$$\begin{aligned} y &= e^{rt} \\ y' &= r e^{rt} \\ y'' &= r^2 e^{rt} \end{aligned}$$

Let's plug these in our equation:

$$\begin{aligned} r^2 e^{rt} + 11r e^{rt} + 24e^{rt} &= 0 \\ e^{rt}(r^2 + 11r + 24) &= 0 \\ r^2 + 11r + 24 &= 0 \\ (r + 8)(r + 3) &= 0 \end{aligned}$$

therefore, $r = -8$ and $r = -3$, which are real numbers and distinct.

we can now come up with the following equations:

$$\begin{aligned} y(t) &= c_1 e^{-3t} + c_2 e^{-8t} \\ y'(t) &= -3c_1 e^{-3t} - 8c_2 e^{-8t} \end{aligned}$$

We can use gaussian elimination to solve for the constants. Our final answer is $y(t) = -\frac{7}{5}e^{-3t} + \frac{7}{5}e^{-8t}$

8.6 Complex Roots

8.6.1 Introduction

Let's again, look at solutions to the differential equations following the form:

$$\begin{aligned} ay'' + by' + cy &= 0 \\ ar^2 + br + c &= 0 \end{aligned}$$

We call roots **complex roots**, if they have the form $r_{1,2} = \lambda \pm \mu i$.

Assuming we're using the two differential equations, $y(t) = e^{rt}$, we say that our solutions to the differential equation is:

$$\begin{aligned} y_1(t) &= e^{(\lambda + \mu i)t} \\ y_2(t) &= e^{(\lambda - \mu i)t} \end{aligned}$$

Having complex roots is great, but it's ideal for our solution to only have real numbers. we use **Euler's formula** for this:

$$e^{i\theta} = \cos\theta + i\sin\theta$$

We can use another variant:

$$e^{-i\theta} = \cos(-\theta) + i\sin(-\theta) = \cos\theta - i\sin\theta$$

We can now split our exponent into two groups:

$$y_1(t) = e^{\lambda t + \mu i t} = e^{\lambda t} e^{\mu i t} \quad y_2(t) = e^{\lambda t - \mu i t} = e^{\lambda t} e^{-\mu i t}$$

We have our equation from above, which we can use:

$$\begin{aligned} y_1(t) &= e^{\lambda t} e^{\mu i t} = e^{\lambda t} (\cos(\mu t) + i\sin(\mu t)) \\ y_2(t) &= e^{\lambda t} e^{-\mu i t} = e^{\lambda t} (\cos(\mu t) - i\sin(\mu t)) \end{aligned}$$

We know that, given two solutions are "nice enough", any solution can be written as:

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

We can ignore constants for now, so

$$\begin{aligned} y(t) &= y_1(t) + y_2(t) \\ y_1(t) + y_2(t) &= e^{\lambda t} \cos(\mu t) + e^{\lambda t} i\sin(\mu t) + e^{\lambda t} \cos(\mu t) - e^{\lambda t} i\sin(\mu t) \\ y_1(t) + y_2(t) &= e^{\lambda t} \cos(\mu t) + e^{\lambda t} \cos(\mu t) \\ y_1(t) + y_2(t) &= 2(e^{\lambda t} \cos(\mu t)) \end{aligned}$$

Now, we can assume the constants, c_1 and c_2 , are $\frac{1}{2}$, which means

$$\frac{1}{2}y_1(t) + \frac{1}{2}y_2(t) = (e^{\lambda t} \cos(\mu t))$$

main idea

We can say, if the roots of the characteristic equation happens to be $r_{1,2} = \lambda \pm \mu i$, the solution to the differential equation is:

$$y(t) = c_1 e^{\lambda t} \cos(\mu t) + c_2 e^{\lambda t} \sin(\mu t)$$

8.6.2 Example

Solve the following IVP:

$$y'' - 4y' + 9y = 0, \text{ where } y(0) = 0 \text{ and } y'(0) = -8$$

our characteristic equation is $r^2 - 4r + 9 = 0$. There's no easy way to solve this, so we need to use the quadratic formula.

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Where $a = 1$, $b = -4$ and $c = 9$, which we can plug in:

$$r = \frac{4 \pm \sqrt{-4^2 - 4(1)(9)}}{2(1)}, \text{ which simplifies to:}$$
$$r = 2 \pm \sqrt{5}i$$

This resembles our complex root equation, $\lambda \pm \sqrt{\mu}i$, where $\lambda = 2$ and $\mu = 5$. We can plug these variables into our equation

$$y(t) = c_1 e^{2t} \cos(\sqrt{5}t) + c_2 e^{2t} \sin(\sqrt{5}t)$$

we can now think about our first initial condition, $y(0) = 0$

$$0 = c_1 e^{2(0)} \cos(\sqrt{5}(0)) + c_2 e^{2(0)} \sin(\sqrt{5}(0))$$
$$0 = c_1 \times 1$$
$$c_1 = 0$$

this cancels out the first part, giving us:

$$y(t) = c_2 e^{2t} \sin(\sqrt{5}t)$$

now, we need to find the derivative, which is done using the product rule:

$$y'(t) = c_2 e^{2t} \sqrt{5} \cos(\sqrt{5}t) + 1 c_2 e^{2t} \sin(\sqrt{5}t)$$

Here, we can use the initial condition, $y'(0) = -8$

$$-8 = c_2 e^{2(0)} \sqrt{5} \cos(\sqrt{5}(0)) + 1 c_2 e^{2(0)} \sin(\sqrt{5}(0))$$
$$-8 = c_2 \sqrt{5}$$
$$c_2 = -\frac{8}{\sqrt{5}}$$

Therefore, our solution is:

$$y(t) = -\frac{8}{\sqrt{5}} e^{2t} \sin(\sqrt{5}t)$$

8.7 Repeated Roots

8.7.1 Introduction

Here, if the roots are the same, or $r_1 = r_2 = r$, we can say the general solution will be $y(t) = c_1 e^{rt} + c_2 t e^{rt}$

8.7.2 Example

Solve the following IVP:

$$y'' - 4y' + 4y = 0, \text{ where } y(0) = 12 \text{ and } y'(0) = -3$$

We can get the characteristic equation

$$\begin{aligned} r^2 - 4r + 4 &= 0 \\ (r - 2)(r - 2) &= 0 \\ r_1 = r_2 &= 2 \end{aligned}$$

We can see that this falls under the Repeated root, therefore, we can plug r into the general solution:

$$y(t) = c_1 e^{2t} + c_2 t e^{2t}$$

Now, we can solve for the constants, first using $y(0) = 12$:

$$\begin{aligned} 12 &= c_1 e^{2(0)} + c_2(0) e^{2(0)} \\ 12 &= c_1 \end{aligned}$$

We can now find the derivative of $y(t)$, and find c_2 with $y'(0) = -3$

$$\begin{aligned} y(t) &= 12e^{2t} + c_2 t e^{2t} \\ y'(t) &= 24e^{2t} + c_2 e^{2t} + 2c_2 t e^{2t} \text{ (used product rule)} \\ -3 &= 24e^{2(0)} + c_2 e^{2(0)} + 2c_2(0) e^{2(0)} \\ -3 &= 24 + c_2 \\ -27 &= c_2 \end{aligned}$$

Therefore, our solution is $y(t) = 12e^{2t} - 27te^{2t}$