

Weighted Kolmogorov-Smirnov Tests in One- and Two-Sample Problems

Veronika Gontscharuk and Helmut Finner

Institute for Biometrics and Epidemiology,
German Diabetes Center (DDZ),
Leibniz Center for Diabetes Research at Heinrich Heine University Düsseldorf

Adaptive Designs and Multiple Testing Procedures
Workshop 2016
April 28 – 29, 2016,
University of Padua - Padua, Italy

One-Sample Weighted Kolmogorov-Smirnov (W-KS) Tests

Let $X_1, \dots, X_m \sim F$ be iid. W.l.o.g. we test

$$H_0 : F(t) = t, \quad t \in [0, 1].$$

For $\nu \in [0, 1]$ one-sample W-KS statistics are defined by

$$KS_m^\nu = \sup_{t \in (0,1)} \sqrt{m} \frac{|\hat{F}_m(t) - t|}{(t(1-t))^\nu}.$$

Reject H_0 if $KS_m^\nu > b_m$, where b_m denotes a suitable critical value.


Let b_m be such that $\mathbb{P}(KS_m^\nu > b_m | H_0) = \alpha$.

For $\nu = 0$ we get the original KS test and for $\nu = 1/2$ we get the supremum version of the Anderson-Darling test, also known as higher criticism (HC) test.

One-Sample W-KS Tests:

Graphical Representation


Ecdf of m iid $U(0, 1)$ -random variables and rejection curves
related to level α W-KS tests with $m = 100$ and $\alpha = 0.05$



One-Sample W-KS Tests:

Graphical Representation


Ecdf of m iid $U(0, 1)$ -random variables and rejection curves
related to level α W-KS tests with $m = 100$ and $\alpha = 0.05$



One-Sample W-KS Tests:

Graphical Representation


Ecdf of m iid $U(0, 1)$ -random variables and rejection curves related to level α W-KS tests with $m = 100$ and $\alpha = 0.05$




One-Sample W-KS Tests:

Graphical Representation

Ecdf of m iid $U(0, 1)$ -random variables and rejection curves related to level α W-KS tests with $m = 100$ and $\alpha = 0.05$



A New Point of View: Local Levels




Local levels α_i , $i = 1, \dots, m$, are defined as probabilities that $X_{i:m}$ leads to the rejection of true H_0 , i.e.,

$$\alpha_i^{low} = \mathbb{P}(X_{i:m} < c_i^\nu | H_0), \quad \alpha_i^{up} = \mathbb{P}(X_{i:m} > d_i^\nu | H_0),$$

$$\alpha_i = \alpha_i^{low} + \alpha_i^{up}.$$

A New Point of View: Local Levels




Local levels α_i , $i = 1, \dots, m$, are defined as probabilities that $X_{i:m}$ leads to the rejection of true H_0 , i.e.,

$$\alpha_i^{low} = \mathbb{P}(X_{i:m} < c_i^\nu | H_0), \quad \alpha_i^{up} = \mathbb{P}(X_{i:m} > d_i^\nu | H_0),$$

$$\alpha_i = \alpha_i^{low} + \alpha_i^{up}.$$

A New Point of View: Local Levels




Local levels $\alpha_i, i = 1, \dots, m$, are defined as probabilities that $X_{i:m}$ leads to the rejection of true H_0 , i.e.,

$$\alpha_i^{low} = \mathbb{P}(X_{i:m} < c_i^v | H_0), \quad \alpha_i^{up} = \mathbb{P}(X_{i:m} > d_i^v | H_0),$$

$$\alpha_i = \alpha_i^{low} + \alpha_i^{up}.$$

A New Point of View: Local Levels




Local levels $\alpha_i, i = 1, \dots, m$, are defined as probabilities that $X_{i:m}$ leads to the rejection of true H_0 , i.e.,

$$\alpha_i^{low} = \mathbb{P}(X_{i:m} < c_i^v | H_0), \quad \alpha_i^{up} = \mathbb{P}(X_{i:m} > d_i^v | H_0),$$

$$\alpha_i = \alpha_i^{low} + \alpha_i^{up}.$$

One-Sample W-KS Local Levels: Examples

Local levels of level α W-KS tests with $\alpha = 0.05$, $m = 100$ and
 $\nu = 0(0.125)0.875$




Asymptotics of One-Sample W-KS Local Levels

The asymptotics of the W-KS statistics is examined in many papers and can be applied to derive the asymptotics of W-KS local levels.

- For $\nu \in [0, 1/2)$ we get $\lim_{m \rightarrow \infty} \alpha_i > 0$ for central i -values, i.e., $\lim_{m \rightarrow \infty} i/m \in (0, 1)$, and $\lim_{m \rightarrow \infty} \alpha_i = 0$ else. The asymptotic sensitivity range is the central range.
- For $\nu = 1/2$ (HC) we get that $\lim_{m \rightarrow \infty} \alpha_i = 0$ for all i and the most HC local levels are asymptotically equal, i.e., $\lim_{m \rightarrow \infty} \alpha_i/\alpha_j = 1$, cf. **Gontscharuk, Landwehr & Finner [2015,2016]**. The asymptotic sensitivity range consists of intermediates.
- For $\nu \in (1/2, 1]$ we get $\lim_{m \rightarrow \infty} \alpha_i > 0$ and $\lim_{m \rightarrow \infty} \alpha_{m-i} > 0$ only for fixed $i \in \mathbb{N}$. The asymptotic sensitivity range coincides with the extreme range.

Asymptotic Local Levels for $\nu < 1/2$

Asymptotic local levels $\lim_{m \rightarrow \infty} \alpha_i$ with $\lim_{m \rightarrow \infty} i/m = \eta$ for level α
 W-KS tests with $\alpha = 0.05$, $\nu = 0(0.05)0.45$ (from top to bottom)



$$\lim_{m \rightarrow \infty} \alpha_i = 2\Phi\left(-\frac{b_{\nu,\alpha}}{(\eta(1-\eta))^{1/2-\nu}}\right) \text{ for } \eta \in (0, 1)$$

$$\lim_{m \rightarrow \infty} \alpha_i = 0 \text{ for } \eta = 0, 1$$

Two-Sample W-KS Tests

Let $X_1, \dots, X_m \sim F$ be iid and let $Y_1, \dots, Y_n \sim G$ be iid with continuous F, G . We test

$$H_0 : F(t) = G(t), \quad t \in \mathbb{R}.$$

For $\nu \in [0, 1]$, two-sample W-KS statistics are defined by

$$KS_{m,n}^\nu = \sup_{t \in \mathbb{R}} \sqrt{\frac{mn}{m+n}} \frac{|\hat{G}_n(t) - \hat{F}_m(t)|}{(\hat{H}_{m+n}(t)(1 - \hat{H}_{m+n}(t)))^\nu}.$$

Reject H_0 if $KS_{m,n}^\nu > b_{m,n}$, where $b_{m,n}$ denotes a suitable critical value.

Two-Sample W-KS Local Levels

Let $t_1 \leq \dots \leq t_{m+n}$ denote the ordered combined sample and let $V_{m,s} = m\hat{F}_m(t_s)$. For suitable $c_s^\nu, d_s^\nu, s = 1, \dots, m+n$, we get

$$KS_{m,n}^\nu \leq b_{m,n} \text{ iff } c_s^\nu \leq V_{m,s} \leq d_s^\nu \text{ for all } s = 1, \dots, m+n.$$

Assumptions:

- Critical values $c_s^\nu, d_s^\nu, s = 1, \dots, m+n$, are proper;
- $\mathbb{P}(KS_{m,n}^\nu > b_{m,n} | H_0) \leq \alpha$;
- $\mathbb{P}(KS_{m,n}^\nu > b_{m,n} | H_0)$ is as large as possible.


Similar as in the one-sample case, W-KS **local levels** are defined by

$$\alpha_s^{low} = \mathbb{P}(V_{m,s} < c_s^\nu | H_0), \quad \alpha_s^{up} = \mathbb{P}(V_{m,s} > d_s^\nu | H_0),$$

$$\alpha_s = \alpha_s^{low} + \alpha_s^{up}.$$

Two-Sample W-KS Local Levels: $m = n$

Local levels of level α W-KS tests with $\alpha = 0.05$, $m = n = 50$ and
 $\nu = 0(0.125)0.875$



Asymptotics of Two-Sample W-KS Local Levels

In contrast to the one-sample case, it seems that the asymptotics of two-sided W-KS test statistics for $\nu \in (0, 1]$ has not been considered in the literature.

- For $\nu \in [0, 1/2)$ the two-sample W-KS asymptotics **coincides** with the one-sample case.
- For $\nu = 1/2$ (HC) the two-sample W-KS asymptotics **coincides** with the related one-sample HC asymptotics with **sample size** $\min(m, n)$.
- For $\nu \in (1/2, 1]$ the two-sample W-KS asymptotics **differs** from the one-sample counterpart if $\lim_{m,n \rightarrow \infty} m/n = \zeta \in (0, \infty)$.

Key tools: Weighted Brownian bridges, weighted standardized Poisson and Binomial processes.

Asymptotics for $\nu < 1/2$

Let \mathbb{B} denote a standard Brownian bridge on $[0, 1]$ and let

$$\mathbb{B}^\nu(t) = \frac{\mathbb{B}(t)}{(t(1-t))^\nu}, \quad t \in (0, 1), \quad \nu \in [0, 1/2].$$

Theorem 1: For $\nu \in [0, 1/2)$ it holds

$$\lim_{m,n \rightarrow \infty} KS_{m,n}^\nu \stackrel{\mathcal{D}}{=} \sup_{t \in (0,1)} |\mathbb{B}^\nu(t)|.$$

Moreover, let $b_{\nu,\alpha}$ be the asymptotic critical value, i.e.,

$\mathbb{P}(\sup_{t \in (0,1)} |\mathbb{B}^\nu(t)| > b_{\nu,\alpha}) = \alpha$, then

$$\lim_{m,n \rightarrow \infty} \alpha_s = 2\Phi\left(-\frac{b_{\nu,\alpha}}{(\eta(1-\eta))^{1/2-\nu}}\right)$$

for s such that $\lim_{m,n \rightarrow \infty} s/(m+n) = \eta \in (0, 1)$.

Asymptotic and Finite Local Levels

Illustration for $\nu = 0$ and $\nu = 0.25$

The W-KS asymptotics seems to be faster for smaller ν -values.

Asymptotics for $\nu = 1/2$

Theorem 2: Let $m = \min(m, n)$ and let $b_{m,\alpha}$ be such that

$$\lim_{m,n \rightarrow \infty} \mathbb{P}\left(\sup_{t \in (1/m, 1-1/m)} |\mathbb{B}^{1/2}(t)| > b_{m,\alpha}\right) = \alpha.$$

Then

$$\lim_{m,n \rightarrow \infty} \mathbb{P}(KS_{m,n}^{1/2} > b_{m,\alpha}) = \alpha.$$

Moreover, almost all HC local levels are asymptotically equal to

$$\alpha_m^* = \frac{-\log(1 - \alpha)}{2 \log(m) \log(\log(m))}$$

in the sense $\lim_{m,n \rightarrow \infty} \alpha_s / \alpha_m^* = 1$.

Asymptotic and Finite HC Local Levels

Illustration for $\nu = 1/2$

The two-sample HC asymptotics seems to be much better than
the related one-sample asymptotics.



Comparison of HC and Minimum p -Value Tests

The HC and minP tests coincide asymptotically under H_0 and differ not much even for a finite setting.

Asymptotics for $\nu > 1/2$

Let $Z_i, i \in \mathbb{N}$, be iid Bernoulli distributed with parameter $p \in (0, 1)$ and let $Y_s = \sum_{i=1}^s Z_i, s \in \mathbb{N}$. Let \tilde{Y}_s be an independent copy of Y_s .

Theorem 3: Let $\lim_{m,n \rightarrow \infty} m/(m+n) = p \in (0, 1)$, then

$$\lim_{m,n \rightarrow \infty} \left(\frac{m+n}{mn} \right)^{\nu-1/2} KS_{m,n}^{\nu} \stackrel{\mathcal{D}}{=} Q_{\nu,p},$$

where

$$Q_{\nu,p} = \frac{1}{(p(1-p))^{\nu}} \max \left(\sup_{s \in \mathbb{N}} \frac{|sp - Y_s|}{s^{\nu}}, \sup_{s \in \mathbb{N}} \frac{|sp - \tilde{Y}_s|}{s^{\nu}} \right).$$

The asymptotic W-KS distribution is typically discrete for $\nu > 1/2$.

Asymptotic and Finite Local Levels

Illustration for $\nu = 0.75$

The W-KS asymptotics is fast at least for larger ν -values.

Summary

- Local levels are a measure for local sensitivity of goodness-of-fit tests.
- We provided asymptotic local levels of one- and two-sample W-KS tests.
- The one-sample HC asymptotics is extremely slow, while the asymptotics is much better for two-sample HC tests.
- HC and minimum p -value tests coincide asymptotically under H_0 .
- In the two-sample case, HC and minimum p -value tests differ not much even for finite samples.

References

- **Berk, R. and Jones, D. (1978).** Relatively optimal combinations of test statistics. *Scand. J. Stat.*, **5**, 158–162.
- **Berk, R. and Jones, D. (1979).** Goodness-of-fit test statistics that dominate the Kolmogorov Statistics. *Z. Wahrscheinlichkeit.*, **47**, 47–59.
- **Gontscharuk, V., Landwehr, S. and Finner, H. (2016).** Goodness of fit tests in terms of local levels with special emphasis on higher criticism tests. *Bernoulli*, accepted for publication.
- **Gontscharuk, V., Landwehr, S. and Finner, H. (2015).** The intermediates take it all: asymptotics of higher criticism statistics and a powerful alternative based on equal local levels. *Biometrical J.*, **57**, 159–180.
- **Gontscharuk, V. and Finner, H. (2016).** Asymptotics of goodness-of-fit tests based on minimum p -value statistics. *Commun. Stat. A-Theor.*, accepted for publication.
- **Jager, L. and Wellner, J. A. (2004).** On the "Poisson boundaries" of the family of weighted Kolmogorov statistics. In: *A Festschrift for Herman Rubin, Institute of Mathematical Statistics*, 319–331.
- **Mason, D. M. (1983).** The asymptotic distribution of weighted empirical distribution functions. *Stoch. Proc. Appl.*, **15**, 99–109.
- ... thousands of papers on Kolmogorov-Smirnov tests