

# Analysis, sample size calculation and recalculation in designs with multiple nested subgroups

Marius Placzek, Tim Friede

Department of Medical Statistics, University Medical Center Göttingen, Germany

BMBF project (BundesMinisterium für Bildung und Forschung) „**BIOSTATISTISCHE METHODEN ZUR  
EFFIZIENTEN EVALUATION VON INDIVIDUALISIERTEN THERAPIEN (BIMIT)**“.

# Multiple Nested Subgroups

Consider the following design


▷ Full population with k nested subgroups  $F = S_0 \supset S_1 \cdots \supset S_k$

▷ Prevalences  $\tau_1, \dots, \tau_k$   
and treatment effects  $\Delta_0, \dots, \Delta_k$

▷ Assume

$X_{ij1} \sim \mathcal{N}(0, \sigma_{S_i}^2)$ ,  $i = 0, \dots, k$ ,  $j = 1, \dots, n^{S_i}$  (control group)

$X_{ij2} \sim \mathcal{N}(\Delta'_i, \sigma_{S_i}^2)$ ,  $i = 0, \dots, k$ ,  $j = 1, \dots, n^{S_i}$  (treatment group)



# Analysis (fixed design)

## Hypotheses and test statistics

- ▶ individual hypotheses  $H_0^{\{F\}} : \Delta_0 = 0$  (no effect in full population)  
 $H_0^{\{S_i\}} : \Delta_i = 0, i = 1, \dots, k$  (no effect in subpopulation i)
- ▶ intersection hypotheses  $H_0^{\cap_{i \in I} S_i} : \Delta_i = 0 \forall i \in I \subseteq \{0, \dots, k\}$
- ▶ standardized test statistics

$$Z^{\{F\}} = \sqrt{\frac{n}{2}} \frac{\bar{X}_F^T - \bar{X}_F^C}{\hat{\sigma}_F}$$

$$Z^{\{S_i\}} = \sqrt{\frac{n\hat{\tau}_i}{2}} \frac{\bar{X}_{S_i}^T - \bar{X}_{S_i}^C}{\hat{\sigma}_{S_i}}, i = 1, \dots, k$$

with

$$\bar{X}_{S_i}^r = \frac{1}{\sum_{j \geq i} n^{S_j}} \sum_{j \geq i} \sum_{k=1}^{n^{S_j}} X_{jk}^r, r \in \{T, C\}$$

# Analysis (fixed design)

- ▶ Test intersection hypotheses using the joint distribution of the standardized test statistics
  - Spiessens and Debois, 2010: normal distributed standardized test statistics with known variances
- ▶ Joint distribution? What do we know about the variances?
- ▶ **Case 1: known variances** (widely used in publications)
  - ▶ Replace  $\hat{\sigma}_{S_i}$  with  $\sigma_{S_i}$  in the standardized test statistics
  - ▶ Then we know that under the intersection hypothesis, e.g. for only one subgroup,  $H_0^{\{F,S\}}$ :

$$\begin{pmatrix} Z^{\{F\}} \\ Z^{\{S\}} \end{pmatrix} \sim MN \left( \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \sqrt{\tau} \\ \sqrt{\tau} & 1 \end{pmatrix} \right)$$

# Analysis (fixed design)

## ▷ Case 2: Unknown but same variances across subpopulations

$$\sigma = \sigma_F = \sigma_{S_1} = \cdots = \sigma_{S_k}$$

- ▷ Then we have to estimate only one variance  $\hat{\sigma}$  and replace  $\hat{\sigma}_{S_i} = \hat{\sigma}$  in the standardized test statistics
- ▷ One can show that the joint distribution is a multivariate t-distribution
  - ▷ degrees of freedom depending on the number of subjects used to estimate the variance
- ▷ e.g. under the global intersection hypothesis  $H_0^{\{\cap_{i=0}^k S_i\}}$

$$\mathbf{Z} = \begin{pmatrix} Z^{\{F\}} \\ Z^{\{S_1\}} \\ \vdots \\ Z^{\{S_k\}} \end{pmatrix} = \begin{pmatrix} \sqrt{\frac{n}{2}} \frac{\hat{\Delta}_F}{\hat{\sigma}} \\ \vdots \\ \sqrt{\frac{\tau_k n}{2}} \frac{\hat{\Delta}_{S_k}}{\hat{\sigma}} \end{pmatrix} \sim MT_{2(n-k-1)}(\mathbf{0}, \mathbf{\Sigma})$$

# Analysis (fixed design)

$$\Sigma = \begin{pmatrix} 1 & \frac{\sqrt{\tau_1}\sigma_{S_1}}{\sigma_F} & \frac{\sqrt{\tau_2}\sigma_{S_2}}{\sigma_F} & \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sigma_F} & \cdots & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sigma_F} \\ \frac{\sqrt{\tau_1}\sigma_{S_1}}{\sigma_F} & 1 & \frac{\sqrt{\tau_2}\sigma_{S_2}}{\sqrt{\tau_1}\sigma_{S_1}} & \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sqrt{\tau_1}\sigma_{S_1}} & \cdots & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sqrt{\tau_1}\sigma_{S_1}} \\ \frac{\sqrt{\tau_2}\sigma_{S_2}}{\sigma_F} & \frac{\sqrt{\tau_2}\sigma_{S_2}}{\sqrt{\tau_1}\sigma_{S_1}} & 1 & \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sqrt{\tau_2}\sigma_{S_2}} & \cdots & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sqrt{\tau_2}\sigma_{S_2}} \\ \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sigma_F} & \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sqrt{\tau_1}\sigma_{S_1}} & \frac{\sqrt{\tau_3}\sigma_{S_3}}{\sqrt{\tau_2}\sigma_{S_2}} & 1 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sqrt{\tau_{k-1}}\sigma_{S_{k-1}}} \\ \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sigma_F} & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sqrt{\tau_1}\sigma_{S_1}} & \cdots & \cdots & \frac{\sqrt{\tau_k}\sigma_{S_k}}{\sqrt{\tau_{k-1}}\sigma_{S_{k-1}}} & 1 \end{pmatrix}$$

## ► Case 3: Unknown and unequal variances

► Here we only have an asymptotic result, namely

$$\text{under } H_0^{\{\cap_{i=0}^k S_i\}}, \quad Z = \begin{pmatrix} Z\{F\} \\ Z\{S_1\} \\ \vdots \\ Z\{S_k\} \end{pmatrix} = \begin{pmatrix} \sqrt{\frac{n}{2}} \frac{\hat{\Delta}_F}{\hat{\sigma}_F} \\ \vdots \\ \sqrt{\frac{\tau_k n}{2}} \frac{\hat{\Delta}_{S_k}}{\hat{\sigma}_{S_k}} \end{pmatrix} \overset{\cdot}{\sim} \mathcal{N}(0, \Sigma)$$

with covariance matrix as described above

# Analysis (fixed design)

## Case 3: Unknown and unequal variances

- ▶ Try to approximate the joint distribution by a multivariate t-distribution
  - ▶ How to choose the degrees of freedom?
  - ▶ E.g. in the case of only one subgroup: under  $H_0^{\{F,S\}}$  (k=1)


$$\begin{pmatrix} Z^F \\ Z^S \end{pmatrix} = \begin{pmatrix} \sqrt{\frac{n}{2}} \frac{\hat{\Delta}_F}{\hat{\sigma}_F} \\ \sqrt{\frac{n^S}{2}} \frac{\hat{\Delta}_S}{\hat{\sigma}_S} \end{pmatrix} \dot{\sim} MT_{df}(\mathbf{0}, \Sigma)$$

- ▶ Choose df of the full population  $df = 2n - 4$  for a more liberal,  
or of the smallest subpopulation  $df = 2n^S - 2$  for a more  
conservative approximation

# Analysis (fixed design)

## Simulations – unknown, unequal variances – t-approximation

▷ Type-I-error rates,  $\sigma_F = 1$ ,  $\sigma_S = 1.3$ ,  $n_{sim} = 100,000$





# Sample Size Calculation

- ▶ under the alternative
 
$$\begin{aligned}
 \mathbf{Z} &\sim MN(\boldsymbol{\delta}, \boldsymbol{\Sigma}) && \text{(known variances)} \\
 \mathbf{Z} &\sim MT_{2n-4}(\boldsymbol{\delta}, \tilde{\boldsymbol{\Sigma}}) && \text{(unknown, same variances)} \\
 \mathbf{Z} &\dot{\sim} MT_{df}(\boldsymbol{\delta}, \tilde{\boldsymbol{\Sigma}}) && \text{(unknown, unequal variances)} \\
 &&& df \in \{2n-4, 2n^S-2\}
 \end{aligned}$$

with 
$$\boldsymbol{\delta} = \begin{pmatrix} \sqrt{\frac{n}{2}} \frac{\Delta_F}{\sigma_F} \\ \sqrt{\frac{n\tau}{2}} \frac{\Delta_S}{\sigma_S} \end{pmatrix}$$

- ▶ let  $\mathbf{G}_{MT_{df}}(\boldsymbol{\delta}, \tilde{\boldsymbol{\Sigma}})$  denote the distribution function of  $MT_{df}(\boldsymbol{\delta}, \tilde{\boldsymbol{\Sigma}})$  and  $z_{MT_{df}(\mathbf{0}, \boldsymbol{\Sigma}), 1-\alpha}$  the  $(1-\alpha)$ -equicoordinate quantile of  $MT_{df}(\mathbf{0}, \boldsymbol{\Sigma})$
- ▶ use estimates of nuisance parameters and effect sizes, e.g. based on previous studies, to calculate the initial sample size via

$$N_{init} = \min n, \text{ s.t. } 1 - \mathbf{G}_{MT_{df}}(\boldsymbol{\delta}, \tilde{\boldsymbol{\Sigma}})(z_{MT_{df}(\mathbf{0}, \boldsymbol{\Sigma}), 1-\alpha}) \geq 1 - \beta$$

# Sample Size Calculation

- ▶ In the case of unknown, unequal variances one has to choose which df to use for the MT-distribution in the final analysis and in the sample size calculation
- ▶ Calculating the sample size via a multivariate T-distribution requires to calculate a new equicoordinate quantile in each step of the search algorithm
  - ▶ Using the MN-distribution only one quantile is calculated
- ▶ Sample size calculation heavily depends on estimates of the nuisance parameters  $\hat{\sigma}_F, \hat{\sigma}_{S_1}, \dots, \hat{\sigma}_{S_k}$   
 $\hat{\tau}_1, \dots, \hat{\tau}_k$
- ▶ Multivariate Normal and t Probabilities, see Genz and Bretz (2009)
  - ▶ R package multcomp


# Sample Size Calculation - Performance

- ▶ Misspecification of the nuisance parameters can lead to substantial under/over estimation of the sample size

$$n_{sim} = 10,000$$

$$\Delta_S = 1$$

$$1 - \beta = 0.9$$




- ▶ Solution: Internal Pilot Study Design (Wittes & Brittain, 1990)

# IPS design with Blinded Review

- ▷ here: nuisance parameters

$$\sigma_F^2, \sigma_{S_1}^2, \dots, \sigma_{S_k}^2$$

$$\tau_1, \tau_2, \dots, \tau_k$$



- ▷ after  $n_1 = p \cdot N_0$  subjects per group (treatment/control):

- ▷ blinded reestimation via „lumped variance“

$$\hat{\sigma}_F^2 = \hat{\sigma}^2 = \frac{1}{2n_1 - 1} \sum_{i=0}^k \sum_{j=1}^{n_1^{S_i}} \sum_{l=1}^2 (X_{ijl} - \bar{X}_{i..})^2$$

$$\hat{\sigma}_{S_i}^2 = \frac{1}{2n_1^{S_i} - 1} \sum_{s=i}^k \sum_{j=1}^{n_1^{S_i}} \sum_{l=1}^2 (X_{sjl} - \bar{X}_{s..})^2, \quad i = 1, \dots, k$$

- ▷ Prevalences

$$\hat{\tau}_i = \frac{n_1^{S_i}}{n_1}, \quad i = 1, \dots, k$$



- ▷ plug in the new nuisance parameter estimates in the sample size calculation method

# BSSR - Simulations

- Simulated power and corresponding recalculated sample sizes depending on the number of subjects in the subgroup at the timepoint of the blinded review

$$1 - \beta = 0.9 \quad \Delta_S = 1 \quad \sigma_S^* \neq \sigma_S \quad n_{sim} = 10,000$$

- Conservative approximation  $df = 2n^S - 2$



# Blinded Sample Size Reestimation



- ▶ minimal number of subjects in the smallest subgroup to get an appropriate approximation
- ▶ when plugging in the new nuisance parameter estimates, we use the same search algorithm to determine the new sample size
  - ▶ Zucker et. al (1999): use df dependent on the number of subjects at the blinded review
  - ▶ Here: try to use df depending on  $n_1^S$  to improve when dealing with small subgroups

# BSSR - Simulations

- Simulated power and corresponding recalculated sample sizes depending on the number of subjects in the subgroup at the timepoint of the blinded review

$$1 - \beta = 0.9 \quad \Delta_S = 1 \quad \sigma_S^* \neq \sigma_S \quad n_{sim} = 10,000$$

- Conservative approximation at final analysis



## Conclusions & Discussion

- ▶ Method to analyze multiple nested subgroup designs via joined distribution of standardized test statistics
- ▶ Approximation for unknown and unequal variances in the subgroups
- ▶ Sample size calculation approach derived using this approx requires minimum 20-30 subjects in smallest subgroup
  - ▶ compare Sandvik et. al (1996)
- ▶ R package in work
- ▶ Next step: use these findings in the combination of BSSR and Adaptive Enrichment Methods



# References

- ▶ Spiessens B, Debois M (2010) *Adjusted significance levels for subgroup analysis in clinical trials*. Contemporary Clinical Trials 31: 647-656
- ▶ Wittes J, Brittain E (1990). *The role of internal pilot studies in increasing the efficiency of clinical trials*. Statistics in Medicine 9: 65–72.
- ▶ Genz A, Bretz F (2009). Computation of Multivariate Normal and t Probabilities. SpringerVerlag, New York.
- ▶ Friede T, Kieser M (2006). Sample size recalculation in internal pilot study designs: a review. *Biometrical Journal* 2006; 48:537–555.
- ▶ Sandvik, L., Erikssen, J., Mowinckel, P. and Rodland, E. A. (1996). A method for determining the size of internal pilot studies. *Statistics in Medicine* 15, 1587–1590.
- ▶ Zucker, D.M., Wittes, J.T., Schabenberger, O., Brittain, E. (1999). Internal pilot studies II: Comparison of various procedures. *Statistics in Medicine* 18, 3493–3509.