

## GAD-7 Anxiety

Over the <u>last two weeks</u> , how often have you been bothered by the following problems?	Not at all	Several days	More than half the days	Nearly every day
1. Feeling nervous, anxious, or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid, as if something awful might happen	0	1	2	3

Column totals      \_\_\_\_\_ + \_\_\_\_\_ + \_\_\_\_\_ + \_\_\_\_\_ =

*Total score* \_\_\_\_\_

If you checked any problems, how difficult have they made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all

Somewhat difficult

### Very difficult

Extremely difficult



## Scoring GAD-7 Anxiety Severity

This is calculated by assigning scores of 0, 1, 2, and 3 to the response categories, respectively, of "not at all," "several days," "more than half the days," and "nearly every day."

GAD-7 total score for the seven items ranges from 0 to 21.

0–4: minimal anxiety

5–9: mild anxiety

## 10–14: moderate anxiety

## 15–21: severe anxiety

## Copyright

© 1999 Pfizer Inc. All rights reserved. Reproduced with permission.

## **Terms of Use**

Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PRIME-MD-PHQ). The PHQ was developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke, and colleagues. For research information, contact Dr. Spitzer at [ris8@columbia.edu](mailto:ris8@columbia.edu). PRIME-MD® is a trademark of Pfizer Inc.

## Reference

Spitzer, R. L., Kroenke, K., Williams, J. B. W., Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, 166(10), 1092-1097.