


# Diet and age effects on Gasping or abnormal rate/depth of breathing (0, 0.5, 1)



Only the following timepoints were used when testing for direct diet and age–diet interaction effects (all timepoints were used when testing for direct age effects): 05 months, 08 months, 10 months, 16 months, 22 months and 28 months. The effects of age, diet, and the age–diet interaction were estimated using mixed linear models and the significance of the effects were assessed with an approximate F–test using the Kenward and Roger (1997) approach. The p–values for the diet effect at each timepoint are: 05 months = NA; 08 months = NA; 10 months = 0.567; 16 months =  $7.94e-05$ ; 22 months = 0.00719 and 28 months = 0.00323. The diet pairs that have significantly different (Tukey p–value < 0.05) means at 16 months are AL–2D, AL–20 and AL–40. The diet pairs that have significantly different (Tukey p–value < 0.05) means at 22 months are AL–40. The diet pairs that have significantly different (Tukey p–value < 0.05) means at 28 months are AL–1D, AL–20 and AL–40. The p–value for the direct effect of age on BreathingRateDepth is  $1.72e-07$ . The p–value for the effect of the interaction between age and diet on BreathingRateDepth is  $1.49e-06$ . The diet pairs that have significantly different (Tukey p–value < 0.05) rates of change with age are AL–1D, AL–2D, AL–20 and AL–40.