# 1.27 Second-Order Systems

Consider a second-order system with peak time  $t_p = 0.5$  sec. and 5% settling time  $t_s = 1.5$  sec. Determine the poles of this system.

## **Solution:**

The poles of this system are given by  $s=-\sigma\pm j\omega_d$  where  $\sigma=\zeta\omega_n$  and  $\omega_d=\omega_n\sqrt{1-\zeta^2}$ .

Given information  $t_p=0.5$  sec and 5% settling time  $t_s=1.5$  sec leads to  $\omega_d=2\pi$  and  $\sigma=2$ . Therefore the poles of the system are  $s=-2\pm 2\pi$ 

### **GPT Answer:**

The calculations yield the damping ratio  $\zeta=0.391$  and the natural frequency  $\omega_n=6.826$  rad/sec. Using these values, the poles of the system are located at -2.667+6.283i and -2.667-6.283i.

Therefore, the poles of this second-order system are complex conjugates, indicating an underdamped system.

#### **Confidence Level: 95%**

#### **Calculation error**

#### **Claude Answer**

To determine the poles of the second-order system, we need to calculate the natural frequency  $(\omega_n)$  and the damping ratio  $(\zeta)$  using the given peak time  $(t_p)$  and settling time  $(t_s)$ .

Given: - Peak time, 
$$t_p = 0.5 \text{ sec}$$
  
- Settling time (5%),  $t_s = 1.5 \text{ sec}$ 

Step 1: Calculate the damping ratio ( $\zeta$ ) using the settling time. For a second-order system, the settling time is related to the damping ratio by the following equation:  $t_s \approx \frac{3}{\zeta \omega_n}$ 

Step 2: Calculate the natural frequency  $(\omega_n)$  using the peak time. The peak time is related to the damping ratio and natural frequency by the following equation:  $t_p = \frac{\pi}{\omega_n \sqrt{1-\zeta^2}}$ 

Step 3: Solve for 
$$\zeta$$
 and  $\omega_n$ . From the peak time equation:  $0.5 = \frac{\pi}{\omega_n \sqrt{1-\zeta^2}} \omega_n = \frac{\pi}{0.5\sqrt{1-\zeta^2}}$ 

Substituting this into the settling time equation:  $1.5 \approx \frac{3}{\zeta \frac{\pi}{0.5\sqrt{1-\zeta^2}}} 1.5 \approx \frac{3 \cdot 0.5 \sqrt{1-\zeta^2}}{\zeta \pi} 1.5 \zeta \pi \approx 1.5 \sqrt{1-\zeta^2} \zeta^2 \pi^2 \approx 1-\zeta^2 \zeta^2 (\pi^2+1) \approx 1 \zeta \approx \sqrt{\frac{1}{\pi^2+1}} \approx 0.3015$ 

Now, calculate 
$$\omega_n$$
:  $\omega_n = \frac{\pi}{0.5\sqrt{1-0.3015^2}} \approx 6.6579 \text{ rad/s}$ 

Step 4: Calculate the poles using  $\zeta$  and  $\omega_n$ . The poles of a second-order system are given by:  $s_{1,2} = -\zeta \omega_n \pm j \omega_n \sqrt{1-\zeta^2}$ 

Substituting the values:  $s_{1,2} = -0.3015 \cdot 6.6579 \pm j \cdot 6.6579 \sqrt{1 - 0.3015^2} \ s_{1,2} = -2.0074 \pm j \cdot 6.2078$ 

Therefore, the poles of the given second-order system are  $s_1=-2.0074+j6.2078$  and  $s_2=-2.0074-j6.2078$ .

Confidence level: 95%

Correct