

# Geometric Patterns and Microstructures in the study of Material Defects and Composites

Silvio Fanzon

supervised by  
Mariapia Palombaro

University of Sussex  
Department of Mathematics



# Presentation Plan

## ① Geometric Patterns of Dislocations

## ② Microstructures in Composites

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations

## ② Microstructures in Composites

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces (Chapter 3)

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*.  
Journal of Nonlinear Science (2017)

## ② Microstructures in Composites

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces (Chapter 3)  
F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*.  
Journal of Nonlinear Science (2017)
- ▶ Linearised polycrystals (Chapter 4)  
F., Palombaro, Ponsiglione. *Linearized Polycrystals from a 2D System of Edge Dislocations*.  
Preprint (2017)

## ② Microstructures in Composites

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces (Chapter 3)  
F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*. Journal of Nonlinear Science (2017)
- ▶ Linearised polycrystals (Chapter 4)  
F., Palombaro, Ponsiglione. *Linearized Polycrystals from a 2D System of Edge Dislocations*. Preprint (2017)

## ② Microstructures in Composites

- ▶ Critical lower integrability (Chapter 5)  
F., Palombaro. *Optimal lower exponent for the higher gradient integrability of solutions to two-phase elliptic equations in two dimensions*. Calculus of Variations and Partial Differential Equations (2017)

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces (Chapter 3)  
F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*. Journal of Nonlinear Science (2017)
- ▶ Linearised polycrystals (Chapter 4)  
F., Palombaro, Ponsiglione. *Linearized Polycrystals from a 2D System of Edge Dislocations*. Preprint (2017)

## ② Microstructures in Composites

- ▶ Critical lower integrability (Chapter 5)  
F., Palombaro. *Optimal lower exponent for the higher gradient integrability of solutions to two-phase elliptic equations in two dimensions*. Calculus of Variations and Partial Differential Equations (2017)
- ▶ Convex integration

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces (Chapter 3)  
F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*. Journal of Nonlinear Science (2017)
- ▶ Linearised polycrystals (Chapter 4)  
F., Palombaro, Ponsiglione. *Linearized Polycrystals from a 2D System of Edge Dislocations*. Preprint (2017)

## ② Microstructures in Composites

- ▶ Critical lower integrability (Chapter 5)  
F., Palombaro. *Optimal lower exponent for the higher gradient integrability of solutions to two-phase elliptic equations in two dimensions*. Calculus of Variations and Partial Differential Equations (2017)
- ▶ Convex integration
- ▶ Proof of the main theorem



# Presentation Plan

## ① Geometric Patterns of Dislocations



- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.



# Edge dislocations


**Dislocations:** topological defects in the otherwise periodic structure of a crystal.



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.

**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .









# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.

**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .




# Screw dislocations

**Screw dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi // \gamma$ .

# Screw dislocations

**Screw dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \parallel \gamma$ .



# Mixed type dislocations

**Mixed dislocations:** Burgers vector  $\xi$  is constant and  $\gamma$  is curved.

# Mixed type dislocations



**Mixed dislocations:** Burgers vector  $\xi$  is constant and  $\gamma$  is curved.

**Dislocation type:** given by the angle between  $\xi$  and  $\dot{\gamma}$ .

# Mixed type dislocations

**Mixed dislocations:** Burgers vector  $\xi$  is constant and  $\gamma$  is curved.


**Dislocation type:** given by the angle between  $\xi$  and  $\dot{\gamma}$ .





# Nonlinear Elasticity


**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded



# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$




# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$



# Nonlinear Elasticity


**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Energy:** associated to a deformation strain  $\beta$

$$E(\beta) := \int_{\Omega} W(\beta) \, dx.$$



# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded


**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Energy:** associated to a deformation strain  $\beta$

$$E(\beta) := \int_{\Omega} W(\beta) \, dx.$$

**Energy Density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  s.t.



# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$


**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Energy:** associated to a deformation strain  $\beta$

$$E(\beta) := \int_{\Omega} W(\beta) \, dx.$$

**Energy Density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  s.t.

- $W$  is continuous



# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$


**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Energy:** associated to a deformation strain  $\beta$

$$E(\beta) := \int_{\Omega} W(\beta) \, dx.$$

**Energy Density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  s.t.

- ▶  $W$  is continuous
- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$ ,  $F \in \mathbb{M}^{3 \times 3}$   
(frame indifferent),



# Nonlinear Elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$


**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Energy:** associated to a deformation strain  $\beta$

$$E(\beta) := \int_{\Omega} W(\beta) \, dx.$$

**Energy Density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  s.t.


- ▶  $W$  is continuous
- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$ ,  $F \in \mathbb{M}^{3 \times 3}$   
(frame indifferent),
- ▶  $W(F) \sim \text{dist}(F, SO(3))^2 \implies W(I) = 0$ .





# Semi-discrete model for dislocations


**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected



# Semi-discrete model for dislocations

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions




# Semi-discrete model for dislocations

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions

**Strain generating  $(\gamma, \xi)$ :** map  $\beta: \Omega \rightarrow \mathbb{M}^{3 \times 3}$  s.t.

$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma \iff \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$



# Semi-discrete model for dislocations

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions


**Strain generating  $(\gamma, \xi)$ :** map  $\beta: \Omega \rightarrow \mathbb{M}^{3 \times 3}$  s.t.

$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma \iff \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$

**Geometric interpretation:** if  $D$  encloses  $\gamma$ , there exists a deformation  $v \in SBV(\Omega; \mathbb{R}^3)$  s.t.

$$Dv = \nabla v \, dx + \xi \otimes n \mathcal{H}^2 \llcorner D, \quad \beta = \nabla v.$$


$v$  has constant jump  $\xi$  across the slip region  $D$ .



# Strains are not $L^2$

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$




# Strains are not $L^2$

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma))$$



# Strains are not $L^2$


Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma))$$

**Proof:** let  $\sigma > \varepsilon$  and  $L := \text{length}(\gamma)$



# Strains are not $L^2$

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and


$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma))$$

**Proof:** let  $\sigma > \varepsilon$  and  $L := \text{length}(\gamma)$

$$\int_{I_\sigma \setminus I_\varepsilon} |\beta|^2 = L \int_\varepsilon^\sigma \int_{\partial B_\rho(\gamma(s))} |\beta|^2 d\mathcal{H}^1 d\rho$$





# Strains are not $L^2$

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$


Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma))$$

**Proof:** let  $\sigma > \varepsilon$  and  $L := \text{length}(\gamma)$

$$\int_{I_\sigma \setminus I_\varepsilon} |\beta|^2 = L \int_\varepsilon^\sigma \int_{\partial B_\rho(\gamma(s))} |\beta|^2 d\mathcal{H}^1 d\rho$$

$$(\text{Jensen}) \geq L \int_\varepsilon^\sigma \frac{1}{2\pi\rho} \left| \int_{\partial B_\rho(\gamma(s))} \beta \cdot t d\mathcal{H}^1 \right|^2 d\rho$$



# Strains are not $L^2$

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have


$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma))$$

**Proof:** let  $\sigma > \varepsilon$  and  $L := \text{length}(\gamma)$

$$\int_{I_\sigma \setminus I_\varepsilon} |\beta|^2 = L \int_\varepsilon^\sigma \int_{\partial B_\rho(\gamma(s))} |\beta|^2 d\mathcal{H}^1 d\rho$$

$$(\text{Jensen}) \geq L \int_\varepsilon^\sigma \frac{1}{2\pi\rho} \left| \int_{\partial B_\rho(\gamma(s))} \beta \cdot t d\mathcal{H}^1 \right|^2 d\rho$$

$$= L \frac{|\xi|^2}{2\pi} \log \frac{\sigma}{\varepsilon} \rightarrow \infty \text{ as } \varepsilon \rightarrow 0$$




# Regularise the problem

**Energy Truncation.** Fix  $p \in (1, 2)$  and assume

$$W(F) \sim \text{dist}(F, SO(3))^2 \wedge (|F|^p + 1).$$

Strains are maps  $\beta \in L^2(\Omega; \mathbb{M}^{3 \times 3})$  such that

$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma.$$



# Regularise the problem

**Energy Truncation.** Fix  $p \in (1, 2)$  and assume

$$W(F) \sim \text{dist}(F, SO(3))^2 \wedge (|F|^p + 1).$$

Strains are maps  $\beta \in L^2(\Omega; \mathbb{M}^{3 \times 3})$  such that

$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma.$$

**Core Radius Approach.** Assume


$$W(F) \sim \text{dist}(F, SO(3))^2.$$

Let  $\varepsilon > 0$  ( $\propto$  atomic distance) and consider

$$\Omega_\varepsilon(\gamma) := \Omega \setminus I_\varepsilon(\gamma).$$

Strains are maps  $\beta \in L^2(\Omega_\varepsilon(\gamma); \mathbb{M}^{3 \times 3})$  such that

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\gamma) = 0, \quad \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$



# Presentation Plan

## ① Geometric Patterns of Dislocations


- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result

# Semi-coherent interfaces


Two different crystalline materials joined at a flat interface:



# Semi-coherent interfaces

Two different crystalline materials joined at a flat interface:


- **Underlayer:** cubic lattice  $\Lambda^-$ , spacing  $b > 0$  (equilibrium  $l$ ),



# Semi-coherent interfaces

Two different crystalline materials joined at a flat interface:

- **Underlayer:** cubic lattice  $\Lambda^-$ , spacing  $b > 0$  (equilibrium  $l$ ),
- **Overlayer:** lattice  $\Lambda^+ = \alpha\Lambda^-$ , with  $\alpha > 1$  (not in equilibrium).






# Semi-coherent interfaces

Two different crystalline materials joined at a flat interface:

- **Underlayer:** cubic lattice  $\Lambda^-$ , spacing  $b > 0$  (equilibrium  $l$ ),
- **Overlayer:** lattice  $\Lambda^+ = \alpha\Lambda^-$ , with  $\alpha > 1$  (not in equilibrium).

**Semi-coherent interface:** small dilation  $\alpha \approx 1$ .




# Semi-coherent interfaces

Two different crystalline materials joined at a flat interface:

- **Underlayer:** cubic lattice  $\Lambda^-$ , spacing  $b > 0$  (equilibrium  $l$ ),
- **Overlayer:** lattice  $\Lambda^+ = \alpha\Lambda^-$ , with  $\alpha > 1$  (not in equilibrium).


**Semi-coherent interface:** small dilation  $\alpha \approx 1$ .

**Equilibrium:**  $\Lambda^+$  has lower density than  $\Lambda^- \implies$  **edge dislocations** at interface.



# Network of dislocations


## Experimentally observed phenomena:



# Network of dislocations

## Experimentally observed phenomena:

- ▶ two non-parallel sets of **edge dislocations** with spacing  $\delta = \frac{b}{\alpha-1}$ ,




D.A. Porter, K.E. Easterling. *Phase transformations in metals and alloys*. CRC Press (2009)

G. Gottstein. *Physical foundations of materials science*. Springer (2013)

# Network of dislocations

## Experimentally observed phenomena:

- ▶ two non-parallel sets of **edge dislocations** with spacing  $\delta = \frac{b}{\alpha-1}$ ,
- ▶ far field stress is completely relieved.



D.A. Porter, K.E. Easterling. *Phase transformations in metals and alloys*. CRC Press (2009)

G. Gottstein. *Physical foundations of materials science*. Springer (2013)

# Goal of the Paper

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces.*  
Journal of Nonlinear Science (2017)

# Goal of the Paper

$R$  is the size of the interface.

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces.*  
Journal of Nonlinear Science (2017)

# Goal of the Paper

$R$  is the size of the interface.

**Goal:** define a **continuum model** such that

- ▶  $\exists$  critical size  $R^*$  such that nucleation of dislocations is energetically more favorable for  $R > R^*$ ,

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces.*  
Journal of Nonlinear Science (2017)



# Goal of the Paper

$R$  is the size of the interface.

**Goal:** define a **continuum model** such that

- ▶  $\exists$  critical size  $R^*$  such that nucleation of dislocations is energetically more favorable for  $R > R^*$ ,
- ▶ as  $R \rightarrow \infty$  the far field stress is relieved,

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces.*  
Journal of Nonlinear Science (2017)

# Goal of the Paper

$R$  is the size of the interface.

**Goal:** define a **continuum model** such that

- ▶  $\exists$  critical size  $R^*$  such that nucleation of dislocations is energetically more favorable for  $R > R^*$ ,
- ▶ as  $R \rightarrow \infty$  the far field stress is relieved,
- ▶ the dislocation spacing tends to  $\delta = \frac{b}{\alpha - 1}$ .

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*.  
Journal of Nonlinear Science (2017)

# Goal of the Paper

$R$  is the size of the interface.

**Goal:** define a **continuum model** such that

- ▶  $\exists$  critical size  $R^*$  such that nucleation of dislocations is energetically more favorable for  $R > R^*$ ,
- ▶ as  $R \rightarrow \infty$  the far field stress is relieved,
- ▶ the dislocation spacing tends to  $\delta = \frac{b}{\alpha - 1}$ .

**Plan:**

- ▶ analysis of a **semi-discrete model** where dislocations are line defects,

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*.  
Journal of Nonlinear Science (2017)

# Goal of the Paper

$R$  is the size of the interface.

**Goal:** define a **continuum model** such that

- ▶  $\exists$  critical size  $R^*$  such that nucleation of dislocations is energetically more favorable for  $R > R^*$ ,
- ▶ as  $R \rightarrow \infty$  the far field stress is relieved,
- ▶ the dislocation spacing tends to  $\delta = \frac{b}{\alpha - 1}$ .

**Plan:**


- ▶ analysis of a **semi-discrete model** where dislocations are line defects,
- ▶ derive the simplified (dislocation density) **continuum model**.

F., Palombaro, Ponsiglione. *A Variational Model for Dislocations at Semi-coherent Interfaces*.  
Journal of Nonlinear Science (2017)

# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha l$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).




# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha l$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).

**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.




# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).

**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),




# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).

**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),
- ▶  $W(F) \sim \text{dist}(F, \alpha SO(3))^2 \wedge (|F|^p + 1)$  for  $1 < p < 2$ .





# Semi-discrete line defect model


**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).

**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),
- ▶  $W(F) \sim \text{dist}(F, \alpha SO(3))^2 \wedge (|F|^p + 1)$  for  $1 < p < 2$ .

**Admissible dislocations:** compatible with cubic lattice.  
 $(\Gamma, B) \in \mathcal{AD}$  if  $\Gamma = \{\gamma_i\}$ ,  $B = \{\xi_i\}$  with



# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).


**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),
- ▶  $W(F) \sim \text{dist}(F, \alpha SO(3))^2 \wedge (|F|^p + 1)$  for  $1 < p < 2$ .

**Admissible dislocations:** compatible with cubic lattice.

$(\Gamma, B) \in \mathcal{AD}$  if  $\Gamma = \{\gamma_i\}$ ,  $B = \{\xi_i\}$  with

- ▶ dislocation line  $\gamma_i \subset \mathcal{G}$  relatively closed,



# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).


**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),
- ▶  $W(F) \sim \text{dist}(F, \alpha SO(3))^2 \wedge (|F|^p + 1)$  for  $1 < p < 2$ .

**Admissible dislocations:** compatible with cubic lattice.

$(\Gamma, B) \in \mathcal{AD}$  if  $\Gamma = \{\gamma_i\}$ ,  $B = \{\xi_i\}$  with

- ▶ dislocation line  $\gamma_i \subset \mathcal{G}$  relatively closed,
- ▶ Burgers vector  $\xi_i \in b(\mathbb{Z} \oplus \mathbb{Z})$ .



# Semi-discrete line defect model

**Reference configuration:**  $\Omega_r := \Omega_r^- \cup S_r \cup \Omega_r^+$ ,  $r > 0$ ,

- ▶  $\Omega_r^+$  overlayer (equilibrium  $\alpha I$ ),
- ▶  $\Omega_r^-$  underlayer (in equilibrium and rigid).

**Energy density:**  $W: \mathbb{M}^{3 \times 3} \rightarrow [0, \infty)$  continuous, s.t.

- ▶  $W(F) = W(RF)$ ,  $\forall R \in SO(3)$  (frame indifference),
- ▶  $W(F) \sim \text{dist}(F, \alpha SO(3))^2 \wedge (|F|^p + 1)$  for  $1 < p < 2$ .


**Admissible dislocations:** compatible with cubic lattice.

$(\Gamma, B) \in \mathcal{AD}$  if  $\Gamma = \{\gamma_i\}$ ,  $B = \{\xi_i\}$  with

- ▶ dislocation line  $\gamma_i \subset \mathcal{G}$  relatively closed,
- ▶ Burgers vector  $\xi_i \in b(\mathbb{Z} \oplus \mathbb{Z})$ .

**Admissible strains:** for a dislocation  $(\Gamma, B)$  are the maps  $\beta \in AS(\Gamma, B)$ , such that  $\beta \in L^p(\Omega_r; \mathbb{M}^{3 \times 3})$  and

$$\beta = I \text{ in } \Omega_r^-, \quad \text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \Gamma.$$



# Scaling properties of the energy

**Energies:** induced by the misfit

# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \operatorname{Curl} \beta = 0 \right\} \quad (\text{Elastic energy})$$

# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \text{Curl } \beta = 0 \right\} \quad (\text{Elastic energy})$$

$$E_{\alpha,r} := \min_{(\Gamma,B) \in \mathcal{AD}} \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \beta \in \mathcal{AS}(\Gamma, B) \right\} \quad (\text{Plastic energy})$$

# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \operatorname{Curl} \beta = 0 \right\} \quad (\text{Elastic energy})$$

$$E_{\alpha,r} := \min_{(\Gamma,B) \in \mathcal{AD}} \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \beta \in \mathcal{AS}(\Gamma, B) \right\} \quad (\text{Plastic energy})$$

**Theorem (F., Palombaro, Ponsiglione '15)**

*The dislocation-free elastic energy scales like  $r^3$ : we have  $E_{\alpha,1}(\emptyset) > 0$  and*

$$E_{\alpha,r}(\emptyset) = r^3 E_{\alpha,1}(\emptyset).$$



# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \text{Curl } \beta = 0 \right\} \quad (\text{Elastic energy})$$

$$E_{\alpha,r} := \min_{(\Gamma,B) \in \mathcal{AD}} \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \beta \in \mathcal{AS}(\Gamma, B) \right\} \quad (\text{Plastic energy})$$

## Theorem (F., Palombaro, Ponsiglione '15)

*The dislocation-free elastic energy scales like  $r^3$ : we have  $E_{\alpha,1}(\emptyset) > 0$  and*

$$E_{\alpha,r}(\emptyset) = r^3 E_{\alpha,1}(\emptyset).$$

*The plastic energy scales like  $r^2$ : there exists  $0 < E_\alpha < +\infty$  such that*

$$E_{\alpha,r} = r^2 E_\alpha + o(r^2).$$

# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \text{Curl } \beta = 0 \right\} \quad (\text{Elastic energy})$$

$$E_{\alpha,r} := \min_{(\Gamma,B) \in \mathcal{AD}} \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \beta \in \mathcal{AS}(\Gamma, B) \right\} \quad (\text{Plastic energy})$$

## Theorem (F., Palombaro, Ponsiglione '15)

*The dislocation-free elastic energy scales like  $r^3$ : we have  $E_{\alpha,1}(\emptyset) > 0$  and*

$$E_{\alpha,r}(\emptyset) = r^3 E_{\alpha,1}(\emptyset).$$

*The plastic energy scales like  $r^2$ : there exists  $0 < E_\alpha < +\infty$  such that*

$$E_{\alpha,r} = r^2 E_\alpha + o(r^2).$$

*Large  $r \implies$  dislocations are energetically favourable.*

# Scaling properties of the energy

**Energies:** induced by the misfit

$$E_{\alpha,r}(\emptyset) := \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \operatorname{Curl} \beta = 0 \right\} \quad (\text{Elastic energy})$$

$$E_{\alpha,r} := \min_{(\Gamma,B) \in \mathcal{AD}} \inf \left\{ \int_{\Omega_r^+} W(\beta) dx : \beta \in \mathcal{AS}(\Gamma, B) \right\} \quad (\text{Plastic energy})$$

## Theorem (F., Palombaro, Ponsiglione '15)

*The dislocation-free elastic energy scales like  $r^3$ : we have  $E_{\alpha,1}(\emptyset) > 0$  and*

$$E_{\alpha,r}(\emptyset) = r^3 E_{\alpha,1}(\emptyset).$$

*The plastic energy scales like  $r^2$ : there exists  $0 < E_\alpha < +\infty$  such that*

$$E_{\alpha,r} = r^2 E_\alpha + o(r^2).$$

*Large  $r \implies$  dislocations are energetically favourable.*


# Upper bound construction

**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .

# Upper bound construction

**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .



- Divide  $S_r$  into  $(r/\delta)^2$  squares of side  $\delta$ .



# Upper bound construction

**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .



- ▶ Divide  $S_r$  into  $(r/\delta)^2$  squares of side  $\delta$ .
- ▶ Above each  $Q_i$  define pyramids  $C_i^1$  (height  $\delta/2$ ) and  $C_i^2$  (height  $\delta$ ).



# Upper bound construction

**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .

- ▶ Divide  $S_r$  into  $(r/\delta)^2$  squares of side  $\delta$ .
- ▶ Above each  $Q_i$  define pyramids  $C_i^1$  (height  $\delta/2$ ) and  $C_i^2$  (height  $\delta$ ).
- ▶ Define deformation  $\mathbf{v} \in SBV(\Omega_r; \mathbb{R}^3)$ , and strain  $\beta := \nabla \mathbf{v}$  (a.c. part of  $D\mathbf{v}$ ).





# Upper bound construction

**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .

- ▶ Divide  $S_r$  into  $(r/\delta)^2$  squares of side  $\delta$ .
- ▶ Above each  $Q_i$  define pyramids  $C_i^1$  (height  $\delta/2$ ) and  $C_i^2$  (height  $\delta$ ).
- ▶ Define deformation  $\mathbf{v} \in SBV(\Omega_r; \mathbb{R}^3)$ , and strain  $\beta := \nabla \mathbf{v}$  (a.c. part of  $D\mathbf{v}$ ).

**Induced dislocations:**  $\text{Curl } \beta = - \sum_{i,j} \xi_{ij} \otimes \dot{\gamma}_{ij} d\mathcal{H}^1 \llcorner \gamma_{ij}$  with

- ▶  $\gamma_{ij} := Q_i \cap Q_j$  admissible dislocation curve ( $\alpha = 1 + 1/n \implies \delta = nb$ )
- ▶  $\xi_{ij} := (\alpha - 1)(x_j - x_i) \in \pm b\{e_1, e_2\}$  Burgers vector





# Upper bound construction



**Goal:** define a square array of edge dislocations with spacing  $\delta := \frac{b}{\alpha - 1}$ .

- ▶ Divide  $S_r$  into  $(r/\delta)^2$  squares of side  $\delta$ .
- ▶ Above each  $Q_i$  define pyramids  $C_i^1$  (height  $\delta/2$ ) and  $C_i^2$  (height  $\delta$ ).
- ▶ Define deformation  $\mathbf{v} \in SBV(\Omega_r; \mathbb{R}^3)$ , and strain  $\beta := \nabla \mathbf{v}$  (a.c. part of  $D\mathbf{v}$ ).

**Induced dislocations:**  $\text{Curl } \beta = - \sum_{i,j} \xi_{ij} \otimes \dot{\gamma}_{ij} d\mathcal{H}^1 \llcorner \gamma_{ij}$  with


- ▶  $\gamma_{ij} := Q_i \cap Q_j$  admissible dislocation curve ( $\alpha = 1 + 1/n \implies \delta = nb$ )
- ▶  $\xi_{ij} := (\alpha - 1)(x_j - x_i) \in \pm b\{e_1, e_2\}$  Burgers vector

**Energy:** in each pyramid is  $c = c(\alpha, b, p) \implies E_{\alpha,r} \leq c \frac{r^2}{\delta^2}$  (as  $W(\alpha I) = 0$ ).




# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction





# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction




# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction



# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction




**Limitations of the considered model:**

- $v(S_r)$  does not match  $S_r \implies$  not appropriate for semi-coherent interfaces,

# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction




## Limitations of the considered model:

- ▶  $v(S_r)$  does not match  $S_r \implies$  not appropriate for semi-coherent interfaces,
- ▶ expected dislocation geometry with spacing  $\frac{b}{\alpha-1}$  is only optimal in scaling.

# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction



## Limitations of the considered model:


- ▶  $v(S_r)$  does not match  $S_r \implies$  not appropriate for semi-coherent interfaces,
- ▶ expected dislocation geometry with spacing  $\frac{b}{\alpha-1}$  is only optimal in scaling.

## What we do now:

- ▶ take a smaller overlayer and enforce match at the interface,

# Remarks on the semi-discrete model

**Deformed configuration:**  $v(S_R)$  with  $v$  from the upper bound construction



## Limitations of the considered model:


- ▶  $v(S_r)$  does not match  $S_r \implies$  not appropriate for semi-coherent interfaces,
- ▶ expected dislocation geometry with spacing  $\frac{b}{\alpha-1}$  is only optimal in scaling.

## What we do now:

- ▶ take a smaller overlay and enforce match at the interface,
- ▶ introduce a simplified continuum (dislocation density) model to better describe true minimisers.




# Heuristic for the continuum model



**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$


# Heuristic for the continuum model



**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_R \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1}$


# Heuristic for the continuum model



**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

# Heuristic for the continuum model




**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

$$L = 2R \frac{r}{\delta} = \frac{2r^2}{b} (\theta^{-2} - \theta^{-1}) \stackrel{(\theta^{-1} \approx 1)}{\approx} \frac{r^2}{b} (\theta^{-2} - 1) = \frac{1}{b} (R^2 - r^2) = \frac{1}{b} \text{Area Gap}$$

# Heuristic for the continuum model




**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

$$\text{Dislocation Length} \approx \frac{1}{b} \text{Area Gap}$$

# Heuristic for the continuum model




**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

Dislocation Length  $\approx \frac{1}{b}$  Area Gap

$$E_{\alpha,r} \approx r^2 E_\alpha$$

# Heuristic for the continuum model




**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

Dislocation Length  $\approx \frac{1}{b}$  Area Gap

$$E_{\alpha,r} \approx r^2 E_\alpha = \sigma \text{ Area Gap} \quad \text{with} \quad \sigma := \frac{E_\alpha}{\theta^{-2} - 1}$$

# Heuristic for the continuum model



**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Upper bound construction:** with  $\theta = \alpha^{-1}$  and  $\delta = \frac{b}{\theta^{-1}-1} \implies$  perfect match

$$\text{Dislocation Length} \approx \frac{1}{b} \text{Area Gap}$$


$$E_{\alpha,r} \approx r^2 E_\alpha = \sigma \text{Area Gap} \quad \text{with} \quad \sigma := \frac{E_\alpha}{\theta^{-2} - 1}$$

**Hypothesis:** Dislocation Energy  $\propto$  Dislocation Length. Then optimise over  $\theta$ .



# Continuum model


**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  
 $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$



# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)




# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$




# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$



# Continuum model


**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$

**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$



# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$


**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$

**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$

**Total Energy:**  $E_{\alpha,R}^{tot}(\theta) := \min_{\theta} \left( E_{\alpha,R}^{el}(\theta) + E_R^{pl}(\theta) \right)$



# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)


**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$

**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$

**Total Energy:**  $E_{\alpha,R}^{tot}(\theta) := \min_{\theta} \left( E_{\alpha,R}^{el}(\theta) + E_R^{pl}(\theta) \right)$

**Question:** behaviour of  $E_{\alpha,R}^{tot}(\theta)$  as  $R \rightarrow \infty$ ?



# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$


**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$

**Total Energy:**  $E_{\alpha,R}^{tot}(\theta) := \min_{\theta} \left( E_{\alpha,R}^{el}(\theta) + E_R^{pl}(\theta) \right)$

**Question:** behaviour of  $E_{\alpha,R}^{tot}(\theta)$  as  $R \rightarrow \infty$ ?

**Energy competition:**

►  $\theta = 1 \implies$  no dislocation energy





# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$

**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$


**Total Energy:**  $E_{\alpha,R}^{tot}(\theta) := \min_{\theta} \left( E_{\alpha,R}^{el}(\theta) + E_R^{pl}(\theta) \right)$

**Question:** behaviour of  $E_{\alpha,R}^{tot}(\theta)$  as  $R \rightarrow \infty$ ?

**Energy competition:**

- ▶  $\theta = 1 \implies$  no dislocation energy
- ▶  $\theta = \alpha^{-1} \implies$  no elastic energy

$$(v := \alpha x, W(\alpha I) = 0)$$



# Continuum model

**Reference configuration:**  $\Omega_{R,r} := \Omega_R^- \cup S_r \cup \Omega_r^+$ , with  $r := \theta R$ ,  $\theta \in [\alpha^{-1}, 1]$

**Deformations:**  $v \in W^{1,2}(\Omega_r^+; \mathbb{R}^3)$  such that  $v = \frac{x}{\theta}$  on  $S_r$   
 $\implies v(S_r) = S_R$  (interface match)

**Energy density:**  $W(F) \sim \text{dist}(F, \alpha SO(3))^2$

**Elastic:**  $E_{\alpha,R}^{el}(\theta) := \inf \left\{ \int_{\Omega_r^+} W(\nabla v) dx : v = x/\theta \text{ on } S_r \right\}$


**Plastic:**  $E_R^{pl}(\theta) := \sigma \text{ Area Gap} = \sigma R^2(1 - \theta^2)$ ,  $\sigma > 0$

**Total Energy:**  $E_{\alpha,R}^{tot}(\theta) := \min_{\theta} \left( E_{\alpha,R}^{el}(\theta) + E_R^{pl}(\theta) \right)$

**Question:** behaviour of  $E_{\alpha,R}^{tot}(\theta)$  as  $R \rightarrow \infty$ ?

**Energy competition:**

- ▶  $\theta = 1 \implies$  no dislocation energy
- ▶  $\theta = \alpha^{-1} \implies$  no elastic energy
- ▶  $\theta \in (\alpha^{-1}, 1) \implies$  both present



$$(v := \alpha x, W(\alpha I) = 0)$$

# Asymptotic for $E_{\alpha,R}^{tot}$

Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser for  $E_{\alpha,R}^{tot}$  and define

$$\mathcal{E}^{el}(R) := \frac{\sigma^2}{\alpha^3 C^{el}} R, \quad \mathcal{E}^{pl}(R) := \sigma R^2 \left(1 - \frac{1}{\alpha^2}\right) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R.$$

# Asymptotic for $E_{\alpha,R}^{tot}$

Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser for  $E_{\alpha,R}^{tot}$  and define

$$\mathcal{E}^{el}(R) := \frac{\sigma^2}{\alpha^3 C^{el}} R, \quad \mathcal{E}^{pl}(R) := \sigma R^2 \left(1 - \frac{1}{\alpha^2}\right) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R.$$

## Theorem (F., Palombaro, Ponsiglione '15)

As  $R \rightarrow +\infty$  we have

$$E_{\alpha,R}^{el}(\theta_R) = \mathcal{E}^{el}(R) + O(R), \quad E_R^{pl}(\theta_R) = \mathcal{E}^{pl}(R) + O(R),$$

and therefore

$$E_{\alpha,R}^{tot} = \mathcal{E}^{el}(R) + \mathcal{E}^{pl}(R) + o(R).$$

# Asymptotic for $E_{\alpha,R}^{tot}$

Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser for  $E_{\alpha,R}^{tot}$  and define

$$\mathcal{E}^{el}(R) := \frac{\sigma^2}{\alpha^3 C^{el}} R, \quad \mathcal{E}^{pl}(R) := \sigma R^2 \left(1 - \frac{1}{\alpha^2}\right) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R.$$

## Theorem (F., Palombaro, Ponsiglione '15)

As  $R \rightarrow +\infty$  we have

$$E_{\alpha,R}^{el}(\theta_R) = \mathcal{E}^{el}(R) + O(R), \quad E_R^{pl}(\theta_R) = \mathcal{E}^{pl}(R) + O(R),$$

and therefore

$$E_{\alpha,R}^{tot} = \mathcal{E}^{el}(R) + \mathcal{E}^{pl}(R) + o(R).$$

In particular, for large  $R$ :

- ▶ *dislocations are energetically more favourable,*

# Asymptotic for $E_{\alpha,R}^{tot}$

Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser for  $E_{\alpha,R}^{tot}$  and define

$$\mathcal{E}^{el}(R) := \frac{\sigma^2}{\alpha^3 C^{el}} R, \quad \mathcal{E}^{pl}(R) := \sigma R^2 \left(1 - \frac{1}{\alpha^2}\right) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R.$$

## Theorem (F., Palombaro, Ponsiglione '15)

As  $R \rightarrow +\infty$  we have

$$E_{\alpha,R}^{el}(\theta_R) = \mathcal{E}^{el}(R) + O(R), \quad E_R^{pl}(\theta_R) = \mathcal{E}^{pl}(R) + O(R),$$

and therefore

$$E_{\alpha,R}^{tot} = \mathcal{E}^{el}(R) + \mathcal{E}^{pl}(R) + o(R).$$

In particular, for large  $R$ :

- ▶ dislocations are energetically more favourable,
- ▶ dislocation spacing (density) tends to  $\delta = \frac{b}{\alpha-1}$ ,

# Asymptotic for $E_{\alpha,R}^{tot}$

Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser for  $E_{\alpha,R}^{tot}$  and define

$$\mathcal{E}^{el}(R) := \frac{\sigma^2}{\alpha^3 C^{el}} R, \quad \mathcal{E}^{pl}(R) := \sigma R^2 \left(1 - \frac{1}{\alpha^2}\right) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R.$$

## Theorem (F., Palombaro, Ponsiglione '15)

As  $R \rightarrow +\infty$  we have

$$E_{\alpha,R}^{el}(\theta_R) = \mathcal{E}^{el}(R) + O(R), \quad E_R^{pl}(\theta_R) = \mathcal{E}^{pl}(R) + O(R),$$

and therefore

$$E_{\alpha,R}^{tot} = \mathcal{E}^{el}(R) + \mathcal{E}^{pl}(R) + o(R).$$

In particular, for large  $R$ :

- ▶ dislocations are energetically more favourable,
- ▶ dislocation spacing (density) tends to  $\delta = \frac{b}{\alpha-1}$ ,
- ▶ far field stress is relieved.

# Idea of the Proof

**Step 1.** Rescale the elastic energy

$$E_{\alpha,R}^{el}(\theta) = R^3 \theta^3 E_{\alpha,1}^{el}(\theta)$$



# Idea of the Proof

**Step 1.** Rescale the elastic energy

$$E_{\alpha,R}^{el}(\theta) = R^3 \theta^3 E_{\alpha,1}^{el}(\theta)$$

**Step 2.** Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser of  $E_{\alpha,R}^{tot}$ . Then, as  $R \rightarrow \infty$

$$E_{\alpha,1}^{el}(\theta_R) \rightarrow 0, \quad \theta_R \rightarrow \alpha^{-1}$$

# Idea of the Proof

**Step 1.** Rescale the elastic energy

$$E_{\alpha,R}^{el}(\theta) = R^3 \theta^3 E_{\alpha,1}^{el}(\theta)$$

**Step 2.** Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser of  $E_{\alpha,R}^{tot}$ . Then, as  $R \rightarrow \infty$

$$E_{\alpha,1}^{el}(\theta_R) \rightarrow 0, \quad \theta_R \rightarrow \alpha^{-1} \implies \text{Linearisation (about } \alpha l \text{)}$$

# Idea of the Proof

**Step 1.** Rescale the elastic energy

$$E_{\alpha,R}^{el}(\theta) = R^3 \theta^3 E_{\alpha,1}^{el}(\theta)$$

**Step 2.** Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser of  $E_{\alpha,R}^{tot}$ . Then, as  $R \rightarrow \infty$

$$E_{\alpha,1}^{el}(\theta_R) \rightarrow 0, \quad \theta_R \rightarrow \alpha^{-1} \implies \text{Linearisation (about } \alpha l)$$

**Step 3.** There exists  $C^{el} > 0$  such that, as  $R \rightarrow \infty$ ,

$$\frac{1}{(\theta_R^{-1} - \alpha)^2} E_{\alpha,1}^{el}(\theta_R) \rightarrow C^{el}$$

Dal Maso, Negri, Percivale. Set-Valued Analysis (2002).

# Idea of the Proof

**Step 1.** Rescale the elastic energy

$$E_{\alpha,R}^{el}(\theta) = R^3 \theta^3 E_{\alpha,1}^{el}(\theta)$$

**Step 2.** Let  $\theta_R \in [\alpha^{-1}, 1]$  be a minimiser of  $E_{\alpha,R}^{tot}$ . Then, as  $R \rightarrow \infty$

$$E_{\alpha,1}^{el}(\theta_R) \rightarrow 0, \quad \theta_R \rightarrow \alpha^{-1} \implies \text{Linearisation (about } \alpha I)$$

**Step 3.** There exists  $C^{el} > 0$  such that, as  $R \rightarrow \infty$ ,

$$\frac{1}{(\theta_R^{-1} - \alpha)^2} E_{\alpha,1}^{el}(\theta_R) \rightarrow C^{el}$$

**Step 4.** Write the elastic energy as a polynomial

$$E_{\alpha,R}^{el}(\theta_R) = R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2 \frac{1}{(\theta_R^{-1} - \alpha)^2} E_{\alpha,1}^{el}(\theta_R) = k_R^{el} R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2$$

where  $k_R^{el} := C^{el} + \varepsilon_R > 0$  and  $k_R^{el} \rightarrow C^{el}$ .

Dal Maso, Negri, Percivale. Set-Valued Analysis (2002).

# Idea of the Proof

**Step 5.** The total energy computed along  $\theta_R$  is equal to

$$E_{\alpha,R}^{tot}(\theta_R) = k_R^{el} R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta_R^2) \quad (1.1)$$

with  $\theta_R \rightarrow \alpha^{-1}$  minimisers and  $k_R^{el} \rightarrow C^{el}$ .

# Idea of the Proof

**Step 5.** The total energy computed along  $\theta_R$  is equal to

$$E_{\alpha,R}^{tot}(\theta_R) = k_R^{el} R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta_R^2) \quad (1.1)$$

with  $\theta_R \rightarrow \alpha^{-1}$  minimisers and  $k_R^{el} \rightarrow C^{el}$ .

**Step 6.** For a fixed parameter  $k > 0$ , introduce the family of polynomials

$$P_{R,k}(\theta) := k R^3 \theta^3 (\theta^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta^2)$$

# Idea of the Proof

**Step 5.** The total energy computed along  $\theta_R$  is equal to

$$E_{\alpha,R}^{\text{tot}}(\theta_R) = k_R^{\text{el}} R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta_R^2) \quad (1.1)$$

with  $\theta_R \rightarrow \alpha^{-1}$  minimisers and  $k_R^{\text{el}} \rightarrow C^{\text{el}}$ .

**Step 6.** For a fixed parameter  $k > 0$ , introduce the family of polynomials

$$P_{R,k}(\theta) := k R^3 \theta^3 (\theta^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta^2)$$

**Step 7.** Show that for  $R \gg 0$  there exists a unique minimiser  $\theta_R^m$  to

$$P_{R,k}(\theta_R^m) = \min_{\theta \in [\alpha^{-1}, 1]} P_{R,k}(\theta).$$

Moreover  $\theta_R^m \rightarrow \alpha^{-1}$ .

# Idea of the Proof

**Step 5.** The total energy computed along  $\theta_R$  is equal to

$$E_{\alpha,R}^{tot}(\theta_R) = k_R^{el} R^3 \theta_R^3 (\theta_R^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta_R^2) \quad (1.1)$$

with  $\theta_R \rightarrow \alpha^{-1}$  minimisers and  $k_R^{el} \rightarrow C^{el}$ .

**Step 6.** For a fixed parameter  $k > 0$ , introduce the family of polynomials

$$P_{R,k}(\theta) := k R^3 \theta^3 (\theta^{-1} - \alpha)^2 + \sigma R^2 (1 - \theta^2)$$

**Step 7.** Show that for  $R \gg 0$  there exists a unique minimiser  $\theta_R^m$  to

$$P_{R,k}(\theta_R^m) = \min_{\theta \in [\alpha^{-1}, 1]} P_{R,k}(\theta).$$

Moreover  $\theta_R^m \rightarrow \alpha^{-1}$ .

**Step 8.** Since  $\theta_R - \theta_R^m \rightarrow 0$ , by using (1.1), minimality, and computing  $P_{R,k}(\theta_R^m)$ , we have the thesis

$$E_{\alpha,R}^{tot}(\theta_R) = \underbrace{\frac{\sigma^2}{\alpha^3 C^{el}} R}_{\text{Elastic}} + \underbrace{\sigma R^2 (1 - \alpha^{-2}) - 2 \frac{\sigma^2}{\alpha^3 C^{el}} R}_{\text{Plastic}} + O(R).$$



# Conclusions and Perspectives

## Conclusions:

- ▶ A basic variational model describing the competition between the plastic energy spent at interfaces, and the corresponding release of bulk energy.

# Conclusions and Perspectives

## Conclusions:

- ▶ A basic variational model describing the competition between the plastic energy spent at interfaces, and the corresponding release of bulk energy.
- ▶ The size of the overlayer is a free parameter  $\implies$  free boundary problem, but only through the scalar parameter  $\theta$ .

# Conclusions and Perspectives


## Conclusions:

- ▶ A basic variational model describing the **competition between the plastic energy** spent at interfaces, and the corresponding release of **bulk energy**.
- ▶ The size of the overlayer is a free parameter  $\implies$  free boundary problem, but only through the scalar parameter  $\theta$ .

## Perspectives:

- ▶ **Grain boundaries**, the misfit between the crystal lattices are described by rotations rather than dilations.

Read, Shockley (1950) - Hirth, Carnahan (1992)



# Conclusions and Perspectives

## Conclusions:

- ▶ A basic variational model describing the **competition between the plastic energy** spent at interfaces, and the corresponding release of **bulk energy**.
- ▶ The size of the overlayer is a free parameter  $\implies$  free boundary problem, but only through the scalar parameter  $\theta$ .


## Perspectives:

- ▶ **Grain boundaries**, the misfit between the crystal lattices are described by rotations rather than dilations.

Read, Shockley (1950) - Hirth, Carnahan (1992)

- ▶ **Optimal geometry** for the dislocation net (square vs hexagonal)

Koslowski, Ortiz (2004)



# Presentation Plan

## ① Geometric Patterns of Dislocations


- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result


# Polycrystals

**Polycrystal:** formed by many grains, having the **same** lattice structure, mutually rotated  $\implies$  interface misfit at **grain boundaries**.



# Polycrystals

**Polycrystal:** formed by many grains, having the **same** lattice structure, mutually rotated  $\implies$  interface misfit at **grain boundaries**.




**Goal:** obtain polycrystalline structures as minimisers of some energy functional.

F., Palombaro, Ponsiglione. *Linearised Polycrystals from a 2D System of Edge Dislocations*. Preprint (2017)

# Tilt grain boundaries




**Tilt boundary:** small angle rotation  $\theta$  between grains





# Tilt grain boundaries



**Tilt boundary:** small angle rotation  $\theta$  between grains  $\Rightarrow$  **edge dislocations**.




# Tilt grain boundaries

**Tilt boundary:** small angle rotation  $\theta$  between grains  $\implies$  **edge dislocations**.

**Boundary structure:** periodic array of edge dislocations with spacing  $\delta = \frac{\varepsilon}{\theta}$ .




$$N_{\varepsilon} \approx \frac{1}{\varepsilon}$$



# Tilt grain boundaries

**Tilt boundary:** small angle rotation  $\theta$  between grains  $\Rightarrow$  **edge dislocations**.

**Boundary structure:** periodic array of edge dislocations with spacing  $\delta = \frac{\varepsilon}{\theta}$ .



Porter, Easterling. CRC Press (2009) - Gottstein. Springer (2013)

# Plan

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Let  $\mathcal{F}_\varepsilon$  be the energy of such system.

# Plan

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Let  $\mathcal{F}_\varepsilon$  be the energy of such system.

**Plan:**

- compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,

# Plan

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Let  $\mathcal{F}_\varepsilon$  be the energy of such system.

## Plan:

- ▶ compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,
- ▶ show that under suitable boundary conditions  $\mathcal{F}$  is minimised by polycrystals.

# Plan

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Let  $\mathcal{F}_\varepsilon$  be the energy of such system.

## Plan:

- ▶ compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,
- ▶ show that under suitable boundary conditions  $\mathcal{F}$  is minimised by polycrystals.


**Linearised polycrystals:** our energy regime will imply

$$N_\varepsilon \ll \frac{1}{\varepsilon}$$

$\implies$  we have less dislocations than tilt grain boundaries. However we still obtain polycrystalline minimisers, but with grains rotated by an infinitesimal angle  $\theta \approx 0$ .

# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.







# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .





# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .



# Setting (linearised planar elasticity)



**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$



# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.



**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$

**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .



# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .



**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$

**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .

**Strains:** inducing  $\mu$  are maps  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\mu) = 0, \quad \int_{\partial B_\varepsilon(x_i)} \beta \cdot t \, ds = \xi_i.$$



# Setting (linearised planar elasticity)

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$



**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .

**Strains:** inducing  $\mu$  are maps  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\mu) = 0, \quad \int_{\partial B_\varepsilon(x_i)} \beta \cdot t \, ds = \xi_i.$$

**Linearised Energy:**  $\mathbb{C}F : F \sim |F^{\text{sym}}|^2$ , then

$$E_\varepsilon(\mu, \beta) := \int_{\Omega_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx = \int_{\Omega} \mathbb{C}\beta : \beta \, dx.$$



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx$$



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx \geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho \end{aligned}$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$

The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$

The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$

**Remark:** let  $s \in (0, 1)$ , then

$$\int_{B_{\varepsilon^s} \setminus B_\varepsilon} |\beta|^2 dx \geq (1-s) \frac{|\xi|^2}{2\pi} |\log \varepsilon|$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$

The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$

**Remark:** let  $s \in (0, 1)$ , then

$$\int_{B_{\varepsilon^s} \setminus B_\varepsilon} |\beta|^2 dx \geq (1-s) \frac{|\xi|^2}{2\pi} |\log \varepsilon|$$

**Self-energy:** is of order  $|\log \varepsilon|$  and concentrated in a small region around  $B_\varepsilon$ .


# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .

# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.






# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$



**Hypothesis on HC Radius:** as  $\varepsilon \rightarrow 0$

►  $\rho_\varepsilon / \varepsilon^s \rightarrow \infty, \quad \forall s \in (0, 1),$  (HC contains almost all the self-energy)

# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$



**Hypothesis on HC Radius:** as  $\varepsilon \rightarrow 0$

►  $\rho_\varepsilon / \varepsilon^s \rightarrow \infty, \quad \forall s \in (0, 1),$

(HC contains almost all the self-energy)

►  $N_\varepsilon \rho_\varepsilon^2 \rightarrow 0.$

(Measure of HC region vanishes)

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|,$$

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|,$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta : \beta \, dx.$$

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|,$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta : \beta \, dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

►  $N_\varepsilon \ll |\log \varepsilon| \leadsto$  Dilute dislocations



# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|,$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta : \beta \, dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

- ▶  $N_\varepsilon \ll |\log \varepsilon| \leadsto$  Dilute dislocations
- ▶  $N_\varepsilon \approx |\log \varepsilon| \leadsto$  Critical regime

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|,$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta : \beta \, dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

►  $N_\varepsilon \ll |\log \varepsilon| \leadsto$  Dilute dislocations

►  $N_\varepsilon \approx |\log \varepsilon| \leadsto$  Critical regime

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

►  $N_\varepsilon \gg |\log \varepsilon| \leadsto$  Super-critical regime

F., Palombaro, Ponsiglione. *Linearised Polycrystals from a 2D System of Edge Dislocations*.  
Preprint (2017)

## Candidate $\Gamma$ -limit

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^N \xi_i \delta_{x_i}$  be such that “ $\text{Curl } \beta = \mu$ ”.

## Candidate $\Gamma$ -limit

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^N \xi_i \delta_{x_i}$  be such that “**Curl**  $\beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^N B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx.$$

## Candidate $\Gamma$ -limit

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^N \xi_i \delta_{x_i}$  be such that “**Curl**  $\beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^N B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx.$$

**$\Gamma$ -limit:**  $S \in L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$ ,  $A \in L^2(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ,  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2)$  with **Curl**  $A = \mu$ ,

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

## Candidate $\Gamma$ -limit

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^N \xi_i \delta_{x_i}$  be such that “**Curl**  $\beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^N B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx.$$

**$\Gamma$ -limit:**  $S \in L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$ ,  $A \in L^2(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ,  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2)$  with **Curl**  $A = \mu$ ,

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Density  $\varphi$ :** the self-energy for a single dislocation core  $\xi \delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \min_{\beta} \left\{ \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi \delta_0 \text{”} \right\}.$$

# Candidate $\Gamma$ -limit

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^N \xi_i \delta_{x_i}$  be such that “ $\text{Curl } \beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^N B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx.$$

**$\Gamma$ -limit:**  $S \in L^2(\Omega; \mathbb{M}_{\text{sym}}^{2 \times 2})$ ,  $A \in L^2(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ,  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2)$  with  $\text{Curl } A = \mu$ ,

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Density  $\varphi$ :** the self-energy for a single dislocation core  $\xi \delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \min_{\beta} \left\{ \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi \delta_0 \text{”} \right\}.$$

Define  $\varphi : \mathbb{R}^2 \rightarrow [0, \infty)$  as the relaxation of  $\psi$  (splitting multiple dislocations)

$$\varphi(\xi) := \min \left\{ \sum_{i=1}^M \lambda_i \psi(\xi_i) : \xi = \sum_{i=1}^M \lambda_i \xi_i, M \in \mathbb{N}, \lambda_i \geq 0, \xi_i \in \mathbb{S} \right\}.$$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$



# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

$$\blacktriangleright \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A \text{ in } L^2(\Omega; \mathbb{M}^{2 \times 2}),$$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A \text{ in } L^2(\Omega; \mathbb{M}^{2 \times 2}),$
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu \text{ in } \mathcal{M}(\Omega; \mathbb{R}^2),$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. " $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ " and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  and  $\text{Curl } A = \mu$ .

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. "**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ " and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  and **Curl**  $A = \mu$ .

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  and **Curl**  $A = \mu$ .

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

### Remark:

- ▶  $S$  and  $A$  live on two different scales with  $S \ll A \implies$  terms in  $\mathcal{F}$  decoupled.

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  and **Curl**  $A = \mu$ .

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

### Remark:

- ▶  $S$  and  $A$  live on two different scales with  $S \ll A \implies$  terms in  $\mathcal{F}$  decoupled.
- ▶ In the critical regime  $N_\varepsilon \approx |\log \varepsilon|$  we have  $S \approx A$  and  $\text{Curl}(S + A) = \mu$ .

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$C \geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n)$$



# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$C \geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho \varepsilon_n}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) dx$$

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$\begin{aligned} C &\geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho_{\varepsilon_n}}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) \, dx \\ &\geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \psi_{\varepsilon_n}(\xi_{n,i}) \end{aligned}$$

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$\begin{aligned} C &\geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho_{\varepsilon_n}}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) dx \\ &\geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \psi_{\varepsilon_n}(\xi_{n,i}) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \psi_{\varepsilon_n} \left( \frac{\xi_{n,i}}{|\xi_{n,i}|} \right) \end{aligned}$$

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$\begin{aligned} C &\geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho_{\varepsilon_n}}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) dx \\ &\geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \psi_{\varepsilon_n}(\xi_{n,i}) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \psi_{\varepsilon_n} \left( \frac{\xi_{n,i}}{|\xi_{n,i}|} \right) \geq \frac{c}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \end{aligned}$$

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$\begin{aligned} C &\geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho \varepsilon_n}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) dx \\ &\geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \psi_{\varepsilon_n}(\xi_{n,i}) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \psi_{\varepsilon_n} \left( \frac{\xi_{n,i}}{|\xi_{n,i}|} \right) \geq \frac{c}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \\ &\geq \frac{c}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \end{aligned}$$

# Compactness of the measures

Let  $\mu_n := \sum_{i=1}^{M_n} \xi_{n,i} \delta_{x_{n,i}}$  and “ $\text{Curl } \beta_n = \mu_n$ ”. We show that

$$\frac{1}{N_{\varepsilon_n}} |\mu_n|(\Omega) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| \leq C, \quad (1.2)$$

so that  $\frac{\mu_n}{N_{\varepsilon_n}} \xrightarrow{*} \nu$ .

$$\begin{aligned} C &\geq \mathcal{F}_{\varepsilon_n}(\mu_n, \beta_n) \geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \frac{1}{|\log \varepsilon_n|} \int_{B_{\rho \varepsilon_n}(x_{n,i}) \setminus B_{\varepsilon_n}(x_{n,i})} W(\beta_n) dx \\ &\geq \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} \psi_{\varepsilon_n}(\xi_{n,i}) = \frac{1}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \psi_{\varepsilon_n} \left( \frac{\xi_{n,i}}{|\xi_{n,i}|} \right) \geq \frac{c}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}|^2 \\ &\geq \frac{c}{N_{\varepsilon_n}} \sum_{i=1}^{M_n} |\xi_{n,i}| = c \frac{|\mu_n|(\Omega)}{N_{\varepsilon_n}} \implies (1.2) \end{aligned}$$

# Compactness of the strains

## Symmetric Part:

$$CN_{\varepsilon_n} |\log \varepsilon_n| \geq CE_{\varepsilon_n}(\mu_n, \beta_n) \geq C \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx$$

# Compactness of the strains

## Symmetric Part:

$$CN_{\varepsilon_n} |\log \varepsilon_n| \geq CE_{\varepsilon_n}(\mu_n, \beta_n) \geq C \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx \implies \frac{\beta_n^{\text{sym}}}{\sqrt{N_{\varepsilon_n} |\log \varepsilon_n|}} \rightharpoonup S.$$



# Compactness of the strains

## Symmetric Part:

$$CN_{\varepsilon_n} |\log \varepsilon_n| \geq CE_{\varepsilon_n}(\mu_n, \beta_n) \geq C \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx \implies \frac{\beta_n^{\text{sym}}}{\sqrt{N_{\varepsilon_n} |\log \varepsilon_n|}} \rightharpoonup S.$$

**Skew Part:** since “ $\text{Curl } \beta_n = \mu_n$ ” we can apply the generalised **Korn inequality**:

$$\int_{\Omega} |\beta_n^{\text{skew}}|^2 dx \leq C \left( \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx + (|\mu_n|(\Omega))^2 \right) \quad (\text{Gen. Korn})$$

$$(N_{\varepsilon} \gg |\log \varepsilon|)$$

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

# Compactness of the strains

## Symmetric Part:

$$CN_{\varepsilon_n} |\log \varepsilon_n| \geq CE_{\varepsilon_n}(\mu_n, \beta_n) \geq C \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx \implies \frac{\beta_n^{\text{sym}}}{\sqrt{N_{\varepsilon_n} |\log \varepsilon_n|}} \rightharpoonup S.$$

**Skew Part:** since “ $\text{Curl } \beta_n = \mu_n$ ” we can apply the generalised **Korn inequality**:

$$\begin{aligned} \int_{\Omega} |\beta_n^{\text{skew}}|^2 dx &\leq C \left( \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx + (|\mu_n|(\Omega))^2 \right) && \text{(Gen. Korn)} \\ &\leq C \left( \sqrt{N_{\varepsilon_n} |\log \varepsilon_n|} + N_{\varepsilon_n}^2 \right) \leq CN_{\varepsilon_n}^2 && (N_{\varepsilon} \gg |\log \varepsilon|) \end{aligned}$$

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

# Compactness of the strains

## Symmetric Part:

$$CN_{\varepsilon_n} |\log \varepsilon_n| \geq CE_{\varepsilon_n}(\mu_n, \beta_n) \geq C \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx \implies \frac{\beta_n^{\text{sym}}}{\sqrt{N_{\varepsilon_n} |\log \varepsilon_n|}} \rightharpoonup S.$$

**Skew Part:** since “ $\text{Curl } \beta_n = \mu_n$ ” we can apply the generalised Korn inequality:

$$\begin{aligned} \int_{\Omega} |\beta_n^{\text{skew}}|^2 dx &\leq C \left( \int_{\Omega} |\beta_n^{\text{sym}}|^2 dx + (|\mu_n|(\Omega))^2 \right) && \text{(Gen. Korn)} \\ &\leq C \left( \sqrt{N_{\varepsilon_n} |\log \varepsilon_n|} + N_{\varepsilon_n}^2 \right) \leq CN_{\varepsilon_n}^2 && (N_{\varepsilon} \gg |\log \varepsilon|) \end{aligned}$$

so that  $\frac{\beta_n^{\text{skew}}}{N_{\varepsilon_n}} \rightharpoonup A.$

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations.*  
J. Eur. Math. Soc. (JEMS) (2010)

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Admissible dislocations:** measures  $\mu$  satisfying

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds. \quad (\text{GND})$$

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Admissible dislocations:** measures  $\mu$  satisfying

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds. \quad (\text{GND})$$

**Admissible strains:**  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  such that “ $\text{Curl } \beta = \mu$ ” and

$$\beta \cdot t = g_\varepsilon \cdot t \quad \text{on } \partial\Omega.$$

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Admissible dislocations:** measures  $\mu$  satisfying

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds. \quad (\text{GND})$$

**Admissible strains:**  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  such that “ $\text{Curl } \beta = \mu$ ” and

$$\beta \cdot t = g_\varepsilon \cdot t \quad \text{on } \partial\Omega.$$

**$\Gamma$ -limit:** the usual energy  $\mathcal{F}_\varepsilon$   $\Gamma$ -converges to

$$\mathcal{F}_{\text{BC}}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) \, ds,$$

such that  $\text{Curl } A = \mu$ , with  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2) \cap H^{-1}(\Omega; \mathbb{R}^2)$ .

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Admissible dislocations:** measures  $\mu$  satisfying

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds. \quad (\text{GND})$$

**Admissible strains:**  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  such that “ $\text{Curl } \beta = \mu$ ” and

$$\beta \cdot t = g_\varepsilon \cdot t \quad \text{on } \partial\Omega.$$

**$\Gamma$ -limit:** the usual energy  $\mathcal{F}_\varepsilon$   $\Gamma$ -converges to

$$\mathcal{F}_{\text{BC}}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) \, ds,$$

such that  $\text{Curl } A = \mu$ , with  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2) \cap H^{-1}(\Omega; \mathbb{R}^2)$ .

**Remark:**  $\beta_\varepsilon^{\text{sym}} \ll \beta_\varepsilon^{\text{skew}} \implies$  BC pass to the limit only for  $A$ .



# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

**Piecewise constant BC:** Fix a piecewise constant BC

$$g_A := \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}, \quad a := \sum_{k=1}^M m_k \chi_{U_k},$$

with  $m_k < m_{k+1}$  and  $\{U_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .

# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

**Piecewise constant BC:** Fix a piecewise constant BC

$$g_A := \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}, \quad a := \sum_{k=1}^M m_k \chi_{U_k},$$

with  $m_k < m_{k+1}$  and  $\{U_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .

## Problem

*Minimise*

$$\mathcal{F}_{\text{BC}}(\mu, 0, A) = \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds,$$

with  $\text{Curl } A = \mu$  and  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2) \cap H^{-1}(\Omega; \mathbb{R}^2)$ .


# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a *piecewise constant boundary condition*  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{\text{BC}}(\mu, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .  
We interpret  $A$  as a *linearised polycrystal*.




# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{\text{BC}}(\mu, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .  
We interpret  $A$  as a *linearised polycrystal*.



**Open Question:** Are all minimisers piecewise constant? Uniqueness?


# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a **piecewise constant** minimiser of  $\mathcal{F}_{BC}(\mu, 0, A)$


$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .  
We interpret  $A$  as a **linearised polycrystal**.



**Open Question:** Are all minimisers piecewise constant? Uniqueness?

**Essential:** that the boundary condition is piecewise affine on the **whole**  $\partial\Omega$ .



# Idea of the proof

**Problem:** given a piecewise constant BC  $g_A$ , consider

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds : \text{Curl } A = \mu \in \mathcal{M} \cap H^{-1} \right\}.$$

# Idea of the proof

**Problem:** given a piecewise constant BC  $g_A$ , consider

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds : \text{Curl } A = \mu \in \mathcal{M} \cap H^{-1} \right\}.$$

Since  $A$  and  $g_A$  are antisymmetric,  $\exists u, a \in L^2(\Omega)$  s.t.

$$A = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad g_A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}.$$



# Idea of the proof

**Problem:** given a piecewise constant BC  $g_A$ , consider

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds : \text{Curl } A = \mu \in \mathcal{M} \cap H^{-1} \right\}.$$

Since  $A$  and  $g_A$  are antisymmetric,  $\exists u, a \in L^2(\Omega)$  s.t.

$$A = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad g_A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}.$$

**Note:**  $\text{Curl } A = Du \in \mathcal{M}(\Omega; \mathbb{R}^2) \implies u \in BV(\Omega)$

# Idea of the proof

**Problem:** given a piecewise constant BC  $g_A$ , consider

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds : \text{Curl } A = \mu \in \mathcal{M} \cap H^{-1} \right\}.$$

Since  $A$  and  $g_A$  are antisymmetric,  $\exists u, a \in L^2(\Omega)$  s.t.

$$A = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad g_A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}.$$

**Note:**  $\text{Curl } A = Du \in \mathcal{M}(\Omega; \mathbb{R}^2) \implies u \in BV(\Omega) \implies$  **Equivalent Problem:**

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{dDu}{d|Du|} \right) d|Du| + \int_{\partial\Omega} \varphi((u - a)\nu) ds : u \in BV(\Omega) \right\}. \quad (1.3)$$

## Idea of the proof

**Problem:** given a piecewise constant BC  $g_A$ , consider

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds : \text{Curl } A = \mu \in \mathcal{M} \cap H^{-1} \right\}.$$

Since  $A$  and  $g_A$  are antisymmetric,  $\exists u, a \in L^2(\Omega)$  s.t.

$$A = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad g_A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}.$$

**Note:**  $\text{Curl } A = Du \in \mathcal{M}(\Omega; \mathbb{R}^2) \implies u \in BV(\Omega) \implies$  **Equivalent Problem:**

$$\inf \left\{ \int_{\Omega} \varphi \left( \frac{dDu}{d|Du|} \right) d|Du| + \int_{\partial\Omega} \varphi((u - a)\nu) ds : u \in BV(\Omega) \right\}. \quad (1.3)$$

**Proof:** let  $\tilde{u}$  be a minimiser for (1.3). By anisotropic Coarea Formula

$$\int_{\Omega} \varphi \left( \frac{dD\tilde{u}}{d|D\tilde{u}|} \right) d|D\tilde{u}| = \int_{\mathbb{R}} \text{Per}_{\varphi}(\{x \in \Omega : \tilde{u}(x) > t\}) dt,$$

we can select the levels with minimal perimeter. This defines the Caccioppoli partition.

# Comparison with classical Read-Shockley formula

**Read-Shockley formula:** Elastic energy =  $E_0\theta(1 + |\log \theta|)$ .

- ▶ This energy corresponds to small rotations  $\theta$  between grains: small rotations but larger than linearised rotations.

# Comparison with classical Read-Shockley formula

**Read-Shockley formula:** Elastic energy =  $E_0\theta(1 + |\log \theta|)$ .

- ▶ This energy corresponds to small rotations  $\theta$  between grains: small rotations but larger than linearised rotations.
- ▶ It is a nonlinear formula that corresponds to a higher energy regime.

# Comparison with classical Read-Shockley formula

**Read-Shockley formula:** Elastic energy =  $E_0\theta(1 + |\log \theta|)$ .

- ▶ This energy corresponds to small rotations  $\theta$  between grains: small rotations but larger than linearised rotations.
- ▶ It is a nonlinear formula that corresponds to a higher energy regime.
- ▶ The density of dislocations to obtain small rotations is

$$\text{Density} \approx \frac{1}{\varepsilon} \gg N_\varepsilon.$$

# Comparison with classical Read-Shockley formula

**Read-Shockley formula:** Elastic energy =  $E_0\theta(1 + |\log \theta|)$ .

- ▶ This energy corresponds to small rotations  $\theta$  between grains: small rotations but larger than linearised rotations.
- ▶ It is a nonlinear formula that corresponds to a higher energy regime.
- ▶ The density of dislocations to obtain small rotations is

$$\text{Density} \approx \frac{1}{\varepsilon} \gg N_\varepsilon.$$

**Question:**  $\Gamma$ -convergence analysis of the Read-Shockley formula?

Lauteri, Luckhaus. *An energy estimate for dislocation configurations and the emergence of Cosserat-type structures in metal plasticity*. Preprint (2017)

# Comparison with classical Read-Shockley formula

**Read-Shockley formula:** Elastic energy =  $E_0\theta(1 + |\log \theta|)$ .

- ▶ This energy corresponds to small rotations  $\theta$  between grains: small rotations but larger than linearised rotations.
- ▶ It is a nonlinear formula that corresponds to a higher energy regime.
- ▶ The density of dislocations to obtain small rotations is

$$\text{Density} \approx \frac{1}{\varepsilon} \gg N_\varepsilon.$$

**Question:**  $\Gamma$ -convergence analysis of the Read-Shockley formula?

Lauteri, Luckhaus. *An energy estimate for dislocation configurations and the emergence of Cosserat-type structures in metal plasticity*. Preprint (2017)

**Question:** Are there some relevant energy regimes in between?



# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Perspectives:

- ▶ **Uniqueness** of piecewise constant minimisers?

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Perspectives:

- ▶ **Uniqueness** of piecewise constant minimisers?
- ▶ Comparison with the **Read-Shockley formula**?  
Lauteri, Luckhaus. Preprint (2017).

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Perspectives:

- ▶ **Uniqueness** of piecewise constant minimisers?
- ▶ Comparison with the **Read-Shockley formula**?
- ▶ **Dynamics** for linearised polycrystals?

Taylor. *Crystalline variational problems*. Bull. Amer. Math. Soc. (1978).

Chambolle, Morini, Ponsiglione. *Existence and Uniqueness for a Crystalline Mean Curvature Flow*. Comm. Pure Appl. Math (2017).

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Perspectives:

- ▶ **Uniqueness** of piecewise constant minimisers?
- ▶ Comparison with the **Read-Shockley formula**?  
Lauteri, Luckhaus. Preprint (2017).
- ▶ **Dynamics** for linearised polycrystals?  
Taylor. *Crystalline variational problems*. Bull. Amer. Math. Soc. (1978).  
Chambolle, Morini, Ponsiglione. *Existence and Uniqueness for a Crystalline Mean Curvature Flow*. Comm. Pure Appl. Math (2017).
- ▶ Supercritical regime analysis starting from a **non-linear energy**?  
Müller, Scardia, Zeppieri. *Geometric rigidity for incompatible fields and an application to strain-gradient plasticity*. Indiana University Mathematics Journal (2014).

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result



# Gradient integrability for solutions to elliptic equations

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

# Gradient integrability for solutions to elliptic equations

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

## Problem

Study the gradient integrability of distributional solutions  $u \in W^{1,1}(\Omega)$  to

$$\operatorname{div}(\sigma \nabla u) = 0, \tag{2.1}$$

when

$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

with  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  constant elliptic matrices,  $\{E_1, E_2\}$  measurable partition of  $\Omega$ .

# Gradient integrability for solutions to elliptic equations

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

## Problem

Study the gradient integrability of distributional solutions  $u \in W^{1,1}(\Omega)$  to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad (2.1)$$

when

$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

with  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  constant elliptic matrices,  $\{E_1, E_2\}$  measurable partition of  $\Omega$ .

## Application to composites:

- ▶  $\Omega$  is a section of a **composite conductor** obtained by mixing two materials with **conductivities**  $\sigma_1$  and  $\sigma_2$ ,
- ▶ the **electric field**  $\nabla u$  solves (2.1),
- ▶ concentration of  $\nabla u$  in relation to the geometry  $\{E_1, E_2\}$ .

# Astala's Theorem



## Theorem (Astala '94)


Let  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  be uniformly elliptic. There exists exponents  $1 < q < 2 < p$  such that if  $u \in W^{1,q}(\Omega)$  solves

$$\operatorname{div}(\sigma \nabla u) = 0,$$

then  $\nabla u \in L^p_{\text{weak}}(\Omega; \mathbb{R}^2)$ .

*Astala. Area distortion of quasiconformal mappings. Acta Mathematica (1994)*

# Astala's Theorem



## Theorem (Astala '94)

Let  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  be uniformly elliptic. There exists exponents  $1 < q < 2 < p$  such that if  $u \in W^{1,q}(\Omega)$  solves

$$\operatorname{div}(\sigma \nabla u) = 0,$$

then  $\nabla u \in L^p_{\text{weak}}(\Omega; \mathbb{R}^2)$ .


## Question

Are the exponents  $q$  and  $p$  optimal among two-phase elliptic conductivities

$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2}?$$

Astala. *Area distortion of quasiconformal mappings*. Acta Mathematica (1994)


# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).


# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).

# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

**Remark:** it is sufficient to prove optimality in the case


$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where

$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$



# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

**Remark:** it is sufficient to prove optimality in the case

$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where


$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$

The corresponding critical exponents for Astala's theorem are

$$q_{\sigma_1, \sigma_2} = \frac{2K}{K+1}, \quad p_{\sigma_1, \sigma_2} = \frac{2K}{K-1}.$$

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).

# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

**Remark:** it is sufficient to prove optimality in the case

$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where


$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$

The corresponding critical exponents for Astala's theorem are

$$q_{\sigma_1, \sigma_2} = \frac{2K}{K+1}, \quad p_{\sigma_1, \sigma_2} = \frac{2K}{K-1}.$$

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).

# Upper exponent optimality



## Theorem (Nesi, Palombaro, Ponsiglione '14)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .


❶ If  $\sigma \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and  $u \in W^{1, \frac{2K}{K+1}}(\Omega)$  solves

$$\text{div}(\sigma \nabla u) = 0 \quad (2.2)$$

then  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

❷ There exists  $\bar{\sigma} \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and a weak solution  $\bar{u} \in W^{1,2}(\Omega)$  to (2.2) with  $\sigma = \bar{\sigma}$ , satisfying affine boundary conditions and such that  $\nabla \bar{u} \notin L^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

# Upper exponent optimality



## Theorem (Nesi, Palombaro, Ponsiglione '14)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .

❶ If  $\sigma \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and  $u \in W^{1, \frac{2K}{K+1}}(\Omega)$  solves

$$\text{div}(\sigma \nabla u) = 0 \quad (2.2)$$


then  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

❷ There exists  $\bar{\sigma} \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and a weak solution  $\bar{u} \in W^{1,2}(\Omega)$  to (2.2) with  $\sigma = \bar{\sigma}$ , satisfying affine boundary conditions and such that  $\nabla \bar{u} \notin L^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

## Question we address

Is the lower exponent  $\frac{2K}{K+1}$  optimal?

# Lower exponent optimality



## Theorem (F., Palombaro '17)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .  
There exist

- ▶ coefficients  $\sigma_n \in L^\infty(\Omega; \{\sigma_1; \sigma_2\})$ ,
- ▶ exponents  $p_n \in \left[1, \frac{2K}{K+1}\right]$ ,
- ▶ functions  $u_n \in W^{1,1}(\Omega)$  such that  $u_n(x) = x_1$  on  $\partial\Omega$ ,

such that

$$\begin{aligned} \operatorname{div}(\sigma_n \nabla u_n) &= 0, \\ \nabla u_n &\in L_{\text{weak}}^{p_n}(\Omega; \mathbb{R}^2), \quad p_n \rightarrow \frac{2K}{K+1}, \quad \nabla u_n \notin L^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2). \end{aligned}$$

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$



# Solving differential inclusions

## Theorem (Approximate solutions for two phases)


Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- rank-one connection allows to laminate in direction  $n$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,
- ▶  $\lambda$  proportion for  $A$ ,  $1 - \lambda$  proportion for  $B$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,
- ▶  $\lambda$  proportion for  $A$ ,  $1 - \lambda$  proportion for  $B$ ,
- ▶ this allows to recover boundary data  $C$ .



Müller. *Variational models for microstructure and phase transitions*.

# Laminates of first order

$\mathcal{L}_{\Omega}^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_{\Omega}^2(B) := |B \cap \Omega|/|\Omega|$ .

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$



# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- ▶ **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.
- ▶ **Boundary condition** since the barycentre of  $\nu$  is  $\bar{\nu} := \int_{\mathbb{M}^{2 \times 2}} M d\nu(M) = C$ .

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- ▶ **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.
- ▶ **Boundary condition** since the barycentre of  $\nu$  is  $\bar{\nu} := \int_{\mathbb{M}^{2 \times 2}} M d\nu(M) = C$ .
- ▶ **Integrability** since for  $p > 1$  we have

$$\frac{1}{|\Omega|} \int_\Omega |\nabla f_\delta|^p dx = \int_{\mathbb{M}^{2 \times 2}} |M|^p d\nu_\delta(M).$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

**New gradient:** apply previous Proposition to the set  $\{x \in \Omega: \nabla f \sim B\}$  to obtain  $\tilde{f}: \Omega \rightarrow \mathbb{R}^2$  such that  $\tilde{f}(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla \tilde{f}, \{A, B_1, B_2\}) < \delta \quad \text{a.e. in } \Omega.$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

**New gradient:** apply previous Proposition to the set  $\{x \in \Omega: \nabla f \sim B\}$  to obtain  $\tilde{f}: \Omega \rightarrow \mathbb{R}^2$  such that  $\tilde{f}(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla \tilde{f}, \{A, B_1, B_2\}) < \delta \quad \text{a.e. in } \Omega.$$

The gradient distribution of  $\tilde{f}$  is given by

$$\nu = \lambda \delta_A + (1 - \lambda)\mu \delta_{B_1} + (1 - \lambda)(1 - \mu) \delta_{B_2}.$$

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.



# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,
- ▶  $f(x) = Ax$  on  $\partial\Omega$ ,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,
- ▶  $f(x) = Ax$  on  $\partial\Omega$ ,
- ▶  $|\{x \in \Omega : |\nabla f(x) - A_i| < \delta\}| = \lambda_i |\Omega|$ .

# Presentation Plan

## ① Geometric Patterns of Dislocations

- ▶ Dislocations
- ▶ Semi-coherent interfaces
- ▶ Linearised polycrystals

## ② Microgeometries in Composites

- ▶ Critical lower integrability
- ▶ Convex integration
- ▶ Proof of our main result

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- ① Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (2.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.



# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- 1 Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (2.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- 2 Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- 1 Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (2.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- 2 Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.
- 3 Convex integration Proposition  $\implies$  construct  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ .  
In this way  $f$  solves (2.3) and

$$\nabla f \in L_{\text{weak}}^q(\Omega; \mathbb{R}^2), \quad q \in \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right], \quad \nabla f \notin L^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2).$$

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- 1 Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (2.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- 2 Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.
- 3 Convex integration Proposition  $\implies$  construct  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ .  
In this way  $f$  solves (2.3) and

$$\nabla f \in L_{\text{weak}}^q(\Omega; \mathbb{R}^2), \quad q \in \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right], \quad \nabla f \notin L^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2).$$

These methods were developed for isotropic conductivities  $\sigma \in L^\infty(\Omega; \{KI, \frac{1}{K}I\})$ .

The adaptation to our case is non-trivial because of the lack of symmetry of the target set  $T$ , due to the anisotropy of  $\sigma_1$  and  $\sigma_2$ .

Astala, Faraco, Székelyhidi. *Convex integration and the  $L^p$  theory of elliptic equations*.

Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2008)

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

## Lemma (F., Palombaro '17)

A function  $u \in W^{1,1}(\Omega)$  is solution to

$$\text{div}(\sigma \nabla u) = 0$$

iff there exists  $v \in W^{1,1}(\Omega)$  such that  $f = (u, v): \Omega \rightarrow \mathbb{R}^2$  satisfies

$$\nabla f(x) \in T_1 \cup T_2 \quad \text{in } \Omega.$$

Moreover  $E_1 = \{x \in \Omega: \nabla f(x) \in T_1\}$  and  $E_2 = \{x \in \Omega: \nabla f(x) \in T_2\}$ .

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

## Lemma (F., Palombaro '17)

A function  $u \in W^{1,1}(\Omega)$  is solution to

$$\text{div}(\sigma \nabla u) = 0$$

iff there exists  $v \in W^{1,1}(\Omega)$  such that  $f = (u, v): \Omega \rightarrow \mathbb{R}^2$  satisfies

$$\nabla f(x) \in T_1 \cup T_2 \quad \text{in } \Omega.$$

Moreover  $E_1 = \{x \in \Omega: \nabla f(x) \in T_1\}$  and  $E_2 = \{x \in \Omega: \nabla f(x) \in T_2\}$ .

**Key Remark:**  $u$  and  $f$  enjoy the **same** integrability properties.

# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \overline{w}, \quad \forall w \in \mathbb{C}.$$

# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \overline{w}, \quad \forall w \in \mathbb{C}.$$

The sets of conformal linear maps and anti-conformal linear maps are

$$E_0 := \{(z, 0) : z \in \mathbb{C}\} \quad (\text{Conformal maps})$$

$$E_\infty := \{(0, z) : z \in \mathbb{C}\} \quad (\text{Anti-conformal maps})$$



# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \bar{w}, \quad \forall w \in \mathbb{C}.$$

The sets of conformal linear maps and anti-conformal linear maps are

$$E_0 := \{(z, 0) : z \in \mathbb{C}\} \quad (\text{Conformal maps})$$

$$E_\infty := \{(0, z) : z \in \mathbb{C}\} \quad (\text{Anti-conformal maps})$$

**Target sets** in conformal coordinates are

$$T_1 = \{(a, d_1(\bar{a})) : a \in \mathbb{C}\}, \quad T_2 = \{(a, -d_2(\bar{a})) : a \in \mathbb{C}\},$$


where the operators  $d_j: \mathbb{C} \rightarrow \mathbb{C}$  are defined as

$$d_j(a) := k \operatorname{Re} a + i s_j \operatorname{Im} a, \quad \text{with} \quad k := \frac{K-1}{K+1} \quad \text{and} \quad s_j := \frac{S_j-1}{S_j+1}.$$

# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$JR_\theta =$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$$JR_\theta = \lambda_1 A_1 + (1 - \lambda_1) P_1$$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned}
 JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\
 &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\
 &\leadsto \nu_1
 \end{aligned}$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$2JR_\theta = \lambda_2 A_2 + (1 - \lambda_2) P_2$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.



# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

►  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$





# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$



# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$



# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^{p(\theta)} d\nu_n(M) \rightarrow \infty$  as  $n \rightarrow \infty$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.


- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^{p(\theta)} d\nu_n(M) \rightarrow \infty$  as  $n \rightarrow \infty$

**Remark:** barycentre  $J$  gives the right growth.



# Constructing approximate solutions


We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that

- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,






# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that

- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,
- ▶  $\nabla f \notin L^{\frac{2K}{K+1}}$ .




# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,
- ▶  $\nabla f \notin L^{\frac{2K}{K+1}}$ .

**Idea:** alternate one step of the staircase laminate with the convex integration Proposition.



# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$



# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$





# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .


**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$





# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$


**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $v_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } v_1 \implies \nabla f_2$  grows like  $p_1$


This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $v_2$  with  $\bar{v}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } v_2 \implies \nabla g$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $v_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } v_1 \implies \nabla f_2$  grows like  $p_1$


This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $v_2$  with  $\bar{v}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } v_2 \implies \nabla g$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $v_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } v_1 \implies \nabla f_2$  grows like  $p_1$


This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $v_2$  with  $\bar{v}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } v_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$



**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

► Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$

► Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



►  $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

► Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$

► Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

► Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$

► Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$

►  $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$



**Step 2.** Define map  $f_3$  by modifying  $f_2$

► Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$

► Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$

**Iterating:**  $\rightsquigarrow f_n$  obtained by successive modifications  
on nested sets going to zero in measure  $\implies f_n \rightarrow f$



# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.



# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (2.4) and s.t.  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .

Modifying staircase laminate?

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (2.4) and s.t.  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .

Modifying staircase laminate?

- Extend these results to **three-phase** conductivities  $\sigma \in \{\sigma_1, \sigma_2, \sigma_3\}$ .

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (2.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- ▶ Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- ▶ We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- ▶ **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (2.4) and s.t.  $\nabla u \in L^{\frac{2K}{K+1}}_{\text{weak}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .

Modifying staircase laminate?

- ▶ Extend these results to **three-phase** conductivities  $\sigma \in \{\sigma_1, \sigma_2, \sigma_3\}$ .
- ▶ **Dimension  $d \geq 3$ ?** Even only in the isotropic case  $\sigma \in \{KI, K^{-1}I\}$  for  $K > 1$ .  
Main difficulty: Astala's Theorem is missing in higher dimensions.

