

# Linearised Polycrystals from a 2D System of Edge Dislocations

Silvio Fanzon


in collaboration with

M. Palombaro and M. Ponsiglione

Graz, 31st January 2018

# Classical Elasticity: ideal crystals



**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded



# Classical Elasticity: ideal crystals

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$





# Classical Elasticity: ideal crystals

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$




# Classical Elasticity: ideal crystals

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Displacement:** map  $u: \Omega \rightarrow \mathbb{R}^3$  s.t.  $v = x + u$



# Classical Elasticity: ideal crystals

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$


**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Displacement:** map  $u: \Omega \rightarrow \mathbb{R}^3$  s.t.  $v = x + u$

**Nonlinear Elasticity:** the energy associated to a deformation strain  $\beta$  is

$$E(\beta) := \int_{\Omega} W(\beta) dx,$$

where  $W(F) \sim \text{dist}(F, SO(3))^2$  (so  $W(I) = 0$ ).



# Classical Elasticity: ideal crystals

**Reference configuration:**  $\Omega \subset \mathbb{R}^3$  open bounded

**Deformations:** regular maps  $v: \Omega \rightarrow \mathbb{R}^3$

**Deformation strain:**  $\beta := \nabla v: \Omega \rightarrow \mathbb{M}^{3 \times 3}$

**Displacement:** map  $u: \Omega \rightarrow \mathbb{R}^3$  s.t.  $v = x + u$

**Nonlinear Elasticity:** the energy associated to a deformation strain  $\beta$  is


$$E(\beta) := \int_{\Omega} W(\beta) dx,$$

where  $W(F) \sim \text{dist}(F, SO(3))^2$  (so  $W(I) = 0$ ).

**Linear Elasticity:** let  $v = x + \varepsilon u$  with  $\varepsilon \approx 0$ . Then


$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \int_{\Omega} W(\beta) dx = \frac{1}{2} \int_{\Omega} \mathbb{C} \nabla^{\text{sym}} u : \nabla^{\text{sym}} u dx,$$

where  $\mathbb{C} = \partial^2 W(I)$  and  $\nabla^{\text{sym}} u := (\nabla u + \nabla u^T)/2$ .



# Edge dislocations



**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





# Edge dislocations





**Dislocations:** topological defects in the otherwise periodic structure of a crystal.



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.





**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Edge dislocations

**Dislocations:** topological defects in the otherwise periodic structure of a crystal.


**Edge dislocation:** pair  $(\gamma, \xi)$  of dislocation line and Burgers vector, with  $\xi \perp \gamma$ .



# Adding dislocations: the semi-discrete model

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions




# Adding dislocations: the semi-discrete model

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions

**Strain generating  $(\gamma, \xi)$ :** map  $\beta: \Omega \rightarrow \mathbb{M}^{3 \times 3}$  s.t.

$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma \implies \int_c \beta \cdot t \, d\mathcal{H}^1 = \xi.$$



# Adding dislocations: the semi-discrete model

**Dislocation lines:** Lipschitz curves  $\gamma \subset \Omega$  such that  $\Omega \setminus \gamma$  is not simply connected

**Burgers vector:**  $\xi \in \mathcal{S}$  set of slip directions

**Strain generating  $(\gamma, \xi)$ :** map  $\beta: \Omega \rightarrow \mathbb{M}^{3 \times 3}$  s.t.


$$\text{Curl } \beta = -\xi \otimes \dot{\gamma} \mathcal{H}^1 \llcorner \gamma \implies \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$

**Geometric interpretation:** if  $D$  encloses  $\gamma$ , there exists a deformation  $v \in SBV(\Omega; \mathbb{R}^3)$  s.t.

$$Dv = \nabla v \, dx + \xi \otimes n \mathcal{H}^2 \llcorner D, \quad \beta = \nabla v.$$

In particular:


- ▶  $D$  = slip region,
- ▶  $v$  has constant jump  $\xi$  across  $D$ ,
- ▶ the absolutely continuous part of  $Dv$  is  $\beta$ .



# Regularise the problem: Core Radius Approach

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$






# Regularise the problem: Core Radius Approach

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma)).$$



# Regularise the problem: Core Radius Approach


Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma)).$$

**CRA:** new ref. conf.  $\Omega_\varepsilon(\gamma) := \Omega \setminus I_\varepsilon(\gamma)$ .



# Regularise the problem: Core Radius Approach

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$


Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma)).$$

**CRA:** new ref. conf.  $\Omega_\varepsilon(\gamma) := \Omega \setminus I_\varepsilon(\gamma)$ .

**New Strains:** maps  $\beta \in L^2(\Omega_\varepsilon(\gamma); \mathbb{M}^{3 \times 3})$  s.t.

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\gamma) = 0, \quad \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$



# Regularise the problem: Core Radius Approach

Let  $\beta$  generate  $(\gamma, \xi)$ . Consider  $\varepsilon > 0$  and

$$I_\varepsilon(\gamma) := \{x \in \mathbb{R}^3 : \text{dist}(x, \gamma) < \varepsilon\}.$$

Then we have

$$|\beta(x)| \sim \frac{1}{\text{dist}(x, \gamma)} \text{ in } I_\varepsilon(\gamma) \implies \beta \notin L^2(I_\varepsilon(\gamma)).$$


**CRA:** new ref. conf.  $\Omega_\varepsilon(\gamma) := \Omega \setminus I_\varepsilon(\gamma)$ .

**New Strains:** maps  $\beta \in L^2(\Omega_\varepsilon(\gamma); \mathbb{M}^{3 \times 3})$  s.t.

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\gamma) = 0, \quad \int_C \beta \cdot t \, d\mathcal{H}^1 = \xi.$$

**Elastic energy** associated to  $\beta$  is

$$E_\varepsilon(\beta) := \int_{\Omega_\varepsilon(\gamma)} W(\beta) \, dx.$$



# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   $\Gamma$ -converges to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   $\Gamma$ -converges to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

- ( $\Gamma$ -liminf inequality) for every  $x \in \mathcal{X}$  and every  $\{x_n\}$  such that  $x_n \rightarrow x$ ,

$$F(x) \leq \liminf_{n \rightarrow \infty} F_n(x_n),$$

# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   **$\Gamma$ -converges** to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

- ▶ ( $\Gamma$ -liminf inequality) for every  $x \in \mathcal{X}$  and every  $\{x_n\}$  such that  $x_n \rightarrow x$ ,

$$F(x) \leq \liminf_{n \rightarrow \infty} F_n(x_n),$$

- ▶ ( $\Gamma$ -limsup inequality) for every  $x \in \mathcal{X}$  there exists a **recovery sequence**  $\{x_n\}$  such that  $x_n \rightarrow x$  and

$$F(x) = \lim_{n \rightarrow \infty} F_n(x_n).$$



# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   **$\Gamma$ -converges** to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

- ▶ ( $\Gamma$ -liminf inequality) for every  $x \in \mathcal{X}$  and every  $\{x_n\}$  such that  $x_n \rightarrow x$ ,

$$F(x) \leq \liminf_{n \rightarrow \infty} F_n(x_n),$$

- ▶ ( $\Gamma$ -limsup inequality) for every  $x \in \mathcal{X}$  there exists a **recovery sequence**  $\{x_n\}$  such that  $x_n \rightarrow x$  and

$$F(x) = \lim_{n \rightarrow \infty} F_n(x_n).$$

## Theorem (Fundamental Theorem of $\Gamma$ -convergence)

If  $F_n \xrightarrow{\Gamma} F$  and  $\{x_n\}$  are (almost) minimisers of  $F_n$  at each fixed  $n$ ,

# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   **$\Gamma$ -converges** to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

- ▶ ( $\Gamma$ -liminf inequality) for every  $x \in \mathcal{X}$  and every  $\{x_n\}$  such that  $x_n \rightarrow x$ ,

$$F(x) \leq \liminf_{n \rightarrow \infty} F_n(x_n),$$

- ▶ ( $\Gamma$ -limsup inequality) for every  $x \in \mathcal{X}$  there exists a **recovery sequence**  $\{x_n\}$  such that  $x_n \rightarrow x$  and

$$F(x) = \lim_{n \rightarrow \infty} F_n(x_n).$$

## Theorem (Fundamental Theorem of $\Gamma$ -convergence)

If  $F_n \xrightarrow{\Gamma} F$  and  $\{x_n\}$  are (almost) minimisers of  $F_n$  at each fixed  $n$ ,

- ▶  $F$  admits minimum in  $\mathcal{X}$  and  $\inf_{\mathcal{X}} F_n \rightarrow \min_{\mathcal{X}} F$

# $\Gamma$ -convergence

Let  $\mathcal{X}$  be a metric space and  $F_n: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ .

## Definition ( $\Gamma$ -convergence)

We say that  $F_n$   **$\Gamma$ -converges** to  $F: \mathcal{X} \rightarrow \mathbb{R} \cup \{\pm\infty\}$  as  $n \rightarrow \infty$  if:

- ▶ ( $\Gamma$ -liminf inequality) for every  $x \in \mathcal{X}$  and every  $\{x_n\}$  such that  $x_n \rightarrow x$ ,

$$F(x) \leq \liminf_{n \rightarrow \infty} F_n(x_n),$$

- ▶ ( $\Gamma$ -limsup inequality) for every  $x \in \mathcal{X}$  there exists a **recovery sequence**  $\{x_n\}$  such that  $x_n \rightarrow x$  and

$$F(x) = \lim_{n \rightarrow \infty} F_n(x_n).$$


## Theorem (Fundamental Theorem of $\Gamma$ -convergence)

If  $F_n \xrightarrow{\Gamma} F$  and  $\{x_n\}$  are (almost) minimisers of  $F_n$  at each fixed  $n$ ,

- ▶  $F$  admits minimum in  $\mathcal{X}$  and  $\inf_{\mathcal{X}} F_n \rightarrow \min_{\mathcal{X}} F$
- ▶ if  $x_n \rightarrow x$  then  $F(x) = \min_{\mathcal{X}} F$ .


# $\Gamma$ -convergence: basic example

Let  $\mathcal{X} = \mathbb{R}$  and define  $F_n(x) := x^2 + \cos(nx)$ .




# $\Gamma$ -convergence: basic example

Let  $\mathcal{X} = \mathbb{R}$  and define  $F_n(x) := x^2 + \cos(nx)$ .



# $\Gamma$ -convergence: basic example


Let  $\mathcal{X} = \mathbb{R}$  and define  $F_n(x) := x^2 + \cos(nx)$ .



We have that  $F_n \xrightarrow{\Gamma} F := x^2 - 1$  as  $n \rightarrow \infty$ .


# Motivation: polycrystals

**Polycrystal:** formed by many grains, having the **same** lattice structure, mutually rotated  $\Rightarrow$  interface misfit at **grain boundaries**.



# Motivation: polycrystals

**Polycrystal:** formed by many grains, having the **same** lattice structure, mutually rotated  $\implies$  interface misfit at **grain boundaries**.



**Goal:** to obtain polycrystalline structures as minimisers of some energy functional.


F., Palombaro, Ponsiglione. *Linearised Polycrystals from a 2D System of Edge Dislocations*. Preprint (2017)



# Structure of Tilt Grain Boundaries

**Tilt boundary:** small angle rotation  $\theta$  between grains  $\Rightarrow$  **edge dislocations**.

**Boundary structure:** periodic array of edge dislocations with spacing  $\delta = \varepsilon/\theta$ .



Porter, Easterling. CRC Press (2009) - Gottstein. Springer (2013)

# Plan of the paper

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

# Plan of the paper

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

**Plan:** let  $\mathcal{F}_\varepsilon$  be the energy of such system.

- We compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,

# Plan of the paper

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

**Plan:** let  $\mathcal{F}_\varepsilon$  be the energy of such system.

- ▶ We compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,
- ▶ we show that under suitable boundary conditions  $\mathcal{F}$  is minimised by polycrystals.

# Plan of the paper

**Setting:** consider a 2D system of  $N_\varepsilon$  edge dislocations, where  $\varepsilon > 0$  is the lattice spacing and

$$N_\varepsilon \rightarrow +\infty \quad \text{as} \quad \varepsilon \rightarrow 0.$$

**Plan:** let  $\mathcal{F}_\varepsilon$  be the energy of such system.

- ▶ We compute  $\mathcal{F}$ , the  $\Gamma$ -limit of  $\mathcal{F}_\varepsilon$  as  $\varepsilon \rightarrow 0$ ,
- ▶ we show that under suitable boundary conditions  $\mathcal{F}$  is minimised by polycrystals.


**Linearised polycrystals:** our energy regime will imply

$$N_\varepsilon \ll \frac{1}{\varepsilon}$$

$\implies$  we have less dislocations than tilt grain boundaries. However we still obtain polycrystalline minimisers, but with grains rotated by an infinitesimal angle  $\theta \approx 0$ .

# Setting: linear planar elasticity


**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.



# Setting: linear planar elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .





# Setting: linear planar elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .





# Setting: linear planar elasticity



**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$



# Setting: linear planar elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.



**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$

**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .



# Setting: linear planar elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .



**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$

**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .

**Strains:** inducing  $\mu$  are maps  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  s.t.  $\beta = 0$  in  $\cup B_\varepsilon(x_i)$  and

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\mu) = 0, \quad \int_{\partial B_\varepsilon(x_i)} \beta \cdot t \, ds = \xi_i.$$



# Setting: linear planar elasticity

**Reference configuration:**  $\Omega \subset \mathbb{R}^2$  open bounded.

**Dislocation lines:** points  $x_0 \in \Omega$  separated by  $2\varepsilon$ .

**Burgers vectors:** finite set  $\mathcal{S} := \{b_1, \dots, b_s\} \subset \mathbb{R}^2$ .

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathcal{S}.$$



**Core radius approach:**  $\Omega_\varepsilon(\mu) := \Omega \setminus \cup B_\varepsilon(x_i)$ .

**Strains:** inducing  $\mu$  are maps  $\beta: \Omega_\varepsilon(\mu) \rightarrow \mathbb{M}^{2 \times 2}$  s.t.  $\beta = 0$  in  $\cup B_\varepsilon(x_i)$  and

$$\text{Curl } \beta \llcorner \Omega_\varepsilon(\mu) = 0, \quad \int_{\partial B_\varepsilon(x_i)} \beta \cdot t \, ds = \xi_i.$$


**Linear Energy:**  $\mathbb{C}F : F \sim |F^{\text{sym}}|^2$ , then

$$E_\varepsilon(\mu, \beta) := \int_{\Omega} \mathbb{C}\beta : \beta \, dx = \int_{\Omega} \mathbb{C}\beta^{\text{sym}} : \beta^{\text{sym}} \, dx.$$



# Self-energy of a single dislocation core


Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”


$$\int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx$$



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”


$$\int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx \geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho$$



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho \end{aligned}$$






# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”


$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$



# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$




The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$



The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$


**Remark:** let  $s \in (0, 1)$ , then

$$\int_{B_{\varepsilon^s} \setminus B_\varepsilon} |\beta|^2 dx \geq (1-s) \frac{|\xi|^2}{2\pi} |\log \varepsilon|.$$

# Self-energy of a single dislocation core

Let  $\beta$  generate  $\xi \delta_0$ , that is “ $\text{Curl } \beta = \xi \delta_0$ ”

$$\begin{aligned} \int_{B_1 \setminus B_\varepsilon} |\beta|^2 dx &\geq \int_\varepsilon^1 \int_{\partial B_\rho} |\beta \cdot t|^2 ds d\rho \geq (\text{Jensen}) \\ &\geq \frac{1}{2\pi} \int_\varepsilon^1 \frac{1}{\rho} \left| \int_{\partial B_\rho} \beta \cdot t ds \right|^2 d\rho = \frac{|\xi|^2}{2\pi} |\log \varepsilon|. \end{aligned}$$



The reverse inequality can be obtained by computing the energy of

$$\beta(x) := \frac{1}{2\pi} \xi \otimes J \frac{x}{|x|^2}, \quad J := \text{clock-wise rotation of } \frac{\pi}{2}.$$

**Remark:** let  $s \in (0, 1)$ , then

$$\int_{B_{\varepsilon^s} \setminus B_\varepsilon} |\beta|^2 dx \geq (1-s) \frac{|\xi|^2}{2\pi} |\log \varepsilon|.$$

**Self-energy:** is of order  $|\log \varepsilon|$  and concentrated in a small region around  $B_\varepsilon$ .


# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.





# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .


**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.


**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$

**Note:** dislocations separation is a technical assumption for energy estimates.



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses


$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$

with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$

**Note:** dislocations separation is a technical assumption for energy estimates.

**Hypothesis on HC Radius:** as  $\varepsilon \rightarrow 0$



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$


with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$

**Note:** dislocations separation is a technical assumption for energy estimates.

**Hypothesis on HC Radius:** as  $\varepsilon \rightarrow 0$

►  $\rho_\varepsilon / \varepsilon^s \rightarrow \infty, \quad \forall s \in (0, 1), \quad (\text{HC region contains almost all the self-energy})$



# The Hard Core assumption

**HC Radius:** fixed scale  $\rho_\varepsilon \gg \varepsilon$  with  $\rho_\varepsilon \rightarrow 0$ .

**Clusters** of dislocations at scale  $\rho_\varepsilon$  are identified with a single **multiple dislocation**.

**Admissible dislocations:** finite sums of Dirac masses

$$\mu := \sum_{i=1}^N \xi_i \delta_{x_i}, \quad \xi_i \in \mathbb{S},$$


with  $\mathbb{S} := \text{Span}_{\mathbb{Z}} \mathcal{S}$  set of multiple Burgers vectors, and

$$|x_i - x_j| > 2\rho_\varepsilon, \quad \text{dist}(x_k, \partial\Omega) > \rho_\varepsilon.$$

**Note:** dislocations separation is a technical assumption for energy estimates.

**Hypothesis on HC Radius:** as  $\varepsilon \rightarrow 0$

- ▶  $\rho_\varepsilon / \varepsilon^s \rightarrow \infty, \quad \forall s \in (0, 1),$  (HC region contains almost all the self-energy)
- ▶  $N_\varepsilon \rho_\varepsilon^2 \rightarrow 0.$  (Measure of HC region vanishes)



# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|.$$

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|.$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta^{\text{sym}} : \beta^{\text{sym}} dx.$$

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|.$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta^{\text{sym}} : \beta^{\text{sym}} dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

►  $N_\varepsilon \ll |\log \varepsilon| \leadsto$  Dilute dislocations



# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|.$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta^{\text{sym}} : \beta^{\text{sym}} dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

- ▶  $N_\varepsilon \ll |\log \varepsilon| \leadsto$  Dilute dislocations
- ▶  $N_\varepsilon \approx |\log \varepsilon| \leadsto$  Critical regime

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

# Energy regimes

**Energy scaling:** each dislocation accounts for  $|\log \varepsilon| \implies$  relevant scaling is

$$E_\varepsilon \approx N_\varepsilon |\log \varepsilon|.$$

**Rescaled energy functionals:**

$$\mathcal{F}_\varepsilon(\mu, \beta) := \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega_\varepsilon(\mu)} \mathbb{C} \beta^{\text{sym}} : \beta^{\text{sym}} dx.$$

**Energy regimes:** the behaviour of  $N_\varepsilon$  determines three different regimes:

►  $N_\varepsilon \ll |\log \varepsilon| \rightsquigarrow$  Dilute dislocations

►  $N_\varepsilon \approx |\log \varepsilon| \rightsquigarrow$  Critical regime

Garroni, Leoni, Ponsiglione. *Gradient theory for plasticity via homogenization of discrete dislocations*.  
J. Eur. Math. Soc. (JEMS) (2010)

►  $N_\varepsilon \gg |\log \varepsilon| \rightsquigarrow$  Super-critical regime

F., Palombaro, Ponsiglione. *Linearised Polycrystals from a 2D System of Edge Dislocations*.  
Preprint (2017)

## Behaviour of $\mathcal{F}_\varepsilon$ as $\varepsilon \rightarrow 0$ (Heuristic)


Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^{N_\varepsilon} \xi_i \delta_{x_i}$  be such that “Curl  $\beta = \mu$ ”.

# Behaviour of $\mathcal{F}_\varepsilon$ as $\varepsilon \rightarrow 0$ (Heuristic)

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^{N_\varepsilon} \xi_i \delta_{x_i}$  be such that “ $\text{Curl } \beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^{N_\varepsilon} B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}.$$




# Behaviour of $\mathcal{F}_\varepsilon$ as $\varepsilon \rightarrow 0$ (Heuristic)

Let  $(\mu, \beta)$  with  $\mu = \sum_{i=1}^{N_\varepsilon} \xi_i \delta_{x_i}$  be such that “ $\text{Curl } \beta = \mu$ ”.

**Energy decomposition:** let  $\text{HC}_\varepsilon(\mu) := \cup_{i=1}^{N_\varepsilon} B_{\rho_\varepsilon}(x_i)$  be the HC region

$$E_\varepsilon(\mu, \beta) = \int_{\Omega \setminus \text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx + \int_{\text{HC}_\varepsilon(\mu)} \mathbb{C}\beta : \beta \, dx = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}.$$



**Idea:** rescaling by  $N_\varepsilon |\log \varepsilon|$ , we have  $E_\varepsilon^{\text{interaction}} \rightarrow E^{\text{elastic}}$  and  $E_\varepsilon^{\text{self}} \rightarrow E^{\text{plastic}}$ .

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “**Curl**  $\beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

$$\blacktriangleright \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A \text{ in } L^2(\Omega; \mathbb{M}^{2 \times 2}),$$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,



# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\text{Curl } A = \mu$  and  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  ( $\implies A \in BV(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ).

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\text{Curl } A = \mu$  and  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  ( $\implies A \in BV(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ).

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\text{Curl } A = \mu$  and  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  ( $\implies A \in BV(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ).

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

### Remark:

- ▶  $S$  and  $A$  live on two different scales with  $S \ll A \implies$  terms in  $\mathcal{F}$  decoupled.

# $\Gamma$ -convergence result for $N_\varepsilon \gg |\log \varepsilon|$

## Theorem (F., Palombaro, Ponsiglione '17)

**Compactness:** consider  $(\mu_\varepsilon, \beta_\varepsilon)$  s.t. “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” and  $\mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C \implies$

- ▶  $\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S$ ,  $\frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A$  in  $L^2(\Omega; \mathbb{M}^{2 \times 2})$ ,
- ▶  $\frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu$  in  $\mathcal{M}(\Omega; \mathbb{R}^2)$ ,
- ▶  $\text{Curl } A = \mu$  and  $\mu \in H^{-1}(\Omega; \mathbb{R}^2)$  ( $\implies A \in BV(\Omega; \mathbb{M}_{\text{skew}}^{2 \times 2})$ ).

**$\Gamma$ -convergence:** the functionals  $\mathcal{F}_\varepsilon$   $\Gamma$ -converge to

$$\mathcal{F}(\mu, S, A) := \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|, \quad \text{with } \text{Curl } A = \mu.$$

## Remark:

- ▶  $S$  and  $A$  live on two different scales with  $S \ll A \implies$  terms in  $\mathcal{F}$  decoupled.
- ▶ In the critical regime  $N_\varepsilon \approx |\log \varepsilon|$  we have  $S \approx A$  and  $\text{Curl}(S + A) = \mu$ .

# The relaxation formula for $\varphi$

**Self-energy** for a single dislocation core  $\xi\delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \min_{\beta} \left\{ \frac{1}{|\log \varepsilon|} \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi\delta_0\text{”} \right\} .$$

# The relaxation formula for $\varphi$

**Self-energy** for a single dislocation core  $\xi\delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \min_{\beta} \left\{ \frac{1}{|\log \varepsilon|} \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi\delta_0\text{”} \right\}.$$

**Plastic density:** the map  $\varphi: \mathbb{R}^2 \rightarrow [0, \infty)$  defined as the relaxation of  $\psi$

$$\varphi(\xi) := \min \left\{ \sum_{i=1}^M \lambda_i \psi(\xi_i) : \xi = \sum_{i=1}^M \lambda_i \xi_i, M \in \mathbb{N}, \lambda_i \geq 0, \xi_i \in \mathbb{S} \right\}.$$

# The relaxation formula for $\varphi$

**Self-energy** for a single dislocation core  $\xi\delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \min_{\beta} \left\{ \frac{1}{|\log \varepsilon|} \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi\delta_0\text{”} \right\}.$$

**Plastic density:** the map  $\varphi: \mathbb{R}^2 \rightarrow [0, \infty)$  defined as the relaxation of  $\psi$

$$\varphi(\xi) := \min \left\{ \sum_{i=1}^M \lambda_i \psi(\xi_i) : \xi = \sum_{i=1}^M \lambda_i \xi_i, M \in \mathbb{N}, \lambda_i \geq 0, \xi_i \in \mathbb{S} \right\}.$$

**Note:** since the energy is quadratic, in the  $\Gamma$ -limit we have  $\varphi$  instead of  $\psi$ .

# The relaxation formula for $\varphi$

**Self-energy** for a single dislocation core  $\xi\delta_0$  is

$$\psi(\xi) := \lim_{\varepsilon \rightarrow 0} \min_{\beta} \left\{ \frac{1}{|\log \varepsilon|} \int_{B_1 \setminus B_\varepsilon} \mathbb{C}\beta : \beta \, dx : \text{“Curl } \beta = \xi\delta_0\text{”} \right\}.$$

**Plastic density:** the map  $\varphi: \mathbb{R}^2 \rightarrow [0, \infty)$  defined as the relaxation of  $\psi$

$$\varphi(\xi) := \min \left\{ \sum_{i=1}^M \lambda_i \psi(\xi_i) : \xi = \sum_{i=1}^M \lambda_i \xi_i, M \in \mathbb{N}, \lambda_i \geq 0, \xi_i \in \mathbb{S} \right\}.$$

**Note:** since the energy is quadratic, in the  $\Gamma$ -limit we have  $\varphi$  instead of  $\psi$ .

**Properties:**  $\varphi$  is convex and positively 1-homogeneous. Moreover  $\exists c > 0$  s.t.

$$c^{-1} |\xi| \leq \varphi(\xi) \leq c |\xi|, \quad \forall \xi \in \mathbb{R}^2.$$



# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$C \geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$C \geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$C \geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

$$(\rho_\varepsilon \gg \varepsilon) \gtrsim \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \psi(\xi_{\varepsilon,i})$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$\begin{aligned} C &\geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \\ (\rho_\varepsilon \gg \varepsilon) &\gtrsim \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \psi(\xi_{\varepsilon,i}) \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) \end{aligned}$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$\begin{aligned} C &\geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \\ (\rho_\varepsilon \gg \varepsilon) &\gtrsim \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \psi(\xi_{\varepsilon,i}) \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \varphi\left(\frac{\xi_{\varepsilon,i}}{|\xi_{\varepsilon,i}|}\right) \end{aligned}$$

# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$\begin{aligned} C &\geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \\ (\rho_\varepsilon \gg \varepsilon) &\gtrsim \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \psi(\xi_{\varepsilon,i}) \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \varphi\left(\frac{\xi_{\varepsilon,i}}{|\xi_{\varepsilon,i}|}\right) \\ \left(c := \min_{|\xi|=1} \varphi(\xi) > 0\right) &\geq \frac{c}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \end{aligned}$$



# Ideas for compactness: measures

Let  $(\mu_\varepsilon, \beta_\varepsilon)$  with  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$  and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”. Assume that

$$\sup_{\varepsilon} \mathcal{F}_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) \leq C.$$

We show that

$$\frac{|\mu_\varepsilon|(\Omega)}{N_\varepsilon} = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \leq C \implies \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu. \quad (1)$$

$$\begin{aligned} C &\geq \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \\ (\rho_\varepsilon \gg \varepsilon) &\gtrsim \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \psi(\xi_{\varepsilon,i}) \geq \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) = \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| \varphi\left(\frac{\xi_{\varepsilon,i}}{|\xi_{\varepsilon,i}|}\right) \\ \left(c := \min_{|\xi|=1} \varphi(\xi) > 0\right) &\geq \frac{c}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} |\xi_{\varepsilon,i}| = \frac{c}{N_\varepsilon} |\mu_\varepsilon|(\Omega) \implies (1) \end{aligned}$$

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx$$

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx \implies \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S.$$

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx \implies \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S.$$

**Skew Part:** we use a **Generalised Korn inequality**: there exists  $C > 0$  s.t. for every  $\beta \in L^1(\Omega; \mathbb{M}^{2 \times 2})$  with  $\text{Curl } \beta \in \mathcal{M}(\Omega; \mathbb{R}^2)$ ,

$$\int_{\Omega} |\beta^{\text{skew}}|^2 \, dx \leq C \left( \int_{\Omega} |\beta^{\text{sym}}|^2 \, dx + |\text{Curl } \beta|(\Omega)^2 \right). \quad (2)$$

Garroni, Leoni, Ponsiglione. J. Eur. Math. Soc. (JEMS) (2010)

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C}\beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx \implies \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S.$$

**Skew Part:** we use a **Generalised Korn inequality**: there exists  $C > 0$  s.t. for every  $\beta \in L^1(\Omega; \mathbb{M}^{2 \times 2})$  with  $\text{Curl } \beta \in \mathcal{M}(\Omega; \mathbb{R}^2)$ ,

$$\int_{\Omega} |\beta^{\text{skew}}|^2 \, dx \leq C \left( \int_{\Omega} |\beta^{\text{sym}}|^2 \, dx + |\text{Curl } \beta|(\Omega)^2 \right). \quad (2)$$

Since “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”, by (2) and assumption  $N_\varepsilon \gg |\log \varepsilon|$  we get

$$\int_{\Omega} |\beta_\varepsilon^{\text{skew}}|^2 \, dx \leq C \left( \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx + |\mu_\varepsilon|(\Omega)^2 \right)$$

Garroni, Leoni, Ponsiglione. J. Eur. Math. Soc. (JEMS) (2010)

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx \implies \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S.$$

**Skew Part:** we use a **Generalised Korn inequality**: there exists  $C > 0$  s.t. for every  $\beta \in L^1(\Omega; \mathbb{M}^{2 \times 2})$  with  $\text{Curl } \beta \in \mathcal{M}(\Omega; \mathbb{R}^2)$ ,

$$\int_{\Omega} |\beta^{\text{skew}}|^2 \, dx \leq C \left( \int_{\Omega} |\beta^{\text{sym}}|^2 \, dx + |\text{Curl } \beta|(\Omega)^2 \right). \quad (2)$$

Since “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”, by (2) and assumption  $N_\varepsilon \gg |\log \varepsilon|$  we get

$$\begin{aligned} \int_{\Omega} |\beta_\varepsilon^{\text{skew}}|^2 \, dx &\leq C \left( \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx + |\mu_\varepsilon|(\Omega)^2 \right) \\ &\leq C (N_\varepsilon |\log \varepsilon| + N_\varepsilon^2) \leq C N_\varepsilon^2. \end{aligned}$$

Garroni, Leoni, Ponsiglione. J. Eur. Math. Soc. (JEMS) (2010)

# Ideas for compactness: strains

**Symmetric Part:** recalling that  $\mathbb{C}F : F \geq C|F^{\text{sym}}|^2$ , we have

$$N_\varepsilon |\log \varepsilon| \geq C \int_{\Omega} \mathbb{C}\beta_\varepsilon : \beta_\varepsilon \, dx \geq C \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx \implies \frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S.$$

**Skew Part:** we use a **Generalised Korn inequality**: there exists  $C > 0$  s.t. for every  $\beta \in L^1(\Omega; \mathbb{M}^{2 \times 2})$  with  $\text{Curl } \beta \in \mathcal{M}(\Omega; \mathbb{R}^2)$ ,

$$\int_{\Omega} |\beta^{\text{skew}}|^2 \, dx \leq C \left( \int_{\Omega} |\beta^{\text{sym}}|^2 \, dx + |\text{Curl } \beta|(\Omega)^2 \right). \quad (2)$$

Since “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”, by (2) and assumption  $N_\varepsilon \gg |\log \varepsilon|$  we get

$$\begin{aligned} \int_{\Omega} |\beta_\varepsilon^{\text{skew}}|^2 \, dx &\leq C \left( \int_{\Omega} |\beta_\varepsilon^{\text{sym}}|^2 \, dx + |\mu_\varepsilon|(\Omega)^2 \right) \\ &\leq C (N_\varepsilon |\log \varepsilon| + N_\varepsilon^2) \leq CN_\varepsilon^2 \implies \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A. \end{aligned}$$

Garroni, Leoni, Ponsiglione. J. Eur. Math. Soc. (JEMS) (2010)



# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightarrow S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightarrow A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightarrow S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightarrow A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightarrow S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightarrow A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightarrow S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightarrow A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

$$\gtrsim \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i})$$

# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx$$

$$\gtrsim \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) = \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \int_{\Omega} \varphi \left( \frac{d\mu_\varepsilon}{d|\mu_\varepsilon|} \right) d|\mu_\varepsilon|$$



# Ideas for $\Gamma$ -liminf

Assume that  $(\mu_\varepsilon, \beta_\varepsilon)$  is such that  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi_{\varepsilon,i} \delta_{x_{\varepsilon,i}}$ , “Curl  $\beta_\varepsilon = \mu_\varepsilon$ ” and

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightarrow S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightarrow A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad \text{with } \text{Curl } A = \mu.$$

We have to show

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|.$$

**Idea:** split the energy  $E_\varepsilon(\mu_\varepsilon, \beta_\varepsilon) = E_\varepsilon^{\text{interaction}} + E_\varepsilon^{\text{self}}$  and use lower semicontinuity:

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega \setminus HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \geq \int_{\Omega} \mathbb{C} S : S \, dx \quad (N_\varepsilon \rho_\varepsilon^2 \rightarrow 0 \implies |HC_\varepsilon| \rightarrow 0)$$

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{HC_\varepsilon} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx &= \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \frac{1}{|\log \varepsilon|} \int_{B_{\rho_\varepsilon}(x_{\varepsilon,i})} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx \\ &\gtrsim \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \sum_{i=1}^{M_\varepsilon} \varphi(\xi_{\varepsilon,i}) = \liminf_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon} \int_{\Omega} \varphi \left( \frac{d\mu_\varepsilon}{d|\mu_\varepsilon|} \right) d|\mu_\varepsilon| \geq \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| \end{aligned}$$

by Reshetnyak's lower semicontinuity Theorem, since  $\varphi$  is 1-homogeneous.

# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ .

# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.

# Ideas for $\Gamma$ -limsup


Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.

**Recovery measures:**



# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.


$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.

## Recovery measures:

- divide  $\Omega$  in  $\approx N_\varepsilon$  squares of side  $2r_\varepsilon = C/\sqrt{N_\varepsilon}$



# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.


$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.

## Recovery measures:

- ▶ divide  $\Omega$  in  $\approx N_\varepsilon$  squares of side  $2r_\varepsilon = C/\sqrt{N_\varepsilon}$
- ▶ the recovery sequence is  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi \delta_{x_{\varepsilon,i}}$ .



# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$


$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.

## Recovery measures:

- ▶ divide  $\Omega$  in  $\approx N_\varepsilon$  squares of side  $2r_\varepsilon = C/\sqrt{N_\varepsilon}$
- ▶ the recovery sequence is  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi \delta_{x_{\varepsilon,i}}$ .

**Recovery strains:** let  $K_\varepsilon$  be the solution to the cell-problem about each  $x_{\varepsilon,i}$ .





# Ideas for $\Gamma$ -limsup

Consider  $(\mu, S, A)$  with  $S$  symmetric,  $A$  skew and  $\text{Curl } A = \mu$ . We have to construct a **recovery sequence**  $(\mu_\varepsilon, \beta_\varepsilon)$  with “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ” s.t.

$$\frac{\beta_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup S, \quad \frac{\beta_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup A, \quad \frac{\mu_\varepsilon}{N_\varepsilon} \xrightarrow{*} \mu, \quad (3)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{N_\varepsilon |\log \varepsilon|} \int_{\Omega} \mathbb{C} \beta_\varepsilon : \beta_\varepsilon \, dx = \int_{\Omega} \mathbb{C} S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu|. \quad (4)$$

For simplicity assume  $\mu = \xi \, dx$  with  $\varphi(\xi) = \psi(\xi)$ . The general is more technical.


## Recovery measures:

- ▶ divide  $\Omega$  in  $\approx N_\varepsilon$  squares of side  $2r_\varepsilon = C/\sqrt{N_\varepsilon}$
- ▶ the recovery sequence is  $\mu_\varepsilon = \sum_{i=1}^{M_\varepsilon} \xi \delta_{x_{\varepsilon,i}}$ .

**Recovery strains:** let  $K_\varepsilon$  be the solution to the cell-problem about each  $x_{\varepsilon,i}$ . Then  $K_\varepsilon / \sqrt{N_\varepsilon |\log \varepsilon|} \rightharpoonup 0$  and

$$\beta_\varepsilon = \sqrt{N_\varepsilon |\log \varepsilon|} S + N_\varepsilon A + K_\varepsilon + O(\sqrt{N_\varepsilon |\log \varepsilon|})$$

satisfies (3), (4) and “ $\text{Curl } \beta_\varepsilon = \mu_\varepsilon$ ”.



# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Dislocations and strains:**  $(\mu, \beta)$  such that “ $\text{Curl } \beta = \mu$ ” and

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds, \quad \beta \cdot t = g_\varepsilon \cdot t \text{ on } \partial\Omega.$$

# Adding boundary conditions

**Dirichlet type BC:** at level  $\varepsilon > 0$  fix a boundary condition  $g_\varepsilon: \Omega \rightarrow \mathbb{M}^{2 \times 2}$  s.t.

$$\frac{g_\varepsilon^{\text{sym}}}{\sqrt{N_\varepsilon |\log \varepsilon|}} \rightharpoonup g_S, \quad \frac{g_\varepsilon^{\text{skew}}}{N_\varepsilon} \rightharpoonup g_A.$$

**Dislocations and strains:**  $(\mu, \beta)$  such that “ $\text{Curl } \beta = \mu$ ” and

$$\mu(\Omega) = \int_{\partial\Omega} g_\varepsilon \cdot t \, ds, \quad \beta \cdot t = g_\varepsilon \cdot t \quad \text{on } \partial\Omega.$$

## Theorem (F., Palombaro, Ponsiglione '17)

*The energy functionals  $\mathcal{F}_\varepsilon$  are equi-coercive and they  $\Gamma$ -converge to*

$$\mathcal{F}_{\text{BC}}(\mu, S, A) := \int_{\Omega} \mathbb{C}S : S \, dx + \int_{\Omega} \varphi \left( \frac{d\mu}{d|\mu|} \right) d|\mu| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) \, ds,$$

*with  $\text{Curl } A = \mu$  and  $\mu \in \mathcal{M}(\Omega; \mathbb{R}^2) \cap H^{-1}(\Omega; \mathbb{R}^2)$ .*

**Remark:**  $\beta^{\text{sym}} \ll \beta^{\text{skew}} \implies$  BC pass to the limit only for  $A$ .

# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

**Piecewise constant BC:** Fix a piecewise constant BC

$$g_A := \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}, \quad a := \sum_{k=1}^M m_k \chi_{U_k},$$

with  $m_k < m_{k+1}$  and  $\{U_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .

# Minimising $\mathcal{F}_{\text{BC}}$ with piecewise constant BC

**Remark:** there are no BC on  $S \implies$  we can neglect elastic energy.

**Piecewise constant BC:** Fix a piecewise constant BC

$$g_A := \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}, \quad a := \sum_{k=1}^M m_k \chi_{U_k},$$

with  $m_k < m_{k+1}$  and  $\{U_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ .

## Problem

*Minimise*

$$\mathcal{F}_{\text{BC}}(\text{Curl } A, 0, A) = \int_{\Omega} \varphi \left( \frac{d \text{Curl } A}{d|\text{Curl } A|} \right) d|\text{Curl } A| + \int_{\partial\Omega} \varphi((g_A - A) \cdot t) ds,$$

with  $\text{Curl } A \in \mathcal{M}(\Omega; \mathbb{R}^2) \cap H^{-1}(\Omega; \mathbb{R}^2)$ .


# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{BC}(\text{Curl } A, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ . We interpret  $A$  as a *linearised polycrystal*.






# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{BC}(\text{Curl } A, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ . We interpret  $A$  as a *linearised polycrystal*.



**Proof Strategy:** We are minimising an anisotropic total variation functional. By Coarea formula we select the levels with minimal perimeter, defining the Caccioppoli partition.


# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{BC}(\text{Curl } A, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$

with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ . We interpret  $A$  as a *linearised polycrystal*.



**Proof Strategy:** We are minimising an anisotropic total variation functional. By Coarea formula we select the levels with minimal perimeter, defining the Caccioppoli partition.

**Open Question:** Are all minimisers piecewise constant? Uniqueness?


# Polycrystals as energy minimisers

## Theorem (F., Palombaro, Ponsiglione '17)

Given a piecewise constant boundary condition  $g_A$ , there exists a *piecewise constant* minimiser of  $\mathcal{F}_{BC}(\text{Curl } A, 0, A)$

$$A = \sum_{k=1}^M A_k \chi_{E_k},$$


with  $A_k \in \mathbb{M}_{\text{skew}}^{2 \times 2}$  and  $\{E_k\}_{k=1}^M$  Caccioppoli partition of  $\Omega$ . We interpret  $A$  as a *linearised polycrystal*.



**Proof Strategy:** We are minimising an anisotropic total variation functional. By Coarea formula we select the levels with minimal perimeter, defining the Caccioppoli partition.

**Open Question:** Are all minimisers piecewise constant? Uniqueness?

**Essential:** that the boundary condition is piecewise affine on the *whole*  $\partial\Omega$ .



# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Open Questions:

- ▶ **Uniqueness** of piecewise constant minimisers?

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Open Questions:

- ▶ **Uniqueness** of piecewise constant minimisers?
- ▶ **Dynamics** for linearised polycrystals?

Taylor. Bull. Amer. Math. Soc. (1978).

Chambolle, Morini, Ponsiglione. Comm. Pure Appl. Math (2017).

# Conclusions and Perspectives

## Conclusions:

- ▶ A variational model for **linearised polycrystals** with infinitesimal rotations between the grains, deduced by  $\Gamma$ -convergence.
- ▶ Networks of dislocations are obtained as the result of **energy minimisation**, under suitable boundary conditions.

## Open Questions:

- ▶ **Uniqueness** of piecewise constant minimisers?
- ▶ **Dynamics** for linearised polycrystals?  
Taylor. Bull. Amer. Math. Soc. (1978).  
Chambolle, Morini, Ponsiglione. Comm. Pure Appl. Math (2017).
- ▶  $\Gamma$ -convergence analysis starting from a **non-linear energy**?  
Namely, considering small deformations  $v = x + \varepsilon u$ . Now the Burgers vectors are  $\varepsilon \xi$  and the equivalent rescaling is  $\varepsilon^2 N_\varepsilon |\log \varepsilon|$ .  
Müller, Scardia, Zeppieri. Indiana University Mathematics Journal (2014).



Thank You!