

# Optimal lower exponent of solutions to two-phase elliptic equations in two dimensions

Silvio Fanzon

(joint work with Mariapia Palombaro)

Karl-Franzens University, Graz  
Department of Mathematics

# Problem

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \sigma^{-1} \xi \cdot \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

# Problem

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \sigma^{-1} \xi \cdot \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

## Problem

Study the gradient integrability of distributional solutions  $u \in W^{1,1}(\Omega)$  to

$$\operatorname{div}(\sigma \nabla u) = 0, \tag{0.1}$$

when

$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

with  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  constant elliptic matrices,  $\{E_1, E_2\}$  measurable partition of  $\Omega$ .

# Problem

$\Omega \subset \mathbb{R}^2$  bounded open domain. A map  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  is **uniformly elliptic** if

$$\sigma \xi \cdot \xi \geq \lambda |\xi|^2, \quad \sigma^{-1} \xi \cdot \xi \geq \lambda |\xi|^2 \quad \forall \xi \in \mathbb{R}^2, x \in \Omega.$$

## Problem

Study the gradient integrability of distributional solutions  $u \in W^{1,1}(\Omega)$  to

$$\operatorname{div}(\sigma \nabla u) = 0, \tag{0.1}$$

when


$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$

with  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  constant elliptic matrices,  $\{E_1, E_2\}$  measurable partition of  $\Omega$ .

## Application to composites:

- ▶  $\Omega$  is a section of a **composite conductor** obtained by mixing two materials with **conductivities**  $\sigma_1$  and  $\sigma_2$
- ▶ the **electric field**  $\nabla u$  solves (0.1)
- ▶ How much can  $\nabla u$  concentrate, given the geometry  $\{E_1, E_2\}$ ?

# Astala's Theorem



## Theorem (Astala '94)

Let  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  be uniformly elliptic. There exists exponents  $1 < q < 2 < p$  such that if  $u \in W^{1,q}(\Omega)$  solves

$$\operatorname{div}(\sigma \nabla u) = 0,$$

then  $\nabla u \in L^p_{\text{weak}}(\Omega; \mathbb{R}^2)$ .

Astala. *Area distortion of quasiconformal mappings*. Acta Mathematica (1994)

# Astala's Theorem



## Theorem (Astala '94)

Let  $\sigma \in L^\infty(\Omega; \mathbb{M}^{2 \times 2})$  be uniformly elliptic. There exists exponents  $1 < q < 2 < p$  such that if  $u \in W^{1,q}(\Omega)$  solves

$$\operatorname{div}(\sigma \nabla u) = 0,$$

then  $\nabla u \in L^p_{\text{weak}}(\Omega; \mathbb{R}^2)$ .

## Question

Are the exponents  $q$  and  $p$  optimal among two-phase elliptic conductivities

$$\sigma = \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2}?$$

Astala. *Area distortion of quasiconformal mappings*. Acta Mathematica (1994)


# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).

# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).



# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.


**Remark:** it is sufficient to prove optimality in the case

$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where

$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$

# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

**Remark:** it is sufficient to prove optimality in the case

$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where


$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$

The corresponding critical exponents for Astala's theorem are

$$q_{\sigma_1, \sigma_2} = \frac{2K}{K+1}, \quad p_{\sigma_1, \sigma_2} = \frac{2K}{K-1}.$$

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

# Astala's exponents for two-phase conductivities



For two-phase conductivities Astala's exponents  $q = q_{\sigma_1, \sigma_2}$  and  $p = p_{\sigma_1, \sigma_2}$  have been characterised.

**Remark:** it is sufficient to prove optimality in the case

$$\sigma_1 = \begin{pmatrix} 1/K & 0 \\ 0 & 1/S_1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} K & 0 \\ 0 & S_2 \end{pmatrix},$$

where


$$K > 1 \quad \text{and} \quad \frac{1}{K} \leq S_j \leq K, \quad j = 1, 2.$$

The corresponding critical exponents for Astala's theorem are

$$q_{\sigma_1, \sigma_2} = \frac{2K}{K+1}, \quad p_{\sigma_1, \sigma_2} = \frac{2K}{K-1}.$$

Nesi, Palombaro, Ponsiglione. Ann. Inst. H. Poincaré Anal. Non Linéaire (2014).

# Upper exponent optimality



## Theorem (Nesi, Palombaro, Ponsiglione '14)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .


(i) If  $\sigma \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and  $u \in W^{1, \frac{2K}{K+1}}(\Omega)$  solves

$$\text{div}(\sigma \nabla u) = 0 \tag{0.2}$$

then  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

(ii) There exists  $\bar{\sigma} \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and a weak solution  $\bar{u} \in W^{1,2}(\Omega)$  to (0.2) with  $\sigma = \bar{\sigma}$ , satisfying affine boundary conditions and such that  $\nabla \bar{u} \notin L^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

# Upper exponent optimality



## Theorem (Nesi, Palombaro, Ponsiglione '14)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .

❶ If  $\sigma \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and  $u \in W^{1, \frac{2K}{K+1}}(\Omega)$  solves

$$\text{div}(\sigma \nabla u) = 0 \tag{0.2}$$


then  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

❷ There exists  $\bar{\sigma} \in L^\infty(\Omega; \{\sigma_1, \sigma_2\})$  and a weak solution  $\bar{u} \in W^{1,2}(\Omega)$  to (0.2) with  $\sigma = \bar{\sigma}$ , satisfying affine boundary conditions and such that  $\nabla \bar{u} \notin L^{\frac{2K}{K-1}}(\Omega; \mathbb{R}^2)$ .

## Question we address

Is the lower exponent  $\frac{2K}{K+1}$  optimal?

# Lower exponent optimality



## Theorem (F., Palombaro '17)

Let  $\sigma_1 = \text{diag}(1/K, 1/S_1)$ ,  $\sigma_2 = \text{diag}(K, S_2)$  with  $K > 1$  and  $S_1, S_2 \in [1/K, K]$ .  
There exist

- ▶ coefficients  $\sigma_n \in L^\infty(\Omega; \{\sigma_1; \sigma_2\})$ ,
- ▶ exponents  $p_n \in \left[1, \frac{2K}{K+1}\right]$ ,
- ▶ functions  $u_n \in W^{1,1}(\Omega)$  such that  $u_n(x) = x_1$  on  $\partial\Omega$ ,

such that

$$\begin{aligned} \operatorname{div}(\sigma_n \nabla u_n) &= 0, \\ \nabla u_n &\in L_{\text{weak}}^{p_n}(\Omega; \mathbb{R}^2), \quad p_n \rightarrow \frac{2K}{K+1}, \quad \nabla u_n \notin L^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2). \end{aligned}$$

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$



# Solving differential inclusions

## Theorem (Approximate solutions for two phases)


Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that

$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**



# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- rank-one connection allows to laminate in direction  $n$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,
- ▶  $\lambda$  proportion for  $A$ ,  $1 - \lambda$  proportion for  $B$ ,



Müller. *Variational models for microstructure and phase transitions*.

# Solving differential inclusions

## Theorem (Approximate solutions for two phases)

Let  $A, B \in \mathbb{M}^{2 \times 2}$ ,  $C := \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$ , and  $\delta > 0$ . Assume that


$$B - A = a \otimes n \quad \text{for some } a \in \mathbb{R}^2, n \in S^1. \quad (\text{Rank-one connection})$$

$\exists$  **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$  and

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Solutions:** built through **simple laminates**

- ▶ rank-one connection allows to laminate in direction  $n$ ,
- ▶  $\nabla f$  oscillates in  $\delta$ -neighbourhoods of  $A$  and  $B$ ,
- ▶  $\lambda$  proportion for  $A$ ,  $1 - \lambda$  proportion for  $B$ ,
- ▶ this allows to recover boundary data  $C$ .



Müller. *Variational models for microstructure and phase transitions*.

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$



# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.
- **Boundary condition** since the barycentre of  $\nu$  is  $\overline{\nu} := \int_{\mathbb{M}^{2 \times 2}} M d\nu(M) = C$ .

# Laminates of first order

$\mathcal{L}_\Omega^2$  is the normalised Lebesgue measure restricted to  $\Omega \rightsquigarrow \mathcal{L}_\Omega^2(B) := |B \cap \Omega|/|\Omega|$ .

## Gradient distribution

Let  $f: \Omega \rightarrow \mathbb{R}^2$  be Lipschitz. The **gradient distribution** of  $f$  is the Radon measure  $\nabla f_\#(\mathcal{L}_\Omega^2)$  on  $\mathbb{M}^{2 \times 2}$  defined by

$$\nabla f_\#(\mathcal{L}_\Omega^2)(V) := \mathcal{L}_\Omega^2((\nabla f)^{-1}(V)), \quad \forall \text{ Borel set } V \subset \mathbb{M}^{2 \times 2}.$$

Let  $f_\delta$  be the map given by the previous Theorem. Then as  $\delta \rightarrow 0$ ,

$$\nu_\delta := (\nabla f_\delta)_\#(\mathcal{L}_\Omega^2) \xrightarrow{*} \nu := \lambda \delta_A + (1 - \lambda) \delta_B \quad \text{in } \mathcal{M}(\mathbb{M}^{2 \times 2}).$$

The measure  $\nu$  is called a **laminate of first order**, and it encodes:

- **Oscillations** of  $\nabla f_\delta$  about  $\{A, B\}$  and their proportions.
- **Boundary condition** since the barycentre of  $\nu$  is  $\overline{\nu} := \int_{\mathbb{M}^{2 \times 2}} M d\nu(M) = C$ .
- **Integrability** since for  $p > 1$  we have

$$\frac{1}{|\Omega|} \int_\Omega |\nabla f_\delta|^p dx = \int_{\mathbb{M}^{2 \times 2}} |M|^p d\nu_\delta(M).$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

**New gradient:** apply previous Proposition to the set  $\{x \in \Omega: \nabla f \sim B\}$  to obtain  $\tilde{f}: \Omega \rightarrow \mathbb{R}^2$  such that  $\tilde{f}(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla \tilde{f}, \{A, B_1, B_2\}) < \delta \quad \text{a.e. in } \Omega.$$

# Iterating the Proposition

Let  $C = \lambda A + (1 - \lambda)B$  with  $\lambda \in [0, 1]$  and  $\text{rank}(B - A) = 1$ . Let  $f: \Omega \rightarrow \mathbb{R}^2$  such that  $f(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla f, \{A, B\}) < \delta \quad \text{a.e. in } \Omega.$$

**Further splitting:**  $B = \mu B_1 + (1 - \mu)B_2$  with  $\mu \in [0, 1]$ ,  $\text{rank}(B_2 - B_1) = 1$ .

**New gradient:** apply previous Proposition to the set  $\{x \in \Omega: \nabla f \sim B\}$  to obtain  $\tilde{f}: \Omega \rightarrow \mathbb{R}^2$  such that  $\tilde{f}(x) = Cx$  on  $\partial\Omega$ ,

$$\text{dist}(\nabla \tilde{f}, \{A, B_1, B_2\}) < \delta \quad \text{a.e. in } \Omega.$$

The gradient distribution of  $\tilde{f}$  is given by

$$\nu = \lambda \delta_A + (1 - \lambda)\mu \delta_{B_1} + (1 - \lambda)(1 - \mu) \delta_{B_2}.$$

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.



# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,
- ▶  $f(x) = Ax$  on  $\partial\Omega$ ,

# Laminates of finite order

**Laminates of finite order:** laminates obtained iteratively through the splitting procedure in the previous slide.

## Proposition (Convex integration)

Let  $\nu = \sum_{i=1}^N \lambda_i \delta_{A_i}$  be a laminate of finite order, s.t.

- ▶  $\bar{\nu} = A$ ,
- ▶  $A = \sum_{i=1}^N \lambda_i A_i$  with  $\sum_{i=1}^N \lambda_i = 1$ .

Fix  $\delta > 0$ .  $\exists$  a **piecewise affine Lipschitz** map  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ , that is,

- ▶  $\text{dist}(\nabla f, \text{supp } \nu) < \delta$  a.e. in  $\Omega$ ,
- ▶  $f(x) = Ax$  on  $\partial\Omega$ ,
- ▶  $|\{x \in \Omega : |\nabla f(x) - A_i| < \delta\}| = \lambda_i |\Omega|$ .

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- ① Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (0.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- 1 Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (0.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- 2 Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.



# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- ① Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (0.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- ② Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.  
③ Convex integration Proposition  $\implies$  construct  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ .  
In this way  $f$  solves (0.3) and

$$\nabla f \in L^q_{\text{weak}}(\Omega; \mathbb{R}^2), \quad q \in \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right], \quad \nabla f \notin L^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2).$$

# Strategy of the Proof

**Strategy:** explicit construction of  $u_n$  by **convex integration methods**.

- ① Rewrite the equation  $\operatorname{div}(\sigma \nabla u) = 0$  as a differential inclusion

$$\nabla f(x) \in T, \quad \text{for a.e. } x \in \Omega \quad (0.3)$$

for  $f: \Omega \rightarrow \mathbb{R}^2$  and an appropriate target set  $T \subset \mathbb{M}^{2 \times 2}$ .

**Note:**  $u$  and  $f$  have the **same** integrability.

- ② Construct a laminate  $\nu$  with  $\operatorname{supp} \nu \subset T$  and the right integrability.  
③ Convex integration Proposition  $\implies$  construct  $f: \Omega \rightarrow \mathbb{R}^2$  s.t.  $\nabla f \sim \nu$ .  
In this way  $f$  solves (0.3) and

$$\nabla f \in L^q_{\text{weak}}(\Omega; \mathbb{R}^2), \quad q \in \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right], \quad \nabla f \notin L^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2).$$

These methods were developed for isotropic conductivities  $\sigma \in L^\infty(\Omega; \{KI, \frac{1}{K}I\})$ .  
The adaptation to our case is non-trivial because of the lack of symmetry of the target set  $T$ , due to the anisotropy of  $\sigma_1$  and  $\sigma_2$ .

Astala, Faraco, Székelyhidi. *Convex integration and the  $L^p$  theory of elliptic equations*.

Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2008)

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$
$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$
$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

## Lemma (F., Palombaro '17)

A function  $u \in W^{1,1}(\Omega)$  is solution to

$$\text{div}(\sigma \nabla u) = 0$$

iff there exists  $v \in W^{1,1}(\Omega)$  such that  $f = (u, v): \Omega \rightarrow \mathbb{R}^2$  satisfies

$$\nabla f(x) \in T_1 \cup T_2 \quad \text{in } \Omega.$$

Moreover  $E_1 = \{x \in \Omega: \nabla f(x) \in T_1\}$  and  $E_2 = \{x \in \Omega: \nabla f(x) \in T_2\}$ .

# Rewriting the PDE as a differential inclusion

Let  $K > 1$ ,  $S_1, S_2 \in [1/K, K]$  and define

$$\sigma_1 := \text{diag}(1/K, 1/S_1), \quad \sigma_2 := \text{diag}(K, S_2), \quad \sigma := \sigma_1 \chi_{E_1} + \sigma_2 \chi_{E_2},$$
$$T_1 := \left\{ \begin{pmatrix} x & -y \\ S_1^{-1} y & K^{-1} x \end{pmatrix} : x, y \in \mathbb{R} \right\}, \quad T_2 := \left\{ \begin{pmatrix} x & -y \\ S_2 y & K x \end{pmatrix} : x, y \in \mathbb{R} \right\}.$$

## Lemma (F., Palombaro '17)

A function  $u \in W^{1,1}(\Omega)$  is solution to

$$\text{div}(\sigma \nabla u) = 0$$

iff there exists  $v \in W^{1,1}(\Omega)$  such that  $f = (u, v): \Omega \rightarrow \mathbb{R}^2$  satisfies

$$\nabla f(x) \in T_1 \cup T_2 \quad \text{in } \Omega.$$

Moreover  $E_1 = \{x \in \Omega: \nabla f(x) \in T_1\}$  and  $E_2 = \{x \in \Omega: \nabla f(x) \in T_2\}$ .

**Key Remark:**  $u$  and  $f$  enjoy the **same** integrability properties.

# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \overline{w}, \quad \forall w \in \mathbb{C}.$$

# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \overline{w}, \quad \forall w \in \mathbb{C}.$$

The sets of conformal linear maps and anti-conformal linear maps are

$$E_0 := \{(z, 0) : z \in \mathbb{C}\} \quad (\text{Conformal maps})$$

$$E_\infty := \{(0, z) : z \in \mathbb{C}\} \quad (\text{Anti-conformal maps})$$

# Targets in conformal coordinates

**Conformal coordinates:** Let  $A \in \mathbb{M}^{2 \times 2}$ . Then  $A = (a_+, a_-)$  for  $a_+, a_- \in \mathbb{C}$ , defined by

$$Aw = a_+ w + a_- \overline{w}, \quad \forall w \in \mathbb{C}.$$

The sets of conformal linear maps and anti-conformal linear maps are

$$E_0 := \{(z, 0) : z \in \mathbb{C}\} \quad (\text{Conformal maps})$$

$$E_\infty := \{(0, z) : z \in \mathbb{C}\} \quad (\text{Anti-conformal maps})$$

**Target sets** in conformal coordinates are

$$T_1 = \{(a, d_1(\overline{a})) : a \in \mathbb{C}\}, \quad T_2 = \{(a, -d_2(\overline{a})) : a \in \mathbb{C}\},$$

where the operators  $d_j: \mathbb{C} \rightarrow \mathbb{C}$  are defined as


$$d_j(a) := k \operatorname{Re} a + i s_j \operatorname{Im} a, \quad \text{with} \quad k := \frac{K-1}{K+1} \quad \text{and} \quad s_j := \frac{S_j-1}{S_j+1}.$$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$JR_\theta =$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$$JR_\theta = \lambda_1 A_1 + (1 - \lambda_1) P_1$$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$2JR_\theta = \lambda_2 A_2 + (1 - \lambda_2) P_2$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$




# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.



# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

►  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$



# Staircase Laminate (F., Palombaro '17)



Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$





# Staircase Laminate (F., Palombaro '17)



Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$



# Staircase Laminate (F., Palombaro '17)


Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .

$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.

- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^{p(\theta)} d\nu_n(M) \rightarrow \infty$  as  $n \rightarrow \infty$



# Staircase Laminate (F., Palombaro '17)

Let  $\theta \in [0, 2\pi]$ ,  $JR_\theta = (0, e^{i\theta})$ .


$$\begin{aligned} JR_\theta &= \lambda_1 A_1 + (1 - \lambda_1) P_1 \\ &= \lambda_1 A_1 + (1 - \lambda_1)(\mu_1 B_1 + (1 - \mu_1) 2JR_\theta) \\ &\leadsto \nu_1 \end{aligned}$$

$$\begin{aligned} 2JR_\theta &= \lambda_2 A_2 + (1 - \lambda_2) P_2 \\ &= \lambda_2 A_2 + (1 - \lambda_2)(\mu_2 B_2 + (1 - \mu_2) 3JR_\theta) \\ &\leadsto \nu_2 \end{aligned}$$

**Lemma:**  $\exists p(\theta) \in [\frac{2S}{S+1}, \frac{2K}{K+1}]$  continuous, with  $p(0) = \frac{2K}{K+1}$  and a sequence  $\nu_n$  of laminates s.t.


- ▶  $\text{supp } \nu_n \subset T_1 \cup T_2 \cup E_\infty$
- ▶  $\bar{\nu}_n = JR_\theta$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^q d\nu_n(M) < \infty, \quad \forall q < p(\theta)$
- ▶  $\int_{\mathbb{M}^{2 \times 2}} |M|^{p(\theta)} d\nu_n(M) \rightarrow \infty$  as  $n \rightarrow \infty$

**Remark:** barycentre  $J$  gives the right growth.



# Constructing approximate solutions


We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,



# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that

- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,
- ▶  $\nabla f \notin L^{\frac{2K}{K+1}}$ .






# Constructing approximate solutions

We want to construct  $f: \Omega \rightarrow \mathbb{R}^2$  such that


- ▶  $\text{dist}(\nabla f, T_1 \cup T_2) < \varepsilon$  a.e. in  $\Omega$ ,
- ▶  $f = Jx$  on  $\partial\Omega$ ,
- ▶  $\nabla f \in L^q_{\text{weak}}, q \in I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ ,
- ▶  $\nabla f \notin L^{\frac{2K}{K+1}}$ .

**Idea:** alternate one step of the staircase laminate with the convex integration Proposition.



# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$




# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$



# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$



# Constructing approximate solutions


Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$





# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .


**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$



**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$



## Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$


**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \leadsto$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
 and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $l_\delta$

### Step 1. One step of the staircase

- ▶ Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- ▶ Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- ▶  $\leadsto$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$


**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$





# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$


This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase


► Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$

► Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$

►  $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

► Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $v_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } v_1 \implies \nabla f_2$  grows like  $p_1$


This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $v_2$  with  $\bar{v}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } v_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$


**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$


**Step 1.** One step of the staircase

- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$



# Constructing approximate solutions

Recall  $I_\delta := \left( \frac{2K}{K+1} - \delta, \frac{2K}{K+1} \right]$ .

**Step A.** Define  $f_1(x) := Jx \implies \theta_1 = 0, p_1 = \frac{2K}{K+1}$

**Step B.** Laminate  $\nu_1$  from  $J$  to  $2J \rightsquigarrow$  growth  $p_1$

**Step C.** Proposition  $\implies \exists$  map  $f_2$  s.t.  $f_2 = Jx$  on  $\partial\Omega$   
and  $\nabla f_2 \sim \text{supp } \nu_1 \implies \nabla f_2$  grows like  $p_1$

This determines the exponent range  $I_\delta$

**Step 1.** One step of the staircase



- Split  $W_1$ . Since  $W_1 \sim 2J \implies$  point  $(2 + \rho)JR_{\theta_2}$  with  $\theta_2, \rho$  small.  $\implies p_2 \in I_\delta$
- Climb from  $(2 + \rho)JR_{\theta_2}$  to  $3JR_{\theta_2}$
- $\rightsquigarrow$  Laminate  $\nu_2$  with  $\bar{\nu}_2 = W_1$  and growth  $p_2$

**Step 2.** Define map  $f_3$  by modifying  $f_2$

- Proposition  $\implies \exists$  map  $g$  s.t.  $g = W_1 x$  on  $\partial\Omega$   
and  $\nabla g \sim \text{supp } \nu_2 \implies \nabla g$  grows like  $p_2$
- Set  $f_3 := g$  in the set  $\{\nabla f_2 \sim W_1\}$  and  $f_3 := f_2$  otherwise  $\implies \nabla f_3$  grows like  $p_2$

**Step 1.** Split  $W_2 \rightsquigarrow$  Laminate  $\nu_3$  with growth  $p_3 \in I_\delta$

**Iterating:**  $\rightsquigarrow f_n$  obtained by successive modifications  
on nested sets going to zero in measure  $\implies f_n \rightarrow f$



# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).



# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- ▶ Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- ▶ We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (0.4) and s.t.  $\nabla u \in L^{\frac{2K}{K+1}}_{\text{weak}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .  
Modifying staircase laminate?

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (0.4) and s.t.  $\nabla u \in L_{\text{weak}}^{\frac{2K}{K+1}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .

Modifying staircase laminate?

- Extend these results to **three-phase** conductivities  $\sigma \in \{\sigma_1, \sigma_2, \sigma_3\}$ .

# Conclusions and Perspectives

**Conclusions:** analysis of critical integrability of distributional solutions to

$$\operatorname{div}(\sigma \nabla u) = 0, \quad \text{in } \Omega, \quad (0.4)$$

when  $\sigma \in \{\sigma_1, \sigma_2\}$  for  $\sigma_1, \sigma_2 \in \mathbb{M}^{2 \times 2}$  elliptic.

- ▶ Optimal exponents  $q_{\sigma_1, \sigma_2}$  and  $p_{\sigma_1, \sigma_2}$  were already characterised and the upper exponent  $p_{\sigma_1, \sigma_2}$  was proved to be optimal.

Nesi, Palombaro, Ponsiglione. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2014).

- ▶ We proved the optimality of the lower critical exponent  $q_{\sigma_1, \sigma_2}$ .

## Perspectives:

- ▶ **Stronger** result for lower critical exponent: showing  $\exists u \in W^{1,1}(\Omega)$  solution to (0.4) and s.t.  $\nabla u \in L^{\frac{2K}{K+1}}_{\text{weak}}(\Omega; \mathbb{R}^2)$  but  $\nabla u \notin L^{\frac{2K}{K+1}}(B; \mathbb{R}^2)$ ,  $\forall$  ball  $B \subset \Omega$ .

Modifying staircase laminate?

- ▶ Extend these results to **three-phase** conductivities  $\sigma \in \{\sigma_1, \sigma_2, \sigma_3\}$ .
- ▶ **Dimension  $d \geq 3$ ?** Even only in the isotropic case  $\sigma \in \{KI, K^{-1}I\}$  for  $K > 1$ .  
Main difficulty: Astala's Theorem is missing in higher dimensions.

Thank You!