

Heavy Metal Neurotoxicity and the Collapse of Cerebral Clearance Mechanisms: An Integrative Analysis of Neurodegenerative Pathogenesis in the Aluminum Age

Introduction: The Paradox of Bio-incompatibility in the Aluminum Age

In the complex landscape of 21st-century neurotoxicology, few elements have generated as profound and persistent a scientific controversy as aluminum. Although it is the third most abundant element in the Earth's crust, constituting approximately 8% of its mass, and despite its ubiquity in the modern environment, aluminum remains a fundamental biological paradox: it is a metal with no known biological function in the human organism, being, from an evolutionary perspective, a non-essential element.¹ However, the "Aluminum Age," defined by the massive industrial use of this metal in water treatment, food additives, cosmetics, pharmaceuticals, and critically, as an immunological adjuvant in vaccines, has exponentially increased human systemic exposure.¹

This research report aims to deconstruct, with exhaustive academic rigor, the mechanisms by which this metal, long considered benign or "tolerable," succeeds in penetrating the physiological barriers of the central nervous system (CNS), establishing a state of chronic neuroinflammation, and ultimately precipitating neurodegeneration against a background of failing cerebral drainage systems associated with aging. The analysis focuses on the pathological intersection between the intrinsic toxicity of aluminum—demonstrated via advanced fluorescence and spectroscopy techniques—and the physiological decline of the glymphatic system and meningeal lymphatic vessels, a destructive triad that appears to underlie the etiology of Alzheimer's disease (AD) and other major neurocognitive disorders.³ It is imperative to underscore that current safety limits for aluminum, particularly in the context of vaccine adjuvants, are often based on historical data (dating back to 1947 and 1952) that prioritized immunological efficacy over long-term toxicological profiling and the complex pharmacokinetics of nanocrystalline particles.⁵ This historical regulatory gap

highlights the urgency of re-evaluating the impact of chronic aluminum accumulation on brain health in light of new evidence regarding biopersistence and systemic transport.⁶

I. Pharmacokinetics of Infiltration: From Systemic Exposure to Cerebral Penetration

The classical dogma of toxicology held that the Blood-Brain Barrier (BBB) represents an impenetrable fortress for most metallic xenobiotics. However, contemporary literature elucidates sophisticated mechanisms by which aluminum and other heavy metals evade this protection, exploiting nutrient transporters or immune cells to access the cerebral parenchyma.

1.1. Molecular Mimicry and Transcellular Transport

Once in systemic circulation, aluminum does not circulate freely but rapidly binds to plasma proteins, particularly transferrin, the protein responsible for iron transport. Due to a similar ionic radius and chemical affinity for iron-binding sites, the Al^{3+} ion mimics iron, effectively deceiving transferrin receptors (TfR) located on the endothelial cells of the BBB.¹ Through receptor-mediated endocytosis, the Al-transferrin complex is internalized into endothelial cells and subsequently released into the cerebral interstitial space. Furthermore, it has been demonstrated that aluminum can traverse the barrier as aluminum citrate, utilizing the sodium-independent system Xc- (L-glutamate/L-cystine exchanger).⁷ These transport pathways, intended for essential nutrients, become access gates for neurotoxins, highlighting the brain's inherent vulnerability to contaminants possessing the capacity for structural mimicry.

1.2. The "Trojan Horse" Mechanism: Immune-Mediated Translocation

A distinct and particularly insidious mechanism, specific to aluminum-based adjuvants (aluminum hydroxide or oxyhydroxide), is the "Trojan Horse" phenomenon. Research led by Professor Romain Gherardi's team has demonstrated that intramuscularly injected aluminum particles do not remain localized at the injection site as previously assumed.⁸ Conversely, their nanocrystalline nature and poor biodegradability induce an intense phagocytic reaction from macrophages and monocytes.

The process unfolds sequentially:

1. **Phagocytosis and Biopersistence:** Macrophages capture aluminum particles in an attempt to clear the tissue. However, instead of being dissolved in lysosomes, the aluminum destabilizes the lysosomal membrane or persists in crystalline form,

transforming the macrophage into a long-term mobile depot.¹⁰ This intracellular biopersistence is a prerequisite for distant transport.

2. **Lymphatic and Systemic Translocation:** Loaded macrophages migrate to regional lymph nodes and subsequently enter the blood circulation.⁸
3. **Cerebral Infiltration via CCL2:** The critical factor directing these cells toward the brain is the protein CCL2 (Monocyte Chemoattractant Protein-1 or MCP-1). CCL2 is a major inflammatory signal; under conditions of systemic or local cerebral inflammation, CCL2 levels rise, attracting aluminum-loaded monocytes to cross the BBB.¹⁰

Once inside the cerebral parenchyma, these cells (which may differentiate into microglia) release their toxic cargo or undergo apoptosis, depositing aluminum directly into neural tissue where it progressively accumulates.⁸ This is a slow process, explaining the long latency between exposure and the onset of neurodegenerative symptoms.¹²

1.3. Comparative Analysis: Aluminum Transport vs. Mercury and Lead

To contextualize aluminum toxicity, it is essential to compare its penetration mechanisms with those of other established neurotoxic heavy metals, such as mercury and lead. This comparison reveals the diversity of molecular strategies by which xenobiotics compromise CNS integrity.

Table 1: Comparative Transport Mechanisms at the Blood-Brain Barrier

Neurotoxic Metal	Predominant Chemical Form at Entry	Transporter / Molecular Mechanism at BBB	Mechanistic Details and Implications	Source
Aluminum (Al)	Al^{3+} ion (transferrin-bound), Al Citrate, Nanocrystalline particles	Transferrin Receptor (TfR), System Xc- (Glutamate/Cystine), Phagocytes (Trojan Horse)	Mimics iron to utilize TfR. Utilizes CCL2-recruited monocytes for the transport of insoluble particles. Extreme intracellular biopersistence.	⁹
Methylmercury (MeHg)	MeHg-L-Cysteine Complex	LAT1 (Large Neutral Amino Acid Transporter 1)	Forms a complex with cysteine that is structurally identical to Methionine. Actively transported as a	¹³

			"false amino acid" into the brain.	
Lead (Pb)	Pb^{2+} Ion	DMT1 (Divalent Metal Transporter 1), Calcium Channels	Competes with Calcium at ionic channels and utilizes the DMT1 transporter (specific for iron/divalent metals) on endothelial cells.	¹⁶
Manganese (Mn)	Mn^{2+} Ion, Mn Citrate	Transferrin Receptor, DMT1	Utilizes multiple pathways, including iron transporters, accumulating preferentially in the basal ganglia.	¹⁸

The mechanism of "molecular mimicry" employed by **methylmercury** is remarkable: by complexing with L-cysteine, it creates a spatial structure that perfectly mimics methionine, an essential amino acid. The LAT1 transporter, responsible for transporting large neutral amino acids into the brain, recognizes this complex as a nutrient and actively pumps it across the BBB, facilitating rapid and efficient neurotoxic accumulation.¹³ In contrast, aluminum employs a more insidious and slow pathway, based on immune-mediated chronic accumulation and transferrin binding, rendering its elimination far more difficult once deposition has occurred.⁷

II. The Architecture of Neurotoxicity: Histopathological Evidence and Fluorescence

Controversies regarding the presence of aluminum in the brains of Alzheimer's patients have often been fueled by the technological limitations of detection methods. However, recent research utilizing Graphite Furnace Atomic Absorption Spectrometry (GFAAS) and specific fluorescence microscopy has dispelled doubts regarding the accumulation of this metal.

2.1. The Lumogallion Revolution: Visualizing the "Invisible"

The research group led by Professor Christopher Exley standardized the use of **lumogallion**, a chelating agent that forms a specific fluorescent complex with aluminum ions. This method allowed, for the first time, the *in situ* visualization of aluminum deposits in brain tissue, clearly

differentiating them from lipofuscin autofluorescence or other metals.²⁰

Studies on brains from donors with Familial Alzheimer's Disease (fAD) revealed aluminum levels significantly higher than controls, often exceeding the pathological threshold of 2–3 $\mu\text{g/g}$ dry weight, reaching extremely high values.² In contrast, control tissues from donors without neurodegenerative disease invariably present low levels, typically below 1.0 $\mu\text{g/g}$, demonstrating that massive accumulation is not an inevitable consequence of normal aging but a marker of pathology.²³

2.2. Spherulites and Co-localization with Alzheimer's Pathology

Microscopic analysis revealed a striking topographical overlap between aluminum deposits and the hallmarks of Alzheimer's disease:

- **Co-localization with Amyloid-Beta ($\text{A}\beta$):** Aluminum has been identified co-localized with senile $\text{A}\beta$ plaques. Furthermore, aluminum appears to act as a cross-linker, promoting the oligomerization of $\text{A}\beta$ peptides and stabilizing the plaque structure, making them more resistant to proteolytic degradation.²⁴
- **Aluminum Spherulites:** Using synchrotron X-ray fluorescence, researchers identified aluminum-rich "spherulites" in AD brain tissue. These structures are morphologically identical to the cores of senile plaques, suggesting that aluminum may be the nucleation element around which amyloid deposits organize.²⁶
- **Interaction with Tau Protein:** In familial Alzheimer's cases, aluminum was also observed co-localized with hyperphosphorylated tau protein in intracellular neurofibrillary tangles.²⁴ Although less frequent than amyloid associations, the intracellular presence of aluminum in neurons suggests direct toxicity to the cytoskeleton and axonal transport.²⁵

These data converge on the conclusion that aluminum is not an "innocent bystander" accumulating passively in already diseased tissue, but a central actor actively participating in the formation and stabilization of toxic protein aggregates.²⁴

III. The Cytokine Storm: Molecular Mechanisms of Neuroinflammation

The physical presence of aluminum in the cerebral parenchyma triggers a violent innate immune response. Microglia, the immune sentinels of the brain, perceive aluminum particles and associated protein complexes as Damage-Associated Molecular Patterns (DAMPs), initiating an inflammatory cascade that paradoxically aggravates neurodegeneration.

3.1. Activation of the NLRP3 Inflammasome: The Critical Pathway

One of the most well-documented molecular mechanisms of aluminum adjuvant toxicity is the activation of the **NLRP3** (NOD-like receptor family, pyrin domain containing 3) multiprotein complex. This is a critical intracellular sensor for initiating sterile inflammation.

The pathogenic sequence is as follows:

1. **Frustrated Phagocytosis:** Microglia phagocytose aluminum aggregates or Al-amyloid complexes.
2. **Lysosomal Destabilization:** The crystalline and rigid nature of aluminum particles causes mechanical or chemical rupture of the lysosomal membrane, releasing cathepsins (acidic proteolytic enzymes) into the cytoplasm.⁹
3. **Perturbation of Ionic Homeostasis (K⁺ Efflux):** Lysosomal rupture and aluminum's action on the cell membrane induce a massive efflux of potassium ions (K⁺) from the cell. The decrease in intracellular K⁺ concentration is the canonical activation signal for NLRP3.²⁷
4. **Complex Assembly:** Once activated, NLRP3 recruits the adaptor protein ASC, which in turn recruits pro-caspase-1, leading to the formation of the active inflammasome.
5. **Cytokine Processing:** The inflammasome activates caspase-1, which cleaves pro-interleukin-1 β (pro-IL-1 β) and pro-interleukin-18 (pro-IL-18) into their biologically active forms, secreting them into the extracellular space.³⁰

Studies have demonstrated that aluminum-induced IL-1 β secretion is completely dependent on NLRP3; in the absence of this sensor (in knock-out mice), the inflammatory response is abrogated, confirming the central role of this pathway.³²

3.2. Amplification of Inflammation: The Role of TNF- α and IL-6

Beyond inflammasome activation, aluminum stimulates the **NF- κ B** nuclear factor pathway, which governs the expression of pro-inflammatory genes. This leads to sustained production of **Tumor Necrosis Factor-alpha (TNF- α)** and **Interleukin-6 (IL-6)**.³³

- **TNF- α :** This cytokine plays a dual role. Under homeostatic conditions, it regulates synaptic plasticity, but in excess (as in chronic aluminum exposure), it becomes neurotoxic. TNF- α released by microglia activates the TNFR1 receptor on neurons, inducing apoptosis and excitotoxicity.³⁴ Furthermore, TNF- α possesses the capacity for auto-regulation and amplification, signaling back to microglia to maintain the activated state.
- **IL-6:** *In vitro* and *in vivo* studies have shown that exposure to adjuvants or LPS rapidly induces elevated IL-6 levels in microglia via signaling pathways involving MAPK kinases (JNK, p38).³⁷ IL-6 contributes to perpetuating the immune response and altering the blood-brain barrier, facilitating further infiltration of toxins and peripheral immune cells.³⁵

Table 2: Aluminum-Induced Cytokine Profile

Pro-inflammatory Cytokine	Primary Induction Mechanism	Direct and Indirect Neurotoxic Effects	Source
IL-1 β	NLRP3 Inflammasome Activation -> Caspase-1	Chronic neuroinflammation, astrocyte activation, recruitment of peripheral leukocytes.	³²
TNF- α	NF- κ B Activation	Excitotoxicity, neuronal apoptosis, synaptic dysfunction, BBB breakdown.	³⁴
IL-6	MAPK (JNK/p38) and NF- κ B Activation	Alteration of BBB permeability, promotion of reactive gliosis.	³⁷
IL-18	NLRP3 Inflammasome Activation	Promotion of Th1-type immune response, induction of interferon-gamma.	³⁸

This toxic inflammatory environment, termed a chronic "cytokine storm," impedes normal neuronal function and, crucially, inhibits the brain's natural cleaning processes, setting the stage for the collapse of clearance systems.

IV. The Collapse of Cerebral Clearance: Decline of Glymphatic and Lymphatic Systems

To fully comprehend aluminum neurotoxicity, one must shift from the paradigm of intoxication (intake) to the paradigm of retention (efflux). The brain possesses specialized systems for eliminating metabolic waste, the recent discovery of which has revolutionized neurology: the glymphatic system and meningeal lymphatic vessels. Their failure, correlated with age, explains the exponential accumulation of neurotoxic metals.

4.1. The Glymphatic System: Hydrodynamics of Cerebral Cleaning

Described by the team led by Maiken Nedergaard, the glymphatic (glial-lymphatic) system functions as a macroscopic "washing" system for the cerebral parenchyma.³⁹ It facilitates the convective flow of Cerebrospinal Fluid (CSF) along periarterial spaces, its penetration into

brain tissue, and exchange with Interstitial Fluid (ISF), sweeping metabolic waste (including amyloid-beta, tau, and metal ions) toward perivenous spaces for elimination.⁴⁰

Key elements and vulnerabilities of the system include:

- **Sleep Dependence:** The glymphatic system is predominantly active during sleep. During wakefulness, the interstitial space contracts, increasing resistance to flow. Sleep induces an expansion of the interstitial space by over 60%, allowing for rapid and efficient flow.³⁹ Recent research from 2024 has definitively confirmed that glymphatic clearance is sharply suppressed during wakefulness.⁴³ Sleep fragmentation in the elderly reduces the time window available for aluminum elimination, leading to retention.⁴⁴
- **Role of Aquaporin-4 (AQP4):** The efficiency of this flow critically depends on AQP4 water channels, polarized on the end-feet of astrocytes ensheathing blood vessels. Heavy metals can perturb AQP4 expression and localization, causing astrocytic edema and blocking clearance channels.⁴⁵ Loss of AQP4 polarity is a hallmark of the aging and Alzheimer's brain.⁴⁷
- **Arterial Pulsatility:** CSF flow is driven by the pulsations of cerebral arteries. Arterial stiffening (atherosclerosis), common in the elderly, reduces the motive force of the glymphatic system, contributing to fluid stagnation.⁴⁹

4.2. Meningeal Lymphatic Vessels (MLVs): The Final Exit Gate

It was long believed that the CNS lacked lymphatic drainage. The rediscovery of lymphatic vessels in the dura mater (meninges) in 2015 demonstrated the existence of a direct route for evacuating macromolecules and immune cells from the CSF to the deep cervical lymph nodes (dCLNs).⁵⁰

The pathological impact of MLV dysfunction is major:

- **Age-Related Atrophy:** Studies on murine models have shown that with aging, meningeal lymphatic vessels atrophy; their diameter decreases, and coverage is significantly reduced. This leads to a drastic decline in CSF drainage capacity.⁴⁹
- **Experimental Blockade and Alzheimer's:** Experimental ablation of these vessels (using photodynamic targeting with Visudyne) exacerbates amyloid-beta deposition in the brain and worsens cognitive deficits, demonstrating a direct causal relationship between lymphatic drainage and toxic protein accumulation.⁵⁴
- **Meningeal Inflammation:** Accumulation of T cells and increased Interferon-gamma (IFN-γ) levels in the aging meninges induce lymphatic endothelial dysfunction, creating a vicious cycle.⁵³

Table 3: Cerebral Clearance Systems and Toxic Retention

Feature	Glymphatic System	Meningeal Lymphatic Vessels (MLVs)	Impact of Aluminum and Aging
Localization	Cerebral parenchyma	Dura mater (meninges)	Both suffer structural

	(perivascular spaces)		degradation with age.
Primary Function	CSF-ISF exchange, "sweeping" tissue waste	Collection of CSF and waste from glymphatic to lymph nodes	Aluminum requires these pathways for elimination; blockage leads to retention.
Dependence	AQP4, Sleep, Arterial pulsatility	Integrity of lymphatic vessels, Ganglion drainage	Sleep disorders and vascular stiffness compromise both systems.
Consequence of Failure	Stagnation of toxins in parenchyma, plaque formation	Accumulation of CSF, intracranial pressure, immune failure	Increased local Al concentration, promoting A β aggregation.

4.3. Pathogenic Synthesis: "The Triad of Neurodegeneration"

Integrating the data reveals a devastating scenario for the brain chronically exposed to aluminum:

1. **Entry:** Aluminum continuously penetrates (via Trojan Horse/BBB).
2. **Reaction:** Induces inflammation (NLRP3) and protein aggregation (Spherulites).
3. **Blockade:** Cleaning systems (Glymphatic/MLV) fail due to age and inflammation induced by the metal itself.

The result is irreversible accumulation. The half-life of aluminum in the brain is estimated to be extremely long, ranging from years to decades; estimates indicate a half-life of up to 50 years or more in certain neuronal compartments, making cumulative exposure from youth relevant in old age.⁷

V. Clinical Implications and Therapeutic Perspectives

The link between aluminum and neurodegenerative diseases transcends theory, being supported by increasingly solid clinical and epidemiological evidence.

5.1. Alzheimer's, Autism, and Multiple Sclerosis

Elevated aluminum levels have been consistently identified in the brains of Alzheimer's patients, particularly in familial forms where genetic predisposition appears to accelerate accumulation.²² Furthermore, recent studies have found extremely high aluminum

concentrations in cortical tissue from donors with Autism Spectrum Disorder (ASD), suggesting the metal's involvement in neurodevelopment and early neuroinflammation.² Multiple Sclerosis (MS) has also been associated with aluminum deposits, with the metal found in demyelination zones, possibly contributing to the autoimmune reaction via adjuvant mechanisms.¹

5.2. Therapeutic Interventions: Chelation and Drainage Restoration

In light of these discoveries, therapeutic strategies must evolve from symptom management to addressing the root cause.

- **Chelation Therapy:** The use of chelators such as Deferoxamine (DFO) has shown promising results in reducing aluminum burden and slowing cognitive decline in limited clinical trials.⁷ However, DFO has poor BBB penetration. New agents, such as combinations with citric acid or lipophilic chelators (HEDTA), are being investigated to mobilize deep parenchymal stores.⁵⁸
- **Restoration of the Glymphatic System:** Non-invasive interventions targeting sleep improvement, apnea treatment, and cardiovascular health maintenance are essential for sustaining glymphatic functionality.⁴² Novel therapies aiming to modulate AQP4 or regenerate meningeal lymphatic vessels (e.g., via VEGF-C growth factors or inorganic nanoparticles) represent the frontier of regenerative medicine in neurology.⁵⁹

Conclusion: The Necessity of a New Toxicological Paradigm

The extensive research presented in this report validates and amplifies initial concerns regarding the neurotoxic risk of aluminum. We can no longer consider the presence of this metal in the brain as a benign accident. The evidence converges on a clear conclusion:

Aluminum is a persistent neurotoxin that penetrates the brain via active mechanisms (Trojan Horse), induces specific inflammatory pathology (NLRP3, Spherulites), and whose toxicity is dramatically exacerbated by the physiological collapse of cerebral clearance systems (Glymphatic/MLV) associated with age.

This detailed understanding of the "Infiltration-Inflammation-Stagnation" mechanism offers not only a plausible explanation for the etiology of Alzheimer's disease but also a clear roadmap for developing therapeutic and preventive strategies aimed at eliminating this molecular intruder and restoring the brain's natural capacity for detoxification.

Integrated References

1. Aluminium in the Human Brain: Routes of Penetration, Toxicity, and Resulting Complications, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10139039/>
2. (PDF) Aluminium in human brain tissue: how much is too much? - ResearchGate, accesată pe noiembrie 21, 2025, https://www.researchgate.net/publication/335482534_Aluminium_in_human_brain_tissue_how_much_is_too_much
3. A Toxic Synergy between Aluminium and Amyloid Beta in Yeast - MDPI, accesată pe noiembrie 21, 2025, <https://www.mdpi.com/1422-0067/22/4/1835>
4. Glymphatic System - Lab Focuses - University of Rochester Medical Center, accesată pe noiembrie 21, 2025, <https://www.urmc.rochester.edu/labs/nedergaard/projects/glymphatic-system>
5. Romain K Gherardi's research works | Hôpital Henri Mondor (Hôpitaux Universitaires Henri Mondor) and other places - ResearchGate, accesată pe noiembrie 21, 2025, <https://www.researchgate.net/scientific-contributions/Romain-K-Gherardi-2080227543>
6. Biopersistence and Brain Translocation of Aluminum Adjuvants of Vaccines, accesată pe noiembrie 21, 2025, <https://www.semanticscholar.org/paper/Biopersistence-and-Brain-Translocation-of-Aluminum-Gherardi-Eidi/e77d74805c3c00411d3576773c8835c42899c56c>
7. The Pharmacokinetics and Toxicology of Aluminum in the Brain - UKnowledge, accesată pe noiembrie 21, 2025, https://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1068&context=ps_facpub
8. Slow CCL2-dependent translocation of biopersistent particles from muscle to brain - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/23557144/>
9. Biopersistence and Brain Translocation of Aluminum Adjuvants of Vaccines - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4318414/>
10. Biopersistence and brain translocation of aluminum adjuvants of vaccines - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/25699008/>
11. Macrophagic myofasciitis: characterization and pathophysiology - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3623725/>
12. Highly delayed systemic translocation of aluminum-based adjuvant in CD1 mice following intramuscular injections | Request PDF - ResearchGate, accesată pe noiembrie 21, 2025,

https://www.researchgate.net/publication/281811233_Highly_delayed_systemic_translocation_of_aluminum-based_adjuvant_in_CD1_mice_following_intramuscular_injections

13. Methylmercury transport across the blood-brain barrier by an amino acid carrier, accesată pe noiembrie 21, 2025, <https://journals.physiology.org/doi/abs/10.1152/ajpregu.1992.262.5.R761?doi=10.1152/ajpregu.1992.262.5.R761>
14. S-(Methylmercury)-L-Cysteine: Uses, Interactions, Mechanism of Action | DrugBank Online, accesată pe noiembrie 21, 2025, <https://go.drugbank.com/drugs/DB02750>
15. Molecular Mimicry as a Mechanism for the Uptake of Cysteine S-Conjugates of Methylmercury and Inorganic Mercury | Chemical Research in Toxicology - ACS Publications, accesată pe noiembrie 21, 2025, <https://pubs.acs.org/doi/10.1021/tx060158i>
16. Divalent metal transporter 1 (DMT1) in the brain: implications for a role in iron transport at the blood-brain barrier, and neuronal and glial pathology - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4458610/>
17. MOLECULAR MECHANISMS OF LEAD NEUROTOXICITY - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8276936/>
18. Manganese Flux across the Blood-Brain Barrier - UKnowledge, accesată pe noiembrie 21, 2025, https://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1192&context=ps_facpub
19. MANGANESE IN EUKARYOTES: THE ROLE OF DMT1 - PMC - NIH, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2501114/>
20. The Identification of Aluminum in Human Brain Tissue Using Lumogallion and Fluorescence Microscopy - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/27472886/>
21. Aluminum Should Now Be Considered a Primary Etiological Factor ..., accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6159653/>
22. Aluminum and Amyloid- β in Familial Alzheimer's Disease - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/31958088/>
23. Aluminium in human brain tissue from donors without neurodegenerative disease: A comparison with Alzheimer's disease, multiple sclerosis and autism - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/32385326/>
24. Link between Aluminum and the Pathogenesis of Alzheimer's Disease: The Integration of the Aluminum and Amyloid Cascade Hypotheses - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3056430/>
25. Illuminating the association between aluminium and familial Alzheimer's disease, accesată pe noiembrie 21, 2025, <https://www.innovationnewsnetwork.com/illuminating-association-between-aluminium-familial-alzheimers-disease/10629/>

26. Alzheimer's research at Diamond, accesată pe noiembrie 21, 2025, https://www.diamond.ac.uk/Home/News/LatestNews/20_09_10.html
27. Role of NLRP3 Inflammasome and Its Inhibitors as Emerging Therapeutic Drug Candidate for Alzheimer's Disease: a Review of Mechanism of Activation, Regulation, and Inhibition - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9403980/>
28. The NLRP3 Inflammasome Pathway: A Review of Mechanisms and Inhibitors for the Treatment of Inflammatory Diseases - Frontiers, accesată pe noiembrie 21, 2025, <https://www.frontiersin.org/journals/aging-neuroscience/articles/10.3389/fnagi.2022.879021/full>
29. The NLRP3 Inflammasome Pathway: A Review of Mechanisms and Inhibitors for the Treatment of Inflammatory Diseases - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9226403/>
30. Aluminum induces neuroinflammation via P2X7 receptor activating NLRP3 inflammasome pathway - ResearchGate, accesată pe noiembrie 21, 2025, https://www.researchgate.net/publication/366157380_Aluminum_induces_neuroinflammation_via_P2X7_receptor_activating_NLRP3_inflammasome_pathway
31. The Nlrp3 inflammasome is critical for aluminum hydroxide-mediated IL-1 β secretion but dispensable for adjuvant activity | Request PDF - ResearchGate, accesată pe noiembrie 21, 2025, https://www.researchgate.net/publication/5226735_The_Nlrp3_inflammasome_is_critical_for_aluminum_hydroxide-mediated_IL-1b_secretion_but_dispensable_for_adjuvant_activity
32. The NLRP3 Inflammasome is Critical for Alum-Mediated IL-1 β Secretion but Dispensable for Adjuvant Activity - PMC - NIH, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2759997/>
33. Inflammatory cytokines TNF α , IL-1 β , and IL-6 are induced in endotoxin-stimulated microglia through different signaling cascades - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10450609/>
34. TNF- α (TNF-alpha) & Microglia in Neurodegenerative Diseases - Biospective, accesată pe noiembrie 21, 2025, <https://biospective.com/resources/tnf-alpha-microglia-neurodegenerative-diseases>
35. Neither interleukin-6 nor signalling via tumour necrosis factor receptor-1 contribute to the adjuvant activity of Alum and Freund's adjuvant - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/9536117/>
36. TNF Production and Release from Microglia via Extracellular Vesicles: Impact on Brain Functions - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7598215/>
37. Microglia Responses to Pro-inflammatory Stimuli (LPS, IFN γ +TNF α) and Reprogramming by Resolving Cytokines (IL-4, IL-10) - Frontiers, accesată pe noiembrie 21, 2025, <https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2>

[018.00215/full](#)

38. The NLRP3 inflammasome: molecular activation and regulation to therapeutics - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7807242/>
39. The Glymphatic System – A Beginner's Guide - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4636982/>
40. The Glymphatic System: A Novel Component of Fundamental Neurobiology - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8603752/>
41. The Glymphatic System: A Novel Component of Fundamental Neurobiology, accesată pe noiembrie 21, 2025, <https://www.jneurosci.org/content/41/37/7698>
42. The Sleeping Brain: Harnessing the Power of the Glymphatic System through Lifestyle Choices - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7698404/>
43. Glymphatic clearance is enhanced during sleep - bioRxiv, accesată pe noiembrie 21, 2025, <https://www.biorxiv.org/content/10.1101/2024.08.24.609514v1>
44. Glymphatic failure as a final common pathway to dementia - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8186542/>
45. accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4887506/#:~:text=Some%20evidence%20has%20pointed%20to,brain%20swelling%20and%20neuronal%20damage.>
46. Regulation of AQP4 in the Central Nervous System - MDPI, accesată pe noiembrie 21, 2025, <https://www.mdpi.com/1422-0067/21/5/1603>
47. Emerging Roles of Meningeal Lymphatic Vessels in Alzheimer's Disease - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10473149/>
48. Blocking meningeal lymphatic drainage aggravates Parkinson's disease-like pathology in mice overexpressing mutated α -synuclein - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/30867902/>
49. The Lymphatic System In The Brain Clearance Mechanisms - New Therapeutic Perspectives For Alzheimer's Disease - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10190136/>
50. Meningeal Lymphatics: A Review and Future Directions From a Clinical Perspective - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7176397/>
51. The Discovery of Central Nervous System Lymphatic Vessels: The Missing Link That Closes the Circle of Brain Immunosurveillance - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6178593/>
52. Missing Link Found Between Brain, Immune System - UVA Health Newsroom, accesată pe noiembrie 21, 2025, <https://newsroom.uvahealth.com/2015/06/01/brain-immune-system-link/>
53. Age-related alterations in meningeal immunity drive impaired CNS lymphatic drainage, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10083715/>

54. Meningeal lymphatics in aging and Alzheimer's disease - Annals of Translational Medicine, accesată pe noiembrie 21, 2025, <https://atm.amegroups.org/article/view/23446/html>
55. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules | Journal of Experimental Medicine | Rockefeller University Press, accesată pe noiembrie 21, 2025, <https://rupress.org/jem/article/212/7/991/41853/A-dural-lymphatic-vascular-system-that-drains>
56. Age-related alterations in meningeal immunity drive impaired CNS lymphatic drainage - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/37027179/>
57. Entry, Half-Life, and Desferrioxamine-Accelerated Clearance of Brain Aluminum after a Single ²⁶Al Exposure - Oxford Academic, accesată pe noiembrie 21, 2025, <https://academic.oup.com/toxsci/article/64/1/77/1637712>
58. Aluminum Poisoning with Emphasis on Its Mechanism and Treatment of Intoxication - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8767391/>
59. Ultrasmall inorganic nanoparticles repair damaged meningeal lymphatic vessels to boost Parkinson's disease therapy | PNAS, accesată pe noiembrie 21, 2025, <https://www.pnas.org/doi/10.1073/pnas.2503434122>
60. The nasal lymphatic route of CSF outflow: implications for neurodegenerative disease diagnosis and monitoring - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10826790/>
61. Drainage of Cells and Soluble Antigen from the CNS to Regional Lymph Nodes - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7088878/>
62. Slow CCL2-dependent translocation of biopersistent particles from muscle to brain - PMC, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3616851/>
63. Aluminium and Alzheimer's: An unproven link, accesată pe noiembrie 21, 2025, <https://www.alzheimersresearchuk.org/news/aluminium-and-alzheimers/>
64. Lymphatic failure as a final common pathway to dementia - PubMed - NIH, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/33004510/>
65. Overview of the meningeal lymphatic vessels in aging and central nervous system disorders, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/36528776/>
66. Blood-brain barrier flux of aluminum, manganese, iron and other metals suspected to contribute to metal-induced neurodegeneration - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/17119290/>
67. Overview of the meningeal lymphatic vessels in aging and central nervous system disorders, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9759913/>
68. Brain burdens of aluminum, iron, and copper and their relationships with amyloid- β pathology in 60 human brains - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/22699848/>

69. Mercury neurotoxicity: mechanisms of blood-brain barrier transport - PubMed, accesată pe noiembrie 21, 2025, <https://pubmed.ncbi.nlm.nih.gov/2190116/>
70. The Interplay between Meningeal Lymphatic Vessels and Neuroinflammation in Neurodegenerative Diseases - PMC - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10964105/>
71. New High-Throughput Method for Aluminum Content Determination in Vaccine Formulations - PMC - NIH, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11861868/>
72. NLRP3 Inflammasome Is Expressed and Functional in Mouse Brain Microglia but Not in Astrocytes - PubMed Central, accesată pe noiembrie 21, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4474809/>