

# Introduction to Plack Stars

Alejandro Hernández A.

Universidad de los Andes, Bogotá, Colombia

November 6, 2015

# Overview

- 1 Introduction
  - Black hole solutions
- 2 Regularized Schwarzschild metric
  - Hayward metric
- 3 Quantum Field Theory
  - Newtonian Potential
  - Quantum cosmology remarks
  - Plank stars

# Preliminaries

## Einstein Field Equations

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi T_{\mu\nu} \quad (1)$$

## Minkowski metric

$$ds^2 = -dt^2 + dr^2 + r^2 d\Omega^2 \quad (2)$$

# Black hole solutions

## Classification of Black holes

	Non-rotating ( $J = 0$ )	Rotating ( $J \neq 0$ )
Uncharged ( $Q = 0$ )	Schwarzschild	Kerr
Charged ( $Q \neq 0$ )	Reissner-Nordström	Kerr-Newman

# Schwarzschild Black hole

## Schwarzschild metric

$$ds^2 = - \left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 d\Omega^2 \quad (3)$$

## Kretschmann invariant

$$\kappa = R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} = \frac{48m^2}{r^6} \quad (4)$$

# Regularized Schwarzschild metric

According to [1], we can find metrics that are:

- Spherically symmetric.
- Static.
- Asymptotically flat (minkowski).
- **Have regular center.**

The resulting stress-energy tensor is physically reasonable, satisfies the weak energy condition and has components that are bounded and fall off appropriately at large distance.

# Regularized Schwarchild metric

Consider a static, spherically symmetric metric of the form:

$$ds^2 = -F(r)dt^2 + \frac{1}{F(r)}dr^2 + r^2d\Omega^2 \quad (5)$$

We demand

$$F(r) \sim 1 - \frac{2m}{r} \quad \text{as } r \rightarrow \infty \quad (6)$$

$$F(r) \sim 1 - \frac{r^2}{l^2} \quad \text{as } r \rightarrow 0 \quad (7)$$

# Hayward metric

The so called Hayward metric [1] satisfies all the required properties and is given by:

$$F(r) = 1 - \frac{2mr^2}{r^3 + 2ml^2} \quad (8)$$

For this metric, the Kretschmann invariant is

$$OJO \quad (9)$$



# Hayward metric

Analyzing the zeros of  $F(r)$ , we get a critical mass  $m_* = \frac{3\sqrt{3}}{4}l$  and a radius  $r_* = \sqrt{3}l$ .




Figure : Behavior<sup>1</sup> of  $g_{tt} = F(r)$  for different values of the parameter  $m$ .

<sup>1</sup>Image taken from [1].

# Quantum Field Theory

Spacetime metric describing 'non-singular' black holes are commonly studied in the literature [2, 3] as effective modification to the Schwarzschild solution that mimic quantum gravity effects removing the central singularity.

To begin with, two insights from quantum cosmology [4]:

- The onset of quantum gravitational effects is when energy density reaches the Planck scale ( $\sim 5.155 \cdot 10^{96} \frac{\text{kg}}{\text{m}^3}$ ).
- The dominant quantum effect at high density is a strong pressure, sufficient to counterbalance weight and reverse gravitational collapse.

# Quantum Field theory

For a black hole, the previous arguments imply that matter's collapse can be stopped before the central singularity is formed, yielding the formation of a central core, called a **Planck star**.

# References



Hayward, S.A.: Formation and Evaporation of regular black holes. Phys. Rev. Lett. **96**, 31103 (2006).



De Lorenzo, T., Pacilio, C., Rovelli, C., Speziale, S.: On the effective metric of a Planck star. Gen. Relativ. Gravit. **47**, 41 (2015).



Rovelli, C., Vidotto, F.: Planck Stars. Int. J. Mod. Phys. D. **23**, 1142026, (2014).



Ashtekar, A., Pawłowski, T., Singh, P., Vandersloot, K.: Loop quantum cosmology of  $k=1$  FRW models. Phys. Rev. D **75**, 24035 (2007).

# The End