



FREEZEING-IN CANNIBAL DARK SECTORS

Andrzej Hryczuk



NATIONAL
CENTRE
FOR NUCLEAR
RESEARCH
ŚWIERK


based on:

E. Cervantes and AH [2407.12104](#) (to appear in JHEP)

Does DM interact (somewhat)
with any particle of the SM?

Does DM interact (somewhat)
with any particle of the SM?

Yes



At high temperature
DM thermalises, has
very high abundance



An **efficient**
depletion is needed
(freeze-out)

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?



An **efficient**
depletion is needed
(freeze-out)

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?

Yes

No

An **efficient
depletion** is needed
(freeze-out)

A **slow production**
is needed
(freeze-in)

„Initial condition”
dependence

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?

Yes

No

An **efficient
depletion** is needed
(freeze-out)

A **slow production**
is needed
(freeze-in)

„Initial condition”
dependence

depletion to SM

„WIMP”

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?

Yes

No

An **efficient
depletion** is needed
(freeze-out)

A **slow production**
is needed
(freeze-in)

„Initial condition”
dependence

depletion to SM

depletion
to other states

„WIMP”

Secluded
dark sector

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?

Yes

No

An **efficient
depletion** is needed
(freeze-out)

A **slow production**
is needed
(freeze-in)

„Initial condition”
dependence

„WIMP”

depletion to SM

to kinetic energy

depletion to other states

Secluded
dark sector
(e.g. cannibal dark sector)

Does DM interact (somewhat)
with any particle of the SM?

Yes

No (or nearly no)

At high temperature
DM thermalises, has
very high abundance

Is initial population (e.g.
from reheating) small?

Yes

No

An **efficient
depletion** is needed
(freeze-out)

A **slow production**
is needed
(freeze-in)

„Initial condition”
dependence

„WIMP”

depletion to SM

depletion
to other states
to kinetic energy

Secluded
dark sector

(e.g. cannibal dark sector)

CANNIBAL DM

Explains depletion of DM solely
through **self number changing**
reactions!

SELF-INTERACTING DARK MATTER

ERIC D. CARLSON

Lyman Laboratory of Physics, Harvard University, Cambridge, MA 02138

MARIE E. MACHACEK

Department of Physics, Northeastern University, Boston, MA 02115

AND


LAWRENCE J. HALL

Department of Physics, University of California; and Theoretical Physics Group, Physics Division,
Lawrence Berkeley Laboratory, 1 Cyclotron Road, Berkeley, CA 94720

Received 1992 March 17; accepted 1992 April 20

..., Hochberg et al.'14; ...

Simplest example, scalar ϕ with interactions (no coupling to SM!):

$$\frac{g}{3!}\phi^3 + \frac{\lambda}{4!}\phi^4 \Rightarrow$$


To obtain correct relic abundance: $m_\phi \sim \mathcal{O}(10 - 100 \text{ MeV})$

CANNIBAL DM

Explains depletion of DM solely
through **self number changing**
reactions!

SELF-INTERACTING DARK MATTER

ERIC D. CARLSON

Lyman Laboratory of Physics, Harvard University, Cambridge, MA 02138

MARIE E. MACHACEK

Department of Physics, Northeastern University, Boston, MA 02115

AND


LAWRENCE J. HALL

Department of Physics, University of California; and Theoretical Physics Group, Physics Division,
Lawrence Berkeley Laboratory, 1 Cyclotron Road, Berkeley, CA 94720

Received 1992 March 17; accepted 1992 April 20

..., Hochberg et al. '14; ...

Simplest example, scalar ϕ with interactions (no coupling to SM!):

$$\frac{g}{3!}\phi^3 + \frac{\lambda}{4!}\phi^4 \Rightarrow$$


To obtain correct relic abundance: $m_\phi \sim \mathcal{O}(10 - 100 \text{ MeV})$

cannibalisation: mass \rightarrow kinetic energy


expansion: kinetic energy redshifts \rightarrow depletion of E of the dark sector

Problem: structure formation...

de Laix et al. '95



CANNIBAL FREEZE-OUT

Self-heating of DM makes it warmer, **erasing the formation of structures!**
Tracking its temperature evolution is essential!



- Initially DM is *relativistic* ($T_{DM} \gg m_{DM}$);
- During freeze-out the dark sector uses its rest mass as *fuel* to keep itself warm;
- The system decouples and behaves as a non-relativistic gas.

PORTALS



EXAMPLE: SCALAR SINGLET DM

The simplest DM model: $\mathcal{L}_{SM} \supset -\frac{m^2}{2}\varphi^2 - \frac{\lambda}{4!}\varphi^4 - \frac{1}{2}\lambda_{h\varphi}\varphi^2 |H|^2$

$\langle\varphi\rangle = 0$ & $\lambda_{h\varphi} \gtrsim 10^{-4}$ „WIMP” (thermalizes, undergoes freeze-out)

$\langle\varphi\rangle = 0$ & $\lambda_{h\varphi} \lesssim 10^{-9}$ FIMP (feebly-interacting, undergoes freeze-in)

$\langle\varphi\rangle \neq 0$ Cannibal DM: spontaneous \mathbb{Z}_2 breaking \rightarrow Higgs mixing terms

Hufnagel, Tytgat '22

EXAMPLE: SCALAR SINGLET DM

The simplest DM model: $\mathcal{L}_{SM} \supset -\frac{m^2}{2}\varphi^2 - \frac{\lambda}{4!}\varphi^4 - \frac{1}{2}\lambda_{h\varphi}\varphi^2 |H|^2$

$\langle\varphi\rangle = 0$ & $\lambda_{h\varphi} \gtrsim 10^{-4}$ „WIMP” (thermalizes, undergoes freeze-out)

$\langle\varphi\rangle = 0$ & $\lambda_{h\varphi} \lesssim 10^{-9}$ FIMP (feebly-interacting, undergoes freeze-in)

$\langle\varphi\rangle \neq 0$ Cannibal DM: spontaneous \mathbb{Z}_2 breaking \rightarrow Higgs mixing terms

Hufnagel, Tytgat '22

Evolution of number density $Y_\varphi := n_\varphi/s$ and 'temperature' $x_\varphi := m_\varphi/T_\varphi$ governed by set of Boltzmann equations:

$$\begin{aligned} \frac{Y'_\varphi}{Y_\varphi} &= \frac{1}{x\tilde{H}} \left(\langle C_{h\rightarrow\varphi\varphi} \rangle + \langle C_{hh\rightarrow\varphi\varphi} \rangle + \langle C_{3\leftrightarrow 2} \rangle \right) \quad \text{Cannibal} \\ -\frac{x'_\varphi}{x_\varphi} &= \frac{1}{x\tilde{H}} \left(\langle C_{h\rightarrow\varphi\varphi} \rangle_2 + \langle C_{hh\rightarrow\varphi\varphi} \rangle_2 + \langle C_{\phi h \leftrightarrow \phi h} \rangle_2 + \langle C_{3\leftrightarrow 2} \rangle_2 \right) - \frac{Y'_\varphi}{Y_\varphi} + \frac{H}{x\tilde{H}} \frac{\langle p^4/E^3 \rangle}{3T_\varphi} + \frac{2s'}{3s} \end{aligned}$$



Freeze-in Freeze-in/out El. scattering

see also Ghosh et al. '22

BOOSTING FREEZE-IN

see also Bernal '05



Consider an initial cold, low populated dark sector: $T_{DM}^i = 10^{-2} T_{SM}^i$



BOOSTING FREEZE-IN

see also Bernal '05

Consider an initial cold, low populated dark sector: $T_{DM}^i = 10^{-2} T_{SM}^i$




only cannibalization


BOOSTING FREEZE-IN

see also Bernal '05

Consider an initial cold, low populated dark sector: $T_{DM}^i = 10^{-2} T_{SM}^i$



only cannibalization




only freeze-in


BOOSTING FREEZE-IN

see also Bernal '05

Consider an initial cold, low populated dark sector: $T_{DM}^i = 10^{-2} T_{SM}^i$



only cannibalization



only freeze-in


cannibal+freeze-in

THIS MODEL HAS AN ISSUE...

$$\mathbb{Z}_2 \text{ breaking} \Rightarrow \varphi \text{ unstable with } \tau_\varphi \propto 1/\lambda_{h\varphi}^2$$

Gray lines match
observed relic density

Colored areas are
excluded



For decaying DM only very
long lifetimes are allowed
 $\Rightarrow \lambda_{h\varphi}$ **needs to be very small**

The impact of our improved
treatment **negligible in the
allowed region**

... but still it is a viable model
(in the asymmetric reheating scenario)

goes to the scenario of Hufnagel, Tytgat '22

MODIFIED MODEL

Natural extension:

a complex (\mathbb{Z}_3 stabilised) **DM candidate** S with a **real singlet mediator** ϕ

$$\mathcal{L} \supset -\frac{1}{3!}g_s(S^3 + (S^*)^3) - \frac{\lambda_s}{4}|S|^4 - A_{\phi s}\phi|S|^2 - \frac{\lambda_{\phi s}}{2}\phi^2|S|^2 - B_{\phi h}\phi|H|^2$$

DM self interactions

thermalisation between ϕ and S

Portal

Portal to the visible sector $B_{\phi h}\phi|H|^2$ **induces mixing** between ϕ and the Higgs post-EWPT:

$$\phi \rightarrow \phi + \theta h$$

$$h \rightarrow h - \theta \phi$$



ϕ couples to matter in a Higgs-like way

Such a model exhibits new dynamics + provides chance of detection

BOLTZMANN EQUATIONS

DM evolution

Freeze-in

DM-mediator interactions

DM self-interactions

$$\frac{Y'_S}{Y_S} = \frac{1}{x \tilde{H}} \left(\langle C_{h \rightarrow \phi SS^*} \rangle + \langle C_{h \rightarrow SS^*} \rangle + \langle C_{\phi \phi \leftrightarrow SS^*} \rangle + \langle C_{3 \leftrightarrow 2} \rangle \right),$$

$$-\frac{x'_S}{x_S} = \frac{1}{x \tilde{H}} \left(\langle C_{h \rightarrow \phi SS^*} \rangle_2 + \langle C_{h \rightarrow SS^*} \rangle_2 + \langle C_{\phi S \leftrightarrow \phi S} \rangle_2 + \langle C_{3 \leftrightarrow 2} \rangle_2 \right) - \frac{Y'_S}{Y_S} + \frac{H}{x \tilde{H}} \frac{\langle p^4/E^3 \rangle}{3T_S} + \frac{2s'}{3s}$$


Mediator evolution

$$\frac{Y'_\phi}{Y_\phi} = \frac{1}{x \tilde{H}} \left(\langle C_{h \rightarrow \phi SS^*} \rangle + \langle C_{\text{sm sm} \rightarrow \text{sm } \phi} \rangle + \langle C_{\phi \phi \leftrightarrow SS^*} \rangle \right),$$

$$-\frac{x'_\phi}{x_\phi} = \frac{1}{x \tilde{H}} \left(\langle C_{h \rightarrow \phi SS^*} \rangle_2 + \langle C_{\text{sm sm} \rightarrow \text{sm } \phi} \rangle_2 + \langle C_{\phi S \leftrightarrow \phi S} \rangle_2 \right) - \frac{Y'_\phi}{Y_\phi} + \frac{H}{x \tilde{H}} \frac{\langle p^4/E^3 \rangle}{3T_\phi} + \frac{2s'}{3s}.$$

BBN & CMB CONSTRAINTS

Mediator is decaying to SM:
different phenomenology for two mass hierarchies between S (the DM) and ϕ




Resulting constraints are significant, but many viable points found in a scan & some with prospects of CMB distortion in reach of PIXIE



(The Primordial Inflation Explorer)

DM & MEDIATOR DETECTION

$m_S > m_\phi$: indirect DM
detection feasible



$m_S < m_\phi$: weaker BBN
limits allow for points
that are potentially
testable in LLP searches



CONCLUSIONS

1. **Frozen-in** Cannibal Dark Matter is a viable scenario, naturally avoiding the large scale structure formation limits plaguing cannibalising DM.
2. It has unique evolution in the Early Universe & potentially **detectable signals in indirect searches**.
3. Temperature (and momentum distribution) can have a **non-trivial impact** in such scenarios and a **need to be studied** carefully.



Thank you!