

AI505 – Optimization

Sheet 08, Spring 2025

Exercises with the symbol $+$ are to be done at home before the class. Exercises with the symbol $*$ will be tackled in class. The remaining exercises are left for self training after the exercise class. Some exercises are from the text book and the number is reported. They have the solution at the end of the book.

Exercise 1⁺ (13.1)

Filling a multidimensional space requires exponentially more points as the number of dimensions increases. To help build this intuition, determine the side lengths of an n -dimensional hypercube such that it fills half of the volume of the n -dimensional unit hypercube.

Exercise 2⁺ (13.2)

Suppose that you sample randomly inside a unit sphere in n dimensions. Compute the probability that a randomly sampled point is within ϵ distance from the surface of the sphere as $n \rightarrow \infty$. Hint: The volume of a sphere is $C(n)r^n$, where r is the radius and $C(n)$ is a function of the dimension n only.

Exercise 3⁺

Generate a full factorial set of design points in Python using `numpy.meshgrid` for two dimensions and plot the generated points. Find a function from the benchmark suite and evaluate the function in those points.

Repeat the same process for a function with $n = 3$ and for $n > 3$ dimensions.

Exercise 4^{*}

Construct in Python a Latin Hypercube design plan with numbers 1 through $m = 4$ for 3 dimensions. How many different Latin Hypercubes are there? How many different uniform projection plans are there for a single Latin Hypercube? How many different uniform projection plans can be generated from all Latin Hypercubes of m numbers in 3 dimensions?

Exercise 5

Consult the documentation of the quasi-Monte Carlo python submodule `scipy.stats.qmc` and repeat the previous exercises using those functions.

Exercise 6^{*} Quasi-random sequences are typically constructed for the unit n -dimensional hypercube, $[0, 1]^n$. Any multidimensional function with bounds on each variable can be transformed into such a hypercube. Show how.

Exercise 7^{*} (13.3)

Suppose we have a sampling plan $X = \{x_1, \dots, x_{10}\}$, where

$$x_i = [\cos(2\pi i/10), \sin(2\pi i/10)]$$

Compute the Morris-Mitchell criterion for X using an L_2 norm when the parameter q is set to 2. In other words, evaluate $\Phi_2(X)$. If we add $[2, 3]$ to each x_i , will $\Phi_2(X)$ change? Why or why not?

Exercise 8 (13.4)

Additive recurrence requires that the multiplicative factor c be irrational. Why can c not be rational?