

Machine Learning

Linear regression with one variable

Gradient descent

Have some function $J(\theta_0, \theta_1)$

Want $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$

Outline:

- Start with some θ_0, θ_1
- Keep changing θ_0, θ_1 to reduce $J(\theta_0, \theta_1)$ until we hopefully end up at a minimum

começamos neste ponto froma random acho eu
depois 'olhamos nossa volta' e escolhemos proximo de forma minimizar função J(x0,x1)




2 pontos começo diferente acabam com 2 minimos locais
J(θ0,θ1)
θ0
θ1
3
2
1
0
-1
-2
-3
1
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0
0
0.2
0.4
0.6
0.8
1

igual a programação

Assignment

$a := b$

$a := a+1$

Truth assertion

$a = b$

$a = a+1$ ✗

Gradient descent algorithm

repeat until convergence {

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) \quad \text{(for } j = 0 \text{ and } j = 1\text{)}$$

}

learning rate

Simultaneously update θ_0 and θ_1

Correct: Simultaneous update

$$\text{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$
$$\text{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$
$$\theta_0 := \text{temp0}$$
$$\theta_1 := \text{temp1}$$

Incorrect:

$$\text{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$
$$\theta_0 := \text{temp0}$$
$$\text{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$
$$\theta_1 := \text{temp1}$$