

Machine Learning

Linear regression with one variable

Cost function intuition II

Hypothesis: $h_\theta(x) = \theta_0 + \theta_1 x$

Parameters: θ_0, θ_1

Cost Function: $J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^{m} \left(h_\theta(x^{(i)}) - y^{(i)} \right)^2$

Goal: $\underset{\theta_0, \theta_1}{\text{minimize}}\ J(\theta_0, \theta_1)$

$h_\theta(x)$

(for fixed θ_0, θ_1, this is a function of x)



$$h_\theta(x) = 50 + 0.06x$$

$J(\theta_0, \theta_1)$

(function of the parameters θ_0, θ_1)



Contour plots

Contour figures -



$J(\theta_0, \theta_1)$

$h_\theta(x)$

(for fixed θ_0, θ_1, this is a function of x)



$J(\theta_0, \theta_1)$

(function of the parameters θ_0, θ_1)

those three have the same Value for J (theta zero, theta one)



θ0, θ1

$h_\theta(x)$

(for fixed θ_0, θ_1, this is a function of x)



$J(\theta_0, \theta_1)$

(function of the parameters θ_0, θ_1)



$\theta_0 = 360$

$\theta_1 = 0$

$h_\theta(x)$

(for fixed θ_0, θ_1, this is a function of x)



$J(\theta_0, \theta_1)$

(function of the parameters θ_0, θ_1)



$h_\theta(x)$

(for fixed θ_0, θ_1, this is a function of x)



$J(\theta_0, \theta_1)$

(function of the parameters θ_0, θ_1)