# EdgeCDSS kit packet: vial strengths to sign

2026-10-28. One page per drug with a signed dose in the bank: 25 drugs. Each lists the strengths the kit file and SMOG CY24 show, a SIGN line per strength, and whether one strength or two is stocked. **Nothing in this packet is signed.**

Why it matters: a millilitre is served only from a signed strength. One strength stocked: the volume is served without asking. Two: the medic is asked which vial first. None signed: mg only.

Kit file: /home/andrew/pi-cloud-cdss/server/drug_concentrations.json. SMOG: SMOG_CY24_REVISION_FINAL.pdf (printed page = PDF page); quotes are checked against the PDF's text when the packet is built; every quote was found.

Drugs: ketamine, epinephrine, rocuronium, succinylcholine, fentanyl, norepinephrine, cefazolin, ertapenem, moxifloxacin, ceftriaxone, tranexamic acid, midazolam, morphine, dextrose, propofol, levetiracetam, calcium chloride, calcium gluconate, naloxone, phytomenadione, lorazepam, artesunate, atropine, sodium chloride 3%, acetaminophen

# ketamine

**Signed doses in the bank** (20): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| post-intubation sedation — repeated bolus (no infusion pump) | adult\|peds | IV | 0.5 mg/kg |
| prolonged sedation infusion | adult | IV | 0.5–2 mg/kg/hr |
| moderate to severe pain / analgesia | adult\|peds | IV | 0.25 mg/kg |
| moderate to severe pain / analgesia | peds | IV | 0.2 mg/kg |
| agitated or violent patient — dissociative sedation | adult | IV | 2 mg/kg |
| agitated or violent patient — dissociative sedation | adult | IM | 4 mg/kg |
| agitated or violent patient — dissociative sedation | peds | IV | 1 mg/kg |
| agitated or violent patient — dissociative sedation | peds | IM | 3 mg/kg |
| RSI induction | adult | IV | 2 mg/kg |
| RSI induction — haemodynamically unstable / in extremis | adult | IV | 1 mg/kg |
| RSI induction | peds | IV | 2 mg/kg |
| ongoing sedation — loading dose (infusion pump available) | adult\|peds | IV | 1 mg/kg |
| background pain — low-dose analgesia (prolonged field care) | adult | IV | 0.1–0.2 mg/kg |
| refractory seizure (benzodiazepine-refractory) | adult | IV | 2 mg/kg |
| refractory seizure (benzodiazepine-refractory) | adult | IM | 3–4 mg/kg |
| refractory seizure (benzodiazepine-refractory) | peds | IV | 1 mg/kg |
| refractory seizure (benzodiazepine-refractory) | peds | IM | 3 mg/kg |
| breakthrough pain — IM/IN (prolonged field care) | adult | IM | 0.5–1 mg/kg |
| breakthrough pain — IM/IN (prolonged field care) | adult | IN | 0.5–0.75 mg/kg |
| moderate to severe pain / analgesia | peds | IM | 0.5 mg/kg |

**The kit file** (drug_concentrations.json)

- 500 mg / 10 mL vial = 50 mg/mL: signed AI-AIM 2026-08-25
- 200 mg / 20 mL vial = 10 mg/mL: declared, unsigned

The kit asks which vial every time (confirm_required).

**SMOG CY24**

1 mg/mL, infusion mix, not a vial: reference only. p.128: “MIX 500 mg/500 mL ... CONCENTRATION 1 mg/mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 500 mg / 10 mL vial (50 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Kit: 200 mg / 20 mL vial (10 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# epinephrine

**Signed doses in the bank** (13): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| anaphylaxis | adult | IM | 0.3 mg |
| anaphylaxis | peds | IM | 0.15 mg |
| cardiac arrest | adult | IV | 1 mg |
| cardiac arrest | peds | IV | 0.01 mg/kg |
| push-dose vasopressor for hypotension / shock (SBP < 90) | adult | IV | 5–20 mcg |
| infusion for refractory shock | adult | IV | 2–10 mcg/min |
| symptomatic bradycardia — infusion | adult | IV | 0.02–0.2 mcg/kg/min |
| symptomatic bradycardia — push dose | adult | IV | 10–20 mcg |
| shock unresponsive to IV fluids — infusion | adult\|peds | IV | 0.05–0.3 mcg/kg/min |
| shock — push dose | peds | IV | 0.01 mg/kg |
| neonatal resuscitation — HR below 60 despite ventilation | peds | IV | 0.01 mg/kg |
| croup with stridor at rest — nebulised | peds | NEB | 5 mg |
| bronchiolitis with severe respiratory distress — nebulised | peds | NEB | 5 mg |

**The kit file** (drug_concentrations.json)

- 1 mg / 1 mL ampoule = 1 mg/mL: declared, unsigned

The kit does not ask which vial.

**SMOG CY24**

1 mg/mL, vial strength. p.113: “Epinephrine 1mg/1ml (1:1,000)”

0.1 mg/mL, vial strength. p.114: “IV: 1mg/10mL (0.1mg/mL) pre-filled 10cc Syringe”

2 mcg/mL, infusion mix, not a vial: reference only. p.113: “MIX 1 mg/500 mL ... CONCENTRATION 2 mcg/mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 1 mg / 1 mL ampoule (1 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| SMOG p.114: 0.1 mg/mL, not in the kit file (declare: ____ mg in ____ mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# rocuronium

**Signed doses in the bank** (2): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| RSI paralytic | adult | IV | 1.2 mg/kg |
| RSI paralytic | peds | IV | 1 mg/kg |

**The kit file** (drug_concentrations.json)

- 100 mg / 10 mL vial = 10 mg/mL: signed AI-AIM 2026-08-25

The kit does not ask which vial.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 100 mg / 10 mL vial (10 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# succinylcholine

**Signed doses in the bank** (3): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| RSI paralytic | adult | IV | 1.5 mg/kg |
| RSI paralytic — under 5 years | peds | IV | 2 mg/kg |
| RSI paralytic — 5 years and above | peds | IV | 1.5 mg/kg |

**The kit file** (drug_concentrations.json)

- 100 mg / 2 mL ampoule = 50 mg/mL: signed AI-AIM 2026-08-25

The kit does not ask which vial.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 100 mg / 2 mL ampoule (50 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# fentanyl

**Signed doses in the bank** (3): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| acute pain / analgesia | peds | IV | 1 mcg/kg |
| acute pain / analgesia | adult\|peds | IN | 1 mcg/kg |
| acute pain / analgesia — JTS PFC fixed dose | adult | IV | 50 mcg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

10 mcg/mL, infusion mix, not a vial: reference only. p.119: “MIX 1 mg/100 mL ... CONCENTRATION 10 mcg/mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# norepinephrine

**Signed doses in the bank** (4): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| vasodilatory/haemorrhagic shock infusion | adult | IV | 0.05–0.5 mcg/kg/min |
| vasodilatory shock infusion | peds | IV | 0.05–0.5 mcg/kg/min |
| symptomatic bradycardia — infusion | adult | IV | 0.02–0.4 mcg/kg/min |
| cardiogenic shock / pulmonary oedema with SBP under 100 | adult | IV | 0.02–2 mcg/kg/min |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

8 mcg/mL, infusion mix, not a vial: reference only. p.147: “MIX 4 mg/500 mL ... CONCENTRATION 8 mcg/mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# cefazolin

**Signed doses in the bank** (8): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| open fracture / wound prophylaxis | adult | IV | 2 g |
| open fracture / wound prophylaxis | peds | IV | 20–30 mg/kg |
| penetrating brain injury (post-injury antimicrobial prophylaxis) | adult | IV | 2 g |
| penetrating spinal cord injury (post-injury antimicrobial prophylaxis) | adult | IV | 2 g |
| penetrating chest injury without esophageal disruption (post-injury prophylaxis) | adult | IV | 2 g |
| open maxillofacial fracture (post-injury prophylaxis) | adult | IV | 2 g |
| single pre-surgical dose for open wounds (Role 1 / Role 2 without surgical support) | adult | IV | 2 g |
| ventriculostomy in place (prophylaxis) | adult | IV | 1 g |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

8 mg/mL, infusion mix, not a vial: reference only. p.103: “War wounds (dirty wounds), 2g in 250 mL NS IV”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# ertapenem

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| penetrating abdominal injury | adult | IV | 1 g |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

4 mg/mL, infusion mix, not a vial: reference only. p.116: “Give 1g in 250 mL NS IV over 5 min”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# moxifloxacin

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| combat wound prophylaxis (oral) | adult | PO | 400 mg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# ceftriaxone

**Signed doses in the bank** (4): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| sepsis | adult | IV | 2 g |
| open or penetrating TBI (CNS-penetrating antimicrobial prophylaxis) | adult | IV | 2 g |
| penetrating TBI / CNS injury (post-injury prophylaxis) | peds | IV | 100 mg/kg |
| severe COVID-19 with possible bacterial pneumonia co-infection | adult | IV | 2 g |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# tranexamic acid

**Signed doses in the bank** (2): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| traumatic haemorrhage — loading dose | adult | IV | 2 g |
| traumatic haemorrhage | peds | IV | 15 mg/kg |

**The kit file** (drug_concentrations.json)

- 1000 mg / 10 mL ampoule = 100 mg/mL: signed AI-AIM 2026-08-25

The kit does not ask which vial.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 1000 mg / 10 mL ampoule (100 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# midazolam

**Signed doses in the bank** (10): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| active seizure | adult | IV | 0.1 mg/kg |
| active seizure | peds | IM | 0.2 mg/kg |
| procedural sedation | adult | IV | 1–4 mg |
| procedural sedation for pacing / cardioversion | adult | IV | 2.5–5 mg |
| agitated or violent patient / behavioural emergency | adult | IV | 5 mg |
| ongoing sedation (prolonged field care) | adult | IV | 0.5–2 mg |
| procedural sedation (incl. cardioversion, transcutaneous pacing) | peds | IV | 0.05–0.1 mg/kg |
| procedural / conscious sedation | peds | IN | 0.2–0.5 mg/kg |
| post-intubation sedation bolus | adult | IV | 1–4 mg |
| sedation / anxiolysis | adult | IM | 2–4 mg |

**The kit file** (drug_concentrations.json)

- 5 mg / 1 mL vial = 5 mg/mL: declared, unsigned
- 5 mg / 5 mL vial = 1 mg/mL: declared, unsigned

The kit asks which vial every time (confirm_required).

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 5 mg / 1 mL vial (5 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Kit: 5 mg / 5 mL vial (1 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# morphine

**Signed doses in the bank** (4): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| acute pain / analgesia | adult | IV | 0.1 mg/kg |
| acute pain / analgesia | adult | IM | 0.1 mg/kg |
| acute pain / analgesia | peds | IV | 0.1 mg/kg |
| acute pain / analgesia | peds | IM | 0.1 mg/kg |

**The kit file** (drug_concentrations.json)

- 10 mg / 1 mL ampoule = 10 mg/mL: declared, unsigned
- 2 mg / 1 mL ampoule = 2 mg/mL: declared, unsigned
- 1 mg / 1 mL ampoule = 1 mg/mL: declared, unsigned

The kit asks which vial every time (confirm_required).

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 10 mg / 1 mL ampoule (10 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Kit: 2 mg / 1 mL ampoule (2 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Kit: 1 mg / 1 mL ampoule (1 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# dextrose

**Signed doses in the bank** (5): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| symptomatic hypoglycaemia | adult | IV | 25 g |
| symptomatic hypoglycaemia | peds | IV | 0.5–1 g/kg |
| neonatal hypoglycaemia | peds | IV | 0.5 g/kg |
| symptomatic hypoglycaemia — alert patient | adult | PO | 15 g |
| symptomatic hypoglycaemia — alert patient | peds | PO | 0.5 g/kg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

500 mg/mL, vial strength. p.105: “10-25 g (40-100 mL of 25% solution or 20-50 mL of 50% solution)”

250 mg/mL, vial strength. p.105: “10-25 g (40-100 mL of 25% solution or 20-50 mL of 50% solution)”

250 mg/mL, infusion mix, not a vial: reference only. p.105: “D25= 25ml NS + 25ml D50 (12.5g in 50ml's solution)”

100 mg/mL, infusion mix, not a vial: reference only. p.105: “Newborns: 5ml/kg D10 (Max 25 G/dose)”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| SMOG p.105: 500 mg/mL, not in the kit file (declare: ____ mg in ____ mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| SMOG p.105: 250 mg/mL, not in the kit file (declare: ____ mg in ____ mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# propofol

**Signed doses in the bank** (2): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| induction of anaesthesia, hypotensive trauma patient (reduced dose) | adult | IV | 0.5–1 mg/kg |
| sedation infusion | adult | IV | 10–75 mcg/kg/min |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

10 mg/mL, infusion mix, not a vial: reference only. p.154: “MIX 500 mg/50 mL ... CONCENTRATION 10 mg/mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# levetiracetam

**Signed doses in the bank** (3): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| severe TBI — seizure prophylaxis, loading dose | adult | IV | 1500 mg |
| status epilepticus, refractory | peds | IV | 60 mg/kg |
| status epilepticus | adult | IV | 2000 mg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# calcium chloride

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| hypocalcaemia after massive transfusion | adult | IV | 1 g |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

100 mg/mL, vial strength. p.101: “solution concentration of 100 mg/mL (10%) containing 1.4 mEq (27 mg) ... (1gram = 10cc of a 10% solution)”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| SMOG p.101: 100 mg/mL, not in the kit file (declare: ____ mg in ____ mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# calcium gluconate

**Signed doses in the bank** (3): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| hyperkalaemia — membrane stabilisation | adult | IV | 1 g |
| hypocalcaemia after massive transfusion | adult | IV | 3 g |
| hyperkalaemia — membrane stabilisation | peds | IV | 100 mg/kg |

**The kit file** (drug_concentrations.json)

- 1000 mg / 10 mL ampoule = 100 mg/mL: declared, unsigned

The kit does not ask which vial.

**SMOG CY24**

100 mg/mL, vial strength. p.102: “Hypocalcemia prophylaxis from massive transfusion 30mL of 10% solution over 5 minutes”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 1000 mg / 10 mL ampoule (100 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# naloxone

**Signed doses in the bank** (4): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| opioid-induced respiratory depression | adult | IV | 0.4–2 mg |
| opioid-induced respiratory depression | adult | IM | 0.4–2 mg |
| opioid-induced respiratory depression | peds | IV | 0.1 mg/kg |
| opioid-induced respiratory depression | adult | IN | 0.4–2 mg |

**The kit file** (drug_concentrations.json)

- 0.4 mg / 1 mL ampoule = 0.4 mg/mL: signed AI-AIM 2026-08-25

The kit does not ask which vial.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 0.4 mg / 1 mL ampoule (0.4 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# phytomenadione

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| warfarin reversal | adult | IV | 10 mg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# lorazepam

**Signed doses in the bank** (2): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| active seizure | adult\|peds | IV | 0.1 mg/kg |
| agitated or violent patient / behavioural emergency | adult | IV | 2 mg |

**The kit file** (drug_concentrations.json)

- 4 mg / 2 mL ampoule = 2 mg/mL: signed AI-AIM 2026-08-25

The kit does not ask which vial.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Kit: 4 mg / 2 mL ampoule (2 mg/mL) | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# artesunate

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| severe malaria | adult | IV | 2.4 mg/kg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# atropine

**Signed doses in the bank** (6): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| symptomatic bradycardia | adult | IV | 1 mg |
| symptomatic bradycardia | peds | IV | 0.02 mg/kg |
| nerve agent or organophosphate poisoning — autoinjector | adult | IM | 2 mg |
| nerve agent poisoning — escalating IV/IO bolus | adult | IV | 2 mg |
| organophosphate or carbamate poisoning | peds | IV | 0.05 mg/kg |
| RSI pre-dose — paediatric intubation | peds | IV | 0.02 mg/kg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

80 mcg/mL, infusion mix, not a vial: reference only. p.100: “IV/IO: (Nerve agent) Atropine 20mg in 250mL”

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# sodium chloride 3%

**Signed doses in the bank** (1): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| severe TBI — signs of herniation, bolus | adult | IV | 7.5 g |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)

# acetaminophen

**Signed doses in the bank** (4): a signed strength turns these into millilitres.

| Indication | Population | Route | Dose |
|---|---|---|---|
| mild to moderate pain / analgesia | adult | PO | 500–1000 mg |
| mild to moderate pain / analgesia | adult | IV | 500–1000 mg |
| mild to moderate pain / analgesia | peds | PO | 10–15 mg/kg |
| mild to moderate pain / analgesia | peds | IV | 15 mg/kg |

**The kit file** (drug_concentrations.json)

No presentation declared.

**SMOG CY24**

SMOG states doses for this drug in mg and no strength, or has no monograph for it.

**Sign**: one line per strength. Sign only what you have read off the vial.

| | | |
|---|---|---|
| Another strength on the vial: _____ mg in _____ mL | ☐ SIGN as stocked ☐ not stocked | Signed _____ Date _____ |

**Stocked** (tick one)

☐ **One strength stocked**: _____. Doses serve mg and mL from it with no vial question; a different strength the medic states still withholds the volume.

☐ **Two strengths stocked**: _____ and _____. The medic is asked which vial every time before any mL.

☐ **Not stocked**: doses serve mg only.

Declared by _____ Date _____ (entered with tools/set_concentration.py; nothing on this page is signed until then)