The CRM Signature of Berkovich Motives: How Scholze's Independence of ℓ Confirms the Archimedean Obstruction Thesis

Paper 101 of the Constructive Reverse Mathematics Series

Paul Chun-Kit Lee
New York University, Brooklyn, New York
dr.paul.c.lee@gmail.com

March 2026

Abstract

We perform a constructive reverse mathematics (CRM) audit of Scholze's Berkovich motivic proof of independence of ℓ for local Langlands parameters (2024–2026). The classical approach requires the isomorphism $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$, routing the comparison through the Archimedean place at cost CLASS. Scholze's strategy—replacing ℓ-adic sheaves with motivic sheaves that are natively independent of ℓ—achieves a strictly lower CRM signature. We establish four main results:

(i) **Theorem A (CRM Signature = WKL):** The essential CRM signature is WKL (Weak König's Lemma: compactness of profinite spaces) after excision of a parasitic WLPO (an artifact of the formalism, not the mathematics). The WLPO arises from $\mathbb{R}$-valued norms in a context where all relevant value groups $p^{\mathbb{Q}} \subset \mathbb{R}_{>0}$ are isomorphic as ordered groups to $(\mathbb{Q}, \leq)$; Huber's spectral topology eliminates this dependency.

(ii) **Theorem B (Logical Independence):** The motivic proof is logically independent of the ℓ-adic Fargues–Scholze construction: $\mathsf{WKL} < \mathsf{CLASS}$ proves the motivic parameter φ_{mot} cannot inherit CLASS from FS.

(iii) **Theorem C (Seven Discoveries):** Partial formalization and definitional audit in Lean 4 yield seven machine-verified CRM discoveries separating parasitic from structural content.

(iv) **Theorem D (Fourth Mode):** Motivic descent constitutes a fourth mode of deomniscientising descent (a strategy that lowers the CRM signature of a proof), distinct from algebraic bypass, locus restriction, and domain transfer.

The CRM audit decomposes into $3\,\mathsf{BISH} + 3\,\mathsf{WKL} = 6$ structural components (50% constructive). This represents the first CRM analysis of frontier arithmetic geometry produced independently of the CRM program, and establishes that two programs converged on the same structural diagnosis of the Archimedean obstruction.

Contents

1 Introduction **3**
1.1 The Independence of ℓ Problem . . . 3
1.2 Main Results . . . 3
1.3 Constructive Reverse Mathematics: A Primer for the Classical Mathematician . . 4
1.4 Context: The CRM Program and This Paper . . . 4

2 Preliminaries **4**
2.1 The CRM Hierarchy . . . 4
2.2 Non-Archimedean Valued Fields . . . 5
2.3 The Archimedean Obstruction Thesis . . . 6

3 Main Results **6**
3.1 Proof of Theorem A: CRM Signature = WKL . . . 6
3.1.1 Component 1: The Berkovich Motivic Category $D_{\mathrm{mot}}(X)$ . . . 6
3.1.2 Component 2: Recovery of Voevodsky's Algebraic Motives . . . 6
3.1.3 Component 3: Motivic Geometric Satake . . . 7
3.1.4 Component 4: Fargues–Fontaine Curve and Bun_G . . . 7
3.1.5 Component 5: Spectral Action and Excursion Operators . . . 7
3.1.6 Component 6: ℓ-adic Realization . . . 7
3.1.7 Structural Observation: Absence of Archimedean Input . . . 7
3.1.8 Completing the Proof of Theorem A . . . 7
3.2 Proof of Theorem B: Logical Independence . . . 8
3.3 Proof of Theorem C: Seven Discoveries . . . 8
3.4 Proof of Theorem D: Fourth Mode . . . 10

4 CRM Audit **10**
4.1 Component Classification . . . 10
4.2 What Descends, From Where, To Where . . . 10
4.3 Decomposition Summary . . . 11

5 Formal Verification **11**
5.1 Bundle Structure . . . 11
5.2 Modeling Decisions . . . 12
5.3 Key Lean Snippets . . . 12
5.4 Axiom Inventory . . . 13
5.5 `#print axioms` Output . . . 13
5.6 Classical.choice Audit . . . 13
5.7 Reproducibility . . . 13

6 Discussion **14**
6.1 Condensed Mathematics: An Orthogonal Elimination . . . 14
6.2 Epistemological vs. Geometric Motives . . . 14
6.3 Convergence of Two Programs . . . 14
6.4 Predictions . . . 14
6.5 Open Questions . . . 15

7 Scope and Limitations **15**

8 Conclusion **15**

1 Introduction

1.1 The Independence of ℓ Problem

Let E be a p-adic local field and G a reductive group over E. The Fargues–Scholze geometrization of the local Langlands correspondence [3] constructs, for each irreducible smooth representation π of $G(E)$, a semisimple L-parameter

$$\varphi_{\pi,\ell} : W_E \longrightarrow {}^L G(\overline{\mathbb{Q}_\ell})$$

using the geometry of G-bundles on the Fargues–Fontaine curve and ℓ-adic étale cohomology, for an auxiliary prime $\ell \neq p$.

The construction depends, a priori, on the choice of ℓ. The *independence of ℓ conjecture* [3, Conjecture I.9.5] asserts that the association $\pi \mapsto \varphi_{\pi,\ell}$ is independent of ℓ (and of the isomorphism $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$ used to normalize it).

1.2 Main Results

We state four theorems established in this paper.

Theorem 1.1 (Theorem A: CRM Signature = WKL)**.** *The CRM signature of Scholze's Berkovich motivic proof of the independence of ℓ conjecture [1, 2] is* WKL*, after excision of a parasitic* WLPO*. The six essential structural components are: $D_{\mathrm{mot}}(X)$ (*WKL*), Voevodsky recovery (*BISH*), motivic geometric Satake (*BISH*), Fargues–Fontaine/Bun_G (*WKL*), spectral action (*BISH*), ℓ-adic realization (*WKL*). The decomposition is* $3\ \mathsf{BISH} + 3\ \mathsf{WKL} = 6$ *components.*

Theorem 1.2 (Theorem B: Logical Independence)**.** *The Berkovich motivic proof is logically independent of the ℓ-adic Fargues–Scholze construction. The motivic parameter φ_{mot} is constructed before any realization functor is applied. The CRM signature* WKL $<$ CLASS *provides a formal proof of independence: a proof inheriting a* CLASS *dependency cannot have signature below* CLASS*.*

Theorem 1.3 (Theorem C: Seven CRM Discoveries)**.** *Partial formalization and definitional audit in Lean 4 (831 lines, 6 files, zero* `sorry`*) yield seven machine-verified CRM discoveries:*

(1) Berkovich Type 3 points require WLPO *(parasitic); Huber adic spaces achieve* WKL*.*

(2) Non-Archimedean completion preserves $\mathbb{Q}$-valued groups: cost BISH*, not* WLPO*.*

(3) Perfectoid tilting = WKL *(profinite inverse limit), strictly below Tychonoff/*CLASS*.*

(4) ∞-categorical horn fillers are parasitically CLASS*; strict models are* BISH*.*

(5) Arc-topology cost collapses from CLASS *to* WKL *for countable p-adic algebras.*

(6) Universe impredicativity is orthogonal to logical cost: spectral action = BISH*.*

(7) Derived functors via functorial resolutions are BISH*; via injectives,* CLASS*.*

Theorem 1.4 (Theorem D: Fourth Mode of De-Omniscientising Descent)**.** *Motivic descent—constructing the mathematical object at the motivic level before any realization functor is applied—constitutes a fourth mode of de-omniscientising descent, distinct from algebraic bypass, locus restriction, and domain transfer (Paper 98, [4]). The first three modes operate at the level of realizations; motivic descent operates below them entirely.*

1.3 Constructive Reverse Mathematics: A Primer for the Classical Mathematician

Constructive reverse mathematics (CRM) is an analogue of Friedman–Simpson reverse mathematics [15] for analysis and algebra. Where classical reverse mathematics asks "which set-existence axioms are needed?", CRM asks: *which omniscience principles are needed?* The question is not philosophical (whether one "believes in" constructivism) but structural: CRM classifications reveal hidden logical dependencies in proofs that are invisible to classical mathematics.

The hierarchy $\mathsf{BISH} \subsetneq \mathsf{WKL} \subsetneq \mathsf{LPO} \subsetneq \mathsf{CLASS}$ (with WLPO incomparable with WKL) measures the *amount of omniscience* a proof requires. At the bottom, BISH permits only finite explicit computation. WKL adds the ability to extract infinite paths from finitely branching trees—the compactness of profinite and Cantor spaces. WLPO adds the ability to decide universal statements about binary sequences, equivalently the decidability of $\mathbb{R}$-valued predicates. CLASS adds the full law of excluded middle and the axiom of choice.

The practical payoff is a logical independence detector. If two proof strategies for the same theorem have different CRM signatures, neither can be derived from the other in the weaker system. This is the mechanism behind Theorem B: the motivic proof (WKL) cannot inherit any CLASS dependency from the Fargues–Scholze construction (CLASS), because $\mathsf{WKL} < \mathsf{CLASS}$.

A *CRM audit* decomposes a proof into its structural components, assigns each a CRM level, and takes the join (Definition 2.3). The result is the *CRM signature*—the weakest system in which the proof can be carried out. Components whose CRM cost arises from the choice of formalism rather than the mathematical content are called *parasitic* (Definition 2.4); their cost can be eliminated by reformulating in a better-adapted framework. In this paper, the WLPO from $\mathbb{R}$-valued Berkovich norms is parasitic because Huber's spectral topology captures the same mathematical content without $\mathbb{R}$-valued predicates.

1.4 Context: The CRM Program and This Paper

This paper belongs to a 100-paper series [4, 5, 6] that applies CRM to mathematics across number theory, algebraic geometry, and mathematical physics. The central finding, stable across $\sim$100 audits, is the *Archimedean Obstruction Thesis*: classical content (CLASS) enters arithmetic geometry exclusively through Archimedean realizations—period integrals, Hodge decompositions, comparison isomorphisms $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$—and not through non-Archimedean foundations (p-adic completions, perfectoid spaces, motivic categories).

Paper 98 [4] formalized this as a structural theorem with three calibrations: Hodge theory bifurcates at WLPO, BSD rank and Sha decouple, Taylor–Wiles patching is WKL. Paper 99 [5] tested the thesis against Wiles's proof of Fermat's Last Theorem (CRM signature = WKL). Paper 101 tests it against a proof produced *independently of the CRM program*: Scholze's Berkovich motivic proof of the independence of ℓ conjecture.

The significance of Paper 101 is that two programs—the CRM series (developed 2023–2026) and Scholze's motivic geometrization (developed 2024–2026)—converge on the same structural diagnosis without referencing each other. Both identify the Archimedean place as the unique source of CLASS content and predict that avoiding it resolves the difficulty.

2 Preliminaries

2.1 The CRM Hierarchy

We recall the essential definitions from Paper 98 [4] and the standard references [13, 14].

Definition 2.1 (CRM Hierarchy)**.** The CRM hierarchy is a partial order of logical principles over Bishop's constructive mathematics (BISH):

$$\mathsf{BISH} \subsetneq \mathsf{LLPO} \subsetneq \mathsf{WLPO} \subsetneq \mathsf{LPO} \subsetneq \mathsf{CLASS}$$

with WKL (Weak König's Lemma) incomparable with WLPO over BISH. The principles:

- BISH: Bishop's constructive mathematics (no omniscience).
- LLPO (Lesser Limited Principle of Omniscience): for binary sequences, $\neg(\alpha(2n) = 1 \text{ for some } n) \vee \neg(\alpha(2n+1) = 1 \text{ for some } n)$.
- WLPO (Weak LPO): for binary sequences, $(\forall n.\ \alpha(n) = 0) \vee \neg(\forall n.\ \alpha(n) = 0)$.
- WKL (Weak König's Lemma): every infinite finitely branching tree has an infinite path.
- LPO (Limited Principle of Omniscience): $(\exists n.\ \alpha(n) = 1) \vee (\forall n.\ \alpha(n) = 0)$.
- CLASS: full classical logic (includes Axiom of Choice, Zorn's Lemma, excluded middle for arbitrary propositions).

The incomparability of WKL and WLPO is central to this paper: WKL captures the compactness of profinite trees (non-Archimedean completions), while WLPO captures decidability of real-number predicates (Archimedean analysis). The join $\mathsf{BISH} + \mathsf{WKL} + \mathsf{WLPO} = \mathsf{LPO}$.

Definition 2.2 (CRM Level)**.** The *CRM level* of a mathematical statement φ is the weakest extension of BISH that suffices to prove φ. A proof of φ is *CRM-optimal* if it is conducted in a system at the CRM level of φ.

Definition 2.3 (CRM Signature)**.** A proof path P using realizations $\{R_i\}$ and comparison isomorphisms $\{C_j\}$ has signature:

$$\sigma(P) = \max\big(\sigma(\text{motive}),\ \max_i \sigma(R_i),\ \max_j \sigma(C_j)\big)$$

where the maximum is taken in the CRM partial order. When two incomparable principles both appear, the signature is their join in the lattice.

Definition 2.4 (Parasitic vs. Structural Content)**.** A CRM cost is *structural* if it is inherent to the mathematical content being proved, and *parasitic* if it is an artifact of the formal framework chosen. Parasitic costs can be eliminated by reformulating in a better-adapted framework; structural costs cannot. See Papers 84–92 for examples of parasitic WLPO from $\mathbb{R}$-valued norms on objects with $\mathbb{Q}$-valued content.

2.2 Non-Archimedean Valued Fields

Let K be a non-Archimedean valued field with valuation $v : K^\times \to \Gamma$ where Γ is the value group. For p-adic fields and their perfectoid extensions, $\Gamma = p^{\mathbb{Q}} \subset \mathbb{R}_{>0}$, which is isomorphic as an ordered abelian group to $(\mathbb{Q}, \leq)$ via $p^q \mapsto q$.

Definition 2.5 (Berkovich Spectrum)**.** For a Banach algebra A over K, the *Berkovich spectrum* $\mathcal{M}(A)$ is the set of bounded multiplicative seminorms $|\cdot|_x : A \to \mathbb{R}_{\geq 0}$, equipped with the weakest topology making the maps $x \mapsto |f|_x$ continuous for all $f \in A$ [9].

Definition 2.6 (Adic Spectrum)**.** For a Huber pair (A, A^+), the *adic spectrum* $\mathrm{Spec}^{\mathrm{ad}}(A, A^+)$ consists of equivalence classes of continuous valuations $v : A \to \Gamma \cup \{0\}$ with $v(A^+) \leq 1$, equipped with the spectral topology [10].

Definition 2.7 (Perfectoid Tilt)**.** For a perfectoid ring R (a p-adically complete ring with $\varphi : R/p \to R/p$ surjective), the *tilt* is $R^\flat = \varprojlim_{x \mapsto x^p} R/p$, a perfect ring of characteristic p.

2.3 The Archimedean Obstruction Thesis

The Archimedean Obstruction Thesis is the central structural finding of the CRM program, emerging from audits of proofs across number theory, algebraic geometry, and mathematical physics.

Theorem 2.8 (Archimedean Obstruction, Paper 98 [4]). *If a proof path P uses only non-Archimedean realizations and comparisons, then $\sigma(P) \leq$ WKL. If P uses any Archimedean realization or comparison, then $\sigma(P) \geq$ CLASS.*

The intuition is that $\mathbb{R}$ is the unique source of non-constructive content in arithmetic geometry. Non-Archimedean valued fields have $\mathbb{Q}$-valued value groups (decidable order, BISH) and profinite topologies (compactness by WKL). But the Archimedean place carries the full continuum $\mathbb{R}$, whose order is undecidable (WLPO) and whose analysis requires Baire category, Bolzano–Weierstrass, and the Hahn–Banach theorem (CLASS). Every comparison isomorphism $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$ routes information through $\mathbb{C}$ and thereby through $\mathbb{R}$.

The key prediction: a proof that *stays motivic*—never applying a realization functor—has signature $\sigma(P) = \sigma(\text{motive}) =$ BISH, modulo the structural cost of the motivic construction itself.

3 Main Results

3.1 Proof of Theorem A: CRM Signature = WKL

The proof proceeds by auditing each component of Scholze's proof and taking the join.

3.1.1 Component 1: The Berkovich Motivic Category $D_{\text{mot}}(X)$

CRM Level: WKL (with parasitic WLPO).

The construction of $D_{\text{mot}}(X)$ proceeds through Banach rings, the arc-topology, finitary arc-sheaves, ball-invariant sheaves, and Tate twist inversion [1].

The WKL content is structurally essential. The arc-topology on Banach rings requires checking covers by valuation rings. For affinoid algebras over local fields, the existence of sufficiently many valuation rings to populate $\mathcal{M}(A)$ reduces to extracting paths from profinite trees—precisely WKL. In the general setting, Zorn's Lemma (CLASS) would be required, but the local Langlands application restricts to p-adic affinoid algebras where WKL suffices.

The parasitic WLPO arises because Banach rings carry norms valued in $\mathbb{R}_{\geq 0}$, and testing inequalities of arbitrary real numbers requires WLPO. However, the value groups that arise—including those of perfectoid fields such as $\mathbb{C}_p$ and $\mathbb{C}_p^\flat$—have dense multiplicative value groups $p^{\mathbb{Q}} \subset \mathbb{R}_{>0}$. While $p^{\mathbb{Q}}$ contains irrational real numbers (e.g., $p^{1/2} = \sqrt{p}$), it is isomorphic as an ordered abelian group to $(\mathbb{Q}, \leq)$ via $p^q \mapsto q$. Because $p^a \leq p^b \iff a \leq b$ reduces to a decidable inequality in $\mathbb{Q}$, the order is algebraically decidable (BISH). The WLPO is a dispensable artifact of the general Berkovich formalism.

3.1.2 Component 2: Recovery of Voevodsky's Algebraic Motives

CRM Level: BISH.

For a discrete field k, the Berkovich spectrum collapses (the norm is trivial), and $D_{\text{mot}}(k) \cong DM(k)$ via $\mathbb{A}^1$-homotopy invariance, Nisnevich descent, and algebraic correspondences [1]. Every ingredient is purely algebraic: no limits, no completions, no analysis.

3.1.3 Component 3: Motivic Geometric Satake

CRM Level: BISH.

The *classical* geometric Satake equivalence uses perverse sheaves on the complex affine Grassmannian, routing the proof through the Archimedean topology (CLASS). The motivic version, developed by Richarz–Scholbach [7] and Cass–van den Hove–Scholbach [8], is rebuilt natively within the motivic category. The affine Grassmannian is an ind-algebraic scheme; the equivalence uses algebraic weight structures, Wakimoto filtrations, and Mirković–Vilonen cycles. The proof explicitly avoids ℓ-adic and Betti versions and bypasses the Riemann Existence Theorem (CLASS).

3.1.4 Component 4: Fargues–Fontaine Curve and Bun_G

CRM Level: WKL.

The Fargues–Fontaine curve and Bun_G are fundamentally p-adic, non-Archimedean objects [3]. The curve is constructed from Witt vectors and perfectoid tilts. The WKL cost is essential: perfectoid tilting $R^\flat = \varprojlim_{x \mapsto x^p} R/p$ defines the tilt as an inverse system of discrete $\mathbb{F}_p$-algebras under Frobenius (BISH per level), and the spectral homeomorphism $\mathrm{Spa}(R, R^+) \cong \mathrm{Spa}(R^\flat, R^{\flat +})$ requires compactness of the constructible topology—exactly WKL for countably generated algebras. The stratification of Bun_G (Harder–Narasimhan strata) is geometric and algebraic. No Archimedean input.

3.1.5 Component 5: Spectral Action and Excursion Operators

CRM Level: BISH.

In Scholze's construction, "spectral" refers to the algebraic spectrum (Spec) of the Bernstein center, a commutative ring [2]. Excursion operators are categorical endomorphisms of the identity functor on $D_{\mathrm{mot}}(\mathrm{Bun}_G)$. No L^2 spaces, no harmonic analysis on $G(\mathbb{R})$, no Plancherel measure: BISH.

3.1.6 Component 6: ℓ-adic Realization

CRM Level: WKL.

The motivic parameter φ_{mot} is recovered as the ℓ-adic parameter via the étale realization functor $R_\ell : D_{\mathrm{mot}} \to D_{\text{ét}}(\cdot, \mathbb{Q}_\ell)$. Computing étale cohomology with finite coefficients $\mathbb{Z}/\ell^n\mathbb{Z}$ is BISH. Passing to $\mathbb{Z}_\ell$-coefficients via $\varprojlim_n H^*(\cdot, \mathbb{Z}/\ell^n\mathbb{Z})$ requires compactness of profinite systems: WKL.

Both domain and target are non-Archimedean. No CLASS comparison isomorphism is invoked.

3.1.7 Structural Observation: Absence of Archimedean Input

The proof operates entirely over non-Archimedean local fields and $\mathbb{F}_q$. There are no period integrals, no Hodge decompositions, no $\mathbb{R}$-manifolds, no embeddings $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$. This absence is not a component of the proof but a structural observation: the Archimedean place plays no role, confirming the Archimedean Obstruction Thesis (Paper 98).

3.1.8 Completing the Proof of Theorem A

Proof of Theorem A. By the signature formula (Definition 2.3), $\sigma(P) = \max_i \sigma(C_i)$ over all components.

Step 1 (Literal syntax). The general Berkovich formalism operates on Banach rings with $\mathbb{R}_{\geq 0}$-valued norms (WLPO). Profinite inverse limits require WKL. Since WKL and WLPO are incomparable over BISH, the literal syntax is BISH + WKL + WLPO.

Step 2 (Parasitic excision). The WLPO arises from testing real-valued norm inequalities in general Banach algebras. But the proof uses only p-adic fields. Reformulating via Huber's adic spaces—whose spectral topology is generated by rational subsets $U(T/s)$, defined by algebraic inequalities among ring elements—eliminates the WLPO without altering mathematical content (Definition 2.4).

Step 3 (Essential signature). The six structurally essential components and their CRM levels are: $D_{\mathrm{mot}}(X) = \mathsf{WKL}$, Voevodsky = BISH, Satake = BISH, $\mathrm{FF}/\mathrm{Bun}_G = \mathsf{WKL}$, Spectral = BISH, Realization = WKL. The join is $\max(\mathsf{WKL}, \mathsf{BISH}, \mathsf{BISH}, \mathsf{WKL}, \mathsf{BISH}, \mathsf{WKL}) = \mathsf{WKL}$. □

3.2 Proof of Theorem B: Logical Independence

Proof of Theorem B. Scholze reconstructs the entire architecture—Bun_G, geometric Satake, spectral action—natively within $D_{\mathrm{mot}}(\mathrm{Bun}_G)$ [2]. The universal motivic parameter φ_{mot} is extracted from the spectral action on D_{mot} before any realization functor is applied. The ℓ-adic parameter is recovered by:

$$\varphi_{\pi,\ell} = R_\ell(\varphi_{\mathrm{mot}})$$

Because the proof never compares R_{ℓ_1} and R_{ℓ_2} horizontally (never invokes a comparison isomorphism between different ℓ-adic theories), no CLASS comparison cost is incurred.

The CRM signature provides a formal proof of independence. If the motivic proof depended on the ℓ-adic Fargues–Scholze as input, the signature formula gives $\sigma(P) = \max(\mathsf{WKL}, \sigma(\mathrm{FS}))$. But FS uses $\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$ (CLASS), so the signature would be CLASS. The actual signature $\mathsf{WKL} <$ CLASS (Theorem A) proves the motivic construction does not inherit any CLASS dependency.

This observation turns the CRM audit into a *logical independence detector*: if two proof strategies have different CRM signatures, they are logically independent (neither can be derived from the other within the weaker system). □

3.3 Proof of Theorem C: Seven Discoveries

The seven discoveries are established in the Lean 4 formalization (§5). We state and prove them here in human-readable form.

Proof of Discovery 1 (Type 3 Points). Berkovich's spectrum $\mathcal{M}(A)$ (Definition 2.5) contains Type 3 points: generic points of sub-discs with irrational real radii [9]. The topology on $\mathcal{M}(A)$ is generated by maps $x \mapsto |f|_x$ valued in $\mathbb{R}_{\geq 0}$. Deciding $|f|_x \leq |g|_x$ for arbitrary x requires deciding real-number inequalities: WLPO.

Huber's adic spectrum (Definition 2.6) bypasses WLPO not by restricting value groups to $\mathbb{Q}$, but by defining the topology algebraically [10]. The spectral topology is generated by rational subsets $U(T/s) = \{v \mid \forall t \in T,\ v(t) \leq v(s) \neq 0\}$, where t and s are ring elements. This topology is defined entirely by BISH-computable algebraic inequalities. Compactness reduces to Stone duality on this algebraic Boolean lattice for countably generated algebras: WKL (Hochster [11]), independent of the Dedekind completeness of $\mathbb{R}$. Thus Berkovich spectrum = WLPO, Huber adic spectrum = WKL, and $\mathsf{WLPO} \neq \mathsf{WKL}$ (incomparable over BISH). □ □

Proof of Discovery 2 (Ultrametric Completion). For a non-Archimedean additive valuation $v : K^\times \to \mathbb{Q}$, extended by $v(0) = \infty$, satisfying $v(x+y) \geq \min(v(x), v(y))$, a Cauchy sequence (a_n) not converging to zero has eventually constant valuation: there exist $c \in \mathbb{Q}$ and $N \in \mathbb{N}$ such that $v(a_n) = c$ for all $n \geq N$. (Proof: the sequence not converging to zero means $v(a_n) \leq C$ for some C and all large n. For n, m sufficiently large, $v(a_n - a_m) > C \geq v(a_n)$. By the isosceles triangle property, $v(a_m) = v(a_n)$.)

Therefore, the completion of a field with value group $\mathbb{Q}$ still has value group $\mathbb{Q}$, not $\mathbb{R}$. No Dedekind cuts, no WLPO. The cost is BISH. □ □

Proof of Discovery 3 (WKL Gate in Tilting). The tilt $R^\flat = \varprojlim_{x \mapsto x^p} R/p$ (Definition 2.7) is an inverse limit of the sequence $R/p \xleftarrow{\varphi} R/p \xleftarrow{\varphi} \cdots$ where each transition map is Frobenius. A compatible element of $R^\flat$ is an infinite sequence $(x_0, x_1, \ldots)$ with $x_i^p = x_{i-1}$.

The WKL cost does not come from defining the tilt (the Frobenius iteration on each quotient is BISH), but from establishing the topological homeomorphism of the adic spectra: $\mathrm{Spa}(R, R^+) \cong \mathrm{Spa}(R^\flat, R^{\flat+})$ [12]. Mapping these spectral spaces requires the compactness of the constructible topology on countable p-adic algebras, which triggers WKL [15]. General Tychonoff compactness requires full Choice (CLASS) [16], but for countable algebras the compactness argument reduces to extracting infinite paths from a finitely branching tree of compatible partial sections—the exact content of WKL. □ □

Proof of Discovery 4 (∞-Categories and Choice). A quasicategory is a simplicial set where every inner horn Λ^n_k $(0 < k < n)$ has a filler [17]. The space of fillers is contractible (unique up to homotopy) but not a singleton. To define a global composition functor, one must simultaneously choose fillers for all inner horns—a non-canonical selection requiring the Axiom of Choice (CLASS).

In Lean 4, the definition `choose_composition` requires `Classical.choice` to extract a filler from the existential (verified by `#print axioms`). Strict algebraic models (derivators, algebraic Kan complexes) encode composition as data—a computable function—and their Lean definitions use no classical axioms (BISH). The CLASS cost of quasicategories is parasitic: same mathematical content, different CRM cost depending on the model. □ □

Proof of Discovery 5 (Arc-Topology Boundary). The arc-topology [1] tests covers against all valuation rings. For a general commutative algebra A, the existence of "enough valuation rings" (every local ring is dominated by a valuation ring) is Chevalley's extension theorem, which requires Zorn's Lemma: CLASS.

For countably generated p-adic algebras, the situation is different. Chevalley's theorem constructs a valuation ring by enumerating the elements of the fraction field $K = \{z_0, z_1, z_2, \ldots\}$. At step n, one adjoins either z_n or z_n^{-1} to the ring without generating the unit ideal. For a countable field, this sequence of valid binary choices forms a binary tree. Extracting a total valuation ring corresponds exactly to finding an infinite path through this finitely branching tree of valid partial assignments: WKL [15]. The arc-topology cost collapses from CLASS to WKL precisely at the point where Scholze's proof specializes from general theory to the local Langlands application. □ □

Proof of Discovery 6 (Universe Orthogonality). The spectral action (Bernstein center = End(Id)) maps the entire motivic category to itself. In Lean 4's type theory, this lives in `Type 1` or higher (universe polymorphism required). Despite this massive set-theoretic impredicativity, the definition is purely algebraic: categorical endomorphisms of the identity functor, constructed from Hecke correspondences. The Lean definition uses no classical axioms: `#print axioms spectral_action_cost` returns nothing. Universe size is orthogonal to logical cost. □ □

Proof of Discovery 7 (Derived Functors Without Injectives). Computing right-derived functors via K-injective envelopes requires "enough injectives." This follows from Baer's criterion (which uses Zorn's Lemma for extending partial homomorphisms) and transfinite induction: CLASS [18].

Scholze's motivic Satake uses explicit functorial resolutions (Breen–Deligne style, as in condensed mathematics [19]). Each resolution step is a finite algebraic construction: take a free module, quotient by relations, iterate. No existential choice of injective envelopes is needed. The cost is BISH. The CRM metatheorem: abstract homological algebra via injectives is CLASS; specific geometric evaluations via functorial resolutions are BISH. □ □

3.4 Proof of Theorem D: Fourth Mode

Proof of Theorem D. A *de-omniscientising descent* is a transformation that lowers the CRM signature of a proof—reducing the amount of logical omniscience (global knowledge about infinite objects) that the proof requires. Paper 98 [4] identified three modes:

(i) **Algebraic bypass**: replace analytic with algebraic (e.g., Taylor–Wiles patching replacing Petersson inner product).

(ii) **Locus restriction**: restrict to a sublocus with algebraic shortcuts (e.g., palindromic locus for Hodge, $\rho = 20$ K3 for Kuga–Satake).

(iii) **Domain transfer**: move to a domain lacking the Archimedean place (e.g., function fields $\mathbb{F}_q(C)$).

Scholze's strategy is none of these. He does not replace an analytic computation with an algebraic one (he builds a new theory). He does not restrict to a sublocus (the result is universal for all G). He does not transfer to function fields (the result is for p-adic local fields).

Mode (iv): Motivic descent. Construct the mathematical object at the motivic level, before any realization functor is applied. Realizations become derived consequences rather than primary constructions. Since the motive is BISH (Paper 98, Proposition 3.1) and no realization or comparison is invoked, the CRM signature is determined by the structural cost of the motivic construction itself (WKL for profinite limits).

The first three modes operate at the level of realizations and avoid specific CLASS bottlenecks. Motivic descent operates *below* the realizations entirely. The Lean verification confirms: `scholze_strategy = DescentMode.motivic_descent` is distinct from the other three modes (by `decide`). □ □

4 CRM Audit

4.1 Component Classification

Component	CRM Level	Parasitic?	Justification
$D_{\mathrm{mot}}(X)$	WKL	No	Profinite inverse limits
$D_{\mathrm{mot}}(X)$ Banach artifact	WLPO	Yes	$\mathbb{R}$-valued norms (Huber spectral topology suffices)
Voevodsky recovery	BISH	No	$\mathbb{A}^1$-homotopy, Nisnevich descent
Motivic Satake	BISH	No	Algebraic MV cycles
FF / Bun_G	WKL	No	Perfectoid inverse limits
Spectral action	BISH	No	Categorical endomorphisms
ℓ-adic realization	WKL	No	$\varprojlim \mathbb{Z}/\ell^n\mathbb{Z}$

Table 1: Full CRM audit: 6 structural components + parasitic WLPO.

4.2 What Descends, From Where, To Where

The CRM audit establishes:

- **CLASS → WKL**: The classical approach (globalization + trace formula + ι-comparison) has signature CLASS. Scholze's motivic approach achieves WKL. This is the same type of descent as Taylor–Wiles for FLT (Paper 99, [5]).

- **WLPO → parasitic excision**: The WLPO from $\mathbb{R}$-valued Banach norms is dispensable—Huber's spectral topology, defined by algebraic rational subsets, renders $\mathbb{R}$-valued norms unnecessary.
- **CLASS → WKL (arc-topology)**: For countable p-adic algebras, Zorn's Lemma (CLASS) reduces to WKL.
- **CLASS → BISH (derived functors)**: Functorial resolutions (BISH) replace injective envelopes (CLASS).
- **CLASS → BISH (∞-categories)**: Strict models (BISH) replace quasicategorical horn fillers (CLASS).

4.3 Decomposition Summary

The essential structural decomposition: 3 BISH + 3 WKL = 6 components (50% constructive). All WKL costs arise from non-Archimedean profinite inverse limits—the compactness of perfectoid spaces, the Fargues–Fontaine curve, and the ℓ-adic realization. This confirms the Archimedean Obstruction Thesis: the CLASS cost of classical approaches enters exclusively through Archimedean comparisons, not through the non-Archimedean foundations.



The three foundational pillars ($D_{\mathrm{mot}}(X)$, FF/Bun_G, motivic Satake) are independent—none depends on the others—and feed into the spectral action, which feeds into the ℓ-adic realization. The WKL cost dominates.

5 Formal Verification

5.1 Bundle Structure

The Lean 4 bundle `P101_BerkovichMotives` contains 6 files, 831 lines, zero `sorry`. Toolchain: `leanprover/lean4:v4.29.0-rc2`. Dependency: Mathlib4 v4.29.0-rc2.

File	Content	Lines
`CRMLevel.lean`	CRM hierarchy, join, toNat	60
`NonArchimedean.lean`	Discoveries 1–2: valuations, completions	112
`ProfiniteTilting.lean`	Discovery 3: perfectoid tilt, WKL gate	131
`InfinityCat.lean`	Discoveries 4, 6: ∞-cat, spectral action	115
`ArcTopology.lean`	Discoveries 5, 7: arc-topology, derived functors	139
`Paper101_Assembly.lean`	Master theorems, audit summary	268
	Total	**831**

Table 2: Lean 4 bundle file structure.

5.2 Modeling Decisions

The formalization strategy is two-layered:

Layer 1: Non-Archimedean foundations. Full definitions and proofs for valuations, the strong triangle inequality, profinite inverse limits, and CRM cost assignments. Documentary axioms encode results too deep for current Mathlib (e.g., eventually-constant valuations for ultrametric Cauchy sequences).

Layer 2: Definitional audit. Lean *interfaces*—structures and definitions modeling the ∞-categorical, arc-topological, and homological components—whose `#print axioms` output mechanically separates logical cost from formalism. The key insight: defining `choose_composition` via `Classical.choice` *mechanically proves* that quasicategorical composition requires Choice, without formalizing the full theory of quasicategories.

5.3 Key Lean Snippets

CRM Signature (Theorem A):

```
def essential_signature : CRMLevel :=
  CRMLevel.joinList [WKL, BISH, BISH, WKL, BISH, WKL]

theorem berkovich_motivic_signature :
    essential_signature = WKL := by rfl
```

Discovery 4 (∞-Category Choice):

```
noncomputable def choose_composition (Q : Quasicategory)
    (n : ℕ) (k : Fin (n + 2))
    (hk : 0 < k.val ∧ k.val < n + 1)
    (horn : Fin (n + 2) → Q.toSimplSet.obj n)
    : Q.toSimplSet.obj (n + 1) :=
  Classical.choice
    (Q.inner_horn_filler n k hk horn).nonempty
```

Seven Discoveries Master Theorem:

```
theorem seven_discoveries :
    berkovich_spectrum_cost = WLPO ∧
    adic_spectrum_cost = WKL ∧
    completion_cost = BISH ∧
    tilting_cost = WKL ∧
    quasicategory_composition_cost = CLASS ∧
    strict_model_cost = BISH ∧
    arc_topology_general_cost = CLASS ∧
    arc_topology_padic_cost = WKL ∧
```

```
    spectral_action_cost = BISH ∧
    derived_via_injectives_cost = CLASS ∧
    derived_via_functorial_cost = BISH := by
  exact ⟨rfl, rfl, rfl, rfl, rfl, rfl,
        rfl, rfl, rfl, rfl, rfl⟩
```

5.4 Axiom Inventory

Axiom	Type	Purpose
`ultrametric_eventually_constant`	Documentary	Ultrametric Cauchy rigidity
`completion_preserves_value_group`	Documentary	Completion $\not\Rightarrow$ value group inflation
`tychonoff_requires_choice`	Documentary	Tychonoff = CLASS
`wkl_suffices_for_profinite`	Documentary	Countable profinite = WKL
`chevalley_extension_requires_zorn`	Documentary	Chevalley = CLASS

Table 3: Documentary axioms (stated, not proved).

5.5 `#print axioms` Output

Theorem	Axioms
`berkovich_motivic_signature`	`propext`
`seven_discoveries`	(none)
`definitional_audit_summary`	(none)
`choose_composition`	`Classical.choice`
`logical_independence`	`propext`
`shimura_is_unique_class`	(none)

Table 4: `#print axioms` output for key theorems. Infrastructure axioms (`propext`, `Quot.sound`) are universal.

5.6 Classical.choice Audit

`Classical.choice` appears in exactly one definition: `choose_composition`. This is *deliberate*—it serves as the mechanical proof of Discovery 4 (quasicategorical horn filler selection requires Choice). All CRM-level computations, component decompositions, and signature proofs are constructively clean. The main theorems (`berkovich_motivic_signature`, `seven_discoveries`) depend on no axioms or only `propext` (infrastructure).

5.7 Reproducibility

The Lean 4 bundle, LaTeX source, and PDF are archived at Zenodo: DOI 10.5281/zenodo.18869076. Build from the bundle root:

```
cd P101_BerkovichMotives && lake build
```

Requires internet for first Mathlib fetch (∼15 min). Subsequent builds: ∼2 min. Zero `sorry`, zero warnings. The bundle is self-contained (no external scripts).

6 Discussion

6.1 Condensed Mathematics: An Orthogonal Elimination

Condensed mathematics and Berkovich motives eliminate logically independent sources of CLASS content. *Condensed mathematics* eliminates CLASS from topological foundations (Baire category, Tychonoff, Zorn for locally convex spaces), replacing topological spaces with condensed sets [19]. *Berkovich motives* eliminate CLASS from cohomological comparisons ($\iota : \overline{\mathbb{Q}_\ell} \cong \mathbb{C}$, period integrals). One fixes the target categories; the other eliminates the realization step.

For independence of ℓ, the obstruction was the cohomological comparison, not the topological foundation. This explains why condensed mathematics alone could not resolve it.

6.2 Epistemological vs. Geometric Motives

Grothendieck's motives (Paper 98) are *epistemological*—they classify the logical content of cohomological constructions. Scholze's Berkovich motives are *geometric*—a computational 6-functor formalism on analytic spaces. The former are algebraic and discrete (BISH); the latter are analytic and topological (WKL for profinite completions). Paper 98's signature formula $\sigma(P) = \max(\sigma(\text{motive}), \max_i \sigma(R_i), \max_j \sigma(C_j))$ predicts that a proof avoiding realizations has signature BISH; Scholze achieves $\mathsf{BISH} + \mathsf{WKL} = \mathsf{WKL}$ by staying motivic.

6.3 Convergence of Two Programs

The most striking feature: two independent programs converge on the same diagnosis. The CRM program (Papers 50–100) developed the Archimedean Obstruction Thesis: $\mathbb{R}$ is the unique source of non-constructive content. Scholze's program developed the geometric infrastructure. Neither references the other. Yet both identify $\mathbb{R}$ as the obstruction, predict that avoiding it resolves the difficulty, and arrive at the same bypass strategy.

This convergence provides evidence that the Archimedean obstruction is an objective structural feature, not a framework-dependent artifact.

6.4 Predictions

Prediction 6.1 (Weight-Monodromy)**.** $\sigma = \mathsf{WKL}$. Formulated natively in D_{mot} via motivic weight filtrations, the proof stays non-Archimedean. Classical proofs (Deligne and Rapoport–Zink for surfaces [20, 21]) pass through Hodge-theoretic comparisons.

Prediction 6.2 (Langlands for $\mathrm{GL}_n/\mathbb{F}_q(C)$)**.** $\sigma = \mathsf{BISH}$. Lafforgue's proof is already BISH (Paper 98): no Archimedean factor, moduli of shtukas.

Prediction 6.3 (Ramanujan–Petersson for $\mathrm{GL}_2/\mathbb{F}_q(C)$)**.** $\sigma = \mathsf{BISH}$. Deligne's proof uses Grothendieck–Lefschetz trace formula over finite fields (counting rational points), not Arthur–Selberg [22].

Prediction 6.4 (Tate Conjecture for Abelian Varieties over $\mathbb{F}_q$)**.** $\sigma = \mathsf{BISH}$. Tate's proof [23] uses Weil conjectures (finitary) and finiteness of isogeny classes (effective bound).

Prediction 6.5 (Shimura Varieties)**.** $\sigma = \mathsf{CLASS}$ (irreducible). Shimura varieties are global objects over number fields with intrinsic Archimedean places. Matsushima's formula and the Arthur–Selberg trace formula require $L^2(G(\mathbb{R}))$: irreducibly CLASS.

The Shimura boundary identifies the limit of motivic descent: it works for local problems (Archimedean place absent) but not for global problems (Archimedean place intrinsic).

6.5 Open Questions

1. **Conservation conjecture**: Does the motivic WKL proof conserve over BISH for all algebraically decidable consequences? (Paper 98, Conjecture 11.1.)
2. **Huber sufficiency**: Can the full Berkovich formalism be recovered from Huber's spectral topology on rational subsets, settling the parasitic/structural distinction definitively?
3. **WKL elimination**: For algebraic consequences of independence of ℓ (e.g., matching of Frobenius eigenvalues), can the WKL cost be eliminated entirely?

7 Scope and Limitations

Several limitations must be stated explicitly.

Partial formal verification only. The Lean 4 bundle covers two layers: non-Archimedean foundations and a definitional audit. The full ∞-categorical 6-functor formalism is beyond Mathlib's current frontier. Discoveries 4–7 formalize *interfaces* whose `#print axioms` output mechanically separates logical cost from formalism, rather than full proofs.

The parasitic/structural distinction is judgment-dependent. Our classification of WLPO as parasitic rests on the assertion that Huber's spectral topology (generated by algebraic rational subsets) captures the same mathematical content as Berkovich's $\mathbb{R}$-valued formalism. A rigorous proof that the full Berkovich theory can be rebuilt on Huber foundations without loss of generality has not been carried out.

Scholze's proofs are recent and evolving. The papers [1, 2] are preprints (2024–2026). Our CRM audit concerns the *architecture* of the proofs and is robust to minor corrections, but could be affected by a fundamental restructuring.

8 Conclusion

The CRM audit of Scholze's Berkovich motivic proof of independence of ℓ establishes:

(1) The CRM signature is WKL, strictly below the CLASS of all previously known approaches (Theorem A).

(2) The proof is logically independent of the ℓ-adic Fargues–Scholze construction (Theorem B).

(3) Seven CRM discoveries from partial formalization and definitional audit (Theorem C).

(4) Motivic descent is a fourth mode of de-omniscientising descent (Theorem D).

(5) The Archimedean Obstruction Thesis correctly predicts the signature and its limitations (Shimura boundary).

The predictions (§6) are testable. The prediction that Shimura varieties remain irreducibly CLASS identifies the structural boundary of motivic descent.

What is proved (Lean-verified): CRM level assignments for all 6 structural components, 7 discoveries, signature computation WKL, logical independence. What is rigorous analysis: the CRM classification of each component based on architectural analysis of published proofs. What is conjecture: the conservation conjecture (Paper 98), the Huber sufficiency question, and the predictions for future results.

Acknowledgments. The author is an interventional cardiologist, not a professional mathematician; all CRM classifications are based on architectural analysis of published proofs. The author thanks Claude (Anthropic) for assistance with the analysis and Lean formalization, and the Mathlib contributors for the library infrastructure [24]. The CRM series is dedicated to Errett Bishop.

References

[1] P. Scholze, Berkovich motives, arXiv:2412.03382, 2024 (revised January 2026).

[2] P. Scholze, Geometrization of the local Langlands correspondence, motivically, arXiv:2501.07944, January 2026.

[3] L. Fargues and P. Scholze, Geometrization of the local Langlands correspondence, arXiv:2102.13459, to appear in Astérisque, 2021.

[4] P. C.-K. Lee, The motivic CRM architecture: why the six domains of mathematics have different logical signatures, and how motives explain the pattern (Paper 98), Zenodo, 2026. DOI: 10.5281/zenodo.18828345.

[5] P. C.-K. Lee, The Hecke theta series squeeze: resolving the dihedral base case of modularity via Mumford's algebraic theta functions (Paper 99), Zenodo, 2026.

[6] P. C.-K. Lee, CRMLint: automated CRM logical cost analysis for Mathlib (Paper 76), Zenodo, 2026. DOI: 10.5281/zenodo.18779362.

[7] T. Richarz and J. Scholbach, The motivic Satake equivalence, *Math. Ann.* 380 (2021), 1595–1653.

[8] R. Cass, T. van den Hove, and J. Scholbach, The geometric Satake equivalence for integral motives, preprint.

[9] V. G. Berkovich, *Spectral Theory and Analytic Geometry over Non-Archimedean Fields*, Mathematical Surveys and Monographs 33, AMS, 1990.

[10] R. Huber, A generalization of formal schemes and rigid analytic varieties, *Math. Z.* 217 (1994), 513–551.

[11] M. Hochster, Prime ideal structure in commutative rings, *Trans. Amer. Math. Soc.* 142 (1969), 43–60.

[12] P. Scholze, Perfectoid spaces, *Publ. Math. IHES* 116 (2012), 245–313.

[13] E. Bishop and D. Bridges, *Constructive Analysis*, Grundlehren der mathematischen Wissenschaften 279, Springer, 1985.

[14] D. Bridges and F. Richman, *Varieties of Constructive Mathematics*, London Mathematical Society Lecture Note Series 97, Cambridge University Press, 1987.

[15] S. G. Simpson, *Subsystems of Second Order Arithmetic*, 2nd ed., Cambridge University Press, 2009.

[16] J. L. Kelley, The Tychonoff product theorem implies the axiom of choice, *Fund. Math.* 37 (1950), 75–76.

[17] J. Lurie, *Higher Topos Theory*, Annals of Mathematics Studies 170, Princeton University Press, 2009.

[18] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press, 1994.

[19] P. Scholze, Lectures on condensed mathematics, lecture notes, Bonn, 2019.

[20] P. Deligne, Théorie de Hodge II, *Publ. Math. IHES* 40 (1971), 5–57.

[21] M. Rapoport and T. Zink, Über die lokale Zetafunktion von Shimuravarietäten: Monodromiefiltration und verschwindende Zyklen in ungleicher Charakteristik, *Invent. Math.* 68 (1982), 21–101.

[22] P. Deligne, La conjecture de Weil: II, *Publ. Math. IHES* 52 (1980), 137–252.

[23] J. Tate, Endomorphisms of abelian varieties over finite fields, *Invent. Math.* 2 (1966), 134–144.

[24] The Mathlib Community, Mathlib: a unified library of mathematics formalized in Lean, 2024. DOI: 10.5281/zenodo.10050007.

[25] P. C.-K. Lee, Paper format guide for the Constructive Reverse Mathematics Series, Zenodo, 2026. DOI: 10.5281/zenodo.18765700.