Algebraic Gauss–Manin via Griffiths Pole Reduction: A CRMLint Squeeze

(Paper 80, Constructive Reverse Mathematics Series)

Paul Chun-Kit Lee*
New York University
dr.paul.c.lee@gmail.com

February 2026

Abstract

We upgrade the CRMLint Squeeze protocol from exact arithmetic over $\mathbb{Q}$ (Papers 77–79) to symbolic algebra over the function field $\mathbb{Q}(t)$. The target is the Legendre family of elliptic curves $y^2 = x(x-1)(x-t)$, parametrized by t.

The Classical Boundary Node is the Ehresmann–Gauss–Manin theorem: the cohomology of a smooth family forms a local system with flat connection, computed via analytic continuation of period integrals over $\mathbb{C}$. This is CLASS: it requires Ehresmann's fibration theorem [23] (continuous topology), multi-valued functions, and transcendental analysis.

The constructive target: Grothendieck and Dwork proved that the differential equation governing the cohomology is fundamentally algebraic. The Picard–Fuchs operator $\mathcal{L} = t(1-t)\frac{d^2}{dt^2} + (1-2t)\frac{d}{dt} - \frac{1}{4}$ can be computed entirely via Griffiths pole reduction in algebraic de Rham cohomology over $\mathbb{Q}[t]$. We carry out this reduction explicitly: a Python CAS performs the pole reduction and emits formal polynomial identities; Lean 4 verifies them via the `ring` tactic (0 sorry, 0 axioms on the constructive path).

This is an infrastructure upgrade for the CRM program. Papers 77–79 used `native_decide` on concrete integers (0-dimensional point verification). Paper 80 uses `ring` on symbolic polynomials (1-dimensional moduli verification). The function-field bridge enables CRMLint to verify identities over parameterized families of varieties rather than isolated points.

CRM descent: CLASS $\to$ BISH.

Contents

1 Introduction **2**
1.1 The function-field gap . . . 2
1.2 The metamathematical pivot . . . 3
1.3 Main results . . . 3
1.4 Position in the atlas . . . 4

*Lean 4 formalization available at https://doi.org/10.5281/zenodo.18791985.

2 Preliminaries **4**
2.1 The CRM hierarchy . . . 4
2.2 Algebraic de Rham cohomology . . . 5
2.3 The Gauss–Manin connection . . . 5
2.4 The Griffiths pole reduction algorithm . . . 6
2.5 The Picard–Fuchs equation . . . 6

3 The Legendre Family **6**
3.1 The defining polynomial . . . 6
3.2 Branch points and discriminant . . . 6

4 Griffiths Pole Reduction **7**
4.1 The Gauss–Manin differentiation . . . 7
4.2 Step 1: Griffiths polynomial division . . . 7
4.3 Step 2: Coboundary reduction . . . 7
4.4 Row 2 . . . 8
4.5 The Gauss–Manin connection matrix . . . 8

5 The Picard–Fuchs Equation **8**
5.1 Derivation from the connection matrix . . . 8
5.2 The hypergeometric recurrence . . . 9
5.3 Closed form . . . 10

6 CRM Audit **10**
6.1 Constructive strength classification . . . 10
6.2 What descends, from where, to where . . . 10
6.3 Comparison with the Squeeze trilogy . . . 10

7 Formal Verification **11**
7.1 File structure . . . 11
7.2 Axiom inventory . . . 11
7.3 Key code snippets . . . 11
7.4 Reproducibility . . . 12

8 Discussion **12**
8.1 The de-omniscientizing descent for families . . . 12
8.2 What the function-field bridge enables . . . 12
8.3 Structural barriers to deeper applications . . . 13
8.4 Relationship to Grothendieck and Dwork . . . 13

9 Conclusion **13**

1 Introduction

1.1 The function-field gap

The CRMLint Squeeze trilogy (Papers 77–79 [14, 15, 16]) established a pipeline for extracting constructive content from classical existence theorems in algebraic geometry and number theory. Each paper targeted a fixed algebraic object over $\mathbb{Q}$:

- Paper 77: explicit Hodge decompositions for E^4 (Hodge theory).
- Paper 78: explicit local Langlands for $\mathrm{GL}_2(\mathbb{Q}_2)$ (Langlands program).
- Paper 79: Standard Conjecture D for generic Weil fourfolds (algebraic cycles).

In every case, the Squeeze reduced a CLASS existence theorem to a BISH computation on concrete integers, verified by Lean 4's compiled kernel (`native_decide` or `decide`).

This pipeline has a structural limitation: it operates on isolated, rigid objects. The Millennium Prize problems—and more generally, the deep conjectures of arithmetic geometry—are universal structural theorems about *families* of varieties, parameterized over moduli spaces. To extract a BSD rank ≥ 2 cycle, one cannot examine a single elliptic curve; higher Heegner cycles live over the function field of the modular curve $X_0(N)$. To falsify the Hodge Conjecture, one cannot test a single variety at bounded degree; one must prove that the cycle class matrix over $\mathbb{Q}(t)$ has no algebraic solution at *any* degree.

The present paper closes this gap by upgrading the CRMLint Squeeze from exact arithmetic ($\mathbb{Q}$) to function-field algebra ($\mathbb{Q}(t)$).

1.2 The metamathematical pivot

The upgrade changes the verification tactic.

Papers 77–79 used `native_decide`: Lean's compiled kernel evaluates a decidable proposition on concrete integers. This cannot handle a symbolic parameter t—there is no concrete integer to evaluate.

Paper 80 uses `ring`: Lean's polynomial normalization tactic verifies that two formal expressions in $\mathbb{Q}[x,t]$ define the same polynomial. This is the natural tactic for function-field identities. It operates syntactically on the polynomial ring, requiring no numerical evaluation, no convergence argument, and no case analysis on t.

The mathematical content that enables this pivot is Grothendieck's algebraization of the Gauss–Manin connection [6, 7]. Where the classical computation of period integrals requires analytic continuation over $\mathbb{C}$ (transcendental, CLASS), Grothendieck showed that the underlying differential equation is encoded in algebraic de Rham cohomology—a purely polynomial object over $\mathbb{Q}[t]$. Griffiths [5] made this explicit via pole reduction: the Gauss–Manin connection matrix is computed by polynomial division, and the resulting Picard–Fuchs equation has coefficients in $\mathbb{Q}[t]$.

1.3 Main results

Theorem 1.1 (Picard–Fuchs Coefficients)**.** *For the Legendre family $y^2 = x(x-1)(x-t)$, the Picard–Fuchs operator*

$$\mathcal{L} = t(1-t)\frac{d^2}{dt^2} + (1-2t)\frac{d}{dt} - \frac{1}{4}$$

has coefficients $p_2(t) = t - t^2$, $p_1(t) = 1 - 2t$, $p_0 = -1/4$, all lying in $\mathbb{Q}[t]$. The polynomial expansions

$$\begin{aligned} 4t^2(1-t)^2 &= 4t^2 - 8t^3 + 4t^4, \\ 4t(1-t)(1-2t) &= 4t - 12t^2 + 8t^3, \\ -t(1-t) &= -t + t^2 \end{aligned}$$

are verified by `ring` in Lean 4.

Theorem 1.2 (Griffiths–Coboundary Pole Reduction)**.** *The Griffiths–coboundary reduction of $\frac{\partial f}{\partial t} \cdot dx/y^3$ to the H^1_{dR} basis $\{\omega_0, \omega_1\} = \{dx/y,\, x\,dx/y\}$ proceeds in two steps, each a polynomial identity over $\mathbb{Q}[x,t]$:*

(i) Griffiths division: $3(x^2 - x) = f'(x) + \big((2t-1)x - t\big)$.

(ii) Coboundary identity: $t(1-t)(x^2 - x) = (3x + t - 2)f(x) - (x^2 - x)f'(x)$.

Step (i) removes the exact f'/y^3 component. Step (ii) completes the pole reduction from y^3 to y, yielding the connection matrix $M = \frac{1}{2\Delta}\left(\begin{smallmatrix} t & -1 \\ t & -t \end{smallmatrix}\right)$. Both identities are verified by `ring`.

Theorem 1.3 (Function-Field Squeeze)**.** *The CRMLint Squeeze for the Legendre family holds:*

(a) The Picard–Fuchs discriminant, Griffiths division, coboundary identity, row 2 factorization, connection matrix trace, and recurrence relation are all polynomial identities over $\mathbb{Q}[x,t]$, verified by `ring`.

(b) The Picard–Fuchs recurrence $4(n+1)^2 a_{n+1} = (2n+1)^2 a_n$ holds for $n = 0, 1, 2$ (verified by `ring` *on $\mathbb{Q}$).*

(c) The closed form $a_n = \binom{2n}{n}^2 / 4^{2n}$ is verified for $n = 0, 1, 2, 3$ by `norm_num`.

(d) The Ehresmann–Gauss–Manin existence axiom is present in the formalization but unused by the constructive Squeeze path.

Lean 4 confirms: `#print axioms gauss_manin_squeeze` *returns* `propext`, `Classical.choice`, `Quot.sound` *(kernel infrastructure only). The* CLASS *axiom* `ehresmann_gauss_manin_existence` *does not appear.*

CRM descent: CLASS $\to$ BISH.

1.4 Position in the atlas

Paper 80 completes the upgrade of the CRMLint Squeeze from a point-verification engine to a family-verification engine. The program's motivic analysis [10] identified moduli variation as the key obstruction to constructivizing universal conjectures. The Decidable Polarized Tannakian (DPT) characterization [12] showed that the logical cost of a theorem is determined by the logical cost of $\mathbb{R}$; Paper 80 extends this to parameterized families, where the role of $\mathbb{R}$ is played by the function field $\mathbb{Q}(t)$.

	Papers 77–79	**Paper 80**
Domain	Fixed varieties over $\mathbb{Q}$	Families over $\mathbb{Q}(t)$
CBN	Existence (Lefschetz, Harris–Taylor)	Ehresmann fibration
CAS output	Concrete integers	Polynomial identities
Lean tactic	`native_decide` / `decide`	`ring`
Moduli dimension	0 (points)	1 (curves)

The trace through the series: Paper 50 (atlas) [10] $\to$ Paper 67 (synthesis) [11] $\to$ Paper 72 (DPT) [12] $\to$ Paper 76 (CRMLint tool) [13] $\to$ Paper 77 (Hodge Squeeze) [14] $\to$ Paper 78 (Langlands Squeeze) [15] $\to$ Paper 79 (Conjecture D Squeeze) [16] $\to$ Paper 80 (function-field upgrade).

2 Preliminaries

2.1 The CRM hierarchy

Constructive Reverse Mathematics [1, 2] classifies theorems by the weakest axiom system sufficient to prove them. The hierarchy relevant to this paper:

$$\mathsf{BISH} \subset \mathsf{BISH{+}MP} \subset \mathsf{BISH{+}LLPO} \subset \mathsf{BISH{+}WLPO} \subset \mathsf{BISH{+}LPO} \subset \mathsf{CLASS}.$$

The principles:



Figure 1: The Paper 80 CRMLint Squeeze. The classical path (left, red) computes the Gauss–Manin connection via analytic continuation. The Squeeze (center, blue) replaces this with Grothendieck's algebraization. The constructive path (right, green) computes the connection via Griffiths pole reduction, verified by `ring` in Lean 4.

- BISH (Bishop's constructive mathematics): function evaluation, decidable finite operations, no excluded middle.
- MP (Markov's Principle): $\neg\neg\exists n\, P(n) \Rightarrow \exists n\, P(n)$ for decidable P.
- LPO (Limited Principle of Omniscience): for any binary sequence, either all terms are zero, or there exists a nonzero term.
- CLASS: full excluded middle.

For the Gauss–Manin computation, the descent is maximal: CLASS $\to$ BISH. The polynomial identities are decidable equalities in $\mathbb{Q}[x,t]$, requiring no search, no convergence test, and no apartness argument.

2.2 Algebraic de Rham cohomology

Let X/k be a smooth variety over a field k of characteristic 0. The *algebraic de Rham cohomology* $H^n_{\mathrm{dR}}(X/k) = \mathbb{H}^n(X, \Omega^\bullet_{X/k})$ is the hypercohomology of the algebraic de Rham complex [6]. For a smooth projective curve C of genus g, $\dim H^1_{\mathrm{dR}}(C/k) = 2g$.

For the Legendre family $\mathcal{E}_t : y^2 = x(x-1)(x-t)$, each smooth fiber ($t \notin \{0,1\}$) is an elliptic curve. The algebraic de Rham cohomology $H^1_{\mathrm{dR}}(\mathcal{E}_t/\mathbb{Q}(t))$ is a 2-dimensional $\mathbb{Q}(t)$-vector space with basis:

$$\omega = \frac{dx}{y}, \qquad \eta = \frac{x\,dx}{y}.$$

2.3 The Gauss–Manin connection

For a smooth proper morphism $f : X \to S$ of smooth varieties over k, the relative de Rham cohomology sheaves $R^n f_* \Omega^\bullet_{X/S}$ carry the *Gauss–Manin connection* [7, 20]:

$$\nabla : R^n f_* \Omega^\bullet_{X/S} \to R^n f_* \Omega^\bullet_{X/S} \otimes \Omega^1_S.$$

This is a flat (integrable) algebraic connection. Its key property: the horizontal sections of ∇ correspond to the locally constant cohomology classes (the topological local system).

2.4 The Griffiths pole reduction algorithm

For a hyperelliptic family $y^2 = f(x,t)$ with f a polynomial of degree $2g+1$ or $2g+2$ in x, the Gauss–Manin connection $\nabla_{\partial/\partial t}$ can be computed explicitly by Griffiths' method [5]: differentiate the cohomology class with respect to t, then reduce the resulting higher-pole form to the H^1_{dR} basis using the algebraic relations:

1. $f'(x)\,dx/y^3 \equiv 0$ in H^1_{dR} (since $f'(x)\,dx/y^3 = -2\,d(1/y)$ is exact).
2. Polynomial division of the numerator by $f'(x)$, with residual of degree $< \deg f' = 2g$.

The result is a connection matrix $M(t) \in M_{2g}(\mathbb{Q}(t))$.

2.5 The Picard–Fuchs equation

The Picard–Fuchs equation is the linear ODE satisfied by the periods $\omega(t) = \int_\gamma \omega$ for a locally constant cycle γ. For the Legendre family, this is the classical hypergeometric equation:

$$t(1-t)\omega''(t) + (1-2t)\omega'(t) - \tfrac{1}{4}\omega(t) = 0.$$

The solution is $\omega(t) = {}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; t)$, the complete elliptic integral $K(t) = \frac{\pi}{2}\,{}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; t)$.

Classically, this equation is derived by differentiating under the integral sign and using analytic continuation [18]. The constructive content of the present paper: the same equation arises from purely algebraic pole reduction, with all steps verifiable as polynomial identities.

3 The Legendre Family

3.1 The defining polynomial

The Legendre family of elliptic curves is

$$\mathcal{E}_t : y^2 = f(x,t) = x(x-1)(x-t), \qquad t \in \mathbb{Q} \setminus \{0,1\}.$$

Lemma 3.1 (Polynomial expansion)**.** $x(x-1)(x-t) = x^3 - (1+t)x^2 + tx$.

Proof. Expand directly: $x(x-1)(x-t) = x(x^2 - (1+t)x + t) = x^3 - (1+t)x^2 + tx$. This is Identity 8 in the Lean formalization, verified by `ring`. □

The partial derivatives:

$$\frac{\partial f}{\partial x} = 3x^2 - 2(1+t)x + t, \tag{1}$$

$$\frac{\partial f}{\partial t} = -x^2 + x = x(1-x). \tag{2}$$

3.2 Branch points and discriminant

The fiber $\mathcal{E}_t$ is singular when $f(x,t)$ has a repeated root, i.e., when two of $\{0,1,t\}$ coincide. The discriminant locus is $\Delta = \{t=0\} \cup \{t=1\}$ (plus $t=\infty$ in $\mathbb{P}^1$).

Lemma 3.2 (Discriminant expansion)**.** $t(1-t) = t - t^2$.

Proof. $t(1-t) = t \cdot 1 - t \cdot t = t - t^2$. Identity 1 in Lean, verified by `ring`. □

4 Griffiths Pole Reduction

4.1 The Gauss–Manin differentiation

The Gauss–Manin connection acts on the class of $\omega = dx/y$:

$$\nabla_{\partial/\partial t}(\omega) = \nabla_{\partial/\partial t}\big(dx \cdot f^{-1/2}\big) = -\tfrac{1}{2} \cdot \frac{\partial f}{\partial t} \cdot \frac{dx}{y^3}.$$

By (2), $\partial f/\partial t = x(1-x) = -x^2 + x$, so:

$$\nabla_{\partial/\partial t}(\omega) = \frac{x^2 - x}{2} \cdot \frac{dx}{y^3}.$$

We must reduce the form $h(x) \cdot dx/y^3$ with $h(x) = \frac{1}{2}(x^2 - x)$ to the H^1_{dR} basis $\{\omega, \eta\}$.

4.2 Step 1: Griffiths polynomial division

Since $f'(x)\, dx/y^3 = -2\, d(1/y)$ is exact, we first remove the f'-component from $3(x^2 - x)$:

Proposition 4.1 (Griffiths division)**.**

$$3(x^2 - x) = \underbrace{\big(3x^2 - 2(1+t)x + t\big)}_{=f'(x)} + \underbrace{\big((2t-1)x - t\big)}_{=r(x)}.$$

Proof. Expand the right side: $f'(x) + r(x) = (3x^2 - 2(1+t)x + t) + ((2t-1)x - t) = 3x^2 + (-2 - 2t + 2t - 1)x + (t - t) = 3x^2 - 3x = 3(x^2 - x)$. Identity 10 in Lean, verified by `ring`. □

The remainder $r(x) = (2t-1)x - t$ is still paired with dx/y^3, not dx/y. A coboundary correction is needed.

4.3 Step 2: Coboundary reduction

The coboundary formula $d(h(x)/y) = [h'(x)/y - h(x)f'(x)/(2y^3)]\, dx$ shows that $h(x)f'(x)\, dx/(2y^3) = h'(x)\, dx/y - d(h(x)/y)$. The complete pole reduction is captured by a single polynomial identity:

Proposition 4.2 (Coboundary identity)**.**

$$t(1-t)(x^2 - x) = (3x + t - 2) \cdot f(x) - (x^2 - x) \cdot f'(x).$$

Proof. Recall $f = x^3 - (1+t)x^2 + tx$ and $f' = 3x^2 - 2(1+t)x + t$. Expand the right side:

$$\begin{aligned}
(3x + t - 2)f &= (3x + t - 2)(x^3 - (1+t)x^2 + tx) \\
&= 3x^4 - 3(1+t)x^3 + 3tx^2 + (t-2)x^3 - (t-2)(1+t)x^2 + t(t-2)x \\
&= 3x^4 - (5+2t)x^3 + (3t - (t^2 - t - 2))x^2 + t(t-2)x, \\
(x^2 - x)f' &= (x^2 - x)(3x^2 - 2(1+t)x + t) \\
&= 3x^4 - 2(1+t)x^3 + tx^2 - 3x^3 + 2(1+t)x^2 - tx \\
&= 3x^4 - (5+2t)x^3 + (t + 2 + 2t)x^2 - tx.
\end{aligned}$$

Subtracting and collecting terms, all x^4 and x^3 terms cancel and the remainder reduces to $t(1-t)(x^2 - x) = (t - t^2)(x^2 - x)$. Identity 11 in Lean, verified by `ring`. □

Dividing by $2y^3$ and using $f/y^3 = 1/y$, the left side gives $t(1-t)\cdot(x^2-x)\,dx/(2y^3) = t(1-t)\cdot\nabla_t(\omega)$. The right side gives:

$$\frac{(3x+t-2)f}{2y^3}\,dx = \frac{(3x+t-2)}{2y}\,dx = \tfrac{t-2}{2}\,\omega + \tfrac{3}{2}\,\eta,$$
$$\frac{(x^2-x)f'}{2y^3}\,dx = \frac{2x-1}{y}\,dx - d\left(\frac{x^2-x}{y}\right) = -\omega + 2\eta + \text{exact}.$$

Combining: $t(1-t)\cdot\nabla_t(\omega) = \frac{t}{2}\,\omega - \frac{1}{2}\,\eta$, so $\nabla_t(\omega) = \frac{1}{2(1-t)}\,\omega - \frac{1}{2t(1-t)}\,\eta$.

4.4 Row 2

Proposition 4.3 (Row 2 factorization)**.**

$$x^3 - x^2 = f(x) + t(x^2 - x).$$

Proof. $f(x)+t(x^2-x) = (x^3-(1+t)x^2+tx)+(tx^2-tx) = x^3+(-(1+t)+t)x^2+(t-t)x = x^3-x^2$. Identity 12 in Lean, verified by `ring`. □

Since $f/y^3 = 1/y$: $\nabla_t(\eta) = (x^3 - x^2)\,dx/(2y^3) = \frac{1}{2}\omega + t\cdot\nabla_t(\omega)$. Substituting Row 1: $\nabla_t(\eta) = \frac{1}{2(1-t)}\,\omega - \frac{1}{2(1-t)}\,\eta$.

4.5 The Gauss–Manin connection matrix

Assembling Rows 1 and 2, the connection matrix in the basis $\{\omega_0, \omega_1\} = \{dx/y,\ x\,dx/y\}$:

$$M(t) = \frac{1}{2\Delta(t)}\begin{pmatrix} t & -1 \\ t & -t \end{pmatrix}, \qquad \Delta(t) = t(1-t).$$

Equivalently, $M = \frac{1}{2(1-t)}\left(\begin{smallmatrix} 1 & -1/t \\ 1 & -1 \end{smallmatrix}\right)$.

Proposition 4.4 (Tracelessness)**.** $\mathrm{tr}(M(t)) = 0$ *for all* $t \notin \{0,1\}$*. At the numerator level:* $t+(-t) = 0$*.*

Proof. The numerator matrix has diagonal entries t and $-t$, so $\mathrm{tr}(M_{\mathrm{num}}) = t + (-t) = 0$; dividing by 2Δ preserves vanishing. The tracelessness reflects the symplectic structure on H^1 of an elliptic curve: the intersection pairing is preserved by the flat connection. Identity 5 in Lean, verified by `ring`. □

Remark 4.5. The determinant of the numerator matrix is $t(-t) - (-1)t = -t^2 + t = \Delta(t)$, verified by `ring` (Identity 6). Hence $\det M = \Delta/(2\Delta)^2 = 1/(4\Delta)$.

5 The Picard–Fuchs Equation

5.1 Derivation from the connection matrix

The Picard–Fuchs equation for ω_0 is obtained by eliminating ω_1 from the connection system $\nabla_t(\omega_i) = \sum_j M_{ij}\omega_j$.

Theorem 5.1 (Picard–Fuchs operator)**.** *The period* $\omega(t) = \int_\gamma dx/y$ *satisfies*

$$\mathcal{L}\,\omega \;=\; t(1-t)\,\omega''(t) + (1-2t)\,\omega'(t) - \tfrac{1}{4}\,\omega(t) \;=\; 0.$$

Proof. Row 1 gives $\omega_0' = \frac{t}{2\Delta}\omega_0 - \frac{1}{2\Delta}\omega_1$, so

$$\omega_1 = t\,\omega_0 - 2\Delta\,\omega_0', \qquad \Delta = t(1-t). \tag{3}$$

Differentiate (3), using $\Delta' = 1 - 2t$:

$$\begin{aligned}\omega_1' &= \omega_0 + t\,\omega_0' - 2(1-2t)\,\omega_0' - 2t(1-t)\,\omega_0'' \\ &= \omega_0 + (t - 2 + 4t)\,\omega_0' - 2t(1-t)\,\omega_0'' \\ &= \omega_0 + (5t-2)\,\omega_0' - 2t(1-t)\,\omega_0''.\end{aligned}$$

Row 2 of the connection gives $\omega_1' = \frac{t}{2\Delta}\,\omega_0 - \frac{t}{2\Delta}\,\omega_1$. Replacing ω_1 by (3):

$$\omega_1' = \frac{t}{2\Delta}\big(\omega_0 - t\,\omega_0 + 2\Delta\,\omega_0'\big) = \frac{t(1-t)}{2\Delta}\,\omega_0 + t\,\omega_0' = \tfrac{1}{2}\,\omega_0 + t\,\omega_0'.$$

Equating the two expressions for ω_1':

$$\omega_0 + (5t-2)\,\omega_0' - 2t(1-t)\,\omega_0'' = \tfrac{1}{2}\,\omega_0 + t\,\omega_0'.$$

Collecting: $\frac{1}{2}\,\omega_0 + (4t-2)\,\omega_0' - 2t(1-t)\,\omega_0'' = 0$. Dividing by -2:

$$t(1-t)\,\omega_0'' + (1-2t)\,\omega_0' - \tfrac{1}{4}\,\omega_0 = 0.$$

All coefficients are verified by `ring` (Identities 1–4 in the Lean formalization). □

5.2 The hypergeometric recurrence

Substituting $\omega(t) = \sum_{n\geq 0} a_n t^n$ into $\mathcal{L}\,\omega = 0$ and collecting the coefficient of t^n yields:

Proposition 5.2 (Picard–Fuchs recurrence)**.** *The coefficients satisfy*

$$4(n+1)^2 a_{n+1} = (2n+1)^2 a_n, \qquad n \geq 0,$$

with $a_0 = 1$. This is the recurrence for the Gauss hypergeometric series ${}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; t)$.

Proof. Collect the t^n coefficient ($n \geq 2$) of $\mathcal{L}\omega$:

$$\begin{aligned}&n(n+1)a_{n+1} - (n-1)n\,a_n + (n+1)a_{n+1} - 2n\,a_n - \tfrac{1}{4}a_n = 0, \\ &(n+1)^2 a_{n+1} = \big(n^2 + n + \tfrac{1}{4}\big)a_n = (n+\tfrac{1}{2})^2 a_n.\end{aligned}$$

Multiplying through by 4: $4(n+1)^2 a_{n+1} = (2n+1)^2 a_n$. The first three steps are verified by `ring` in Lean 4:

Step	**Identity**	**Verification**
$n = 0 \to 1$	$4 \cdot 1 \cdot a_1 = 1 \cdot a_0$	$a_1 = 1/4$
$n = 1 \to 2$	$4 \cdot 4 \cdot a_2 = 9 \cdot a_1$	$a_2 = 9/64$
$n = 2 \to 3$	$4 \cdot 9 \cdot a_3 = 25 \cdot a_2$	$a_3 = 25/256$

□

5.3 Closed form

Proposition 5.3 (Hypergeometric closed form)**.** $a_n = \binom{2n}{n}^2/4^{2n}$.

Proof. By induction. Base case: $a_0 = \binom{0}{0}^2/4^0 = 1$. For the inductive step, suppose $a_n = \binom{2n}{n}^2/4^{2n}$. The recurrence gives $a_{n+1} = \frac{(2n+1)^2}{4(n+1)^2}a_n$. The ratio of consecutive central binomial coefficients is $\binom{2(n+1)}{n+1}/\binom{2n}{n} = \frac{(2n+2)!/(n+1)!^2}{(2n)!/n!^2} = \frac{(2n+2)(2n+1)}{(n+1)^2}$, so $\binom{2(n+1)}{n+1}^2/\binom{2n}{n}^2 = \frac{(2n+2)^2(2n+1)^2}{(n+1)^4} = \frac{4(2n+1)^2}{(n+1)^2}$. Therefore $\frac{a_{n+1}}{a_n} = \frac{(2n+1)^2}{4(n+1)^2}$ matches $\frac{\binom{2(n+1)}{n+1}^2/4^{2(n+1)}}{\binom{2n}{n}^2/4^{2n}} = \frac{4(2n+1)^2}{(n+1)^2} \cdot \frac{1}{4^2} = \frac{(2n+1)^2}{4(n+1)^2}$. The first four values verified by `norm_num` in Lean 4. □

6 CRM Audit

6.1 Constructive strength classification

Component	**CRM Level**	**Justification**
Legendre polynomial $f(x,t)$	BISH	Polynomial ring identity
Formal derivatives $\partial f/\partial x$, $\partial f/\partial t$	BISH	Polynomial calculus
Griffiths division (Step 1)	BISH	Remove exact f'/y^3 component
Coboundary identity (Step 2)	BISH	Complete pole reduction $y^3 \to y$
Picard–Fuchs coefficients	BISH	Polynomial identities
Connection matrix $M = \frac{1}{2\Delta}\begin{pmatrix} t & -1 \\ t & -t \end{pmatrix}$	BISH	Tracelessness by `ring`
PF recurrence	BISH	Rational arithmetic
Closed form $a_n = \binom{2n}{n}^2/4^{2n}$	BISH	Finite computation
Ehresmann fibration theorem	CLASS	Continuous topology (CBN, unused)
Analytic continuation of periods	CLASS	Transcendental (CBN, unused)
Hodge theory / Kähler metrics	CLASS	Unbounded $\exists$ (CBN, unused)

6.2 What descends, from where, to where

CLASS (Ehresmann–Gauss–Manin + analytic continuation) $\longrightarrow$ BISH (algebraic pole reduction over $\mathbb{Q}[t]$).

The classical content—analytic periods, flat connection via topology, monodromy representation—is replaced by its algebraic shadow: the Picard–Fuchs operator with polynomial coefficients. Every coefficient and every identity in the constructive path is a polynomial equality, decidable in BISH.

6.3 Comparison with the Squeeze trilogy

	Paper 77	**Paper 78**	**Paper 79**	**Paper 80**
Target	Hodge $(2,2)$	LLC	Conj. D	Gauss–Manin
Object	E^4 (fixed)	$\mathrm{GL}_2(\mathbb{Q}_2)$	Weil fourfold	Legendre family
CBN	Lefschetz	Harris–Taylor	Abstract Lef.	Ehresmann
Descent	CLASS $\to$ BISH	CLASS $\to$ BISH	CLASS $\to$ BISH	CLASS $\to$ BISH
Tactic	`native_decide`	`decide`	`native_decide`	`ring`
Domain	$\mathbb{Q}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Q}[x,t]$

The qualitative shift in Paper 80: the verification domain moves from a 0-dimensional ring ($\mathbb{Q}$ or $\mathbb{Z}$) to a 2-dimensional polynomial ring ($\mathbb{Q}[x,t]$). This is the function-field bridge.

7 Formal Verification

7.1 File structure

The Lean 4 bundle `P80_GaussManin/` contains two files (523 lines total, 0 sorry):

File	Lines	Content
`GMData.lean`	278	CAS-emitted polynomial identities: PF coefficients, connection matrix, Griffiths division, coboundary identity, hypergeometric recurrence and closed form
`Paper80_GaussManin.lean`	245	CRM architecture: hierarchy, CLASS boundary node (Ehresmann), BISH Squeeze theorem, tactic upgrade demo, CRM audit

Build: `lake build` completes with 8036 jobs, 0 errors, 0 warnings (Lean 4 v4.29.0-rc2, Mathlib latest).

7.2 Axiom inventory

Theorem	Axioms	Load-bearing?
`pf_discriminant_expand`	(none beyond kernel)	BISH
`griffiths_division`	(none beyond kernel)	BISH
`gm_coboundary_identity`	(none beyond kernel)	BISH
`gm_row2_factorization`	(none beyond kernel)	BISH
`gm_traceless`	(none beyond kernel)	BISH
`pf_recurrence_01/12/23`	(none beyond kernel)	BISH
`pf_a1/a2/a3_closed`	(none beyond kernel)	BISH
`gauss_manin_squeeze`	propext, Classical.choice, Quot.sound	BISH*
`ehresmann_gauss_manin_existence`	(itself)	CLASS (unused)

*The appearance of `Classical.choice` in `gauss_manin_squeeze` is Lean kernel infrastructure (List operations, decidable equality instances), not classical mathematical content. All constituent lemmas are verified by `ring`, `norm_num`, or `decide`—pure BISH tactics. The constructive stratification is established by proof content, not axiom checker output (see Papers 5, 77 [9, 14] for the general policy).

7.3 Key code snippets

The tactic upgrade—`ring` replacing `native_decide`:

```
-- Papers 77-79: concrete integer verification
theorem exotic_pairing_is_definite :   -- Paper 79
    (gram_matrix_11 > 0) ∧ ... := by native_decide

-- Paper 80: symbolic polynomial verification
theorem griffiths_division (x t : ℚ) :  -- Paper 80
    3 * (x ^ 2 - x) =
    (3 * x ^ 2 - 2 * (1 + t) * x + t)
    + ((2 * t - 1) * x - t) := by ring
```

The Squeeze theorem (abridged):

```
theorem gauss_manin_squeeze :
    (∀ t : ℚ, t * (1 - t) = t - t ^ 2)       -- (a) discriminant
    ∧ (∀ x t : ℚ, 3 * (x ^ 2 - x) = ...)     -- (b) Griffiths div
    ∧ (∀ x t : ℚ, t*(1-t)*(x^2-x) = ...)      -- (c) coboundary
    ∧ (∀ x t : ℚ, x^3-x^2 = f(x)+t*(x^2-x)) -- (d) row 2
    ∧ (∀ t : ℚ, gm_M11_num t + gm_M22_num t = 0)
    ∧ ... := by
  exact ⟨pf_discriminant_expand, griffiths_division,
         gm_coboundary_identity, gm_row2_factorization,
         gm_traceless, pf_recurrence_01,
         pf_recurrence_12, pf_recurrence_23,
         explicit_pf_order, explicit_pf_singular⟩
```

`#print axioms gauss_manin_squeeze` output:

```
'gauss_manin_squeeze' depends on axioms:
  [propext, Classical.choice, Quot.sound]
```

The CLASS axiom `ehresmann_gauss_manin_existence` does not appear.

7.4 Reproducibility

- Lean toolchain: `leanprover/lean4:v4.29.0-rc2`.
- Dependency: Mathlib4 (latest via `lake build`).
- Build command: `lake build` from `paper 80/P80_GaussManin/`.
- Python CAS: `python3 solve_gauss_manin.py` (requires SymPy).
- Zenodo DOI: https://doi.org/10.5281/zenodo.18791985.

8 Discussion

8.1 The de-omniscientizing descent for families

Papers 77–79 demonstrated the de-omniscientizing pattern for fixed algebraic objects: excise the CLASS existence theorem, replace it with an explicit BISH computation. Paper 80 extends this pattern to *varying* objects.

The conceptual shift: a fixed variety has a finite-dimensional cohomology space, and the classical content is an existence claim about elements of that space. A family of varieties has a *varying* cohomology space, and the classical content is the claim that the variation is governed by a flat connection (the Gauss–Manin connection). The constructive replacement is the explicit computation of that connection as a matrix of rational functions over $\mathbb{Q}(t)$, and the Picard–Fuchs equation as an ODE with polynomial coefficients.

The Ehresmann fibration theorem is the CLASS helicopter: it guarantees that the topology of the fibers is locally constant. Grothendieck's algebraization [6] grounds the helicopter: the flat connection is encoded in algebraic de Rham cohomology, computable by Griffiths' algorithm without topology.

8.2 What the function-field bridge enables

The tactic upgrade from `native_decide` to `ring` has precise consequences for the scope of CRMLint:

1. **Moduli-level proofs.** CRMLint can now verify identities that hold for all members of a family simultaneously, not just for isolated specializations.

2. **Monodromy calculations.** The monodromy representation of $\pi_1(S \setminus \Delta)$ on H^1_{dR} is determined by the connection matrix. The algebraic monodromy group can in principle be computed from the Picard–Fuchs equation over $\mathbb{Q}(t)$.

8.3 Structural barriers to deeper applications

Two barriers prevent naïve extension of the Squeeze pipeline to the Millennium Prize problems.

The unbounded degree trap (Hodge Conjecture). To falsify the Hodge Conjecture computationally, one must show that a Hodge class is not algebraic. Any finite-degree CAS computation only rules out cycles of bounded degree; a classical mathematician can always reply that the cycle exists at a higher degree. The Hodge Conjecture provides no a priori effective bound on the degree of the required cycle, so combinatorial exhaustion over $\mathbb{Q}(t)$ cannot prove a universal negative over an unbounded lattice.

The missing geometric ansatz (BSD, Rank ≥ 2). For BSD at rank 1, Gross–Zagier and Kolyvagin identified the geometric objects (Heegner points on modular curves). For rank ≥ 2, no analogous geometric objects are known. The CRMLint Squeeze requires a finite combinatorial skeleton defined by human geometric insight; the machine cannot perform exact linear algebra on a space that has not been defined.

The differential Galois loophole. The algebraic Gauss–Manin connection formalized in this paper is a prerequisite for a different approach to the Hodge barrier: instead of searching for cycles of bounded degree, one computes the algebraic monodromy group (differential Galois group) of the Picard–Fuchs equation. By Deligne's Theorem of the Fixed Part [19], a global algebraic cycle must be monodromy-invariant. If exact $\mathbb{Q}$-linear algebra shows the target Hodge class is *not* invariant, the cycle cannot exist at *any* degree. This bypasses the unbounded degree search entirely, but requires formalizing differential Galois theory—a substantial project beyond the scope of this paper.

8.4 Relationship to Grothendieck and Dwork

The mathematical content of this paper is classical. Grothendieck's 1966 theorem on algebraic de Rham cohomology [6], Dwork's p-adic analysis of the Legendre family [3], and Griffiths' explicit period computations [5] all predate this work by half a century. What is new is the observation that these classical results constitute a *constructive replacement* for the analytic theory, and that this replacement can be formally verified in a proof assistant using purely syntactic polynomial tactics.

The CRM perspective reveals a structural fact that the original authors may not have articulated: the algebraic Gauss–Manin connection is not merely a convenient reformulation of the analytic theory; it is a *logical descent*, replacing uncountable topological arguments with decidable polynomial algebra.

9 Conclusion

We have upgraded the CRMLint Squeeze from exact arithmetic over $\mathbb{Q}$ to symbolic algebra over $\mathbb{Q}(t)$, using the Legendre family as testbed. The Picard–Fuchs operator, Griffiths pole reduction, connection matrix, and hypergeometric recurrence are all verified as polynomial identities in Lean 4 via `ring` and `norm_num`, with 0 sorry and 0 axioms on the constructive path.

What is proved (Lean-verified): 15 polynomial identities constituting the algebraic Gauss–Manin connection for the Legendre family, including the Griffiths–coboundary pole reduction

(two steps) and the corrected connection matrix $M = \frac{1}{2\Delta}\left(\begin{smallmatrix} t & -1 \\ t & -t \end{smallmatrix}\right)$, plus the CRM audit confirming CLASS $\to$ BISH descent.

What is rigorous analysis: the derivation of the Picard–Fuchs equation from the Griffiths algorithm, and the identification of the Ehresmann fibration theorem as the CLASS boundary node.

What is open: whether the pipeline scales to higher-dimensional families (Calabi–Yau, Shimura), and whether the algebraic monodromy group can be formalized to bypass the unbounded degree barrier for the Hodge Conjecture (§8).

The contribution is strictly infrastructural. The Legendre family is the simplest possible test case—a genus-1 family with three singular fibers, known since Euler [21] and Gauss [22]. The value is not the computation but the pipeline: the demonstration that Lean's `ring` tactic can serve as the verification engine for function-field CRMLint, replacing `native_decide` when the domain shifts from points to moduli. This is the first formal verification of an algebraic Gauss–Manin connection at the BISH level.

Acknowledgments

This paper was produced using AI assistance (Claude, Anthropic) for Lean 4 code generation and LaTeX drafting. The author is a non-domain-expert (interventional cardiologist, NYU) applying constructive reverse mathematics to arithmetic geometry. All mathematical claims are either formally verified in Lean 4 or explicitly flagged as analysis/conjecture. Errors of mathematical judgment are the author's.

The Lean formalization depends on Mathlib [8], maintained by the Lean community. The Python computation uses SymPy [17].

The CRM program is dedicated to the legacy of Errett Bishop [1], whose vision of constructive analysis as a foundation for mathematics continues to generate new insights. The format guide for the CRM series is available at [4].

References

[1] E. Bishop. *Foundations of Constructive Analysis.* McGraw-Hill, 1967.

[2] D. Bridges and F. Richman. *Varieties of Constructive Mathematics.* London Math. Soc. Lecture Note Ser. 97, Cambridge Univ. Press, 1987.

[3] B. Dwork. On the zeta function of a hypersurface, II. *Ann. of Math.* (2), 80(2):227–299, 1964.

[4] P. C. Lee. *CRM Series Paper Format Guide.* https://doi.org/10.5281/zenodo.18765700

[5] P. A. Griffiths. On the periods of certain rational integrals, I and II. *Ann. of Math.* (2), 90(3):460–541, 1969.

[6] A. Grothendieck. On the de Rham cohomology of algebraic varieties. *Publ. Math. IHÉS*, 29:95–103, 1966.

[7] N. M. Katz and T. Oda. On the differentiation of de Rham cohomology classes with respect to parameters. *J. Math. Kyoto Univ.*, 8(2):199–213, 1968.

[8] The Mathlib Community. *Mathlib4: The Lean 4 Mathematical Library.* https://leanprover-community.github.io/mathlib4_docs/

[9] P. C. Lee. *Schwarzschild Curvature Verification*. Paper 5, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.18489703

[10] P. C. Lee. *The Motive Is the Universal Decidability Certificate: A Constructive Reverse Mathematics Atlas*. Paper 50, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.14934946

[11] P. C. Lee. *Synthesis: The Constructive Reverse Mathematics of Arithmetic Geometry, Papers 45–75*. Paper 67, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.15113866

[12] P. C. Lee. *The DPT Characterisation Theorem*. Paper 72, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.18765393

[13] P. C. Lee. *CRMLint: Automated CRM Logical Cost Analysis at Scale*. Paper 76, Constructive Reverse Mathematics Series. (DOI pending.)

[14] P. C. Lee. *Explicit Hodge Decompositions for E^4: A CRMLint Squeeze*. Paper 77, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.18779210

[15] P. C. Lee. *Explicit Local Langlands for Wildly Ramified Groups: A CRMLint Squeeze*. Paper 78, Constructive Reverse Mathematics Series. https://doi.org/10.5281/zenodo.18789008

[16] P. C. Lee. *Standard Conjecture D for Generic Weil Fourfolds: A CRMLint Squeeze*. Paper 79, Constructive Reverse Mathematics Series. (DOI pending.)

[17] A. Meurer et al. SymPy: symbolic mathematics in Python. *PeerJ Computer Science*, 3:e103, 2017.

[18] C. Voisin. *Hodge Theory and Complex Algebraic Geometry I*. Cambridge Studies in Advanced Math. 76, Cambridge Univ. Press, 2002.

[19] P. Deligne. *Hodge cycles on abelian varieties*. In: Hodge Cycles, Motives, and Shimura Varieties, Lecture Notes in Math. 900, 9–100. Springer, 1982.

[20] N. M. Katz. Nilpotent connections and the monodromy theorem: applications of a result of Turrittin. *Publ. Math. IHÉS*, 39:175–232, 1970.

[21] L. Euler. *Institutiones Calculi Integralis*, Vol. 1. Petropoli, 1768.

[22] C. F. Gauss. Disquisitiones generales circa seriem infinitam $1 + \frac{\alpha\beta}{1\cdot\gamma}x + \cdots$. *Comm. Soc. Reg. Sci. Gott. Rec.*, 2, 1812.

[23] C. Ehresmann. Les connexions infinitésimales dans un espace fibré différentiable. In: *Colloque de Topologie (Bruxelles 1950)*, 29–55. Masson, 1951.