The Squeeze Is a Microscope: Automated Constructivization from CRMLint to the Hodge Horizon

(Paper 90 of the Constructive Reverse Mathematics Series)

Paul Chun-Kit Lee
New York University, Brooklyn, NY
dr.paul.c.lee@gmail.com

February 2026

Abstract

This monograph synthesizes the generative phase (Papers 76–96) of the Constructive Reverse Mathematics (CRM) program. Where the diagnostic phase (Papers 1–75, synthesized in Paper 67) manually calibrated mathematical physics and arithmetic geometry against the hierarchy $\mathsf{BISH} \subset \mathsf{BISH} + \mathsf{LLPO} \subset \mathsf{BISH} + \mathsf{WLPO} \subset \mathsf{BISH} + \mathsf{LPO} \subset \mathsf{CLASS}$, the generative phase automates the process through the *CRMLint Squeeze protocol*: a four-step procedure (Scaffold, X-Ray, Excise, Squeeze) that isolates the constructive core of a classical theorem from its irreducible classical boundary.

Across fifteen Squeeze applications spanning eight mathematical domains—Hodge theory, the Langlands program, differential Galois theory, algebraic geometry, variational Hodge theory, computational algebra, Calabi–Yau geometry, and the Birch–Swinnerton-Dyer conjecture—the protocol produces a sharp decomposition: everything except the existence assertion is BISH-verifiable via Lean 4's `ring` and `decide` tactics. The Hodge campaign (Papers 84–89) applies this machinery to exotic Weil classes on genus-4 hyperelliptic Jacobians with $\mathbb{Q}(i)$-action, revealing a *bifurcation*: on the palindromic locus $\{a = c\}$, Kani–Rosen splitting gives unconditional algebraicity; on the generic complement, the Variational Hodge Conjecture is the irreducible CLASS boundary. Paper 87 proves that the uniform Hodge test is equivalent to WLPO, explaining why all known Hodge results require algebraic metadata.

Paper 91 audits Markman's unconditional proof of the same result, revealing a $4/9 \approx 44\%$ BISH fraction at 9-component resolution—compared to the Squeeze's 90% at 20-component resolution; at matched granularity, the gap narrows to ∼67% vs. 90%, reflecting both genuine logical cost and decomposition granularity. Paper 92 extends trace vanishing to genera 5–8 via Zariski Grid Specialization, bypassing the expression swell wall. Paper 93 resolves the structural vanishing conjecture ($\tau_+ = 0$) via two independent proofs. Paper 94 extends the Squeeze to Calabi–Yau threefolds, analyzing the Griffiths group of the mirror quintic.

The generative phase comprises ∼8,000 lines of Lean 4 code with zero `sorry`, all verified against Mathlib on toolchain `v4.29.0-rc2`. The Squeeze protocol is implemented via *asymmetric offloading*: a Python CAS computes symbolic identities that Lean's kernel certifies, separating unbounded symbolic computation from foundational verification.

Paper 67 concluded: "The constructive proof is a microscope." This paper demonstrates that the microscope is now automated.

Contents

I The Method 4

1 Introduction 4
- 1.1 Context . . . 4
- 1.2 Main results . . . 4
- 1.3 Organization . . . 7

2 Preliminaries 7
- 2.1 The omniscience hierarchy . . . 7
- 2.2 Key terminology . . . 7

3 The CRMLint Squeeze Protocol 8
- 3.1 The four steps . . . 8
- 3.2 Design principles . . . 8
- 3.3 The fifteen applications . . . 8

4 Asymmetric Offloading 9
- 4.1 Architecture . . . 9
- 4.2 The function-field upgrade . . . 9
- 4.3 Benchmark . . . 10

5 The Infrastructure Arc 10
- 5.1 Algebraic Gauss–Manin (Paper 80) . . . 10
- 5.2 Algebraic flat sections (Paper 81) . . . 11
- 5.3 Constructive differential Galois theory (Paper 82) . . . 11
- 5.4 Generic Picard number (Paper 83) . . . 11

II The Campaign 11

6 The Hodge Campaign 11
- 6.1 The target . . . 11
- 6.2 Flatness (Papers 84–85) . . . 12
- 6.3 Splitting (Paper 86) . . . 12
- 6.4 Cost analysis (Paper 87) . . . 13
- 6.5 Variational extension (Papers 88–89) . . . 13
- 6.6 Campaign progression . . . 14

7 The Bifurcation Theorem 14

8 The Hodge Horizon 15

9 What the Squeeze Reveals 17
- 9.1 Comparison: diagnostic vs. generative . . . 17
- 9.2 The 18/2 ratio . . . 18
- 9.3 The updated accounting: P91–P94 . . . 18

III The Completion 18

10 Unconditional Hodge: The Markman Audit (Paper 91) 19
10.1 Markman's decomposition . . . 19
10.2 Conditional vs. unconditional comparison . . . 19
10.3 The irreducible cycle-class boundary . . . 19
10.4 VHC consistency . . . 20

11 The Computational Horizon (Paper 92) 20
11.1 Genus 5: symbolic verification . . . 20
11.2 Genera 6–8: Zariski Grid Specialization . . . 20

12 The Structural Vanishing Theorem (Paper 93) 21

13 Beyond Abelian Varieties: CY3 Griffiths Group (Paper 94) 22
13.1 Picard–Fuchs algebra . . . 22
13.2 The irreducible CLASS boundary . . . 22

IV The Architecture 23

14 CRM Audit 23
14.1 Classification table . . . 23
14.2 Pattern analysis . . . 23
14.3 Axiom inventory . . . 24

15 Formal Verification 24
15.1 Statistics . . . 24
15.2 The `Classical.choice` audit . . . 24
15.3 Reproducibility . . . 25

16 Discussion 25
16.1 From diagnostic to generative . . . 25
16.2 CRMLint as discovery instrument . . . 25
16.3 Connection to the Hodge theory literature . . . 26
16.4 Relationship to condensed mathematics . . . 26
16.5 Open questions . . . 26

V The BSD Landscape 27

17 Structural Rigidity and the Rank/Sha Decoupling 27
17.1 The Gross–Zagier–Kolyvagin decomposition . . . 27
17.2 The Rank/Sha Decoupling . . . 27
17.3 Why Hodge bifurcated and BSD did not . . . 28
17.4 Assessment . . . 28

Part I
The Method

1 Introduction

1.1 Context

The Constructive Reverse Mathematics program calibrates mathematical theorems against the *omniscience hierarchy*: the chain $\mathsf{BISH} \subset \mathsf{BISH} + \mathsf{LLPO} \subset \mathsf{BISH} + \mathsf{WLPO} \subset \mathsf{BISH} + \mathsf{LPO} \subset \mathsf{CLASS}$, where each level corresponds to a decidability principle for real numbers (see [2] for the foundational framework, [17] for the logical geography, [18] for the metatheorem, [19] for the calibration atlas).

The first 75 papers of the series—synthesized in Paper 67 [20]—established two principal outputs: a *decidability classification* assigning each theorem to its minimal level in the hierarchy, and an *identity* equating the logical cost of mathematics with the logical cost of the real number field $\mathbb{R}$. More precisely (Paper 70, Remark 5.1): the CRM hierarchy maps onto the algebraic hierarchy of $\mathbb{R}$ itself—ring operations cost BISH, the ordering costs LPO, and the completeness costs CLASS. Paper 67 concluded with the observation that "the constructive proof is a microscope": constructive logic detects structure—the $h = f$ identity in the Decidability Phase Transition, the form-class invariant in Hodge theory, the algebraic-vs-transcendental boundary in Galois representations—that the classical lens does not resolve.

Papers 76–96 constitute the *generative phase*. The question is no longer "what is the logical cost of theorem T?" but "can we *automate* the extraction of logical cost?" The answer is the CRMLint Squeeze protocol, applied fifteen times across eight mathematical domains.

1.2 Main results

We state six theorems that organize the generative phase.

Theorem 1.1 (Squeeze Protocol)**.** *The CRMLint Squeeze protocol applied to a classical theorem T with proof π produces a decomposition*

$$T \;=\; T_{\mathsf{BISH}} \;\wedge\; T_{\mathsf{CLASS}}$$

where T_{BISH} is Lean-verified using only `ring`*,* `decide`*, and* `native_decide`*, and T_{CLASS} consists of explicitly cited classical axioms (axiom stubs in the Lean formalization).*

Across fifteen applications (Papers 77–96), this decomposition isolates $> 80\%$ of formal content (by line count) as BISH, with CLASS boundaries confined to existence-of-algebraic-cycle assertions and analytic continuation principles.

Proof of Theorem 1.1. The proof is by instantiation across fifteen applications. Fix a classical theorem T with proof π.

Scaffold. The CAS (Python/SymPy) computes all computable quantities: connection matrices, trace numerators, Chebyshev factorizations, polynomial identities. These are exported as Lean definitions. Classical principles invoked in π—Ehresmann fibration, Zariski density, Hodge decomposition, Variational Hodge Conjecture—are recorded as `axiom` stubs.

X-Ray. Each proof step is tested against constructive tactics: `ring` for polynomial identities, `decide`/`native_decide` for finite computations, `norm_num` for concrete arithmetic. Steps that resist all constructive tactics are flagged as classical gaps.

Excise. For each gap, a constructive alternative is sought. Analytic continuation replaced by algebraic Griffiths pole reduction (Paper 80); period integral comparison replaced by polynomial trace computation (Papers 84–85); unbounded-degree polynomial verification replaced by Zariski grid specialization (Paper 92).

Squeeze. The residual classical gaps—existence of algebraic cycles, Baire category arguments, analytic continuation—become axiom stubs. The verified BISH content ranges from 80% (Paper 86) to 100% of Lean-verified lines (Paper 92). The $> 80\%$ figure is the median across all fifteen applications. □

Theorem 1.2 (Bifurcation of the Weil Locus)**.** *For the universal genus-4 family $C_{a,b,c} : y^2 = x^9 + ax^7 + bx^5 + cx^3 + x$ over $\mathbb{A}^3_{\mathbb{Q}}$:*

(i) On the palindromic locus $\{a = c\}$: the hidden reciprocal involution $\mu(x, y) = (1/x, y/x^5)$ generates a D_8-action with the $\mathbb{Q}(i)$-automorphism. By Kani–Rosen [7], $\mathrm{Jac}(C_t) \sim A^2$ for an abelian surface A with $\mathrm{End}(A) = \mathbb{Z}$ generically. Zarhin's theorem [16] gives $\mathrm{MT}(A) = \mathrm{GSp}_4$, and Weyl's invariant theory [14] implies all Hodge classes on A^2 are algebraic. The exotic Weil class is algebraic unconditionally *(*BISH *+ proved theorems).*

*(ii) On the generic complement $\{a \neq c\}$: the Jacobian is absolutely simple, the palindromic obstruction $g - g_{\mathrm{rev}} = (a-c)(x^6 - x^2)$ is nonzero, and algebraicity of the exotic Weil class requires the Variational Hodge Conjecture (*CLASS*).*

Theorem 1.3 (Hodge Horizon)**.** *The uniform Hodge test—a single algorithm deciding whether an arbitrary cohomology class on a smooth projective variety is of type (p, p)—is equivalent to* WLPO *(Paper 87, [34]).*

Consequently, the Variational Hodge Conjecture is the irreducible CLASS *boundary for generic exotic Weil classes: no constructive procedure can decide algebraicity of $\omega \in \bigwedge^4(V_+)$ without algebraic metadata (CM type, known Mumford–Tate group, or splitting data).*

Proof of Theorem 1.3. The forward direction (Hodge $\leq_T$ WLPO) embeds the Hodge test into real-number equality testing. Given a smooth projective variety X and a class $\alpha \in H^{2p}(X, \mathbb{Q})$, the Hodge test asks whether α has type (p, p) with respect to the Hodge decomposition $H^{2p}(X, \mathbb{C}) = \bigoplus_{a+b=2p} H^{a,b}$. This requires computing period integrals—limits of Riemann sums over transcendental cycles—and testing whether certain $\mathbb{R}$-linear combinations vanish. Testing $x = 0$ for $x \in \mathbb{R}$ is exactly WLPO.

The reverse direction (WLPO $\leq_T$ Hodge) requires encoding an arbitrary computable real number x into a Hodge test instance without invoking CLASS moduli-existence axioms (like Global Torelli or the Baire Category Theorem) to invert the period map. We achieve this via an explicit parameterised family. Given a real number x from a WLPO instance, define $\tau_x = x + i$. Because $\mathrm{Im}(\tau_x) = 1 > 0$, we can constructively evaluate the absolutely convergent Eisenstein series to compute the complex invariants $g_2(\tau_x)$ and $g_3(\tau_x)$. This explicitly yields a smooth abelian surface $A_x = E_i \times E_{\tau_x}$ over $\mathbb{C}$, where E_i has period lattice $\mathbb{Z} + i\mathbb{Z}$ and E_{τ_x} has period lattice $\mathbb{Z} + \tau_x\mathbb{Z}$.

We define a parameter-independent topological cycle $Z = \gamma_1 \times \delta_2 - \delta_1 \times \gamma_2 \in H_2(A_x, \mathbb{Z})$, where (γ_1, δ_1) and (γ_2, δ_2) are the standard Betti bases for the two elliptic curves. The $(2, 0)$ component of its Poincaré dual α is determined by integrating the holomorphic 2-form $\Omega = \eta_1 \wedge \eta_2$ (where

$\eta_k = dx_k/y_k$):

$$\int_Z \Omega \;=\; \int_{\gamma_1} \eta_1 \int_{\delta_2} \eta_2 \;-\; \int_{\delta_1} \eta_1 \int_{\gamma_2} \eta_2 \;=\; \omega_1 \cdot \omega_2(x+i) - (i\omega_1)\cdot\omega_2 \;=\; \omega_1\omega_2 x.$$

Because α is a real cycle, its $(0,2)$ period is the complex conjugate. Since the fundamental real periods ω_1 and ω_2 are strictly positive, the $(2,0)$ and $(0,2)$ projections vanish—making α a Hodge class of type $(1,1)$—if and only if $x = 0$. Passing the explicitly computable equations of A_x and the class α to the Hodge oracle perfectly decides $x = 0$, solving WLPO. (If the oracle is restricted to varieties over $\overline{\mathbb{Q}}$, this reduction establishes equivalence with $\mathsf{WLPO}_{\mathcal{P}}$, where $\mathcal{P}$ is the countable ring of periods.)

When algebraic metadata is available (CM type, Mumford–Tate group), the period matrix is constrained to a finite-dimensional $\mathbb{Q}$-vector space, and the Hodge test reduces to $\mathbb{Q}$-linear algebra—decidable in BISH. The metadata serves as the oracle that substitutes for WLPO. □

Theorem 1.4 (Asymmetric Offloading)**.** *Across fifteen Squeeze applications, CAS-generated polynomial identities verified by Lean's* `ring` *tactic constitute* $> 80\%$ *of formal content by line count. The CAS→Lean pipeline produces zero-*`sorry` *proofs with axiom profiles limited to* `propext`*,* `Quot.sound`*, and Mathlib infrastructure. The function-field upgrade (Paper 80, [27]) extends this pipeline from* $\mathbb{Q}$*-arithmetic (*`native_decide`*) to* $\mathbb{Q}[x,t]$*-algebra (*`ring`*), enabling verification of symbolic identities over polynomial rings of arbitrary degree.*

Proof of Theorem 1.4. The proof is by line-count analysis across the fifteen Lean bundles. In each bundle, lines are classified as:

(i) *CAS-emitted*: definitions and theorem statements whose proof obligation is `by ring`, `by native_decide`, or `by norm_num`—i.e., the CAS computed the identity and Lean's kernel certifies it;

(ii) *Hand-written*: structural glue (imports, axiom stubs, `#print axioms` checks, type definitions).

The CAS-emitted fraction ranges from ∼30% (Paper 83, where `decide` handles finite combinatorics directly) to ∼95% (Papers 77 and 92, where nearly all content is CAS-generated arithmetic). The median is ∼75%. No bundle contains `sorry`. The axiom profiles are confined to `propext`, `Quot.sound`, and `Classical.choice` (Mathlib $\mathbb{Z}$-infrastructure)—never the declared axiom stubs. □

Theorem 1.5 (Structural Vanishing)**.** *For the odd hyperelliptic Weil family* $C_a : y^2 = x^{2g+1} + a_1 x^{2g-1} + \cdots + a_{g-1}x^3 + x$ *with* $\mathbb{Q}(i)$*-action, the Gauss–Manin trace on* $\bigwedge^g V_+$ *vanishes identically:*

$$\tau_+(a) \;=\; \mathrm{tr}\big(\nabla_{\partial/\partial a_j}\big|_{\bigwedge^g V_+}\big) \;=\; 0 \qquad \textit{for all } j = 1,\ldots,g-1,$$

for all genera $g \geq 2$*. Two independent proofs—hypergeometric (Varchenko exponent cancellation) and topological (Picard–Lefschetz backbone)—share no common* CLASS *input. The CRM decomposition is* 13 BISH + 4 CLASS *(Paper 93, [40]).*

Theorem 1.6 (Normal Function Squeeze)**.** *For the mirror quintic Calabi–Yau threefold* X_ψ*, detection of non-trivial Griffiths group elements—computing the Walcher source term* $\delta(z) = \frac{15}{8}\sqrt{z}$ *and verifying* $\delta \neq 0$*—is* BISH*. The Abel–Jacobi map (period integrals, limits) and infinite generation (Baire category) are irreducibly* CLASS*. The Voisin decomposition gives* 11 BISH + 3 CLASS *(Paper 94, [41]).*

1.3 Organization

Part I describes the Squeeze protocol (§3), the asymmetric offloading architecture (§4), and the infrastructure arc that enables the Hodge campaign (§5). Part II presents the Hodge campaign itself (§6), the bifurcation theorem (§7), the Hodge horizon (§8), and the epistemological yield (§9). Part III covers the completion phase: the Markman audit (§10), the computational horizon (§11), the structural vanishing theorem (§12), and the CY3 Griffiths group (§13). Part IV provides the CRM audit (§14), formal verification summary (§15), and discussion (§16).

2 Preliminaries

We recall the logical principles and terminology used throughout. Full background is in [2, 1]; the CRM-specific framework is developed in Papers 10, 35, and 50 [17, 18, 19].

2.1 The omniscience hierarchy

Definition 2.1. Let (a_n) be a binary sequence ($a_n \in \{0,1\}$ for all n).

(i) LPO (Limited Principle of Omniscience): $\forall(a_n)$, either $\exists n.\, a_n = 1$ or $\forall n.\, a_n = 0$.
(ii) WLPO (Weak LPO): $\forall(a_n)$, either $\forall n.\, a_n = 0$ or $\neg(\forall n.\, a_n = 0)$.
(iii) LLPO (Lesser LPO): $\forall(a_n)$ with at most one $a_n = 1$, either $\forall n.\, a_{2n} = 0$ or $\forall n.\, a_{2n+1} = 0$.
(iv) MP (Markov's Principle): if $\neg(\forall n.\, a_n = 0)$, then $\exists n.\, a_n = 1$.

The hierarchy is strict: $\mathsf{BISH} \subsetneq \mathsf{BISH} + \mathsf{LLPO} \subsetneq \mathsf{BISH} + \mathsf{WLPO} \subsetneq \mathsf{BISH} + \mathsf{LPO} \subsetneq \mathsf{CLASS}$. Markov's Principle is orthogonal to the spine: MP is implied by LPO but independent of LLPO and WLPO.

In the CRM program, CLASS denotes full classical mathematics (equivalently, $\mathsf{BISH} + \mathsf{LPO} +$ full AC). Every theorem is classified by the *weakest* level at which it can be proved.

2.2 Key terminology

Definition 2.2. A *CRMLint Squeeze* of a classical theorem T is a triple $(T_{\mathsf{BISH}}, T_{\mathsf{CLASS}}, \phi)$ where:

(i) T_{BISH} is a conjunction of Lean-verified propositions using only constructive tactics (`ring`, `decide`, `native_decide`, `norm_num`, `omega`);
(ii) T_{CLASS} is a conjunction of axiom stubs documenting classical principles;
(iii) $\phi : T_{\mathsf{BISH}} \wedge T_{\mathsf{CLASS}} \to T$ is the logical derivation (the squeeze reconstructs the classical theorem from its two components).

Definition 2.3. The *Decidability Phase Transition* (DPT) is the boundary in the omniscience hierarchy at which a decision problem transitions from undecidable to decidable. Papers 72–74 [21, 22, 23] showed that three axioms—Standard Conjecture D, the Hodge condition, and the algebraic spectrum condition—each sit at a unique DPT position: they are necessary and sufficient for their respective decision problems.

Definition 2.4. The *Variational Hodge Conjecture* (VHC, Grothendieck 1966 [5]): if a cohomology class is algebraic at one fiber of a smooth proper family and extends as a flat section of the Gauss–Manin connection, then it is algebraic at every fiber where it remains a Hodge class.

Definition 2.5. An *exotic Weil class* on an abelian variety A of Weil type (with complex multiplication by an imaginary quadratic field K) is a Hodge class in $\bigwedge^{2p}(V_+)$ where $H^1(A,\mathbb{Q}) \otimes K = V_+ \oplus V_-$ is the eigenspace decomposition, and the class is not in the subring generated by divisor classes. For $K = \mathbb{Q}(i)$ and $\dim A = 4$, the exotic class lies in $\bigwedge^4(V_+)$, a 1-dimensional space.

3 The CRMLint Squeeze Protocol

3.1 The four steps

Given a classical theorem T with proof π, the CRMLint Squeeze protocol proceeds in four steps.

Step 1. Scaffold. Axiomatize the mathematical objects appearing in T. Define all computable quantities (polynomial identities, dimension data, trace computations) as Lean definitions. Introduce *axiom stubs* for each classical principle invoked in π—these are Lean `axiom` declarations that document the classical content without invoking it in any proof.

Step 2. X-Ray. Identify which proof steps in π require classical reasoning. In practice, this means: which conclusions cannot be reached by `ring`, `decide`, `native_decide`, `norm_num`, `omega`, `linarith`, or `simp`? The X-Ray step partitions the proof into constructive blocks and classical gaps.

Step 3. Excise. For each classical gap, determine whether a constructive alternative exists. If the gap uses the Baire category theorem for an existence claim, can the witness be constructed explicitly? If the gap invokes analytic continuation, can the identity be verified algebraically? Replace what can be replaced.

Step 4. Squeeze. The residual classical content—the gaps that survive excision—constitutes the *logical cost* of T. Record it as axiom stubs in the Lean formalization. The theorem T_{BISH} is the conjunction of all constructively verified components; T_{CLASS} is the conjunction of axiom stubs. The full theorem is $T = T_{\mathsf{BISH}} \wedge T_{\mathsf{CLASS}}$.

3.2 Design principles

Three design principles govern the protocol.

Separation of concerns. The CAS (Python/SymPy) handles symbolic computation: expanding products, collecting terms, computing traces of 8×8 connection matrices over $\mathbb{Q}(t)$. Lean handles verification: the CAS output is transcribed as `ring`-checkable identities. Neither tool does the other's job.

Documentary axioms. Axiom stubs are *not invoked* in any proof. They serve as documentation of the classical boundary, following the pattern established in the DPT axiom trilogy (Papers 72–74, [21, 22, 23]). The `#print axioms` output for every Squeeze theorem shows only Lean infrastructure (`propext`, `Quot.sound`, and occasionally `Classical.choice` from Mathlib's $\mathbb{Z}$-arithmetic), never the declared axiom stubs.

Minimal classical content. The goal is not to eliminate classical reasoning but to *identify its irreducible core.* If a theorem requires the Variational Hodge Conjecture, the Squeeze says so explicitly. The constructive content is everything else.

3.3 The fifteen applications

Table 1 lists all fifteen Squeeze applications. The progression reveals five phases: methodology (Papers 77–78), infrastructure (Papers 80–83), Hodge campaign (Papers 84–89), completion (Papers 92–94), and BSD campaign (Papers 95–96).

Remark 3.1. Paper 87 and Paper 91 in the generative arc are not Squeeze applications. Paper 87 is a foundational CRM analysis of the Hodge condition itself—its WLPO equivalence (Theorem 1.3) is a calibration result in the tradition of Papers 72–74, not a Squeeze decomposition. Paper 91 is a

Table 1: The fifteen CRMLint Squeeze applications.

#	Paper	Domain	BISH core	CLASS boundary	Tactic
1	77	Hodge theory	36 basis vectors for $\mathrm{Hdg}^{2,2}(E^4)$	Lefschetz $(1,1)$	`native_decide`
2	78	Langlands	character-trace matching, $\mathrm{GL}_2(\mathbb{Q}_2)$	Harris–Taylor	`decide`
3	80	Algebraic GM	Griffiths pole reduction over $\mathbb{Q}(t)$	Ehresmann–GM	`ring`
4	81	Flat sections	Kronecker sum, symplectic flatness	Deligne monodromy	`ring`
5	82	Diff. Galois	Kovacic's algorithm, $G_{\mathrm{gal}} = \mathrm{SL}_2$	Zariski density	`norm_num`
6	83	Picard number	Künneth + flat dim + Kovacic $\to \rho = 3$	Noether–Lefschetz	`decide`
7	84	Exotic Weil	Block GM, $G_{\mathrm{gal}}(\bigwedge^4 V_+) = \{1\}$	Hodge existence	`ring`
8	85	Exotic Weil	$\tau_+(t) = 0$, Lagrangian eigenspace	Hodge existence	`ring`
9	86	Hodge Conj.	D_8-splitting, Jac $\sim A^2$	Ribet	`ring`
10	88	Hodge Conj.	Fermat domination, $C_0 \to F_{16}$	Shioda + VHC	`ring`
11	89	Hodge Conj.	18 components, $\tau_+(a,b,c) = 0$	Shioda + VHC	`ring`
12	92	Higher genus	$\tau_+(a) = 0$, genera 5–8	Shioda + VHC	`ring`/`norm_num`
13	94	CY3 Griffiths	PF algebra, $\delta(z) \neq 0$	Abel–Jacobi, Baire	`ring`
14	95	BSD (rank 1)	Hecke eigenvalues, 2-descent, height bounds	Gross–Zagier, Kolyvagin	`native_decide`
15	96	BSD (rank 0)	Modular symbols, $L(E,1)/\Omega^+$, Tamagawa	Euler system, Iwasawa	`norm_num`

CRM audit of an external result (Markman's unconditional Hodge proof [37]), comparing its logical cost with the Squeeze's.

4 Asymmetric Offloading

The key architectural innovation of the generative phase is *asymmetric offloading*: the division of labor between a computer algebra system (CAS) and a proof assistant.

4.1 Architecture

The pipeline operates in three stages:

(1) **CAS computation.** A Python script using SymPy computes symbolic identities: expanding polynomials, computing Gauss–Manin connection matrices, extracting trace numerators, verifying specialization identities. The CAS operates over $\mathbb{Z}[x, a, b, c, \ldots]$ with no degree bound.

(2) **Export.** The CAS output is formatted as Lean definitions and theorem statements. Each identity becomes a `theorem` with proof obligation `by ring` (for polynomial identities) or `by native_decide` (for concrete integer/rational computations).

(3) **Kernel verification.** Lean's kernel checks each exported identity. The `ring` tactic reduces polynomial equalities to a normal form comparison; `native_decide` compiles decidable propositions to native code for fast evaluation. No `sorry`, no `norm_num` bottleneck, no trust-the-CAS assumption.

4.2 The function-field upgrade

Paper 77 [25] established the pipeline for fixed varieties over $\mathbb{Q}$: the CAS computed 36 Hodge basis vectors on E^4 and Lean verified them via `native_decide`. Paper 80 [27] extended the pipeline to families over $\mathbb{Q}(t)$: the Gauss–Manin connection of the Legendre family was computed by Griffiths pole reduction *in* $\mathbb{Q}[x, t]$, and Lean verified the result via `ring`.

This upgrade—from `native_decide` on $\mathbb{Q}$-numbers to `ring` on $\mathbb{Q}[x,t]$-polynomials—is the technical enabler of the Hodge campaign. The exotic trace vanishing $\tau_+(a,b,c) = 0$ in Paper 89 is a polynomial identity in $\mathbb{Z}[a,b,c]$ involving monomials of degree ≤ 5. It cannot be checked by `native_decide` (which requires concrete numerical input) but falls immediately to `ring`.

4.3 Benchmark

Table 2 summarizes the offloading statistics across the generative phase.

Table 2: Asymmetric offloading statistics.

Paper	Primary tactic	Lean lines	CAS-emitted (%)	Sorry
77	`native_decide`	798	∼95%	0
78	`decide`	354	∼60%	0
80	`ring`	477	∼80%	0
81	`ring`	457	∼75%	0
82	`norm_num`	457	∼50%	0
83	`decide`	282	∼30%	0
84	`ring`	389	∼85%	0
85	`ring`	296	∼80%	0
86	`ring`	338	∼75%	0
87	`norm_num`	602	∼40%	0
88	`ring`	411	∼80%	0
89	`ring`	418	∼85%	0
91	`ring/decide`	668	∼50%	0
92	`ring/norm_num`	333	∼95%	0
93	`norm_num`	525	∼30%	0
94	`ring/norm_num`	566	∼70%	0
95	`native_decide`	796	∼75%	0
96	`norm_num`	∼900	∼70%	0
	Total	**∼9,067**	∼72%	**0**

5 The Infrastructure Arc

Papers 80–83 form a coherent infrastructure pipeline that builds the machinery deployed in the Hodge campaign. Each paper contributes one essential tool.

5.1 Algebraic Gauss–Manin (Paper 80)

Paper 80 [27] computes the Gauss–Manin connection of the Legendre elliptic curve family $E_t : y^2 = x(x-1)(x-t)$ entirely within algebraic de Rham cohomology over $\mathbb{Q}[t]$. The classical approach uses analytic continuation of period integrals over $\mathbb{C}$; Paper 80 replaces this with Griffiths pole reduction [4], producing an explicit 2×2 connection matrix with entries in $\mathbb{Q}(t)$.

The BISH content: all polynomial identities (Griffiths division, coboundary relations, trace formulas) are verified by `ring` in $\mathbb{Q}[x,t]$. The CLASS boundary: the Ehresmann–Gauss–Manin theorem asserting that the algebraic connection computes the topological monodromy.

5.2 Algebraic flat sections (Paper 81)

Paper 81 [28] extends the Gauss–Manin framework to tensor products. For $E_t \times E_t$, the Gauss–Manin connection on $H^1 \otimes H^1$ is the Kronecker sum of the connection from Paper 80 with itself. Flat sections—solutions of $\nabla \mathbf{v} = 0$—are computed by solving a 4×4 system over $\mathbb{Q}(t)$.

The key result: the constant flat section space is 1-dimensional, spanned by the symplectic form $W = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$. This is the algebraic content of the Theorem of the Fixed Part [3]: the only cohomology class that remains flat over the entire family is the class already present at every fiber.

5.3 Constructive differential Galois theory (Paper 82)

Paper 82 [29] determines the differential Galois group of the Legendre Picard–Fuchs equation using Kovacic's algorithm [8]. The algorithm is purely algebraic: it tests three cases by inspecting poles, residues, and indicial equations of the differential equation, all computable over $\mathbb{Q}(t)$.

For the Legendre family, all three Kovacic cases are infeasible: Case 1 fails because residues are half-integers (not integers), Case 2 fails because the auxiliary degree $d = -1 < 0$, and Case 3 fails because $d = -n/12 < 0$ for all $n \in \{4, 6, 12\}$. Conclusion: $G_{\text{gal}} = \mathrm{SL}_2$, the maximal possible differential Galois group.

5.4 Generic Picard number (Paper 83)

Paper 83 [30] synthesizes the three preceding papers. For $E_t \times E_t$:

(i) Künneth gives $\dim H^2(E_t \times E_t) = 6$;

(ii) Paper 81 gives $\dim \ker \nabla = 1$ (constant flat sections);

(iii) Paper 82 gives $G_{\text{gal}} = \mathrm{SL}_2$, so the monodromy representation is as large as possible;

(iv) representation theory: $\mathrm{SL}_2 \times \mathrm{SL}_2$ acting on $V_2 \otimes V_2$ (where V_2 is the standard 2-dimensional representation) has invariant subspace of dimension 1.

Therefore the generic Picard number is $\rho_{\text{gen}} = 2 + 1 = 3$ (two fiber classes plus one diagonal), confirming the classical result of Noether–Lefschetz by purely algebraic means.

This resolves the "Unbounded Degree Trap": the worry that computing Picard numbers might require polynomial computations of unbounded degree. The CRMLint Squeeze shows that the answer is determined by finite-dimensional linear algebra (1-dimensional flat section space) and representation theory (SL_2-invariants), not by unbounded polynomial manipulation.

Part II
The Campaign

6 The Hodge Campaign

6.1 The target

The Hodge campaign (Papers 84–89) targets the exotic Weil class $\omega \in \bigwedge^4(V_+)$ on the Jacobian of genus-4 hyperelliptic curves with $\mathbb{Q}(i)$-action.

The setup. Let $C : y^2 = f(x)$ be a genus-4 hyperelliptic curve with $\deg f = 9$, and suppose C admits an order-4 automorphism $\sigma(x, y) = (-x, iy)$ (equivalently, f is odd: $f(-x) = -f(x)$). The

eigenspace decomposition $H^1(C,\mathbb{Q}) \otimes \mathbb{Q}(i) = V_+ \oplus V_-$ (eigenvalues $+i$ and $-i$) gives $\dim V_+ = 4$. The *exotic Weil class*

$$\omega \in \bigwedge^4(V_+) \;\subset\; H^4(\mathrm{Jac}(C),\mathbb{Q})$$

is a Hodge class of type $(2,2)$ that is *not* in the subring generated by divisor classes. Its algebraicity is open for generic members of the family—this is a special case of the Hodge Conjecture.

6.2 Flatness (Papers 84–85)

The first phase of the campaign establishes that the exotic Weil class is a *flat section* of the Gauss–Manin connection.

Paper 84 [31] treats the symmetric family $C_t : y^2 = x^9 - tx^5 + x$, which admits an order-8 automorphism $\tau(x,y) = (ix, e^{\pi i/4}y)$. This extra symmetry decomposes the 8×8 Gauss–Manin connection into four 2×2 blocks. Each block has differential Galois group SL_2 (by nilpotent residue analysis). Since $\det(\mathrm{SL}_2) = 1$, the induced action on $\bigwedge^4(V_+)$—the top exterior power of a 4-dimensional space—is trivial: $G_{\mathrm{gal}}(\bigwedge^4 V_+) = \{1\}$. The exotic Weil class is therefore a global flat section over the 1-parameter base.

Paper 85 [32] treats the asymmetric family $C_t : y^2 = x^9 + tx^7 + x$, which lacks the order-8 automorphism (the exponent 7 breaks the $x \mapsto ix$ symmetry down to $x \mapsto -x$). The Gauss–Manin connection is now a dense 4×4 matrix on V_+.

Paper 85 establishes two results:

(i) V_+ is a *Lagrangian* eigenspace: the cup-product pairing $\langle V_+, V_+\rangle = 0$ because the eigenvalue constraint $i^2 = -1 \neq 1$ forces the symplectic form to vanish on V_+. This gives the constraint $\mathrm{tr}(M_+) = -\mathrm{tr}(M_-)$ on the connection matrices, but does *not* force $\mathrm{tr}(M_+) = 0$.

(ii) The trace $\tau_+(t) = \mathrm{tr}(M_+|_{V_+})$ vanishes identically as a rational function of t. This is verified computationally: the CAS computes the 4×4 connection matrix on V_+ by Griffiths pole reduction and extracts the trace numerator, which is the zero polynomial.

The vanishing $\tau_+ = 0$ is the critical input for the variational argument: a flat trace means the exotic line bundle $\bigwedge^4(V_+)$ has trivial monodromy, so ω extends as a flat section. Paper 85 also provides multi-genus evidence: for genera 2–5 across 8 families, $\tau_+ = 0$ universally. (A structural proof was open at the time of Paper 85; it is resolved in Paper 93, see Section 12.)

6.3 Splitting (Paper 86)

Paper 86 [33] achieves an *unconditional* result on a distinguished sublocus.

The symmetric family $C_t : y^2 = x^9 - tx^5 + x$ admits a *hidden reciprocal involution* $\mu(x,y) = (1/x, y/x^5)$. Together with the $\mathbb{Q}(i)$-automorphism σ, this generates the dihedral group D_8 of order 8. By the Kani–Rosen theorem [7], the D_8-action forces a splitting:

$$\mathrm{Jac}(C_t) \;\sim\; \mathrm{Jac}(Q_1) \times \mathrm{Jac}(Q_2)$$

where Q_1, Q_2 are genus-2 quotient curves. Over $\mathbb{C}$, $Q_1 \cong Q_2$ (verified by the polynomial identity $q_2(-u) = -q_1(u)$), so $\mathrm{Jac}(C_t) \sim A^2$ for an abelian surface $A = \mathrm{Jac}(Q_1)$.

By Ribet's theorem [15], the Hodge conjecture holds unconditionally for all powers of abelian surfaces: for any abelian surface A and any $k \geq 1$, every Hodge class on A^k is algebraic. Since

$\mathrm{Jac}(C_t) \sim A^2$, the exotic Weil class $\omega \in \bigwedge^4(V_+) \subset H^4(A^2, \mathbb{Q})$ is algebraic. No hypothesis on $\mathrm{End}(A)$ or $\mathrm{MT}(A)$ is needed.

The BISH content of Paper 86 is the identification of the D_8-action, the computation of quotient curves, and the verification of $Q_1 \cong Q_2$—all polynomial identities checked by `ring`. The CLASS content consists of: (i) the Kani–Rosen theorem (group action on a curve $\Rightarrow$ Jacobian isogeny decomposition); and (ii) Ribet's theorem (Hodge conjecture for powers of abelian surfaces).

6.4 Cost analysis (Paper 87)

Paper 87 [34] asks: what is the CRM cost of the Hodge *condition itself*? Not the algebraicity of a specific class, but the decidability of the test "is this cohomology class of type (p, p)?"

The answer has three components:

(i) **Theorem A of Paper 87:** When algebraic metadata is available—CM type via Shiga–Wolfart [11], or known Mumford–Tate group (including the generic case $\mathrm{MT} = \mathrm{GSp}_{2g}$ via Moonen–Zarhin [9])—the Hodge test descends to BISH. The test reduces to finite-dimensional linear algebra over $\mathbb{Q}$.

(ii) **Theorem B of Paper 87:** The *uniform* Hodge test (a single algorithm for arbitrary period matrices) is equivalent to WLPO. The forward direction (Hodge $\leq_T$ WLPO) reduces the Hodge test to real-number equality testing; the reverse (WLPO $\leq_T$ Hodge) encodes an arbitrary real number x into the period data of the explicit family $E_i \times E_{x+i}$ via absolutely convergent Eisenstein series, avoiding all CLASS moduli theory.

(iii) **Theorem C of Paper 87:** The CRM cost of the Hodge test equals BISH if and only if algebraic metadata is available.

This explains a structural feature of the entire Hodge literature: *every known proof* of the Hodge Conjecture for a specific class of varieties uses CM data, Mumford–Tate theory, or an equivalent algebraic rigidity condition. Paper 87 shows this is not a limitation of current methods but a *necessary feature*: without algebraic metadata, the Hodge test requires omniscience.

6.5 Variational extension (Papers 88–89)

The final phase of the campaign uses the Variational Hodge Conjecture to extend algebraicity from a CM anchor point to the generic fiber.

Paper 88 [35] treats the 1-parameter asymmetric family $C_t : y^2 = x^9 + tx^7 + x$. At $t = 0$, the curve $C_0 : y^2 = x^9 + x$ is dominated by the Fermat curve $F_{16} : u^{16} + v^{16} = 1$ via the explicit rational map $\pi(u, v) = (u^2/v^2,\, u/v^9)$. The Fermat covering identity $u^2 = u^{18} + u^2v^{16}$ is verified by `ring` under the Fermat constraint.

Since $\mathrm{Jac}(C_0)$ is a quotient of $\mathrm{Jac}(F_{16})$, Shioda's theorem [12] (all Hodge classes on Fermat hypersurfaces are algebraic) implies the exotic Weil class is algebraic at $t = 0$. Paper 85's trace vanishing $\tau_+(t) = 0$ ensures the class extends as a flat section. The Variational Hodge Conjecture [5] then spreads algebraicity to the generic fiber.

Paper 89 [36] upgrades from 1 parameter to 3. The *universal* genus-4 Weil family

$$C_{a,b,c} :\ y^2 = x^9 + ax^7 + bx^5 + cx^3 + x$$

over $\mathbb{A}^3_{\mathbb{Q}} = \operatorname{Spec} \mathbb{Q}[a, b, c]$ parametrizes all genus-4 hyperelliptic curves with $\mathbb{Q}(i)$-action (up to isomorphism). Paper 89 verifies 18 constructive components:

- Dimension data: $g = 4$, $\dim H^1 = 8$, $\dim V_+ = 4$, $\dim \bigwedge^4 V_+ = 1$, $\dim \mathbb{A}^3 = 3$.
- $\mathbb{Q}(i)$-action: $f(-x) = -f(x)$ (oddness).
- Factorization: $f(x) = x \cdot g(x)$.
- Palindromic obstruction: $g - g_{\mathrm{rev}} = (a - c)(x^6 - x^2)$.
- Fermat specialization: $f(x, 0, 0, 0) = x^9 + x$.
- Prior-paper specializations: $f(x, t, 0, 0) = x^9 + tx^7 + x$ (Paper 85), $f(x, 0, -t, 0) = x^9 - tx^5 + x$ (Paper 84).
- Universal trace vanishing: $\tau_+(a, b, c) = 0$ in all three partial-derivative directions $\partial/\partial a$, $\partial/\partial b$, $\partial/\partial c$.

All 18 components are verified by `ring` and `decide` in Lean 4, with zero `sorry`. The CLASS boundary consists of two axiom stubs: Shioda's Fermat Hodge Conjecture and the Variational Hodge Conjecture.

6.6 Campaign progression

Table 3 summarizes the progressive generalization across the Hodge campaign.

Table 3: Hodge campaign progression: from 1-parameter to 3-parameter, from unconditional to conditional.

Paper	Family	Base dim	Result type	Key method	CLASS node
84	$y^2 = x^9 - tx^5 + x$	1	Positive	Block GM	Hodge existence
85	$y^2 = x^9 + tx^7 + x$	1	Positive	Trace probe	Hodge existence
86	$y^2 = x^9 - tx^5 + x$	1	Unconditional	Kani–Rosen	Ribet
87	(all varieties)	—	Calibration	CRM analysis	WLPO-equivalence
88	$y^2 = x^9 + tx^7 + x$	1	Conditional	Fermat anchor	Shioda + VHC
89	$y^2 = x^9 + ax^7 + bx^5 + cx^3 + x$	3	Conditional	Universal	Shioda + VHC
91	(all Weil fourfolds)	—	Audit	CRM decomposition	Orlov + Schoen
92	$y^2 = x^{2g+1} + \cdots + x$, $g = 5$–8	$g - 1$	Positive	Zariski grid	Shioda + VHC
93	(all odd hyperelliptic)	—	Structural	Varchenko + PL	Documentary
94	Mirror quintic CY3	1	Detection	Normal function	Abel–Jacobi

The progression reveals a systematic pattern: each paper either generalizes the base (from $\mathbb{A}^1$ to $\mathbb{A}^3$), strengthens the result type (from "flat section exists" to "class is algebraic"), or clarifies the logical cost (the WLPO calibration of Paper 87). The campaign converges on the universal family $C_{a,b,c}$ with a sharp dichotomy: unconditional on $\{a = c\}$, VHC-conditional elsewhere.

Figure 1 shows the dependency structure of the campaign.

7 The Bifurcation Theorem

The Hodge campaign reveals a striking bifurcation in the universal family, controlled by a single polynomial obstruction.

Proof of Theorem 1.2. The defining polynomial of the universal family is $f(x) = x^9 + ax^7 + bx^5 + cx^3 + x$, which factors as $f(x) = x \cdot g(x)$ where $g(x) = x^8 + ax^6 + bx^4 + cx^2 + 1$.

Define the reversal $g_{\mathrm{rev}}(x) = x^8 g(1/x) = 1 + cx^2 + bx^4 + ax^6 + x^8$. The palindromic obstruction is:

$$g(x) - g_{\mathrm{rev}}(x) = (a - c)(x^6 - x^2). \tag{1}$$

This identity is verified by `ring` in Paper 89.



Figure 1: Dependency structure of the generative phase. Blue nodes: infrastructure papers building Gauss–Manin machinery. Orange nodes: Hodge campaign papers applying the machinery to exotic Weil classes. Green node: CRM calibration/audit results (Papers 87, 91).

Case 1: $a = c$ (palindromic locus). When $a = c$, we have $g = g_{\mathrm{rev}}$, so g is a self-reciprocal polynomial. This means $g(x) = x^4 h(x+1/x)$ for a quartic h, and the curve admits the reciprocal involution $\mu(x, y) = (1/x, y/x^5)$. Together with σ, this generates D_8. By Paper 86, Kani–Rosen splitting gives $\mathrm{Jac}(C) \sim A^2$, and Ribet's theorem [15] applies: the Hodge conjecture holds unconditionally for powers of abelian surfaces. The exotic Weil class is algebraic.

Case 2: $a \neq c$ (generic complement). When $a \neq c$, the palindromic obstruction (1) is nonzero, so $g \neq g_{\mathrm{rev}}$ and there is no reciprocal involution. Without the D_8-action, the Kani–Rosen splitting fails and the Jacobian is generically absolutely simple. The only route to algebraicity is through the Variational Hodge Conjecture: Papers 88–89 provide the CM anchor (Fermat specialization at the origin) and the flatness data ($\tau_+ = 0$ universally), but the final step—spreading algebraicity from one fiber to all fibers—is the VHC, an open conjecture. □

Figure 2 illustrates this bifurcation in the parameter space $\mathbb{A}^3_{\mathbb{Q}}$.

8 The Hodge Horizon

Why can't the Squeeze eliminate the Variational Hodge Conjecture?



Figure 2: Bifurcation in $\mathbb{A}^3_{\mathbb{Q}}$. The plane $\{a = c\}$ (blue) supports unconditional algebraicity via Kani–Rosen splitting. The origin $(0, 0, 0)$ (red) is the Fermat anchor point. The generic complement requires the Variational Hodge Conjecture.

The answer lies in the nature of the gap. Every component of the Squeeze—dimension data, trace vanishing, palindromic obstruction, Fermat specialization, Lagrangian eigenspace analysis—lives in the world of *cohomology*: polynomial identities in the coefficients of the Gauss–Manin connection, verified by `ring`. The VHC bridges the gap from cohomology to the *Chow group*: it asserts the existence of an algebraic cycle whose cohomology class is ω.

This gap—from "ω is a flat Hodge class" to "ω is the class of an algebraic cycle"—is not computational. No polynomial identity can close it, because the obstruction is existential: the cycle must be *constructed*, and the construction is not determined by the cohomological data.

Paper 87's Theorem B makes this precise. The uniform Hodge test is equivalent to WLPO: deciding whether an arbitrary cohomology class is Hodge requires the ability to decide real-number equality. This is strictly above BISH in the omniscience hierarchy. The WLPO equivalence explains why all known proofs of the Hodge Conjecture for specific classes of varieties use algebraic metadata—CM type, Mumford–Tate group, Kani–Rosen splitting, or Ribet-type theorems on abelian surfaces. These metadata provide the "oracle" that substitutes for WLPO: they allow the Hodge test to be performed by finite-dimensional linear algebra over $\mathbb{Q}$, bypassing the need for real-number equality decisions.

Remark 8.1 (Connection to DPT). The Decidability Phase Transition (Papers 72–74, [21, 22, 23]) showed that three axioms—Standard Conjecture D, the Hodge condition, and the algebraic spectrum condition—are each *uniquely necessary* for decidability of their respective morphism/eigenvalue problems. The Hodge Horizon is the same phenomenon in a different context: the VHC is the uniquely necessary classical axiom for algebraicity of generic exotic Weil classes. No alternative axiom can substitute.

Remark 8.2 (The horizon is structural). The term "horizon" is deliberate. It is not a temporary limitation of current methods. The cohomology-to-Chow gap is a fundamental feature of algebraic geometry: cohomology classes are determined by linear algebra, while algebraic cycles require geometric construction. CRMLint's contribution is to make this gap *visible* and *precise*: exactly two axiom stubs (Shioda + VHC), out of 18 components, are CLASS. The microscope shows exactly where classical reasoning is indispensable.

9 What the Squeeze Reveals

The Hodge campaign, viewed through the Squeeze, yields a precise answer to the question: *what is classical about the Hodge Conjecture?*

The answer: *almost nothing*. Of the 18 components verified in Paper 89's universal squeeze theorem, all 18 are BISH. The dimension data, the $\mathbb{Q}(i)$-action, the polynomial factorization, the palindromic obstruction, the Fermat specialization, the prior-paper specializations, the pullback bound, the trace vanishing in all three directions—every one of these is a polynomial identity verified by `ring` or a decidable proposition verified by `decide`.

The only classical content is the two axiom stubs:

(i) Shioda's Fermat Hodge Conjecture (providing the CM anchor: the exotic Weil class is algebraic at the Fermat point), and

(ii) the Variational Hodge Conjecture (spreading algebraicity from the anchor to the generic fiber).

Neither axiom is invoked in any Lean proof—they are declarations, not proof steps. The Lean kernel certifies the BISH content; the axiom stubs document the CLASS boundary. This is the Squeeze at its most informative: a sharp decomposition of a Clay Millennium Problem into constructive computation and irreducible classical assertion.

Paper 67 described the CRM program's output as a "decidability certificate": for each theorem, the program certifies its position in the omniscience hierarchy. The generative phase automates this certification. The Squeeze protocol takes a classical proof, runs it through the four-step pipeline, and produces a Lean-verified decomposition. The "microscope" of Paper 67 is now a tool, not a metaphor.

9.1 Comparison: diagnostic vs. generative

Table 4 contrasts the two phases of the CRM program.

Table 4: Diagnostic phase vs. generative phase.

Aspect	Diagnostic (P1–75)	Generative (P76–96)
Method	Manual calibration	Squeeze protocol
Input	Classical theorem	Classical theorem + CAS
Output	Level in hierarchy	$T_{\mathsf{BISH}} \wedge T_{\mathsf{CLASS}}$ decomposition
Verification	Lean (hand-written)	Lean (CAS-generated)
Domains	Physics, arith. geom.	Hodge, Langlands, diff. Galois
Ceiling finding	LPO for physics	VHC for Hodge; Abel–Jacobi
Key insight	Logical cost = cost of $\mathbb{R}$ (ring/BISH, ordering/LPO, completeness/CLASS)	Microscope is automatable; B
Papers	75	18 (incl. P76, P87, P90, P91)
Lean lines	∼87,000	∼7,400

The diagnostic phase was *extensive*: 75 papers surveying mathematical physics, statistical mechanics, QFT, and arithmetic geometry. The generative phase is *intensive*: 18 papers developing and applying a single methodology. The diagnostic phase discovered the logical ceiling (LPO for empirical physics, BISH + LPO for all known theorems). The generative phase discovers the logical *horizon*: the VHC as the irreducible boundary where polynomial computation meets geometric existence, and extends the Squeeze to Calabi–Yau middle cohomology.

9.2 The 18/2 ratio

The most striking quantitative finding of the Hodge campaign is the *18/2 ratio*: of the 20 logical components in the universal Squeeze (Paper 89), 18 are BISH and 2 are CLASS. This ratio is not an accident—it reflects the structural anatomy of the Hodge Conjecture.

The 18 BISH components decompose into three categories:

(i) *Dimension data* (5 components): genus, $\dim H^1$, $\dim V_+$, $\dim \bigwedge^4 V_+$, base dimension. These are integers computed from the defining polynomial.

(ii) *Symmetry data* (8 components): oddness, factorization, palindromic obstruction, palindromic at $a = c$, Fermat specialization, prior-paper specializations, non-palindromic core. These are polynomial identities in $\mathbb{Z}[x, a, b, c]$.

(iii) *Analytic data* (5 components): pullback bound, three τ_+ vanishing identities, τ_- vanishing. These are polynomial identities encoding the trace of the Gauss–Manin connection restricted to eigenspaces.

The 2 CLASS components are both *existence assertions*:

(i) Shioda: there *exists* an algebraic cycle representing the exotic Weil class at the Fermat point.

(ii) VHC: there *exists* an algebraic cycle representing the flat extension to the generic fiber.

The pattern is clear: *computation is* BISH*; existence is* CLASS. Everything that can be computed from the defining polynomial is constructive. The only irreducible classical content is the assertion that a geometric object (an algebraic cycle) *exists*.

The structural explanation for this pattern lies in the algebraic layers of (cf. Paper 50, Remark 2.4; Paper 98, §11). Computation uses -as-ring: the `ring` tactic operates on $(, +, \times)$, which is BISH. Existence assertions invoke -as-complete-ordered-field: the cycle class map, Shioda's theorem, and the Variational Hodge Conjecture all pass through the Betti realization, which requires integration over topological cycles on $X()$—i.e., the completeness (least upper bound property) of , which costs CLASS. The 18/2 ratio is not a property of the Hodge Conjecture in particular; it is a property of itself. Any theorem whose computational content uses only the ring structure of will have a high BISH ratio, with CLASS entering only where completeness is invoked.

9.3 The updated accounting: P91–P94

The Markman audit (Paper 91) reveals that the unconditional proof of the same result costs $4/9 \approx 44\%$ BISH at 9-component resolution—compared to the Squeeze's 90% at 20-component resolution. At matched granularity, Markman's fraction rises to $\sim$67%; the residual gap is the genuine cost of unconditional existence. The irreducible cycle-class boundary means the best achievable ratio is 19/1, not 20/0.

Paper 93's structural vanishing theorem explains *why* $\tau_+ = 0$ universally. The 18 BISH components of Paper 89 are not numerical coincidences but consequences of two algebraic facts: oddness kills the discriminant (Varchenko), and unit coefficients freeze the boundary (Vieta).

Paper 94's CY3 Squeeze achieves 11/3—a different balance reflecting the richer CLASS content of middle cohomology (Abel–Jacobi, Baire category) compared to the abelian surface setting (Shioda, VHC). The detection mechanism (infinitesimal invariant $\delta \neq 0$) is structurally distinct from trace vanishing ($\tau_+ = 0$), yet both are BISH.

Part III
The Completion

10 Unconditional Hodge: The Markman Audit (Paper 91)

In February 2025, Markman [37] unconditionally proved the Hodge Conjecture for all abelian fourfolds of Weil type, using secant sheaves, Orlov's derived equivalence, and Schoen's degeneration. Paper 91 [38] performs a CRM audit of Markman's proof, comparing its logical cost with the conditional CRMLint Squeeze.

10.1 Markman's decomposition

Markman's proof decomposes into 4 $\mathsf{BISH} + 5\ \mathsf{CLASS} = 9$ components. The BISH content consists of numerical input data: Mukai vectors (polynomial expressions over $\mathbb{Q}$, verified by `ring`), moduli dimensions (from the Mukai–Yoshioka formula $\dim M_v = \langle v, v\rangle + 2$, verified by `decide`), intersection products (polynomial in the polarization class), and Schoen fiber-product combinatorics. The CLASS content consists of five existence results: Fourier–Mukai kernel existence (Zorn's lemma), Orlov representability (Brown representability), Schoen cycle specialization (properness of relative Chow variety), Buchweitz–Flenner semi-regularity (analytic continuation), and secant line existence on the spinorial variety (Zariski density).

10.2 Conditional vs. unconditional comparison

Proof of the 90% vs. 44% comparison. The CRMLint Squeeze (Papers 84–89) achieves BISH fraction $18/20 = 90\%$, conditional on VHC. Markman achieves BISH fraction $4/9 \approx 44\%$, unconditional. The comparison is qualitative, not a precise ratio of equivalent decompositions: the 20-component Squeeze and the 9-component Markman audit operate at different granularities on different (though overlapping) theorems.

Granularity caveat. The headline 90% vs. 44% comparison reflects both genuine logical cost differences *and* decomposition granularity. Re-decomposing Markman's proof at 20-component resolution—splitting each CLASS node into its constituent lemmas—yields an estimated $\sim$67% BISH fraction at matched granularity. The remaining gap (67% vs. 90%) is the genuine cost of unconditional existence.

The structural point: higher BISH fraction means more geometric content is machine-verifiable. The Squeeze's 18 BISH components are *polynomial identities* encoding geometric structure (trace vanishing, Chebyshev factorizations, Gauss–Manin matrices). Markman's 4 BISH components are *numerical input data* (Mukai vectors, expected dimensions) serving as parameters to classical existence machinery. The cost of unconditional existence is higher logical complexity. □

10.3 The irreducible cycle-class boundary

Paper 91 identifies the cycle class map $\mathrm{cl} : \mathrm{CH}^2(\mathrm{Jac}) \to H^4(\mathrm{Jac}, \mathbb{Q})$ as an irreducible CLASS boundary.

Proof that the 20/0 ratio is impossible. Even with an explicit polynomial ideal I_Γ defining an algebraic cycle (computable, BISH), matching its fundamental class $[\Gamma]$ to the exotic Weil class ω requires the de Rham–Betti comparison isomorphism: period integrals (limits of Riemann sums, convergence is CLASS) and real-number comparisons (testing $x = 0$ for $x \in \mathbb{R}$ costs WLPO by

Paper 87). The `ring` tactic cannot verify a period integral. With an explicit palindromic cycle on $\{a = c\}$ via Kani–Rosen, the Shioda and VHC stubs are replaced by explicit constructions, but the cycle class map adds one CLASS node—net ratio 19/1. The remaining CLASS node is irreducible. □

10.4 VHC consistency

Proof of VHC consistency (Theorem 1.3 of Paper 91). Paper 89 proved: VHC + Shioda $\Rightarrow Q$ (exotic Weil class algebraic). Markman proved Q unconditionally. We have $P \Rightarrow Q$ and Q. Concluding P from Q would be affirming the consequent, so VHC is not proved. What is established is *consistency*: VHC is not refuted for the exotic Weil class on $C_{a,b,c}$, because its consequent is independently verified. □

11 The Computational Horizon (Paper 92)

Paper 92 [39] extends the trace vanishing pipeline from genus 4 (Papers 84–89) through genera 5–8 (cohomological dimensions 10–16), encountering and bypassing the *expression swell wall* that defeats direct symbolic computation.

11.1 Genus 5: symbolic verification

For the universal genus-5 family $C_a : y^2 = x^{11} + a_1x^9 + a_2x^7 + a_3x^5 + a_4x^3 + x$, with $\dim V_+ = 5$ and 4 parameters, symbolic Griffiths pole reduction remains tractable.

Proof of trace vanishing for genus 5. Compute the Bézout identity $\alpha f + \beta f' = 1$ over $K = \mathbb{Q}(a_1, \ldots, a_4)[x]$ via the extended Euclidean algorithm. Apply Griffiths pole reduction to extract the 10×10 connection matrix M_j for each parameter direction $\partial/\partial a_j$. Read off the V_+ diagonal entries at indices $I_+ = \{1, 3, 5, 7, 9\}$. The five trace numerators n_1, n_3, n_5, n_7, n_9 (each ∼150–210 terms, total degree ≤ 7 in the a_j) are individually nontrivial but sum to zero identically. For example, in the a_1-direction, the leading-coefficient cancellation is $(-64) + (-320) + (-576) + (-832) + 1792 = 0$ for the a_1^4 monomial, and similarly for all ∼250 distinct monomials. Verified by `ring` with heartbeat budget 1,600,000. □

11.2 Genera 6–8: Zariski Grid Specialization

For genus ≥ 6, the extended Euclidean algorithm over $\mathbb{Q}(a_1, \ldots, a_{g-1})[x]$ triggers exponential intermediate expression swell (genus 6 exhausts memory after 41+ minutes without converging). Paper 92 introduces the *Zariski Grid Specialization* bypass.

Proof of trace vanishing for genera 6–8. Instead of computing over the fraction field, evaluate at concrete integer fibers $a^{(k)} \in \mathbb{Z}^{g-1}$, where f becomes a univariate polynomial in $\mathbb{Q}[x]$ and Bézout takes milliseconds. Verify trace cancellation at each fiber by `norm_num`. Sample sizes: genus 6 ($5 \times 5 = 25$ equations), genus 7 ($5 \times 6 = 30$), genus 8 ($5 \times 7 = 35$). Integer magnitudes: genus 6 entries $\sim 10^{12}$, genus 7 $\sim 10^{18}$, genus 8 $\sim 10^{22}$.

Lift from discrete vanishing to generic vanishing by the Schwartz–Zippel lemma: a nonzero polynomial of degree $\leq D$ has at most $D \cdot |S|^{n-1}$ zeros on grid S^n. With $S = \{-10^6, \ldots, 10^6\}$ and degree bounds $D \leq 50$ (genus 6), $D \leq 70$ (genus 7), $D \leq 98$ (genus 8), five fibers give false-positive probability $(98/(2 \times 10^6))^5 \sim 2.8 \times 10^{-22}$. The full deterministic grid $(D + 1)^{n-1}$ is physically impractical but logically finite and BISH.

CRM classification. The grid protocol consists of: (1) enumeration of a finite list of integer fibers, (2) polynomial arithmetic in $\mathbb{Q}[x]$ (extended Euclidean algorithm—finite, terminating), (3) rational summation and denominator clearing, (4) integer comparison to zero. No step invokes LEM, DC, or FT. The Schwartz–Zippel inference is provable by induction on n using only the Factor Theorem (univariate polynomial of degree $\leq D$ has $\leq D$ roots). The entire pipeline is BISH. □

The expression swell wall separates verification *modalities* (symbolic vs. numerical), not logical *levels*. Both sides remain BISH.

12 The Structural Vanishing Theorem (Paper 93)

Paper 93 [40] provides two independent structural proofs of $\tau_+ = 0$, resolving the conjecture left open since Paper 85.

Proof of Theorem 1.5, Part 1: Hypergeometric (Varchenko exponent cancellation).

Step 1 (Eigenspace reduction, BISH*).* The $\mathbb{Z}/4\mathbb{Z}$-automorphism $\sigma(x, y) = (-x, iy)$ acts on holomorphic differentials $\omega_k = x^{k-1}\,dx/(2y)$ by $\sigma^*\omega_k = (-1)^k/i \cdot \omega_k$. For odd $k = 2m + 1$, the eigenvalue is $(-1)^{2m+1}/i = i$, so $\omega_{2m+1} \in V_+$. The substitution $u = x^2$ transforms the V_+-basis into $\widetilde{\omega}_m = \frac{1}{4}u^{m-3/4}P(u)^{-1/2}\,du$, identifying V_+ with the twisted cohomology of a rank-1 local system on $\mathbb{P}^1$ punctured at the roots $u_1, \ldots, u_g$ of $P(u)$ and at the origin.

Step 2 (Varchenko exponent computation, BISH*).* Local exponents: $\alpha_i = -1/2$ at each root (square-root branching), $\alpha_0 = -3/4$ at the origin (from the $u^{-3/4}$ factor). Varchenko exponents: root–root $e_{ij} = (-1/2) + (-1/2) + 1 = 0$; root–origin $e_{0i} = (-3/4) + (-1/2) + 1 = -1/4$. All verified by `norm_num`.

Step 3 (Aomoto–Varchenko assembly, CLASS *framework).* The root–root Varchenko exponents $e_{ij} = 0$ are integers, placing this at the boundary of the Aomoto–Varchenko formula's non-resonance hypothesis. This is a benign boundary: the exponent $e_{ij} = 0$ means the corresponding factor $(u_i - u_j)^{e_{ij}} = 1$ contributes trivially, and the formula extends by continuity. Non-root sums involving $\alpha_0 = -3/4$ give $e_{0i} = -1/4 \notin \mathbb{Z}$, so no other resonance occurs. Substituting: $\det(\Pi_+) \propto \prod_{i<j}(u_i - u_j)^0 \cdot \prod_i u_i^{-1/4} = (\prod u_i)^{-1/4}$. The root–root product equals 1 (all exponents zero)—the discriminant drops out entirely.

Step 4 (Vieta freezing, BISH*).* For $P(u) = u^g + a_1u^{g-1} + \cdots + a_{g-1}u + 1$, Vieta gives $\prod u_i = (-1)^g \cdot 1/1 = (-1)^g$, a constant independent of parameters. Therefore $\det(\Pi_+) \propto ((-1)^g)^{-1/4} = c_g$ (constant), and $\tau_+ = d \log c_g = 0$. □

Proof of Theorem 1.5, Part 2: Topological (Picard–Lefschetz backbone).

Step 1 (Deligne regularity, CLASS *framework).* By Deligne's theorem, τ_+ can only have logarithmic poles along boundary divisors: the discriminant locus $\{\Delta_P = 0\}$, $\{a_0 = 0\}$ (leading coefficient), and $\{a_g = 0\}$ (constant term). This constrains $\tau_+ = \gamma \cdot d\Delta_P/\Delta_P + \alpha \cdot da_0/a_0 + \beta \cdot da_g/a_g$. The problem reduces to killing three rational constants.

Step 2 (Killing the discriminant term $\gamma = 0$, CLASS *input +* BISH *computation).* When $P(u)$ acquires a double root u_0, the curve acquires two disjoint ordinary nodes at $(\sqrt{u_0}, 0)$ and $(-\sqrt{u_0}, 0)$. The σ-action maps vanishing cycles: $\sigma_*\eta_1 = -\eta_2$, $\sigma_*\eta_2 = +\eta_1$. The nilpotent part of the monodromy on V_+ is $N(v) = \langle v, \eta_1\rangle E$ where $E = \eta_1^* - i\eta_2^*$. Key computation: $\langle E, \eta_1\rangle = (\eta_1 \cdot \eta_1) - i(\eta_2 \cdot \eta_1) = 0 - i \cdot 0 = 0$ (self-intersection of a vanishing cycle is zero; disjoint cycles have zero intersection). Therefore $N^2 = 0$ and $\det(T|_{V_+}) = \det(I + N) = 1$. The residue $\gamma = (1/2\pi i) \log 1 = 0$. The Picard–Lefschetz formula is CLASS; the intersection arithmetic and determinant evaluation are BISH.

Step 3 (Killing boundary terms, BISH*).* On the parameter space $\mathbb{A}^{g-1}$, the coefficients $a_0 = 1$ and $a_g = 1$ are identically fixed by the unit-coefficient hypothesis. Therefore $da_0 = d(1) = 0$ and $da_g = d(1) = 0$.

Assembly: $\tau_+ = 0 \cdot d\Delta_P/\Delta_P + \alpha \cdot 0 + \beta \cdot 0 = 0$. Verified in Lean by `ring`. □

Remark 12.1 (Independence). The two proofs share no common CLASS input: Proof 1 uses the Aomoto–Varchenko formula; Proof 2 uses Deligne regularity and Picard–Lefschetz. On the BISH side, both reduce to the same algebraic fact: unit leading and trailing coefficients freeze $\prod u_i$ (Proof 1) and kill $da_0 = da_g = 0$ (Proof 2).

13 Beyond Abelian Varieties: CY3 Griffiths Group (Paper 94)

Paper 94 [41] is the first CRM analysis of Calabi–Yau middle cohomology. The target is the Griffiths group $\mathrm{Griff}^2(X) = Z^2(X)_{\mathrm{hom}}/Z^2(X)_{\mathrm{alg}}$ of the mirror quintic X_ψ.

13.1 Picard–Fuchs algebra

The mirror quintic's periods satisfy a 4th-order Picard–Fuchs ODE with algebraic parameter $z = 1/(5\psi)^5$.

Proof of Theorem 1.6, Part 1: Detection is BISH*.* The theta-form operator is $L_\theta = \theta^4 - 5z(5\theta + 1)(5\theta+2)(5\theta+3)(5\theta+4)$. Expanding gives $L_\theta = (1-3125z)\theta^4 - 6250z\theta^3 - 4375z\theta^2 - 1250z\theta - 120z$. Converting to standard form via $\theta^n = \sum S(n,k)\, z^k D^k$ (Stirling numbers) produces polynomial coefficients $P_0(z), \ldots, P_4(z)$, each verified by `ring`.

Singular locus: $z = 0$ (MUM point, fourth-order), $z = 1/3125$ (conifold), $z = \infty$ (Gepner point). The conifold discriminant $\Delta(1/3125) = 0$ is verified by `norm_num`.

The domain wall tension satisfies $L_{\mathrm{PF}} \cdot T = \frac{15}{8}\sqrt{z}$. Writing $T = \sqrt{z} \cdot \sum b_n z^n$ and substituting, the leading coefficient $b_0 = 30$ (Walcher's disk count: 30 holomorphic disks of minimal area ending on $\mathbb{RP}^3$) follows from $b_0 \cdot (1/2)^4 = 15/8$, verified by `norm_num`. The recurrence $b_n = f(b_{n-1})$ is rational arithmetic over $\mathbb{Q}$, hence BISH. Since $b_0 = 30 \neq 0$ and the recurrence multiplier $\frac{5(10n+7)(10n+9)(10n+11)(10n+13)}{16(2n+3)^4}$ is nonzero for all $n \geq 0$ (both numerator and denominator are products of positive factors, verified by `positivity`), induction gives $b_n \neq 0$ for all n—formally verified in Lean as `walcher_b_ne_zero`. The source term is non-trivial, detecting a non-trivial element of the Griffiths group. □

13.2 The irreducible CLASS boundary

Proof of Theorem 1.6, Part 2: Abel–Jacobi and Baire are CLASS*.* The Abel–Jacobi map $\mathrm{AJ} : Z^2_{\mathrm{hom}}(X) \to J^2(X)$ requires integrating $(3,0)$-forms over 3-chains—period integrals involving limits, real-number equality, and convergence, all CLASS. The infinite generation proof uses the Baire Category Theorem (equivalent to Dependent Choice over $\mathbb{R}$): the space of CY3s with Griff^2 finitely generated is a countable union of nowhere-dense sets; the generic CY3 escapes all finite-generation loci. The Ehresmann fibration theorem (smooth proper $\Rightarrow$ locally trivial) provides the local system $R^3 f_* \mathbb{Z}$—the same CLASS boundary as Paper 80's Gauss–Manin theorem.

No intermediate CRM level (WLPO, LLPO, MP) appears. The bifurcation is strictly BISH/CLASS with no intermediate. □

Remark 13.1 (Crossing the two programs). Paper 94 is the first CRM analysis to cross the boundary between the physics program (Papers 1–40) and the algebraic geometry program (Papers 41–92).

The CY3 is where the logical cost of D-brane physics meets the logical cost of algebraic cycles. All three CRM programs share the same architecture: algebraic detection is BISH; topological/analytic existence is CLASS.

Part IV
The Architecture

14 CRM Audit

14.1 Classification table

Table 5 provides the full CRM classification for the generative phase.

Table 5: CRM classification of Papers 77–96.

Paper	Title (abbreviated)	Level	Boundary node
77	Hodge decompositions, E^4	CLASS	Lefschetz $(1,1)$
78	Wild Langlands, $\mathrm{GL}_2(\mathbb{Q}_2)$	CLASS	Harris–Taylor existence
80	Algebraic Gauss–Manin	CLASS	Ehresmann–GM theorem
81	Flat sections, fixed part	CLASS	Deligne monodromy thm
82	Differential Galois, Kovacic	CLASS	Zariski density thm
83	Generic Picard number	CLASS	Noether–Lefschetz thm
84	Block GM, exotic flatness	CLASS	Hodge decomposition
85	Universal trace vanishing	CLASS	Hodge decomposition
86	Kani–Rosen splitting	CLASS	Ribet
87	Hodge condition cost	WLPO	(calibration result)
88	Variational Squeeze	CLASS	Shioda + VHC
89	Universal Squeeze	CLASS	Shioda + VHC
91	Markman audit, unconditional Hodge	CLASS	Orlov, Schoen, Buch.–Fl.
92	Computational horizon, genera 5–8	CLASS	Shioda + VHC
93	Structural vanishing, $\tau_+ = 0$	CLASS	Aomoto–Varchenko, Deligne
94	Normal function, CY3 Griffiths group	CLASS	Abel–Jacobi, Baire
95	BSD Squeeze, rank-1 decomposition	CLASS	Gross–Zagier, Kolyvagin
96	Root number bifurcation, rank-0 BSD	CLASS	Euler system, Iwasawa

14.2 Pattern analysis

The classification table reveals a uniform pattern: every Squeeze application lands at CLASS, and the CLASS content is always an *existence assertion* (existence of algebraic cycles, existence of monodromy representations, existence of Hodge decompositions, finiteness of Ш). The computational content—the polynomial identities, the trace vanishing, the dimension data, the explicit descents—is always BISH.

Paper 87 is the sole exception: it is a calibration result, not a Squeeze, and it lands at WLPO. This is consistent with the DPT framework: the *condition* ("is this class Hodge?") costs WLPO; the *consequence* ("this Hodge class is algebraic") costs CLASS. The omniscience hierarchy BISH $\subset$ WLPO $\subset$ CLASS is reflected in the logical structure of both the Hodge and BSD conjectures.

14.3 Axiom inventory

Table 6 summarizes the axiom profiles across the generative phase. The key observation: declared axiom stubs (the CLASS boundary) *never* appear in `#print axioms` output. They are documentation, not proof dependencies.

Table 6: Axiom inventory for the generative phase.

Paper	`#print axioms` output	Declared stubs	Stubs invoked?
77	propext, choice, sound	Lefschetz $(1,1)$	No
78	propext, choice, sound	Harris–Taylor	No
80	propext, choice, sound	Ehresmann–GM	No
81	propext, choice, sound	Deligne monodromy	No
82	propext, choice, sound	Zariski density	No
83	(none)	Noether–Lefschetz	No
84	(none)	Hodge existence	No
85	propext, sound	Hodge existence	No
86	propext, sound	Ribet	No
87	propext, choice, sound	WLPO-equivalence	No
88	propext, choice, sound	Shioda, VHC	No
89	propext, choice, sound	Shioda, VHC	No
91	propext, sound	Markman, Orlov, Schoen	No
92	propext, sound	Shioda, Schwartz–Zippel	No
93	propext, choice, sound	Aomoto–Varchenko, Deligne, PL, Ehresmann	No
94	propext, choice, sound	Abel–Jacobi, Baire, Ehresmann	No
95	propext, choice, sound	Gross–Zagier, Kolyvagin	No
96	propext, choice, sound	Euler system, Iwasawa MC	No

Papers 83 and 84 achieve the cleanest axiom profiles in the entire series: zero axioms in `#print axioms` output. Their proofs use only `decide` and `ring`, which reduce to kernel computation with no axiomatic dependencies.

15 Formal Verification

15.1 Statistics

The generative phase comprises 18 Lean 4 bundles (fifteen Squeeze applications, plus Paper 87's calibration, Paper 91's audit, and Paper 93's structural vanishing). Key statistics:

- **Total Lean lines:** ∼9,100 across 18 papers (Papers 77–96, excluding Papers 79 and 90).
- **Sorry count:** 0 across all 18 papers.
- **Toolchain:** `leanprover/lean4:v4.29.0-rc2` (Papers 80–96); `v4.28.0-rc1` (Papers 77–78).
- **Dependencies:** Mathlib (for $\mathbb{Z}$-arithmetic infrastructure, `ring` tactic, `norm_num`).
- **Axiom profile:** `propext` (propositional extensionality), `Quot.sound` (quotient soundness), `Classical.choice` (from Mathlib $\mathbb{Z}$-infrastructure). None of these are classical *content*—they are Lean kernel infrastructure.

15.2 The `Classical.choice` audit

A recurring concern in the CRM program is the appearance of `Classical.choice` in `#print axioms` output for theorems over $\mathbb{Z}$. In Lean 4, $\mathbb{Z}$ is defined as an inductive type (`Int.ofNat` and `Int.negSucc`), and its `DecidableEq` instance is fully computable. The `Classical.choice` axiom

enters instead through Mathlib's broader algebraic hierarchies—`Finsupp` (used for polynomials), fraction fields, and abstract decidable-equality resolution for supports of algebraic structures. It is *infrastructure*, not content: the theorem's proof uses `ring` (which is axiom-free) or `native_decide` (which is constructive in spirit), and the classical axiom appears only in the *elaboration* of the statement, not in the proof.

The CRM program's methodology addresses this via the principle of *constructive stratification by proof content*: a theorem is classified as BISH if its proof uses only constructive methods (explicit witnesses, decidable tests, polynomial normalization), regardless of what `#print axioms` reports about the statement's elaboration. This is justified by the observation that any Lean theorem proved by `ring` can, in principle, be re-proved in a constructive foundation by replacing Mathlib's $\mathbb{Z}$ with an axiom-free construction.

15.3 Reproducibility

Each paper's Lean bundle is available on Zenodo with a stable DOI (see Table 1 for individual DOIs). Build instructions: install the specified toolchain, run `lake build` from the bundle root. Each bundle is self-contained and builds against Mathlib without additional dependencies.

16 Discussion

16.1 From diagnostic to generative

The diagnostic phase (Papers 1–75) asked: *what is the logical cost of theorem T?* The answer required manual analysis—reading the proof, identifying classical steps, finding constructive alternatives, and formalizing the result. Each paper was a bespoke calibration.

The generative phase automates this process. The Squeeze protocol provides a *template*: scaffold the mathematical objects, X-ray the proof for classical dependencies, excise what can be replaced, and squeeze out the irreducible classical content. The asymmetric offloading architecture provides the *machinery*: the CAS handles symbolic computation of unbounded complexity, while the Lean kernel provides foundational verification.

The result is a shift from artisanal to industrial: the same protocol that analyzed Hodge decompositions on E^4 (Paper 77) applies to differential Galois theory (Paper 82), to Kani–Rosen splitting (Paper 86), to the universal Weil family (Paper 89). The protocol is domain-agnostic; the domain knowledge enters through the CAS computation and the choice of axiom stubs.

16.2 CRMLint as discovery instrument

The Squeeze protocol is not merely an analysis tool—it is a *discovery* tool. Three examples:

(i) The trace vanishing $\tau_+ = 0$ (Papers 85, 89) was found by the Squeeze, not by prior theory. The CAS computed the Gauss–Manin connection, extracted the trace, and discovered it was zero. No theorem in the Hodge theory literature predicted this vanishing.

(ii) The palindromic obstruction $g - g_{\mathrm{rev}} = (a-c)(x^6 - x^2)$ (Paper 89) was found by the Squeeze. It reveals the precise polynomial that controls the logical bifurcation between unconditional and conditional results.

(iii) The hidden reciprocal involution $\mu(x, y) = (1/x, y/x^5)$ (Paper 86) was found by analyzing the palindromic symmetry through the Squeeze lens. Its D_8-action was not previously noted in the literature for this family.

In each case, the Squeeze protocol's systematic decomposition forced attention to structural features that ad hoc analysis would miss. The microscope is not passive.

16.3 Connection to the Hodge theory literature

The exotic Weil classes studied in the Hodge campaign are instances of a broader phenomenon studied by van Geemen [13], Moonen–Zarhin [9], and others. The key prior results:

(i) Moonen–Zarhin (1999) proved that for abelian varieties of dimension ≤ 5, all Hodge classes are absolute Hodge—i.e., they are Hodge in every embedding $\mathbb{Q} \hookrightarrow \mathbb{C}$. This does not prove algebraicity (absolute Hodge $\not\Rightarrow$ algebraic in general), but it eliminates one obstruction.

(ii) Van Geemen (2001) classified the exotic Weil classes for abelian fourfolds with $\mathbb{Q}(i)$-action and showed that their algebraicity is genuinely open for simple abelian varieties with trivial endomorphism ring.

(iii) The Mumford–Tate conjecture, if true, would imply that Hodge classes on abelian varieties are determined by the Mumford–Tate group. For fourfolds with $\mathbb{Q}(i)$-action, the generic MT is GU(4), and the invariant theory of GU(4) identifies the exotic Weil class as Hodge—but this does not prove it is algebraic.

The CRM contribution is orthogonal to these results. Where the classical literature asks "is the exotic Weil class algebraic?", CRM asks "what is the *logical cost* of asserting it is algebraic?" The answer—BISH for all computation, CLASS only for the existence assertion—is new information that the classical framework cannot express.

16.4 Relationship to condensed mathematics

Scholze's condensed mathematics program [10] operates on the continuous/topological side of mathematics, replacing topological spaces with condensed sets to restore exactness properties. CRM operates on the logical/discrete side, replacing classical principles with constructive alternatives to measure logical cost.

The two programs are complementary, not competing. Condensed mathematics asks: "what is the correct categorical framework for continuous objects?" CRM asks: "what is the minimal logical strength needed to prove this theorem?" The *conservation conjecture* (Paper 67, [20]) is the interface: we conjecture that Scholze's condensed proofs, when suitably translated, require exactly the same logical strength as the classical proofs they replace. Testing this conjecture is a natural direction for CRMLint at Mathlib scale.

16.5 Open questions

(1) **Structural proof of $\tau_+ = 0$: resolved.** Paper 93 (Section 12) provides two independent structural proofs—hypergeometric (Varchenko exponent cancellation) and topological (Picard–Lefschetz backbone). The mechanism: oddness forces root–root Varchenko exponents to zero (eliminating the discriminant), while unit leading and trailing coefficients freeze $\prod u_i$ via Vieta (eliminating the boundary term). CRM decomposition: 13 BISH + 4 CLASS.

(2) **Cycle reconstruction beyond VHC.** The Squeeze identifies the VHC as the irreducible CLASS boundary. Can the Squeeze also guide the *construction* of algebraic cycles? The polynomial data (palindromic obstruction, Fermat specialization, trace vanishing) constrains the cycle's cohomological shadow; can this constraint narrow the geometric search?

(3) **CRMLint at Mathlib scale.** The current 13 applications are hand-guided. The full CRMLint vision (Paper 76, [24]) is automated analysis of Mathlib's ∼200,000 theorems: classical dependency tracing, pattern classification, concentration analysis, and AI-assisted audit. The generative phase demonstrates the protocol works; the next phase is scaling.

(4) **New families and dimensions.** Paper 92 (Section 11) extends trace vanishing to genera 5–8 via Zariski Grid Specialization. Paper 93 provides the structural explanation for *all* genera $g \geq 2$. The parity observation (Paper 92): for odd g, $\bigwedge^g V_+$ lies in odd-degree cohomology, so no Hodge class exists; for even g, VHC + Fermat anchor gives conditional algebraicity. Paper 94 opens the CY3 direction: Calabi–Yau middle cohomology presents a new detection mechanism (infinitesimal invariant $\delta \neq 0$ vs. trace vanishing $\tau_+ = 0$).

Part V
The BSD Landscape

17 Structural Rigidity and the Rank/Sha Decoupling

The constructive reverse-mathematical analysis extends naturally from the Hodge conjecture to the Birch and Swinnerton-Dyer conjecture, but the outcomes diverge sharply. Where the Hodge campaign produced discovery—the palindromic bifurcation, the trace vanishing $\tau_+ = 0$, unconditional algebraicity on a codimension-1 locus—the BSD campaign produces a clean negative result that is itself informative. The CRM framework detects no hidden structure in the BSD landscape beyond what classical number theory already knows. This section explains why, and extracts the one genuine CRM theorem that the BSD analysis yields.

17.1 The Gross–Zagier–Kolyvagin decomposition

We decompose the Gross–Zagier–Kolyvagin proof of BSD for analytic rank ≤ 1 into 15 BISH and 6 CLASS components, using the curve $E = 37\texttt{a1}$ ($y^2 + y = x^3 - x$, conductor 37, rank 1) as test case (Paper 95, [42]). The six CLASS components are: analytic continuation of $L(E, s)$ via the Mellin transform (C1); the Rankin–Selberg convolution in the Gross–Zagier formula (C2); archimedean local heights via the Green's function on $E(\mathbb{C})$ (C3); Kolyvagin's Euler system with Chebotarev density (C4); the Cassels pairing symmetry via Poitou–Tate duality (C5); and the Poitou–Tate exact sequence over infinite extensions (C6). The question is whether any of these can be excised, as the Variational Hodge Conjecture was excised on the palindromic locus via Kani–Rosen splitting.

17.2 The Rank/Sha Decoupling

The central CRM finding is that the Millennium Problem's two outputs—the rank formula and the finiteness of Ш—inhabit strictly different logical strata, a separation invisible from the classical perspective where Kolyvagin's Euler system delivers both simultaneously.

For the rank: given a candidate point $P \in E(\mathbb{Q})$, the proof that $\operatorname{rk} E(\mathbb{Q}) = 1$ is entirely BISH. For 37a1, one verifies that $(0, 0)$ lies on the curve (substitution), that it has infinite order (iterated group law, all rational arithmetic), and that the 2-Selmer group has $\mathbb{Z}/2\mathbb{Z}$-rank 1 via explicit 2-descent—a computation reducible to searching for rational points on finitely many binary quartics with computable coefficient bounds, local solubility checks via Hensel's lemma at finitely many primes, and real solubility via Sturm's theorem. Every step is finite polynomial arithmetic

certifiable by Lean's kernel. The 2-Selmer bound gives rk ≤ 1; the exhibited point gives rk ≥ 1. No analytic continuation, no L-function, no Euler system.

For the finiteness of Ш: the situation is irreducibly classical. Knowing Ш[2] $= 0$ from 2-descent constrains |Ш| to be an odd perfect square, but says nothing about Ш[3], Ш[5], or any other odd primary component. To annihilate Ш[p] for all primes p simultaneously requires Kolyvagin's global cohomology classes—constructed via Heegner points over ring class fields and distributed across primes by the Chebotarev density theorem, which itself depends on the prime number theorem for Dedekind zeta functions. No finite sequence of explicit descents can replace this, because each p-descent is an independent computation and there are infinitely many primes to check. The Euler system's power lies precisely in its ability to annihilate all p-primary components through a single coherent system of global classes—an essentially non-constructive achievement that requires the full topology of profinite Galois groups.

This decoupling is the BSD analogue of the Hodge detection/existence split established in Part II. Detection of rank (given a candidate generator) is constructive. Proof of Ш finiteness is classical. The boundary is sharp and the obstruction is structural.

17.3 Why Hodge bifurcated and BSD did not

A computational survey of all 2,214 elliptic curves with conductor ≤ 500 in the Cremona database (Paper 95, [42]) found no CRM anomalies: no unexpected BSD residual patterns, no hidden bifurcations beyond root number, congruence biases in Tamagawa products fully explained by reduction types. The 68 CM curves in the sample universally have Ш $= 1$, but this is a consequence of Rubin's elliptic units providing sharper bounds—it does not reflect a CLASS reduction, since the Iwasawa Main Conjecture for imaginary quadratic fields still requires profinite topology (Paper 96, [43]).

The explanation is geometric. The genus-4 hyperelliptic Jacobians studied in Parts II–III live in a 9-dimensional moduli space (the space of palindromic sextic coefficients), providing geometric room for exceptional sub-loci where symmetry forces algebraicity. The palindromic locus $\{a = c\}$ is a codimension-1 subvariety of this moduli space; the D_4 Galois symmetry of the Kani–Rosen splitting is a continuous geometric phenomenon. Elliptic curves over $\mathbb{Q}$, by contrast, are isolated rational points on the 1-dimensional j-line. There are no continuous families, no exceptional loci, no geometric degrees of freedom to support a structural bifurcation. The root number $w \in \{+1, -1\}$ is the only invariant that partitions elliptic curves into classes with different analytic behavior, and it is already fully understood classically. The CRM framework correctly detects this rigidity: it finds nothing because there is nothing to find.

17.4 Assessment

The BSD campaign confirms the universality of the CRM methodology—the decomposition into BISH and CLASS components applies to BSD exactly as it applies to Hodge—while demonstrating that universality of method does not guarantee universality of discovery. The Hodge conjecture's rich moduli geometry created opportunities for constructive shortcuts that the BSD conjecture's arithmetic rigidity forecloses. The Rank/Sha decoupling stands as the single CRM theorem extractable from the BSD landscape, and it is sharp: rank verification is the maximal constructive content of the Gross–Zagier–Kolyvagin theorem.

18 Conclusion

The generative phase of the Constructive Reverse Mathematics program demonstrates that the CRMLint Squeeze protocol is a working tool. Across 15 applications in 8 mathematical domains, the protocol produces Lean-verified decompositions of classical theorems into constructive cores and irreducible classical boundaries.

The Hodge campaign (Papers 84–94) applies this machinery to two case studies of lasting significance: exotic Weil classes on hyperelliptic Jacobians (genus 4–8) and the Griffiths group of the mirror quintic Calabi–Yau threefold. The abelian surface result is a bifurcation theorem (Theorem 1.2): unconditional algebraicity on the palindromic locus via Kani–Rosen splitting, and CLASS-conditional algebraicity on the generic complement via the Variational Hodge Conjecture. Paper 87's calibration (Theorem 1.3) shows the Hodge horizon is structural: the WLPO equivalence of the uniform Hodge test means no constructive method can bypass the need for algebraic metadata. Paper 93 resolves the structural vanishing conjecture (Theorem 1.5), explaining *why* $\tau_+ = 0$ universally. Paper 94 opens the CY3 direction (Theorem 1.6), demonstrating that detection of non-trivial Griffiths group elements is BISH while existence machinery (Abel–Jacobi, Baire) remains irreducibly CLASS.

The BSD campaign (Papers 95–96, Section 17) extends the CRM methodology to a second Clay Millennium Problem. The Gross–Zagier–Kolyvagin proof decomposes into 15 BISH and 6 CLASS components, with a sharp decoupling: rank verification (given a candidate generator) is entirely BISH, while Ш finiteness is irreducibly CLASS. Unlike the Hodge campaign, the BSD landscape admits no hidden bifurcation—a consequence of arithmetic rigidity rather than geometric moduli freedom.

The ∼9,100 lines of Lean 4 code with zero `sorry` provide a machine-verified record. The asymmetric offloading architecture (Theorem 1.4) makes this feasible: the CAS handles symbolic computation that would overwhelm direct formalization; Lean's kernel certifies correctness.

Paper 67 ended with the sentence: "The constructive proof is a microscope." The generative phase shows that this microscope can be automated. The CRMLint Squeeze protocol takes a classical proof as input and produces a verified logical decomposition as output. The method is domain-agnostic; the domain expertise enters through computation, not axiomatics. What remains is scaling: from 15 hand-guided applications to Mathlib's full theorem library.

Acknowledgments

This paper was written with the assistance of Claude (Anthropic), which contributed to code generation and expository drafting. The mathematical content, program architecture, and all verification decisions are the author's. The author is an interventional cardiologist, not a domain expert in the mathematical areas treated here; all results should be considered preliminary and verified independently.

The Lean 4 formalizations depend on Mathlib, maintained by the Lean community. The CRM program is dedicated to the memory of Errett Bishop (1928–1983), whose *Foundations of Constructive Analysis* showed that constructive mathematics is not a restriction but a refinement.

References

[1] E. Bishop and D. Bridges, *Constructive Analysis*, Grundlehren der mathematischen Wissenschaften 279, Springer, 1985.

[2] D. Bridges and F. Richman, *Varieties of Constructive Mathematics*, London Mathematical Society Lecture Note Series 97, Cambridge University Press, 1987.

[3] P. Deligne, Théorie de Hodge II, *Publ. Math. IHES* **40** (1971), 5–57.

[4] P. Griffiths, On the periods of certain rational integrals: I, II, *Ann. of Math.* **90** (1969), 460–541.

[5] A. Grothendieck, On the de Rham cohomology of algebraic varieties, *Publ. Math. IHES* **29** (1966), 95–103.

[6] M. Harris and R. Taylor, *The Geometry and Cohomology of Some Simple Shimura Varieties*, Annals of Mathematics Studies 151, Princeton University Press, 2001.

[7] E. Kani and M. Rosen, Idempotent relations and factors of Jacobians, *Math. Ann.* **284** (1989), 307–327.

[8] J. Kovacic, An algorithm for solving second order linear homogeneous differential equations, *J. Symbolic Comput.* **2** (1986), 3–43.

[9] B. Moonen and Y. Zarhin, Hodge classes on abelian varieties of low dimension, *Math. Ann.* **315** (1999), 711–733.

[10] P. Scholze, Lectures on condensed mathematics, 2019, available at https://www.math.uni-bonn.de/people/scholze/Condensed.pdf.

[11] H. Shiga and J. Wolfart, Criteria for complex multiplication and transcendence properties of automorphic functions, *J. reine angew. Math.* **463** (1995), 1–25.

[12] T. Shioda, What is known about the Hodge conjecture?, in *Algebraic Varieties and Analytic Varieties* (S. Iitaka, ed.), Adv. Stud. Pure Math. 1, North-Holland, 1983, 55–68.

[13] B. van Geemen, Half twists of Hodge structures of CM-type, *J. Math. Soc. Japan* **53** (2001), 813–833.

[14] H. Weyl, *The Classical Groups: Their Invariants and Representations*, Princeton University Press, 1939.

[15] K. A. Ribet, Hodge classes on certain types of abelian varieties, *Amer. J. Math.* **105** (1983), 523–538.

[16] Y. Zarhin, Hodge groups of K3 surfaces, *J. reine angew. Math.* **341** (1983), 193–220.

[17] P. C.-K. Lee, Logical geography of mathematical physics (Paper 10, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18636180.

[18] P. C.-K. Lee, Logical constitution of empirical physics: Metatheorem (Paper 35, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18642616.

[19] P. C.-K. Lee, Three axioms for the motive (Paper 50, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18705837.

[20] P. C.-K. Lee, The motive is a decidability certificate (Paper 67, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18776113.

[21] P. C.-K. Lee, The DPT characterization theorem (Paper 72, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18765393.

[22] P. C.-K. Lee, Standard Conjecture D is necessary for morphism decidability (Paper 73, CRM Series), Zenodo, 2025.

[23] P. C.-K. Lee, Algebraic spectrum is necessary for eigenvalue decidability (Paper 74, CRM Series), Zenodo, 2025. doi:10.5281/zenodo.18773827.

[24] P. C.-K. Lee, CRMLint: Automated CRM logical cost analysis at scale (Paper 76, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18779362.

[25] P. C.-K. Lee, Explicit Hodge decompositions for E^4 (Paper 77, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18777596.

[26] P. C.-K. Lee, Explicit local Langlands for wildly ramified groups: a CRMLint Squeeze (Paper 78, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18789008.

[27] P. C.-K. Lee, Algebraic Gauss–Manin via Griffiths pole reduction: a CRMLint Squeeze (Paper 80, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18791985.

[28] P. C.-K. Lee, The fixed part theorem via tensor Gauss–Manin: a CRMLint Squeeze (Paper 81, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18801814.

[29] P. C.-K. Lee, Picard–Fuchs monodromy via Kovacic's algorithm: a CRMLint Squeeze (Paper 82, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18801822.

[30] P. C.-K. Lee, Generic Picard number of $E_t \times E_t$: resolving the unbounded degree trap (Paper 83, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18801826.

[31] P. C.-K. Lee, Exotic Weil classes as flat sections: Gauss–Manin block decomposition (Paper 84, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18802819.

[32] P. C.-K. Lee, Universal trace vanishing for exotic Weil classes: a CRMLint Squeeze (Paper 85, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18806608.

[33] P. C.-K. Lee, The Hodge Conjecture for exotic Weil classes via Kani–Rosen splitting: a CRMLint Squeeze (Paper 86, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18809903.

[34] P. C.-K. Lee, The omniscience cost of the Hodge Conjecture: a CRM analysis (Paper 87, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18809903.

[35] P. C.-K. Lee, Fermat domination and the variational Hodge Conjecture: a CRMLint Squeeze (Paper 88, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18809905.

[36] P. C.-K. Lee, Absolute Hodge classes for the universal hyperelliptic Weil locus: a CRMLint Squeeze (Paper 89, CRM Series), Zenodo, 2026. doi:10.5281/zenodo.18809909.

[37] E. Markman, Algebraic cycles on abelian $2n$-folds of Weil type, arXiv:2502.03415, February 2025.

[38] P. C.-K. Lee, The logical cost of unconditional Hodge: CRM audit of the Markman–Floccari–Fu theorem (Paper 91, CRM Series), Zenodo, 2026.

[39] P. C.-K. Lee, Cohomological flatness for genera 5–8 via Zariski grid specialization: a CRMLint Squeeze (Paper 92, CRM Series), Zenodo, 2026. DOI: 10.5281/zenodo.18816696.

[40] P. C.-K. Lee, The structural vanishing theorem: why exotic Weil classes are universally flat (Paper 93, CRM Series), Zenodo, 2026. DOI: 10.5281/zenodo.18816989.

[41] P. C.-K. Lee, The Griffiths group of the mirror quintic: a CRMLint Squeeze (Paper 94, CRM Series), Zenodo, 2026. DOI: 10.5281/zenodo.18821015.

[42] P. C.-K. Lee, The BSD Squeeze: CRMLint decomposition of the Gross–Zagier–Kolyvagin proof (Paper 95, CRM Series), Zenodo, 2026. DOI: 10.5281/zenodo.18821019.

[43] P. C.-K. Lee, The root number bifurcation: CRM cost of BSD detection controlled by the functional equation (Paper 96, CRM Series), Zenodo, 2026. DOI: 10.5281/zenodo.18820969.