The Motivic CRM Architecture: Why the Six Domains of Mathematics Have Different Logical Signatures, and How Motives Explain the Pattern

Paper 98 of the Constructive Reverse Mathematics Series

A Capstone Synthesis of Papers 50–97

Paul Chun-Kit Lee
New York University, Brooklyn, New York
dr.paul.c.lee@gmail.com

March 2026

DOI: 10.5281/zenodo.18828345

Abstract

We formalize the observation that the six mathematical domains connected by the Langlands program—number theory, algebraic geometry, representation theory, harmonic analysis, topology, and mathematical physics—exhibit systematically different constructive reverse mathematics (CRM) signatures. We show that Grothendieck's theory of motives provides the structural explanation: the motive itself is BISH (Bishop-constructive), each realization functor has a definite CRM cost, and the comparison isomorphisms between realizations are CLASS (fully classical) if and only if they pass through the Archimedean place. This *Archimedean Obstruction Thesis* transcends mere redescription by generating precise structural predictions—most notably, that function field arithmetic (lacking an Archimedean place) must be uniformly constructive. It further explains why algebra is constructive, analysis is classical, algebraic geometry bifurcates, and the Taylor–Wiles repair of Fermat's Last Theorem was a CLASS $\to$ WKL_0 excision. We state three calibration theorems (Hodge detection $\leftrightarrow$ WLPO; TW patching $\leftrightarrow$ WKL_0; BSD rank/III decoupling) and a signature preservation theorem for the Langlands correspondence. A companion 0-`sorry` Lean 4 formalization mechanically verifies the core logical framework, representing the geometric functors as axiomatic stubs. This paper synthesizes the findings of Papers 50–97 in the CRM program. A companion monograph (*The Logical Cost of Everything*) provides a book-length accessible exposition of the full program.

Contents

1 Introduction **4**

2 The CRM Hierarchy **5**

3 Motives: The Universal Constructive Core **6**

4 Realizations and Their CRM Cost **6**

4.1 Betti Realization (CLASS) 6

4.2 de Rham Realization (BISH) 7

4.3 Étale Realization (BISH, with a caveat) 7

4.4 Crystalline Realization (BISH) . . . 7
4.5 Hodge Realization (CLASS) . . . 7
4.6 Automorphic Realization (CLASS) . . . 8
4.7 Summary Table . . . 8

5 Comparison Isomorphisms: The Archimedean Obstruction **8**
5.1 Betti $\leftrightarrow$ de Rham: The Period Map (CLASS) . . . 8
5.2 Crystalline $\leftrightarrow$ de Rham: p-adic Hodge Theory (BISH) . . . 8
5.3 Étale $\leftrightarrow$ Betti: Artin Comparison (CLASS) . . . 9
5.4 de Rham $\leftrightarrow$ Hodge: The $\partial\bar{\partial}$-Lemma (CLASS) . . . 9
5.5 Crystalline $\leftrightarrow$ Étale: Fontaine–Laffaille (BISH) . . . 9
5.6 The Archimedean Obstruction Theorem . . . 9

6 Why the Six Domains Have Different Signatures **10**
6.1 Algebra and Representation Theory: BISH . . . 10
6.2 Harmonic Analysis: CLASS . . . 10
6.3 Number Theory: Sharp Classical Bottlenecks . . . 11
6.4 Algebraic Geometry: Bifurcation . . . 11
6.5 Topology: Mixed . . . 11
6.6 Mathematical Physics . . . 11

7 Calibration Theorem I: Hodge Detection $\leftrightarrow$ WLPO **11**

8 Calibration Theorem II: TW Patching $\leftrightarrow$ WKL_0 **12**
8.1 The Three Tiers . . . 12
8.2 The 1993 $\rightarrow$ 1995 Excision . . . 12

9 Calibration Theorem III: BSD Rank/Ш Decoupling **13**

10 Signature Preservation Under Langlands **14**

11 The Function Field Test **14**

12 Physics Calibrations **15**
12.1 Classical Mechanics: BISH . . . 15
12.2 Quantum Mechanics: BISH + LPO gap . . . 15
12.3 General Relativity: BISH + CLASS gap . . . 15
12.4 Lattice vs. Continuum QFT . . . 15

13 The Archimedean Thesis: A Synthesis **16**

14 CRM Audit **17**

15 Formal Verification **17**
15.1 File Structure . . . 17
15.2 Axiom Inventory . . . 17
15.3 Reproducibility . . . 18

16 Discussion **18**
16.1 Relation to the De-Omniscientising Descent . . . 18
16.2 Open Problems . . . 18
16.3 The Archimedean Stratification . . . 19

17 Conclusion **19**

A The Motivic Reinterpretation: Papers 2–97 Through the Archimedean Lens **20**
A.1 Phase 0: Mathematical Physics (Papers 2–44) . . . 20
A.2 Phase 1: Arithmetic Geometry (Papers 45–66) . . . 21
A.3 Phase 2–4: Proof Methods, Function Fields, Synthesis (Papers 67–75) . . . 22
A.4 Phase 5–7: Generative Phase (Papers 76–97) . . . 23

B Controversies Resolved **24**
B.1 Classical.choice in $\mathbb{R}$-theorems . . . 24
B.2 DPT vs. Fargues–Scholze . . . 25
B.3 FLT vs. BSD: why their CRM profiles diverge . . . 25
B.4 Hodge bifurcation vs. BSD rigidity . . . 25

C New Opportunities **26**
C.1 Constructive p-adic Langlands . . . 26
C.2 Condensed mathematics as realization functor . . . 26
C.3 Motivic Iwasawa theory . . . 26
C.4 Trace formula excision . . . 27
C.5 Higher-dimensional Hodge bifurcations . . . 27
C.6 Explicit Langlands beyond GL_2 . . . 27

D Opportunity Summary **28**

E Feasibility Assessment and Roadmap **28**
E.1 Prime targets (immediately formalizable) . . . 28
E.2 Technically bottlenecked (sound logic, missing infrastructure) . . . 28
E.3 Confirmed irreducible . . . 29
E.4 Candidate Papers 99–100 . . . 29

1 Introduction

The Langlands program connects six areas of mathematics: number theory, algebraic geometry, representation theory, harmonic analysis, topology, and mathematical physics. Each area has its own methods, its own culture, and—as this paper documents—its own *logical signature*. Number theory is algebraic at its core but hits sharp classical bottlenecks at L-functions. Algebraic geometry bifurcates: some theorems are constructive, others irreducibly classical, with the boundary controlled by moduli dimension. Representation theory (of algebraic groups, Hecke algebras) is almost entirely constructive. Harmonic analysis—spectral decomposition on locally symmetric spaces—is the principal factory of classical content. And mathematical physics recapitulates the pattern, with the quantum/classical divide mirroring the BISH/CLASS boundary.

Why? What structural feature of mathematics produces these differences, and what connects them?

The answer, we argue, is Grothendieck's theory of *motives*. The motive of a variety is its universal cohomological invariant—an algebraic object, defined by correspondences, living in a category built from finite-dimensional linear algebra over $\mathbb{Q}$. It is constructive in its syntax. But to *use* a motive, one must apply a *realization functor*: Betti (singular cohomology), de Rham (algebraic differential forms), étale (ℓ-adic Galois representations), crystalline (over $\mathbb{F}_p$), Hodge (filtration and decomposition), or automorphic (via Langlands). Each realization has a definite CRM cost. And the *comparison isomorphisms* between realizations—the period map (Betti $\leftrightarrow$ de Rham), the Artin comparison (étale $\leftrightarrow$ Betti), the Hodge decomposition (de Rham $\leftrightarrow$ Hodge)—are classical if and only if they pass through the topology of the complex manifold, i.e., through the *Archimedean place* $\mathbb{R}$.

This is the **Archimedean Obstruction Thesis**: the Archimedean place—specifically, the completeness and ordering of $\mathbb{R}$, not its ring structure—is the unique source of non-constructive content in arithmetic geometry.

Explanation vs. Redescription. One might ask if this thesis is merely a tautology: does "the classical content comes from $\mathbb{R}$" simply redescribe the truism that "analysis requires analysis"? The explanatory power of the motivic framework lies in its *falsifiable structural predictions*. If the thesis holds, then entirely removing the Archimedean place from the global arithmetic machinery must yield a strictly BISH proof architecture, even for the most profound theorems. This prediction is spectacularly confirmed by the **Function Field Langlands program** (§11). Over $\mathbb{F}_q(C)$, where $\mathbb{A}_F$ lacks an Archimedean factor, Lafforgue's proof utilizes moduli of shtukas (algebraic) rather than the trace formula (spectral analysis), remaining entirely within the constructive envelope.

The thesis yields precise logical consequences, all verified across our program:

(i) **Function field Langlands is BISH** (Papers 69, 78): over $\mathbb{F}_q(C)$, the adèle ring $\mathbb{A}_F$ has no Archimedean factor. All realizations are étale/crystalline. The trace formula uses shtukas instead of spectral theory. The entire correspondence is constructive.

(ii) **The Taylor–Wiles fix was a logical excision** (Papers 68–71, §8): Wiles's 1993 attempt computed the congruence ideal $\eta_{\mathbb{T}}$ via the Petersson inner product (integration on the complex upper half-plane: CLASS). Taylor–Wiles patching replaced this with an inverse limit argument, cleanly excising the Archimedean computation and dropping the logical cost to WKL_0.

(iii) **Hodge detection bifurcates at WLPO** (Papers 86–90, §7): on the generic fiber, detecting Hodge classes requires the Hodge decomposition (CLASS). On special loci, extra algebraic symmetries provide shortcuts (BISH). The gap is precisely WLPO.

(iv) **BSD decouples at the rank/Ш boundary** (Papers 95–96, §9): rank verification is BISH (algebraic descent), Ш finiteness is CLASS (Euler systems, Chebotarev).

(v) **Physics parallels the pattern** (§12): classical mechanics (BISH) vs. quantum spectral theory (CLASS), lattice QFT (BISH) vs. continuum QFT (CLASS), explicit GR solutions (BISH) vs. general existence theorems (CLASS).

A companion monograph, *The Logical Cost of Everything*, provides a book-length exposition of the full 98-paper program for a general mathematical audience, including worked examples, analogies, and the complete paper list. It is included in the Zenodo package alongside this paper.

2 The CRM Hierarchy

We work in the framework of constructive reverse mathematics (CRM), as developed by Ishihara, Bridges, Diener, and others.

Definition 2.1 (CRM Signature)**.** A *CRM signature* of a mathematical theorem T is the weakest level $\sigma \in \{\mathsf{BISH}, \mathsf{MP}, \mathsf{LLPO}, \mathsf{WKL}_0, \mathsf{WLPO}, \mathsf{LPO}, \mathsf{CLASS}\}$ such that T is provable in $\mathsf{BISH} + \sigma$. We write $\sigma(T) = \sigma$.

The hierarchy forms a partial order. The principal principles are strictly ordered by logical strength ($\mathsf{BISH} \subsetneq \mathsf{LLPO} \subsetneq \mathsf{WLPO} \subsetneq \mathsf{LPO} \subsetneq \mathsf{CLASS}$). We also include Weak König's Lemma (WKL_0). Note that WKL_0 (compactness) and WLPO (analytic decidability) are logically incomparable over BISH.

The key principles, for reference:

- BISH: Bishop's constructive mathematics. Intuitionistic logic + dependent choice. Every existential proof provides a computable witness.
- WKL_0: Every infinite, finitely branching tree has an infinite path. Equivalent to: every bounded sequence in $[0, 1]$ has a convergent subsequence.
- WLPO: For every real number x, either $x = 0$ or $x \neq 0$. Decidability of equality for reals.
- CLASS: Full classical logic. Law of excluded middle for all propositions.



Figure 1: The CRM hierarchy. Solid arrows: strict logical inclusion. WKL_0 (compactness) is incomparable with WLPO (analytic decidability) over BISH. Annotations indicate where each level appears in the motivic architecture. The physics ceiling at LPO reflects the maximum logical cost of empirical physics (Paper 70).

Definition 2.2 (CRM Signature of a Proof Path)**.** A *proof path* P consists of a set of realizations $\{R_i\}$ and comparison isomorphisms $\{C_j\}$. Its signature is:

$$\sigma(P) \;=\; \max\bigl(\sigma(\text{motive}),\ \max_i \sigma(R_i),\ \max_j \sigma(C_j)\bigr)$$

3 Motives: The Universal Constructive Core

Let k be a field. Grothendieck's category of pure Chow motives $\mathsf{Mot}_{\mathrm{rat}}(k)$ has:

Objects: Triples (X, p, n) where X is a smooth projective variety over k, $p \in \mathrm{CH}^{\dim X}(X \times X) \otimes \mathbb{Q}$ is an idempotent correspondence, and $n \in \mathbb{Z}$ is a Tate twist.

Morphisms: Algebraic correspondences modulo rational equivalence, with $\mathbb{Q}$-coefficients. Composition is given by intersection theory.

The essential point is that everything here is *algebraic*: correspondences are formal sums of subvarieties, composition is a finite computation in the Chow ring, and all linear algebra takes place over $\mathbb{Q}$.

Proposition 3.1. *The category $\mathsf{Mot}_{\mathrm{rat}}(k)$ is a $\mathbb{Q}$-linear, rigid, symmetric monoidal category. Its objects and formal morphisms are definable in* BISH.

Proof sketch. Equality of morphisms modulo rational equivalence is algebraically decidable in principle, as rational equivalence is explicitly generated by cycles parameterized by $\mathbb{P}^1$ (reducible to solving polynomial systems via elimination theory). All operations—tensor product, internal Hom, duality—are defined by algebraic correspondences and computed by intersection theory, which is a finite algebraic algorithm. No limits, no completions, no analysis. □

Remark 3.2 (Decidability vs. Tractability)**.** While rational equivalence is algebraically decidable *in principle* (thus strictly satisfying the logical requirements of BISH), we emphasize the vast gap between decidability-in-principle and effective computability. Deep problems like Bloch's conjecture demonstrate that bounded geometric constructions can be computationally intractable in practice. The BISH classification reflects logical finiteness, not algorithmic feasibility. (Note: Passing to numerical equivalence $\mathsf{Mot}_{\mathrm{num}}(k)$ guarantees finite-dimensional Hom-spaces via Jannsen's theorem, but testing numerical triviality requires the Standard Conjectures to be computable, necessitating LPO as shown in Paper 73. Rational equivalence cleanly avoids this classical boundary.)

Lean: Motive signature

```
def motive_signature : CRMSignature :=
  ⟨.BISH, "Algebraic cycles and intersection theory modulo rational equivalence"⟩

theorem motive_is_bish : motive_signature.level = .BISH := rfl
```

4 Realizations and Their CRM Cost

A realization functor is an exact, faithful, $\mathbb{Q}$-linear functor from $\mathsf{Mot}(k)$ to a linear category. Each carries a definite CRM cost determined by the analytic or algebraic nature of its target category.

4.1 Betti Realization (CLASS)

For $k \hookrightarrow \mathbb{C}$, the Betti realization sends M to $H^*_{\mathrm{sing}}(X(\mathbb{C}), \mathbb{Q})$. The critical obstruction is forming the complex manifold $X(\mathbb{C})$, which invokes $\mathbb{R}$ and the Archimedean completion.

Calibration Theorem 4.1 (Betti Realization)**.** $\sigma(\mathsf{Real}_{\mathrm{Betti}}) = \mathsf{CLASS}$. The obstruction is the formation of $X(\mathbb{C})$ and integration of differential forms over topological cycles.

4.2 de Rham Realization (BISH)

The algebraic de Rham cohomology $H^*_{\mathrm{dR}}(X/k)$ is computed by the Čech–de Rham spectral sequence on an affine cover. All differentials are algebraic polynomials. No integration, no topology.

Calibration Theorem 4.2 (de Rham Realization)**.** $\sigma(\mathsf{Real}_{\mathrm{dR}}) = \mathsf{BISH}$. Purely algebraic: differential forms modulo exact forms.

4.3 Étale Realization (BISH, with a caveat)

The ℓ-adic étale cohomology $H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Q}_\ell)$ is:

(a) $H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Z}/\ell^n\mathbb{Z})$ for each n: finite Galois cohomology, computed by finite covers. BISH.

(b) $H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Z}_\ell) = \varprojlim_n H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Z}/\ell^n\mathbb{Z})$: a projective limit of finite modules.

(c) $H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Q}_\ell) = H^*_{\text{ét}}(X_{\bar{k}}, \mathbb{Z}_\ell) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$.

Remark 4.3 (The "Smooth Projective" Heavy Lifting)**.** We emphasize that the *smooth projective* hypothesis does heavy lifting here. For pure motives, these are finitely generated modules, and the transition maps satisfy the Mittag-Leffler condition with effectively bounded torsion (via Deligne/Gabber). Thus, the limit is constructively bounded and only requires Dependent Choice (BISH), avoiding WKL_0. Furthermore, while equality testing in the completion $\mathbb{Q}_\ell$ generally requires WLPO, the essential motivic data (Frobenius traces) lie in $\mathbb{Z}$ by Deligne's Weil II, so computations descend to exact $\mathbb{Q}$-arithmetic.

If one moves to singular or open varieties (mixed motives), the Mittag-Leffler bounds on torsion may fail, potentially elevating the projective limit to genuinely require WKL_0. The BISH signature is a strict feature of pure motives.

Calibration Theorem 4.4 (Étale Realization)**.** $\sigma(\mathsf{Real}_{\text{ét}}) = \mathsf{BISH}$ (for pure motives). The limit is effectively bounded, and spectral data descend to $\mathbb{Z}$.

4.4 Crystalline Realization (BISH)

Crystalline cohomology $H^*_{\mathrm{cris}}(X/W(\mathbb{F}_q))$ is a finite-dimensional $W(\mathbb{F}_q)$-module. The Frobenius φ is an explicit matrix whose characteristic polynomial descends to $\mathbb{Z}$, keeping the computational output safely in BISH.

Calibration Theorem 4.5 (Crystalline Realization)**.** $\sigma(\mathsf{Real}_{\mathrm{cris}}) = \mathsf{BISH}$. Linear algebra over Witt vectors.

4.5 Hodge Realization (CLASS)

The Hodge realization associates the decomposition $H^n(X(\mathbb{C}), \mathbb{C}) = \bigoplus H^{p,q}(X)$. The Hodge *filtration* is algebraic (BISH), but the *decomposition* requires harmonic representatives via elliptic regularity and the $\partial\bar{\partial}$-lemma.

Calibration Theorem 4.6 (Hodge Realization)**.** $\sigma(\mathsf{Real}_{\mathrm{Hodge}}) = \mathsf{CLASS}$. Requires elliptic PDE theory on $X(\mathbb{C})$.

4.6 Automorphic Realization (CLASS)

The automorphic realization (via Langlands) requires the spectral decomposition of $L^2(\mathrm{GL}_n(\mathbb{Q})\backslash\mathrm{GL}_n(\mathbb{A}_\mathbb{Q}))$—analysis on the adelic quotient, inherently including the Archimedean factor $\mathrm{GL}_n(\mathbb{R})$.

Calibration Theorem 4.7 (Automorphic Realization)**.** $\sigma(\mathsf{Real}_{\mathrm{aut}}) = \mathsf{CLASS}$. L^2 spectral theory on $G(\mathbb{R})$.

Lean: Realization signatures

```
def realization_signature : Realization -> CRMSignature
  | .Betti       => ⟨.CLASS, "Integration over topological cycles"⟩
  | .deRham      => ⟨.BISH,  "Algebraic differential forms"⟩
  | .etale       => ⟨.BISH,  "Bounded limits and finite Galois"⟩
  | .crystalline => ⟨.BISH,  "Linear algebra over Witt vectors"⟩
  | .Hodge       => ⟨.CLASS, "Hodge decomposition: elliptic PDE"⟩
  | .automorphic => ⟨.CLASS, "L2 spectral decomposition"⟩
```

4.7 Summary Table

Realization	CRM Level	Obstruction
de Rham (algebraic)	BISH	None
Étale ($\mathbb{Q}_\ell$)	BISH	None (pure motives)
Crystalline	BISH	None
Betti (singular)	CLASS	$X(\mathbb{C})$, integration
Hodge (decomposition)	CLASS	Elliptic PDE
Automorphic	CLASS	L^2 spectral theory

Table 1: CRM signatures of realization functors. The BISH/CLASS boundary coincides exactly with the algebraic/analytic boundary.

5 Comparison Isomorphisms: The Archimedean Obstruction

The power of the motivic framework comes from comparison isomorphisms between realizations. These are the "bridges" that allow information to flow between cohomology theories. Their CRM cost is the key structural invariant.

5.1 Betti ↔ de Rham: The Period Map (CLASS)

The comparison is integration:

$$H^n_{\mathrm{dR}}(X/\mathbb{Q}) \otimes_\mathbb{Q} \mathbb{C} \xrightarrow{\sim} H^n_{\mathrm{sing}}(X(\mathbb{C}), \mathbb{Q}) \otimes_\mathbb{Q} \mathbb{C}$$

given by $\omega \mapsto \left(\gamma \mapsto \int_\gamma \omega\right)$. The period matrix consists of transcendental numbers computed by integration over $\mathbb{R}$. This is irreducibly CLASS.

5.2 Crystalline ↔ de Rham: p-adic Hodge Theory (BISH)

Berthelot's comparison isomorphism and Fontaine's functors D_{cris}, D_{dR} operate entirely within p-adic algebra. The filtered φ-module associated to a crystalline Galois representation is computed by explicit linear algebra over $\mathbb{Q}_p$. No $\mathbb{R}$, no integration: BISH.

5.3 Étale ↔ Betti: Artin Comparison (CLASS)

The Artin comparison relates algebraic covers over $\bar{k}$ to topological covers over $\mathbb{C}$ via the Riemann Existence Theorem (RET). This fundamentally requires fixing an embedding $\bar{k} \hookrightarrow \mathbb{C}$. Constructing this embedding invokes the Archimedean continuum and the Axiom of Choice. Furthermore, RET relies on complex analysis (GAGA, elliptic PDEs) to promote analytic sections to global algebraic functions. Thus, the comparison is irreducibly CLASS, even for finite coefficients.

5.4 de Rham ↔ Hodge: The $\partial\bar{\partial}$-Lemma (CLASS)

The comparison between the algebraic Hodge filtration and the analytic Hodge decomposition requires the $\partial\bar{\partial}$-lemma, which in turn requires elliptic regularity on the compact Kähler manifold $X(\mathbb{C})$. This is CLASS.

5.5 Crystalline ↔ Étale: Fontaine–Laffaille (BISH)

For crystalline representations with bounded Hodge–Tate weights, the Fontaine–Laffaille functor gives an explicit, algebraic equivalence between filtered φ-modules and Galois representations. It is BISH.

Comparison	**CRM Level**	**Passes through $\mathbb{R}$?**
Crystalline ↔ de Rham	BISH	No
Crystalline ↔ Étale	BISH	No
Étale ↔ Betti	CLASS	Yes (Embedding $\bar{k} \hookrightarrow \mathbb{C}$, RET)
Betti ↔ de Rham	CLASS	Yes (period integrals)
de Rham ↔ Hodge	CLASS	Yes (elliptic PDE)

Table 2: CRM signatures of comparison isomorphisms. The pattern is sharp: CLASS iff Archimedean.

5.6 The Archimedean Obstruction Theorem



Figure 2: The motivic CRM architecture. Solid blue arrows: BISH (constructive). Dashed red arrows: CLASS (classical). The motive is BISH; the BISH/CLASS boundary coincides exactly with the non-Archimedean/Archimedean boundary.

Theorem 5.1 (Archimedean Obstruction)**.** *Let P be a proof path using only non-Archimedean realizations (de Rham, étale, crystalline) and comparisons between them. Then $\sigma(P) = \mathsf{BISH}$.*

Conversely, if P uses any Archimedean realization (Betti, Hodge, automorphic) or any comparison passing through the Archimedean place, then $\sigma(P) \geq \mathsf{CLASS}$.

Proof. For the forward direction, recall from Definition 2.2 that

$$\sigma(P) = \max\big(\sigma(\text{motive}),\ \max_i \sigma(R_i),\ \max_j \sigma(C_j)\big).$$

By Proposition 3.1, $\sigma(\text{motive}) = \mathsf{BISH}$. By Table 1, each non-Archimedean realization has $\sigma(R_i) = \mathsf{BISH}$ (de Rham: algebraic differential forms; étale: finite Galois cohomology with bounded limits; crystalline: linear algebra over Witt vectors). By Table 2, every comparison between non-Archimedean realizations has $\sigma(C_j) = \mathsf{BISH}$ (Berthelot, Fontaine–Laffaille: explicit algebraic equivalences). Since $\mathsf{BISH} = \max(\mathsf{BISH}, \ldots, \mathsf{BISH})$, we conclude $\sigma(P) = \mathsf{BISH}$.

The converse is immediate: if any component has $\sigma = \mathsf{CLASS}$, the maximum is $\geq \mathsf{CLASS}$. In fact, every Archimedean realization (Betti, Hodge, automorphic) has $\sigma = \mathsf{CLASS}$ by Table 1, and every comparison passing through the Archimedean place has $\sigma = \mathsf{CLASS}$ by Table 2. Hence $\sigma(P) \geq \mathsf{CLASS}$. □

Lean: Archimedean Obstruction

```
theorem archimedean_obstruction (p : ProofPath)
    (h_real : forall r in p.realizations,
              is_non_archimedean r = true)
    (h_comp : forall c in p.comparisons,
              comparison_is_non_archimedean c = true) :
    proof_signature p = .BISH := by
  unfold proof_signature
  -- All realization/comparison signatures are BISH
  -- by helper lemmas + list induction on foldl join
  have hr := map_all_bish_realizations ..
  have hc := map_all_bish_comparisons ..
  exact foldl_join_bish_acc _ .BISH rfl
    (append_all_bish _ _ (append_all_bish ..) hc)
```

6 Why the Six Domains Have Different Signatures

The Archimedean obstruction theorem immediately explains the empirical pattern observed across Papers 50–97.

6.1 Algebra and Representation Theory: BISH

Hecke algebras, Galois deformation rings, modular representation theory—these operate in the étale and de Rham world. No passage through $\mathbb{R}$.

6.2 Harmonic Analysis: CLASS

The trace formula—Selberg, Arthur, Langlands—decomposes $L^2(G(\mathbb{Q})\backslash G(\mathbb{A}))$. The adelic group includes the Archimedean factor $G(\mathbb{R})$. The continuous spectral decomposition at infinity is irreducibly CLASS, making the trace formula the universal classical bottleneck of the Langlands program.

6.3 Number Theory: Sharp Classical Bottlenecks

L-functions are Dirichlet series defined for $\mathrm{Re}(s) \gg 0$ by an algebraic formula. The analytic continuation to all of $\mathbb{C}$ is an Archimedean construction. The critical values $L(k, f)$ involve the Archimedean Γ-factor and period integrals: CLASS. But the core algebraic objects (Selmer groups) are étale: BISH.

6.4 Algebraic Geometry: Bifurcation

The Hodge conjecture sits at the intersection of algebraic and analytic: Hodge classes live in the Betti realization (CLASS), but algebraic classes live in the motive (BISH). Generically, the detection problem is WLPO (Paper 87). But algebraic geometry has moduli. On the palindromic locus of genus-4 curves, the Kani–Rosen decomposition $J(C_t) \sim A_1 \times A_2$ provides an algebraic shortcut (BISH). Moduli dimension controls the availability of such bifurcations.

6.5 Topology: Mixed

Étale cohomology with finite coefficients is BISH. Singular (Betti) cohomology is CLASS via the Archimedean place.

6.6 Mathematical Physics

Classical mechanics (finite-dimensional Hamiltonian ODEs) is BISH. Quantum mechanics introduces the spectral theorem on $L^2(\mathbb{R}^3)$—CLASS in general, but BISH for exactly solvable Hamiltonians with algebraic spectra.

7 Calibration Theorem I: Hodge Detection $\leftrightarrow$ WLPO

We summarize the Hodge calibration from Papers 86–90.

Theorem 7.1 (Hodge Detection Calibration, Papers 87–90)**.** *There exists a family of Hodge classes $\xi_t \in H^4(J(C_t)^2, \mathbb{Q})$ over the genus-4 Torelli locus such that:*

(i) On the palindromic locus: ξ_t is algebraic, verified by BISH methods (Kani–Rosen splitting, Ribet's theorem).

(ii) On the generic fiber: deciding whether ξ_t is algebraic is equivalent to WLPO (decidability of equality on the Mumford–Tate group).

Proof sketch. (i) On the palindromic locus $\{a = c\}$ of the family $C_t : y^2 = x^9 + ax^7 + bx^5 + cx^3 + x$, the hidden reciprocal involution $\mu(x, y) = (1/x, y/x^5)$ generates a dihedral group D_8 with the hyperelliptic involution. By the Kani–Rosen theorem [5], $J(C_t) \sim A_1 \times A_2$ with $A_1 \cong A_2$ over $\mathbb{C}$, so $J(C_t) \sim A^2$. For generic $\mathrm{End}(A) = \mathbb{Z}$, the Mumford–Tate group is GSp_4 (Zarhin), and all Hodge classes on powers of abelian surfaces are algebraic by Ribet [4]. This chain uses only étale data and algebraic correspondences: BISH.

(ii) On the generic fiber, detecting whether ξ_t is a Hodge class requires the Hodge decomposition $H^4(J(C_t)^2, \mathbb{C}) = \bigoplus H^{p,q}$, which is CLASS (elliptic PDE). Deciding whether $\xi_t \in H^{2,2}$ reduces to deciding equality $\mathrm{MT}(J(C_t)) \stackrel{?}{=} \mathrm{GSp}_{2g}$, which is precisely WLPO: decidability of equality for a real algebraic group (Paper 87, Theorem B). The gap between (i) and (ii) is therefore WLPO. □

Remark 7.2 (The Generality of the Bifurcation)**.** The choice of genus 4 in Papers 86–90 was not an ad hoc quirk; it is the minimal dimension where the generic fiber has unforced Hodge classes (unlike curves or surfaces) while still possessing an explicitly computable continuous exceptional

locus (the palindromic locus). The WLPO bifurcation is a general structural phenomenon for any moduli space of dimension ≥ 2.



Figure 3: The Hodge bifurcation (Papers 86–89). On the palindromic locus $\{a = c\}$, algebraic splitting provides BISH Hodge class detection. On the generic fiber, the Hodge decomposition requires CLASS. The gap is exactly WLPO.

8 Calibration Theorem II: TW Patching $\leftrightarrow$ WKL_0

We analyze the CRM structure of the Wiles–Taylor proof of modularity.

8.1 The Three Tiers

Tier 1 (BISH): The algebraic core. Galois deformation rings R, Hecke algebras $\mathbb{T}$, Fontaine–Laffaille, Wiles's numerical criterion—all étale/de Rham.

Tier 2 (WKL_0): Taylor–Wiles patching. The patching argument constructs an inverse limit $R_\infty = \varprojlim R_{Q_n}$. Extracting a compatible system from this inverse limit of finite local rings is precisely Weak König's Lemma.

Calibration Theorem 8.1 (TW Patching $\leftrightarrow$ WKL_0)**.** The Taylor–Wiles patching argument, as a logical principle, is equivalent to WKL_0: extracting a compatible sequence from an inverse system of finite sets is equivalent to finding an infinite path through a finitely branching tree.

Proof. The patching argument constructs an inverse system of surjections $R_{Q_{n+1}} \twoheadrightarrow R_{Q_n}$ between finite local rings. The desired compatible sequence $(x_n)_{n\geq 1}$ with $x_n \in R_{Q_n}$ and $\pi_n(x_{n+1}) = x_n$ exists if and only if the inverse limit $\varprojlim R_{Q_n}$ is nonempty. The tree T whose level-n nodes are elements of R_{Q_n}, with edges given by the transition maps, is infinite (each R_{Q_n} is nonempty) and finitely branching (each R_{Q_n} is finite). By WKL_0, T has an infinite path, which is exactly a compatible sequence. Conversely, if every such inverse system of finite sets has a compatible sequence, then every infinite finitely branching tree has an infinite path (WKL_0). The equivalence is standard (cf. Simpson's subsystems of second-order arithmetic). $\square$

Tier 3 (CLASS): The analytic bridge. The Langlands–Tunnell theorem uses the trace formula. Faltings's isogeny theorem uses Arakelov heights and complex integration: CLASS.

8.2 The 1993 $\to$ 1995 Excision

Wiles's original (1993) strategy attempted to compute the congruence ideal $\eta_{\mathbb{T}}$ via the Petersson inner product, which requires integration over the complex upper half-plane (CLASS). The Taylor–Wiles fix replaced this with the patching argument, which bypasses the inner product entirely.

Theorem 8.2 (CLASS $\to$ WKL_0 Excision)**.** *The Taylor–Wiles (1995) repair of Wiles (1993) is a logical excision: it replaces a CLASS computation (Petersson inner product) with a WKL_0 argument (inverse limit patching), strictly reducing the logical strength of the proof at the patching step.*

Proof. Wiles's 1993 strategy [6] required computing the congruence ideal $\eta_{\mathbb{T}}$ of the Hecke algebra $\mathbb{T}$ acting on $S_2(\Gamma_0(N))$. His method: the Petersson inner product $\langle f, f\rangle = \int_{\Gamma_0(N)\backslash\mathfrak{H}} |f(z)|^2\, y^2\, \frac{dx\,dy}{y^2}$, an integral over the non-compact modular curve, requiring real integration on the upper half-plane. This is irreducibly CLASS (Archimedean: integration over $\mathbb{R}^2$).

Taylor–Wiles [7] replaced this computation with the patching argument: choose auxiliary primes Q_n and construct a system of augmented deformation rings R_{Q_n} whose inverse limit R_∞ is a power series ring, from which the $R = \mathbb{T}$ isomorphism follows by a dimension argument. The patching step requires only WKL_0 (Calibration Theorem 8.1). Since $\mathsf{WKL}_0 <$ CLASS in the CRM hierarchy, the logical strength strictly decreases. □

Remark 8.3 (Historical Intent vs. Logical Reality)**.** To be explicitly clear, this logical excision is a retrospective structural observation, not a historical claim about mathematical intent. Wiles and Taylor invented patching to bypass a profound mathematical gap (an inadequate bound on the Selmer group), not to deliberately reduce the formal logical strength of their proof. Nevertheless, the structural consequence of their historic repair was a strict reduction in the CRM signature.

9 Calibration Theorem III: BSD Rank/Ш Decoupling

Theorem 9.1 (BSD Decoupling)**.** *The two components of the BSD conjecture have different CRM signatures:*

(i) ***Rank verification****: Given a candidate point, computing the rank via algebraic 2-descent is BISH.*

(ii) Ш ***finiteness****: Proving $|Ш(E/\mathbb{Q})| < \infty$ is CLASS. Kolyvagin's Euler system requires annihilating $Ш[p^\infty]$ for all primes simultaneously, necessitating the analytic topology of the Euler system.*

Proof. (i) Given $E/\mathbb{Q}$ and a candidate point $P \in E(\mathbb{Q})$, verifying $\hat{h}(P) > 0$ (hence P has infinite order) requires only explicit computation of the canonical height, which is an algebraic formula involving local heights at each prime (Néron functions) and the Archimedean height $\lambda_\infty(P)$ (the naïve height). For E defined over $\mathbb{Q}$ and $P \in E(\mathbb{Q})$, these are rational computations bounded by explicit Silverman constants (Paper 95). The 2-Selmer group $\mathrm{Sel}_2(E/\mathbb{Q})$ is computed by finite Galois cohomology $H^1(\mathbb{Q}, E[2])$ with local conditions, which is a finite algebraic computation: BISH.

(ii) Kolyvagin's proof [8] constructs an Euler system: a compatible family of cohomology classes $c_n \in H^1(\mathbb{Q}(\mu_n), E[p])$ indexed by square-free integers n, with norm-compatibility relations. The key step—proving $Ш(E/\mathbb{Q})[p^\infty] = 0$—requires showing that the Euler system classes annihilate all of Ш simultaneously for all primes p. This uses Chebotarev's density theorem (distributional statement about Frobenius elements, requiring CLASS) and the Gross–Zagier formula [9] (relating $L'(E, 1)$ to the height of a Heegner point via period integrals: CLASS). Both pass through the Archimedean place. □

Why no continuous bifurcation? Unlike the Hodge conjecture, BSD does not exhibit a continuous constructive/classical bifurcation. Logical bifurcation requires a continuous parameter space where an algebraic constraint defines a positive-dimensional sublocus, allowing one to continuously bypass a generic equality test. For elliptic curves, the moduli space (the j-line) has

dimension 1. A proper algebraic subvariety of a 1-dimensional space is 0-dimensional (isolated CM points). Without positive-dimensional intermediate loci to tune continuous parameters, the WLPO-type structural bifurcation cannot exist.

10 Signature Preservation Under Langlands

Theorem 10.1 (Signature Preservation)**.** *The Langlands correspondence for* $\mathrm{GL}_2/\mathbb{Q}$ *preserves CRM signatures:*

(i) The algebraic data on both sides (Hecke eigenvalues, Galois traces) are BISH.

(ii) Verifying the correspondence for a given pair (E, f) *is BISH.*

(iii) Establishing the correspondence generally is CLASS: the proof uses the trace formula and cannot avoid the Archimedean place.

Proof. (i) The Hecke eigenvalues $a_\ell(f)$ of a weight-2 newform $f \in S_2(\Gamma_0(N))$ are algebraic integers in a number field K_f. For an elliptic curve $E/\mathbb{Q}$ of conductor N, the Galois trace $a_\ell(E) = \ell + 1 - |E(\mathbb{F}_\ell)|$ is computed by counting points on a finite field: a finite algebraic computation. Both sides of the correspondence produce algebraic data by finite arithmetic.

(ii) Given a conjectured pair (E, f), the verification $a_\ell(E) = a_\ell(f)$ for all $\ell \nmid N$ reduces to checking equality of two algebraic integers at each prime. Each check is a finite computation, and by Faltings' isogeny theorem, only finitely many primes suffice (bounded by the Sturm bound $k \cdot [\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(N)]/12$). The entire verification is a finite sequence of finite computations: BISH.

(iii) The proof of modularity (that every $E/\mathbb{Q}$ *has* a corresponding f) uses the trace formula: $\sum_\pi m(\pi) \operatorname{tr} \pi(f) = \sum_\gamma \mathrm{vol}(\dots) \cdot O_\gamma(f)$. The spectral side requires L^2-decomposition of automorphic forms on $\mathrm{GL}_2(\mathbb{A}_\mathbb{Q})$, which passes through the Archimedean component $\mathrm{GL}_2(\mathbb{R})$. The L^2 spectral theory at this place is irreducibly CLASS (Baire category, spectral projections). The algebraic data is preserved—only the *method of establishing the correspondence* incurs classical cost. □

11 The Function Field Test

The ultimate predictive test of the Archimedean Obstruction Thesis lies in function fields $\mathbb{F}_q(C)$, where the adèle ring $\mathbb{A}_F$ lacks an Archimedean factor.

Theorem 11.1 (Function Field Constructivity)**.** *The Langlands correspondence for* GL_n *over function fields* $\mathbb{F}_q(C)$*, as established by Lafforgue, is provable in BISH (with at most* WKL_0 *for algebraic limits).*

Proof. Over a function field $F = \mathbb{F}_q(C)$, the adèle ring $\mathbb{A}_F = \prod'_v F_v$ has *no Archimedean place*: every completion F_v is a non-Archimedean local field (isomorphic to $\mathbb{F}_q(\!(t)\!)$ or a finite extension thereof). By the Archimedean Obstruction Theorem (Theorem 5.1), any proof path whose realizations and comparisons are all non-Archimedean has signature BISH.

Concretely, Lafforgue's proof [10] of the Langlands correspondence for GL_n/F proceeds via the moduli of shtukas, which are algebraic objects over $\mathbb{F}_q$. The cohomology used is étale (ℓ-adic, $\ell \neq \operatorname{char} \mathbb{F}_q$): a non-Archimedean realization. The comparison isomorphisms involved are crystalline–étale (Fontaine–Messing over p-adic fields): non-Archimedean. In the Lean formalization, the proof path `function_field_langlands_path` uses only étale and crystalline realizations with crystalline–étale comparison. `native_decide` confirms $\sigma = \mathsf{BISH}$, and alternatively the result follows from Theorem A applied to the all-non-Archimedean path.

The only logical cost beyond BISH is WKL_0 if one uses inverse limits in the patching arguments (as in Taylor–Wiles style constructions adapted to function fields), which is strictly below CLASS. □

12 Physics Calibrations

The Archimedean obstruction pattern extends beyond pure mathematics. The CRM program has documented logical signatures for the principal domains of mathematical physics (Papers 2, 5–8, 33).

12.1 Classical Mechanics: BISH

Hamiltonian systems on finite-dimensional phase spaces are governed by ODEs with polynomial right-hand sides. For algebraically specified initial data, the Picard–Lindelöf iteration is constructive and convergent. The formal Lean bundle records $\sigma(\text{classical mechanics}) = \mathsf{BISH}$.

12.2 Quantum Mechanics: BISH + LPO gap

The spectral theorem for self-adjoint operators on $L^2(\mathbb{R}^3)$ is CLASS in full generality—it requires the Baire category theorem and the existence of spectral projections. But for exactly solvable systems (hydrogen atom, harmonic oscillator), the spectrum is algebraic and the eigenfunctions are explicit. The gap is LPO: deciding whether a given energy lies in the continuous spectrum.

12.3 General Relativity: BISH + CLASS gap

The Choquet-Bruhat local existence theorem for the Einstein equations requires elliptic regularity and Sobolev embedding: CLASS [22]. But explicit solutions (Schwarzschild, Kerr, FLRW) are algebraically specified and verified by direct substitution: BISH. The gap parallels the Hodge bifurcation—generic existence requires analysis, while explicit instances descend to algebra.

12.4 Lattice vs. Continuum QFT

Lattice gauge theories (Wilson action on a finite lattice) are BISH: the partition function is a finite sum. Continuum QFT requires renormalization, which invokes the operator product expansion and regularization of divergent integrals: CLASS (or worse—some procedures are not even well-defined classically). The confinement problem (Paper 33) sits at this boundary.

In each case, the constructive/classical boundary coincides with the algebraic/analytic boundary. Discretized, finite, algebraic physics is BISH. Continuum, infinite-dimensional, analytic physics is CLASS. The real numbers mediate the transition—but the cost depends on which algebraic layer of $\mathbb{R}$ is invoked. Explicit physics (Schwarzschild by substitution, scattering amplitudes, lattice models) uses $\mathbb{R}$-as-ring: BISH. Spectral physics (gap decidability, phase transitions) uses $\mathbb{R}$-as-ordered-field: LPO. Generic existence theorems (Choquet-Bruhat, continuum spectral decomposition) use $\mathbb{R}$-as-complete-ordered-field: CLASS. The physics ceiling at LPO (Paper 70) reflects the fact that empirical physics never invokes completeness—only the ordering.



Figure 4: The six-domain signature map. Domains that avoid the Archimedean place are BISH (green). Domains that pass through $\mathbb{R}$ acquire CLASS content (red/orange). The Archimedean Thesis: the real numbers are the unique source of non-constructive content.

13 The Archimedean Thesis: A Synthesis

The Archimedean Thesis

The Archimedean place—the completeness and ordering of $\mathbb{R}$, not its ring structure—is the unique source of non-constructive content in arithmetic geometry, and more broadly in the mathematics connected by the Langlands program.
Formally: the CRM signature of a theorem about a motive M is determined by:

$$\sigma(T) \;=\; \max\Big(\underbrace{\sigma(M)}_{=\mathsf{BISH}},\ \underbrace{\sigma(\text{realizations})}_{=\mathsf{BISH}\text{ or }\mathsf{CLASS}},\ \underbrace{\sigma(\text{comparisons})}_{=\mathsf{BISH}\text{ iff non-Archimedean}}\Big)$$

The CRM cost of each realization and comparison is determined by which algebraic layer of $\mathbb{R}$ it invokes:

Layer of $\mathbb{R}$	**Structure**	**CRM cost**	**What it enables**
Ring	$(\mathbb{R}, +, \times)$	BISH	Polynomial identity, algebraic de Rham, `ring`
Ordered field	$(\mathbb{R}, +, \times, <)$	LPO/WLPO	Zero-testing, spectral gaps, Hodge detection
Complete ordered field	$(\mathbb{R}, +, \times, <, \mathrm{LUB})$	CLASS	Integration, L^2 spectral theory, period maps

Consequences:

(1) Remove the Archimedean place (function fields): everything is BISH.

(2) Keep $\mathbb{R}$ but use only its ring structure (algebraic proofs): BISH.

(3) Invoke the ordering (spectral sign decisions): LPO. This is the physics ceiling.

(4) Invoke completeness (analytic proofs, comparison isomorphisms): CLASS enters.

(5) The boundary is sharp: WLPO for Hodge detection (ordering layer), WKL_0 for patching.

(6) The trace formula is the universal CLASS bottleneck (completeness layer).

14 CRM Audit

This paper is itself a CRM object. Its formal content decomposes as follows.

Component	CRM Level	Justification
CRM hierarchy (total order)	BISH	Decidable finite type
Realization signatures (6 functors)	BISH	Definitional classification
Comparison signatures (36 pairs)	BISH	Definitional classification
Archimedean obstruction theorem	BISH	List induction, no LEM
Calibration theorems (Hodge, BSD, TW)	BISH	`native_decide` on finite data
Excision theorem	BISH	`decide` on 2 constants
Langlands signature preservation	BISH	Definitional
Physics parallels	BISH	Definitional

Table 3: CRM audit of Paper 98. The meta-theorem: the logical cost of *classifying* logical cost is BISH.

The entire formal content is BISH. No classical axiom is used in any proof. The Lean bundle's axiom inventory shows only `propext` and `Quot.sound` (Lean infrastructure) for the main theorem, and no axioms at all for the excision and physics gap theorems.

15 Formal Verification

The companion Lean 4 bundle `P98_MotivicCRM/` consists of 6 source files (607 lines), building against Mathlib v4.29.0-rc2 with zero `sorry`, zero warnings. The geometric realization functors are represented as axiomatic stubs (constructors of an inductive type with definitional CRM costs). The Lean verifies the *logical framework*: that if the realizations have the claimed signatures, then the obstruction theorem and calibration theorems follow. The geometric content itself is verified by the papers in the series.

15.1 File Structure

1. `CRMLevel.lean` (94 lines): The 7-level CRM hierarchy as a decidable total order with join (max) operation.
2. `Realizations.lean` (128 lines): Six realization functors, their CRM signatures, all 36 comparison pairs classified, non-archimedean lemmas.
3. `ArchimedeanObstruction.lean` (94 lines): Proof paths, the `proof_signature` function, and Theorem A.
4. `Calibration.lean` (97 lines): Calibration theorems for Hodge (I), BSD (II), and TW excision (III).
5. `Langlands.lean` (66 lines): Signature preservation, function field constructivity (two proofs: `native_decide` and via Theorem A).
6. `Physics.lean` (57 lines): Documentary CRM signatures for classical mechanics, QM, GR.

15.2 Axiom Inventory

The absence of `Classical.choice` is noteworthy: this is a paper *about* the logical cost of classical reasoning, and its own proofs use none. The entire verification is constructive.

Theorem	Axioms
`archimedean_obstruction`	`propext`, `Quot.sound`
`hodge_generic_is_class`	`native_decide`
`bsd_rank_is_bish`	`native_decide`
`modularity_core_is_bish`	`native_decide`
`excision_1993_to_1995`	(none)
`function_field_langlands_is_bish'`	`propext`, `Quot.sound`
`physics_gap_qm`, `physics_gap_gr`	(none)

Table 4: Axiom inventory. No `Classical.choice`. No mathematical axioms.

15.3 Reproducibility

Toolchain: `leanprover/lean4:v4.29.0-rc2`. **Dependency:** Mathlib (fetched automatically by `lake`).

```
cd P98_MotivicCRM && lake build
```

16 Discussion

16.1 Relation to the De-Omniscientising Descent

The central narrative of the CRM program is the *de-omniscientising descent*: the systematic identification and removal of unnecessary omniscience principles from mathematical proofs. The motivic architecture provides the structural explanation for why this descent is possible at all. The motive is BISH; classical content enters only through realization functors and comparison isomorphisms; and these enter only at the Archimedean place. Therefore, any proof that can be rerouted through non-Archimedean realizations automatically descends.

The descent has three modes, all documented across Papers 50–97:

(i) **Algebraic bypass**: Replace an analytic computation with an algebraic one (Taylor–Wiles patching, 2-descent for BSD rank). This excises CLASS and replaces it with WKL_0 or BISH.

(ii) **Locus restriction**: Restrict to a special sublocus where extra symmetry provides algebraic shortcuts (palindromic locus for Hodge, CM points for BSD). This gives BISH on the restricted locus while the generic fiber remains CLASS.

(iii) **Domain transfer**: Move to a domain that lacks the Archimedean place entirely (function fields). This gives uniform BISH without restriction.

16.2 Open Problems

Several structural questions remain open:

(1) **Jacquet–Langlands signature preservation**: Does the Jacquet–Langlands correspondence preserve CRM signatures for all inner forms of GL_n? The algebraic data should be BISH; the proof method is CLASS via the trace formula. But can the proof be rerouted?

(2) **Motivic Langlands for GL_n**: The current analysis covers $\mathrm{GL}_2/\mathbb{Q}$ and function fields. Extending to GL_n requires understanding the CRM cost of the stable trace formula (Arthur's endoscopic classification).

(3) **Positive mass calibration**: The positive mass theorem in general relativity uses Witten's spinor argument or Schoen–Yau minimal surfaces. Both are CLASS. Is there an algebraic proof for specific asymptotically flat manifolds?

(4) **Conservation conjecture**: Can every theorem proved with CLASS that has a BISH-stateable conclusion be re-proved at a strictly lower CRM level? The motivic architecture suggests yes for theorems about motives, but the general case remains open.

16.3 The Archimedean Stratification

A refinement implicit throughout the program but not previously stated as a unified observation. The real number field $\mathbb{R}$ carries three nested algebraic structures, and the CRM hierarchy maps onto them:

Layer	Structure	CRM cost	What it enables	Programme evidence
Ring	$(\mathbb{R}, +, \times)$	BISH	Polynomial identity, `ring`	Papers 80–89, 92–96
Ordered field	$(\mathbb{R}, +, \times, <)$	LPO/WLPO	Sign decisions, spectral gaps	Papers 29, 36, 87
Complete o.f.	$(\mathbb{R}, +, \times, <, \mathrm{LUB})$	CLASS	Integration, L^2, period maps	Papers 45–49, 95–96

Table 5: The Archimedean Stratification: the CRM hierarchy as the algebraic hierarchy of $\mathbb{R}$.

The physics ceiling at LPO (Papers 2–44) reflects physics's use of $\mathbb{R}$-as-ordered-field (the spectral theorem requires sign decisions; phase transitions require Fekete's lemma, which is LPO-equivalent) without invoking completeness. The arithmetic geometry ceiling at CLASS (Papers 45–98) reflects the use of $\mathbb{R}$-as-complete-ordered-field through comparison isomorphisms: period integrals require integration (LUB for Riemann sums), the Hodge decomposition requires elliptic regularity (Sobolev embedding, which uses LUB), and L^2 spectral decomposition requires the Baire category theorem (which requires LUB). The algebra floor at BISH (every `ring` proof in the program) reflects the use of $\mathbb{R}$-as-ring alone.

The u-invariant $u(\mathbb{R}) = \infty$ (Paper 50) is a property of the *ordered field* layer: positive-definiteness requires $<$ but not LUB. This is why it governs the physics ceiling (LPO) rather than the arithmetic ceiling (CLASS).

The three descent modes (§16) can be restated in terms of layers: algebraic bypass descends from the completeness layer to the ring layer; locus restriction provides extra algebraic data that makes the ordering layer sufficient (Hodge detection drops from CLASS to WLPO with CM metadata); domain transfer removes the Archimedean place entirely, eliminating all three layers.

This stratification explains why the natural world—which evolves differential equations (ring operations), propagates waves (ring), and traces geodesics (ring)—operates at BISH+LPO rather than CLASS: physics uses the ring and ordering layers of $\mathbb{R}$ but never invokes its completeness.

17 Conclusion

The motivic CRM architecture provides a unified explanation for why different areas of mathematics feel qualitatively different: they have different logical signatures, determined entirely by their interaction with the real numbers. Motives are the Rosetta Stone. The CRM signature of each domain is the CRM cost of the realization functor it uses. The trace formula is the universal classical bottleneck because it introduces the Archimedean place into the automorphic machinery. Function field arithmetic is fully constructive because it lacks an Archimedean place entirely. And the Taylor–Wiles fix of Fermat's Last Theorem was, at its logical core, an excision of an Archimedean computation.

The companion 0-`sorry` Lean 4 formalization mechanically verifies the abstract logical framework (the CRM hierarchy, signature propagation, and the Archimedean obstruction theorem), while the geometric realization functors are represented as axiomatic stubs. The calibration theorems for Hodge, BSD, and modularity are mechanically checked to have the claimed signatures. The excision theorem (CLASS $\to$ WKL_0) is proved by `decide`.

This paper synthesizes the findings of Papers 50–97 in the CRM program. It does not resolve the open conjectures—Jacquet–Langlands signature preservation, positive mass calibration, spectral theorem stratification, motivic Langlands for GL_n—but it formulates them precisely, within a framework that makes them mechanically verifiable once the mathematical content is supplied.

Acknowledgments. The author is an interventional cardiologist, not a domain expert in the mathematical areas treated here; all results should be considered preliminary and verified independently. The Lean 4 formalization uses the Mathlib library [23]. The Lean code and paper were developed with assistance from Claude (Anthropic) and Gemini (Google). The paper format follows the CRM Series format guide [24].

A The Motivic Reinterpretation: Papers 2–97 Through the Archimedean Lens

This appendix systematically reinterprets the entire CRM program through the framework of Paper 98. The Archimedean Obstruction Thesis (Theorem 5.1) assigns every mathematical object a CRM signature determined by which realization functors it uses and whether its proof path passes through the Archimedean place. We now ask: does this framework retrodict the findings of Papers 2–97? Where does it sharpen them? And where does it reveal new opportunities?

A.1 Phase 0: Mathematical Physics (Papers 2–44)

The physics program established the *empirical ceiling*: $\mathsf{BISH} + \mathsf{LPO}$ suffices for all empirically accessible physics. The motivic framework explains *why*.

Retroactive explanation. Physics operates in two regimes:

(i) **Finite-dimensional algebra** (classical mechanics, lattice models, scattering amplitudes): These are purely algebraic computations—matrix algebra, polynomial evaluation, finite sums—that never invoke the analytic topology of $\mathbb{R}$. By analogy with the motivic framework, they operate in the "de Rham" regime (algebraic identities, no integration). Prediction: BISH. Confirmed by Papers 28, 8–9, 34.

(ii) **Spectral theory on $L^2(\mathbb{R}^n)$** (quantum mechanics, continuum QFT, GR): The spectral decomposition of $L^2(\mathbb{R}^n)$ and the representation theory of the Poincaré group are the exact analytic analogues of the Archimedean component of the automorphic realization. (The Hodge realization is a functor from motives of algebraic varieties to Hodge structures; physics manifolds are generally not algebraic, so the correct parallel is automorphic, not Hodge.) The spectral theorem requires LPO (Paper 36 stratified Cubitt et al.'s spectral gap undecidability as LPO). Prediction: CLASS in general, BISH for exactly solvable systems where the spectrum is algebraic. Confirmed by Papers 4, 6–7, 14, 36–37.

New insight. The six realizations of §4 are defined for smooth projective algebraic varieties, not for arbitrary $\mathbb{R}$-manifolds. Nevertheless, the pattern extends by analogy: physics uses spectral analysis on $\mathbb{R}$-manifolds (the analogue of the automorphic realization at the Archimedean place), but never the inter-realization comparisons (Betti $\leftrightarrow$ étale, period maps) that characterize arithmetic geometry. The physics ceiling LPO is thus a *structural consequence*, not a contingent empirical finding: it is the cost of the Archimedean spectral analysis *alone*, without the additional cost of inter-realization comparisons that push arithmetic geometry to CLASS.

Sharpened predictions.

- Papers 30–31 proved Fan Theorem and Dependent Choice are physically dispensable. The motivic explanation: FT and DC are properties of Brouwerian continua and infinite sequences, but physics uses $\mathbb{R}$ only through its Archimedean metric, not through its Brouwerian topology. FT/DC are orthogonal to the Archimedean obstruction.
- Paper 39 (physical stratification) found that the *epistemic mode*—exact vs promise-gapped vs empirical—determines the arithmetic level of a physical observable, not its thermodynamic scaling. The discrete/continuous distinction in physics parallels the algebraic/analytic distinction in arithmetic geometry: discrete systems (finite sums, combinatorics) are BISH; continuum limits (real integration, spectral theory) are CLASS. The thermodynamic limit is, in motivic language, a change of realization regime—from algebraic to analytic.
- Paper 13 (event horizon as logical boundary): The Schwarzschild event horizon separates the algebraic exterior (Kruskal coordinates: rational functions, BISH) from the singular interior (Penrose diagram: conformal compactification requiring CLASS). By analogy with the motivic framework (the Schwarzschild metric is not an algebraic variety, so the parallel is structural rather than literal), the event horizon acts as a classical boundary node—the passage from algebraic (BISH) to topological (CLASS) description of the spacetime.

Opportunity. Paper 41 (AdS/CFT) and Paper 42 (cosmological constant) found CLASS logical signatures. The motivic framework suggests a specific target: if the bulk/boundary duality can be expressed in terms of étale or crystalline cohomology of the boundary CFT's moduli space (avoiding real integration), the logical cost should drop. This is a *constructive holography* program: translate the holographic dictionary from Betti to étale realization.

A.2 Phase 1: Arithmetic Geometry (Papers 45–66)

This phase established the DPT (Decidable Polarized Tannakian) framework and the $h = f$ discovery. The motivic framework provides the structural explanation for both.

The five conjectures (Papers 45–49). Papers 45–49 calibrated Hodge, Tate, BSD, Fontaine–Mazur, and Weight–Monodromy against the CRM hierarchy. Each was found to require LPO or CLASS in full generality, but to descend to BISH with algebraic metadata.

Motivic reinterpretation. Each conjecture asserts a relationship between realizations:

Conjecture	Comparison involved	Passes through $\mathbb{R}$?
Hodge	Betti $\leftrightarrow$ de Rham (period map)	Yes
Tate	Étale $\leftrightarrow$ algebraic cycles (ℓ-adic cycle class)	Statement: No; proof (Faltings): Yes
BSD	Automorphic $\leftrightarrow$ algebraic (L-function)	Yes
Fontaine–Mazur	Étale $\leftrightarrow$ de Rham (p-adic Hodge)	No (!)
Weight–Monodromy	Étale $\leftrightarrow$ crystalline (local)	No

Note on Tate: The algebraic Tate conjecture—surjectivity of the ℓ-adic cycle class map $\mathrm{CH}^i(X) \otimes \mathbb{Q}_\ell \to H^{2i}_{\text{ét}}(X_{\bar{k}}, \mathbb{Q}_\ell(i))^{\mathrm{Gal}(\bar{k}/k)}$—is defined purely via étale Chern classes and does not pass through $\mathbb{R}$. The "Yes" in the table refers to Faltings's *proof* of the Tate conjecture for abelian varieties, which uses Arakelov geometry (CLASS). Statement and proof must be distinguished: the statement is non-Archimedean, but all known proofs in the number field case invoke the Archimedean place. (If one means the Beilinson–Tate conjecture on L-function poles, the automorphic realization enters directly.)

Key prediction confirmed: Fontaine–Mazur and Weight–Monodromy involve only non-Archimedean comparisons (crystalline–étale, p-adic Hodge theory). The Archimedean Thesis predicts they should be "closer to BISH" than Hodge or BSD. Paper 47 confirmed: Fontaine–Mazur requires only LPO (for the Galois representation side), versus CLASS for Hodge and BSD. The residual

LPO comes not from the comparison isomorphism (which is BISH by Fontaine–Laffaille) but from the *construction* of the Galois representation itself (requiring ℓ-adic limits).

New opportunity: The motivic framework isolates a precise target for Fontaine–Mazur. The comparison (crystalline $\leftrightarrow$ étale) is BISH. The only CLASS content is the construction of the global Galois representation from local data (patching). If a purely p-adic construction existed (avoiding the need for complex embeddings), Fontaine–Mazur could in principle be proved in $\mathsf{BISH} + \mathsf{WKL}_0$. This is a concrete research target.

The DPT framework (Papers 50–53). The three DPT axioms—Standard Conjecture D (morphism decidability), algebraic spectrum (eigenvalue decidability), Archimedean polarization (cycle search)—were discovered empirically. The motivic framework explains their necessity.

Motivic reinterpretation. Each DPT axiom corresponds to a specific comparison isomorphism:

(i) Standard Conjecture D $\leftrightarrow$ cycle class map (Betti $\to$ algebraic): tests whether a Betti class is algebraic. Passes through $\mathbb{R}$ (period integrals). Cost: CLASS without oracle, BISH with D as axiom.

(ii) Algebraic spectrum $\leftrightarrow$ Hodge realization (eigenvalues of Frobenius/Hecke): tests whether spectral data is algebraic. Cost: CLASS without oracle (requires spectral theorem on L^2), BISH with algebraic spectrum as axiom.

(iii) Archimedean polarization $\leftrightarrow$ Hodge $\to$ Betti comparison (positivity of intersection form): the Hodge index theorem requires real integration. Cost: CLASS without oracle, BISH with polarization data.

The DPT axioms are *exactly the data needed to bypass the three Archimedean comparison isomorphisms.* This structural explanation was unavailable when Papers 50–53 were written; the motivic architecture provides it retrospectively.

The $h = f$ discovery (Papers 56–58, 65–66). The exotic Weil class self-intersection computation ($h = f$ for CM abelian fourfolds) was the program's first positive constructive result in Hodge theory.

Motivic reinterpretation. CM abelian varieties have $\mathrm{End}(A) \supsetneq \mathbb{Z}$. The extra endomorphisms provide an *algebraic substitute for the period map*: the CM type determines the Hodge decomposition without integration. In motivic language: CM data provides a splitting of the Betti $\leftrightarrow$ de Rham comparison into algebraic pieces, bypassing the Archimedean obstruction. This is why CM always reduces logical cost—it is a *non-Archimedean path through the Archimedean comparison.*

A.3 Phase 2–4: Proof Methods, Function Fields, Synthesis (Papers 67–75)

FLT: BISH or CLASS? A contested classification (Paper 68). The modularity lifting theorem ($R \cong \mathbb{T}$) routes through étale cohomology (Galois representations: BISH), crystalline cohomology (Fontaine–Laffaille: BISH), and Taylor–Wiles patching (WKL_0). Paper 68 audited the full proof of FLT in five stages:

- **Stages 2–5** (deformation theory, Fontaine–Laffaille, patching, Ribet level-lowering): BISH+ WKL_0. Uncontested.
- **Stage 1** (base case): The residual representation $\bar{\rho}_{E,2}$ must be shown modular. Wiles's 1995 proof used the Langlands–Tunnell theorem (S_4 case, trace formula: CLASS). The Kisin/Khare–Wintenberger route [20] at $p = 2$ replaces this with the D_3 (dihedral) case, where modularity follows from Hecke theta series.

The contested node is whether Hecke theta series are BISH or CLASS. Paper 68 argues BISH: the q-expansion $\theta_\chi(q) = \sum_{n\geq 0} r_\chi(n)q^n$ is a formal power series whose coefficients $r_\chi(n)$ are finite character sums (decidable, no integration). On this reading, CRM(FLT) = BISH. The motivic framework identifies the tension: the classical construction of θ_χ as a function on the upper half-plane uses Poisson summation—an identity of distributions on $\mathbb{R}$ (automorphic realization: CLASS). Whether Poisson summation is *essential Archimedean content* or *dispensable scaffolding* (replaceable by the formal q-expansion) is the precise locus where the constructive/classical judgment must be made.

Note: Paper 68 does *not* use Faltings's isogeny theorem. The FLT proof proceeds by contradiction (Ribet level-lowering to $N = 2$, no such form exists), never requiring an isogeny construction. The Faltings bridge is relevant to the *positive* modularity theorem (associating abelian varieties to modular forms) but not to the FLT application.

Status: Paper 68's classification stands as stated. Resolution requires a formal CRMLint Squeeze of the dihedral base case—a natural target for future work.

Function fields are BISH (Paper 69). This is the strongest confirmation of the Archimedean Thesis. Over $\mathbb{F}_q(C)$, the adèle ring has no Archimedean factor. The Langlands correspondence (Lafforgue [10]) uses moduli of shtukas (algebraic) and étale cohomology (non-Archimedean). The motivic prediction: BISH. Confirmed exactly.

New opportunity: Paper 69 treated only GL_2 and GL_n. The motivic framework predicts that *all* function-field arithmetic should be BISH, including the geometric Langlands program (Fargues–Scholze over function fields), motivic cohomology of function-field varieties, and Iwasawa theory over $\mathbb{F}_q(\!(t)\!)$. Each is a concrete test of the Archimedean Thesis.

The Archimedean Principle (Paper 70). This capstone paper stated the central claim: the logical cost of mathematics is the logical cost of $\mathbb{R}$. Paper 98 upgrades this from an empirical observation to a *structural theorem*: the Archimedean Obstruction Theorem (Theorem 5.1) proves that the cost is determined by the interaction with $\mathbb{R}$ via realization functors and comparison isomorphisms.

DPT reversal (Papers 72–74). These papers proved the DPT axioms are *necessary*, not merely sufficient. The motivic framework explains why: each DPT axiom encodes a comparison isomorphism. Since the comparison isomorphisms are the *only* entry points for CLASS content (by the Archimedean Obstruction Theorem), any axiom scheme that makes cycle-search BISH must neutralize exactly these comparisons. The DPT axioms are the unique minimal such scheme because each addresses exactly one comparison.

A.4 Phase 5–7: Generative Phase (Papers 76–97)

The generative phase tested the Archimedean Thesis as a *predictive instrument*. We review each application through the motivic lens.

CRMLint (Paper 76). The automated analyzer implements the motivic decomposition mechanically: Scaffold = define the motive and realizations; X-Ray = identify which comparisons pass through $\mathbb{R}$; Excise = replace Archimedean comparisons with algebraic substitutes; Squeeze = verify the residual is BISH.

Infrastructure arc (Papers 80–83). The Gauss–Manin connection (P80), flat sections (P81), Kovacic's algorithm (P82), and generic Picard number (P83) all operate in the de Rham realization. The motivic prediction: BISH for the algebraic content, with CLASS entering only when the topological monodromy interpretation (Betti realization) is invoked. Confirmed: the CLASS boundary nodes are exactly Ehresmann–Gauss–Manin (de Rham $\to$ Betti comparison), Deligne's monodromy theorem (Betti realization), and Noether–Lefschetz (Hodge realization).

Hodge campaign (Papers 84–93).

Trace vanishing $\tau_+ = 0$ (Papers 85, 89, 92, 93). The trace τ_+ is computed in the de Rham realization (Gauss–Manin connection, Griffiths pole reduction: all algebraic). The motivic prediction: $\tau_+ = 0$ should be BISH. Confirmed: proved by `ring` in $\mathbb{Z}[a, b, c]$. The structural explanation (Paper 93, Varchenko exponent cancellation) is also BISH: it uses only the hypergeometric structure of the de Rham connection.

Kani–Rosen splitting (Paper 86). The Kani–Rosen isogeny $\mathrm{Jac}(C) \sim A^2$ is algebraic (idempotent relation in the endomorphism algebra). Ribet's theorem asserts Hodge classes are algebraic for powers of abelian surfaces. Ribet's proof uses the Mumford–Tate group, which is Hodge-theoretic (Betti-side). However, on the palindromic locus, the splitting $\mathrm{Jac} \sim A^2$ with $\mathrm{End}(A) = \mathbb{Z}$ determines $\mathrm{MT}(A) = \mathrm{GSp}_4$ by Zarhin's theorem—an algebraic group-theoretic computation (BISH)—and Weyl's theorem on invariants then constructs the algebraic cycles without integration. The Archimedean obstruction is bypassed because the splitting provides algebraic access to the Hodge classes.

Hodge horizon = WLPO (Paper 87). The motivic explanation: the uniform Hodge test requires the Betti $\leftrightarrow$ de Rham comparison (period map: integration over $\mathbb{R}$). Testing whether a period integral vanishes is a real-number equality test = WLPO. With CM metadata, the period map factorizes through algebraic data, bypassing $\mathbb{R}$: cost drops to BISH.

Palindromic bifurcation (Papers 86–89). The motivic framework identifies the bifurcation as a *change of realization functor*: on $\{a = c\}$, the D_8-symmetry provides an algebraic (de Rham/étale) path to algebraicity; off $\{a = c\}$, the only available path goes through the Betti realization (VHC, period integrals), incurring CLASS.

BSD campaign (Papers 95–96).

Rank/Ш decoupling. The motivic explanation is sharp: rank verification (2-descent) operates in the étale realization ($H^1(\mathbb{Q}, E[2])$: non-Archimedean). Ш finiteness requires the automorphic realization (L-function, analytic continuation through the Archimedean Γ-factor) and the Betti realization (Heegner points, complex multiplication, period integrals). Motivic prediction: rank = BISH, Ш = CLASS. Confirmed exactly by Papers 95–96.

Root number bifurcation (Paper 96). For $w = +1$ (even functional equation), detection via modular symbols $L(E, 1)/\Omega^+$ avoids the Betti realization (the ratio is algebraic: BISH). For $w = -1$ (odd functional equation), detection requires $L'(E, 1)$ via the Gross–Zagier formula, which passes through the Betti realization (Heegner point heights involve real integration: CLASS). The motivic prediction matches the root number bifurcation perfectly: w controls whether the L-value computation can stay in the de Rham realization ($w = +1$) or must pass through the Betti realization ($w = -1$).

CY3 Griffiths group (Paper 94). Detection of non-trivial normal functions via the Walcher source term $\delta(z) \neq 0$ operates in the de Rham realization (Picard–Fuchs differential equation: algebraic). Existence of the Abel–Jacobi image requires the Betti realization (integration over real cycles: CLASS). The motivic framework perfectly predicts the BISH/CLASS split.

B Controversies Resolved

The motivic framework resolves several issues that arose during the program.

B.1 Classical.choice in $\mathbb{R}$-theorems

Issue. Every Lean theorem mentioning $\mathbb{R}$ shows `Classical.choice` in `#print axioms`, even when the proof uses only `ring` or `norm_num`. This created confusion about whether the theorems were genuinely constructive.

Resolution. `Classical.choice` is a *Lean infrastructure artifact* from Mathlib's construction of $\mathbb{R}$ (Cauchy completion, quotient field, decidable equality on supports). The CRM signature is determined by the *proof content* (which realization functors and comparison isomorphisms

are used), not by the Lean axiom checker. A theorem proved by `ring` operates in the de Rham realization (algebraic polynomial identity): BISH, regardless of what `#print axioms` reports. This distinction—proof content vs. infrastructure artifact—is the central methodological insight of the program.

B.2 DPT vs. Fargues–Scholze

Issue. The DPT axiom scheme (Papers 50–53, 72–74) and the Fargues–Scholze geometrization of the local Langlands correspondence appear to address similar problems (decidability in arithmetic geometry) from different directions. Are they competing frameworks?

Resolution. They are *complementary and logically forced.* The DPT axioms neutralize the three Archimedean comparison isomorphisms (cycle class, spectral, polarization). Fargues–Scholze operates in the non-Archimedean world (p-adic Hodge theory, diamonds, perfectoid spaces), avoiding $\mathbb{R}$ entirely. In motivic language: DPT provides algebraic oracles that bypass the Archimedean comparisons; Fargues–Scholze provides a framework where those comparisons never arise. They partition the problem along the same boundary—the Archimedean place—from opposite sides.

However, Fargues–Scholze does not achieve BISH for free. It trades the analytic cost of $\mathbb{R}$ for the topological cost of non-Archimedean completeness: completing the algebraic closure of $\mathbb{Q}_p$ to form $\mathbb{C}_p$, and performing functional analysis on Banach and Fréchet spaces over $\mathbb{C}_p$, requires WKL_0 (for compactness and completions of p-adic spaces) and potentially Dependent Choice (for sequential completions and topological limits). The Archimedean CLASS obstruction is eliminated, but a WKL_0 residue remains. The Conservation Conjecture (Paper 67) predicts that this WKL_0 residue is optimal: the non-Archimedean completeness cost is the irreducible minimum for results that require passing to infinite extensions.

B.3 FLT vs. BSD: why their CRM profiles diverge

Issue. Both FLT and BSD concern elliptic curves over $\mathbb{Q}$, yet their CRM profiles are qualitatively different. FLT has an enormous BISH core with a single contested node (Section A.3); BSD is structurally CLASS with CLASS content distributed across multiple components (Paper 95). Why?

Resolution. FLT asserts a *non-existence* result: the proof proceeds by contradiction (Ribet level-lowering to $N = 2$, no such form exists). The entire argument from Frey curve to contradiction routes through étale and crystalline realizations (BISH), with CLASS entering only at the base case anchor (Hecke theta series—contested, see Section A.3). BSD asserts an *equality* between analytic and algebraic invariants ($\mathrm{ord}_{s=1} L(E, s) = \mathrm{rk}\, E(\mathbb{Q})$). The L-function arises from the automorphic realization via the Mellin transform at the Archimedean place, and the finiteness of Ш requires Kolyvagin's Euler system (Chebotarev density: CLASS). The CLASS content is distributed across both the analytic and the Galois-cohomological sides. The structural difference: for FLT, the potential CLASS content is *concentrated in a single node* that Paper 68 argues can be bypassed; for BSD, the CLASS content is *distributed and structural* (the L-function *is* the Archimedean realization). This explains why Paper 96 found no bifurcation for BSD: there is no single node to excise.

B.4 Hodge bifurcation vs. BSD rigidity

Issue. The Hodge conjecture admits a continuous constructive/classical bifurcation (palindromic locus: Papers 86–89), while BSD admits no such bifurcation (Paper 96). Why?

Resolution. The explanation is geometric. Bifurcation requires a continuous family of motives parametrized by a moduli space of dimension ≥ 2, with a positive-dimensional sublocus

where extra symmetry provides an algebraic bypass. For Hodge: the universal hyperelliptic family has 3 parameters (a, b, c), and the palindromic locus $\{a = c\}$ has codimension 1 (dimension 2). For BSD: the moduli of elliptic curves is the j-line (dimension 1), and proper algebraic subvarieties are isolated points (CM curves). No continuous sublocus exists to support a bifurcation. The motivic framework suggests a necessary condition: bifurcation requires dim(moduli) ≥ 2 and a positive-dimensional symmetry sublocus providing an algebraic bypass. Whether this condition is also sufficient remains open.

C New Opportunities

The systematic reinterpretation reveals several concrete research targets.

C.1 Constructive p-adic Langlands

Prediction. The p-adic Langlands correspondence (Breuil, Colmez, Emerton) operates entirely in non-Archimedean realizations: p-adic Galois representations $\leftrightarrow$ p-adic Banach space representations of $\mathrm{GL}_2(\mathbb{Q}_p)$. The motivic framework predicts: $\mathsf{BISH} + \mathsf{WKL}_0$ (WKL for inverse limits in Banach space completions). No CLASS content should arise. *This is untested.* A CRMLint Squeeze of Colmez's proof would be a natural Paper 99 or beyond.

C.2 Condensed mathematics as realization functor

Prediction (revised). Scholze's condensed mathematics replaces topological spaces with condensed sets (sheaves on extremally disconnected profinite sets). A naïve hope is that this avoids the Betti realization's dependence on point-set topology over $\mathbb{R}$, bypassing the Archimedean obstruction. This is *too optimistic*: the category of condensed abelian groups has enough projectives because extremally disconnected spaces are projective in compact Hausdorff spaces (Gleason's theorem), whose proof requires the Boolean Prime Ideal Theorem (CLASS). Condensed mathematics trades analytic CLASS axioms (real integration) for set-theoretic CLASS axioms (infinite choice). The CRM cost is not reduced.

However, Clausen–Scholze's *light condensed sets*—sheaves on totally disconnected spaces of countable cofinality—avoid the Boolean Prime Ideal Theorem entirely. The residual cost is sequential completions, which require either WKL_0 (compactness for inverse limits of finite modules) or DC (dependent choice for sequential constructions). Since WKL_0 and DC are independent over BISH, determining which governs light condensed cohomology is itself a CRM question. The revised prediction: light condensed de Rham $\to$ light condensed Betti has cost $\mathsf{BISH} + \mathsf{WKL}_0$ or $\mathsf{BISH} + \mathsf{DC}$, strictly below the CLASS cost of the classical comparison. The Conservation Conjecture (Paper 67) becomes testable: is this residual cost optimal?

C.3 Motivic Iwasawa theory

Prediction. Classical Iwasawa theory operates over the cyclotomic $\mathbb{Z}_p$-extension (non-Archimedean). The Iwasawa Main Conjecture relates p-adic L-functions to Selmer groups (étale: BISH). The standard construction of p-adic L-functions via interpolation of complex L-values passes through the Archimedean place (CLASS). To achieve $\mathsf{BISH} + \mathsf{WKL}_0$, one must sever this tether to $\mathbb{C}$ entirely.

The correct approach: Kato's Euler system [19] constructs zeta elements z_n purely algebraically in K_2 of modular curves and maps them to Galois cohomology via the p-adic regulator. The p-adic L-function is defined cohomologically as the element generated by this system, and the Selmer group bound is algebraic. The WKL_0 node enters precisely at the inverse limit over the cyclotomic tower $\mathbb{Q}(\mu_{p^\infty})$ to form the Iwasawa module over $\mathbb{Z}_p[\![T]\!]$ (compactness for inverse limits of finite modules). The motivic prediction: $\mathsf{BISH} + \mathsf{WKL}_0$ for the algebraic Iwasawa Main

Conjecture via Kato, with CLASS entering only if one insists on the interpolation definition of p-adic L-functions. Paper 96's rank-0 test case (11a1, $w = +1$) is suggestive: the modular symbol ratio $L(E, 1)/\Omega^+$ is already algebraic.

C.4 Trace formula excision

Assessment: irreducible CLASS. The Arthur–Selberg trace formula is the universal CLASS bottleneck (Section 10). The motivic framework identifies the precise obstruction: the spectral decomposition of $L^2(G(\mathbb{Q})\backslash G(\mathbb{A}))$ at the Archimedean place $G(\mathbb{R})$. Over function fields $\mathbb{F}_q(C)$, the geometric trace formula (Frenkel–Langlands–Ngô) replaces spectral analysis with the Hitchin fibration over the base curve C, and the Archimedean place is absent. Over number fields, this bypass *cannot* work: the "base curve" is $\mathrm{Spec}(\mathcal{O}_K) \cup \{\infty\}$, and the Archimedean places $\{\infty\}$ are structurally required by Arakelov geometry to compactify the arithmetic curve. There is no purely algebraic "arithmetic Hitchin fibration" that ignores infinity. The global trace formula over $\mathbb{Q}$ is an irreducible CLASS node.

The remaining avenue is *local*: the Fargues–Scholze geometrization of the *local* Langlands correspondence uses the Fargues–Fontaine curve (purely p-adic, no Archimedean place). This achieves BISH + WKL_0 for local results (Section B.2), but does not excise the Archimedean place from *global* automorphic arguments.

C.5 Higher-dimensional Hodge bifurcations

Prediction. The palindromic bifurcation (Papers 86–89) requires moduli dimension ≥ 2. For higher-dimensional families:

- Abelian varieties of dimension g have moduli dimension $g(g+1)/2$. For $g \geq 3$, positive-dimensional symmetry loci should exist.
- Calabi–Yau threefolds: the mirror quintic family (Paper 94) has 1 modulus (ψ). The full quintic family has $h^{2,1} = 101$ moduli, but seeking palindromic loci in 101-dimensional parameter space is computationally intractable (Picard–Fuchs ideal expression swell). The viable target is the *Dwork pencil* $x_1^5 + \cdots + x_5^5 - 5\psi x_1x_2x_3x_4x_5 = 0$, where the discrete symmetry group $(\mathbb{Z}/5\mathbb{Z})^3 \rtimes S_5$ plays the role of D_8 from the genus-4 case, allowing algebraic cycle isolation on symmetric resolutions.
- K3 surfaces: moduli dimension 20. The Kuga–Satake construction (Paper 55) builds an abelian variety A_{KS} from the even Clifford algebra of the intersection lattice—pure finite-dimensional quadratic algebra (BISH). By the Shioda–Inose theorem, singular K3 surfaces ($\rho = 20$) have rank-2 transcendental lattice, and A_{KS} splits into CM elliptic curves. The CRM cost of Hodge detection drops to BISH on this locus. The bifurcation is structurally parallel to Paper 86 (Kani–Rosen splitting at the palindromic locus).

C.6 Explicit Langlands beyond GL_2

Prediction. Paper 78 (wild Langlands for $\mathrm{GL}_2(\mathbb{Q}_2)$) already established that the local correspondence is BISH for character-trace matching. The motivic framework predicts this extends to GL_n: the local Langlands correspondence for GL_n over p-adic fields involves only non-Archimedean data (Bernstein–Zelevinsky classification, Bushnell–Kutzko types, compact induction from representations of compact open subgroups). *Caution:* The Harris–Taylor proof [21] globalizes the local field to exploit global Shimura varieties and the Arthur–Selberg trace formula, immediately injecting Archimedean CLASS content. To verify the BISH prediction, one must use the purely local Bushnell–Kutzko construction, which classifies supercuspidal representations via compact induction from finite-dimensional representations of compact open subgroups

(matrices over finite quotients $\mathcal{O}_K/\mathfrak{p}^m$). This is algebraic and BISH, but extends Paper 78 only in combinatorial complexity, not in CRM phenomena.

D Opportunity Summary

Target	Current level	Predicted level	Method
p-adic Langlands (Colmez)	Unaudited	$\mathsf{BISH} + \mathsf{WKL}_0$	CRMLint Squeeze
Light condensed comparison	Unaudited	$\mathsf{BISH} + \mathsf{WKL}_0/\mathsf{DC}$	Gleason bypass
p-adic BSD (Kato Euler system)	CLASS	$\mathsf{BISH} + \mathsf{WKL}_0$	Avoid complex L-values
Geometric trace formula	CLASS	CLASS (irreducible)	Arakelov compactification
CY3 Hodge ($h^{2,1} = 101$)	CLASS	Bifurcation	Palindromic loci
K3 Kuga–Satake	CLASS	Bifurcation	Algebraic metadata
GL_n local Langlands	CLASS	BISH (local)	Bushnell–Kutzko
Fontaine–Mazur (purely p-adic)	LPO	$\mathsf{BISH} + \mathsf{WKL}_0$	Avoid complex embeddings

Table 6: Research targets identified by the motivic reinterpretation. "Current level" is the CRM cost of existing proofs; "predicted level" is the motivic framework's prediction for the optimal achievable cost.

The motivic CRM architecture transforms the program from a catalog of individual results into a *predictive theory*. Each entry in Table 6 is a falsifiable prediction: either the target theorem can be proved at the predicted level, confirming the Archimedean Thesis, or it cannot, revealing a new obstruction beyond the Archimedean place. Either outcome advances the program.

E Feasibility Assessment and Roadmap

Not all targets in Table 6 are equally feasible. We classify them by readiness for formalization.

E.1 Prime targets (immediately formalizable)

1. **K3 Kuga–Satake bifurcation.** The even Clifford algebra construction is finite-dimensional quadratic algebra (BISH). The Shioda–Inose theorem at Picard rank $\rho = 20$ splits A_{KS} into CM elliptic curves, reducing Hodge detection to BISH. This directly extends the Hodge campaign (Papers 84–93) to a new geometric setting and is the strongest validation target.

2. **Iwasawa Main Conjecture via Kato.** The algebraic K-theory zeta elements and p-adic regulator avoid complex L-values entirely. The WKL_0 node (cyclotomic inverse limit) matches the Conservation Conjecture prediction. Clean target for the CAS $\to$ Lean pipeline.

3. **The Hecke theta series resolution.** The sharpest open problem within the existing program. Paper 68 classifies $\mathsf{CRM}(\mathrm{FLT}) = \mathsf{BISH}$ via the Kisin $p = 2$ dihedral bypass. The contested node—whether Hecke theta series require Poisson summation (CLASS) or can be defined via the q-expansion principle on $Y_0(N)$ as a scheme (BISH)—is resolvable by formal methods (Katz, Deligne–Rapoport). Either outcome is a theorem.

E.2 Technically bottlenecked (sound logic, missing infrastructure)

1. **p-adic Langlands (Colmez).** Conceptually $\mathsf{BISH} + \mathsf{WKL}_0$, but formalization requires constructive functional analysis on p-adic Banach and Fréchet spaces (absent from Math-

lib). The WKL_0 boundary at spherical completeness of $\mathbb{C}_p$ is precise; the infrastructure to formalize it does not yet exist.

2. **CY3 Hodge (Dwork pencil).** The discrete symmetry group $(\mathbb{Z}/5\mathbb{Z})^3 \rtimes S_5$ provides the algebraic bypass mechanism, but the Picard–Fuchs differential equation for the mirror quintic is a fourth-order ODE with 101 parameters in the general case. Only the 1-parameter Dwork pencil is computationally tractable.

3. **Fontaine–Mazur (purely p-adic).** Constructing global Galois representations without complex embeddings requires Emerton's completed cohomology, which is defined via p-adic limits of *Betti* cohomology of Shimura varieties. No purely local method currently exists to bypass this.

E.3 Confirmed irreducible

1. **Geometric trace formula over $\mathbb{Q}$.** The Arakelov compactification of $\mathrm{Spec}(\mathcal{O}_K)$ structurally requires Archimedean places. No "arithmetic Hitchin fibration" that ignores infinity exists or can exist over number fields. The global trace formula is an irreducible CLASS node.

2. **Full condensed mathematics.** Gleason's theorem (projective covers in compact Hausdorff spaces) requires the Boolean Prime Ideal Theorem (CLASS). Only Clausen–Scholze's *light* condensed sets avoid this, trading full CLASS for WKL_0/DC.

E.4 Candidate Papers 99–100

Several roadmaps have been proposed. We record two principal options with assessment.

Option A: Resolution papers.

- *Paper 99:* The Hecke Theta Series Squeeze—resolve whether the dihedral base case of modularity is BISH or CLASS via the q-expansion principle. Settles a disputed classification within the program.

- *Paper 100:* The Kuga–Satake Bifurcation—extend the Hodge campaign to K3 surfaces via Shioda–Inose at $\rho = 20$. Geometric capstone concluding with the Conservation Conjecture as the program's principal open problem.

Option B: Verification papers.

- *Paper 99:* p-adic Squeeze—CRMLint Squeeze of the Bertolini–Darmon–Prasanna p-adic Gross–Zagier formula, or of Colmez's p-adic local Langlands for $\mathrm{GL}_2(\mathbb{Q}_p)$ (the latter requires p-adic Banach space infrastructure not yet in Mathlib).

- *Paper 100:* Conservation Law and CRMLint 1.0—formally state the Conservation Conjecture and release an automated CRM audit tool as a Lean 4 metaprogram.

Assessment. Option A produces theorems with definite outcomes (the Hecke theta series question resolves either way; the Kuga–Satake bifurcation is a concrete new result). Option B confirms predictions (the p-adic squeeze verifies Table 6) and builds tooling (CRMLint automation). However, the CRMLint 1.0 ambition must be tempered: the entire program demonstrates that CRM classification requires mathematical judgment—`Classical.choice` appears in BISH theorems, and a naïve dependency tracer would misclassify every $\mathbb{R}$-theorem as CLASS. A fully automated `#crm_audit` macro that correctly outputs BISH/CLASS decompositions is beyond current capability; a *semi*-automated tool that flags Archimedean dependencies for human review is realistic.

References

[1] E. Bishop, *Foundations of Constructive Analysis*, McGraw-Hill, 1967.

[2] D. Bridges and F. Richman, *Varieties of Constructive Mathematics*, LMS Lecture Notes 97, Cambridge, 1987.

[3] H. Ishihara, Reverse mathematics in Bishop's constructive mathematics, *Phil. Sci. Cahier Spécial* 6 (2006), 43–59.

[4] K. Ribet, Hodge classes on certain types of abelian varieties, *Amer. J. Math.* 105 (1983), 523–538.

[5] E. Kani and M. Rosen, Idempotent relations and factors of Jacobians, *Math. Ann.* 284 (1989), 307–327.

[6] A. Wiles, Modular elliptic curves and Fermat's Last Theorem, *Ann. of Math.* 141 (1995), 443–551.

[7] R. Taylor and A. Wiles, Ring-theoretic properties of certain Hecke algebras, *Ann. of Math.* 141 (1995), 553–572.

[8] V. Kolyvagin, Euler systems, in: *The Grothendieck Festschrift*, vol. II, Birkhäuser, 1990, 435–483.

[9] B. Gross and D. Zagier, Heegner points and derivatives of L-series, *Invent. Math.* 84 (1986), 225–320.

[10] L. Lafforgue, Chtoucas de Drinfeld et correspondance de Langlands, *Invent. Math.* 147 (2002), 1–241.

[11] A. Grothendieck, Standard conjectures on algebraic cycles, in: *Algebraic Geometry, Bombay 1968*, Oxford, 1969, 193–199.

[12] U. Jannsen, Motives, numerical equivalence, and semi-simplicity, *Invent. Math.* 107 (1992), 447–452.

[13] J.-M. Fontaine and G. Laffaille, Construction de représentations p-adiques, *Ann. Sci. ENS* 15 (1982), 547–608.

[14] J. Arthur, An introduction to the trace formula, in: *Harmonic Analysis, the Trace Formula, and Shimura Varieties*, Clay Math. Proc. 4, AMS, 2005, 1–263.

[15] B. Mazur, An introduction to the deformation theory of Galois representations, in: *Modular Forms and Fermat's Last Theorem*, Springer, 1997, 243–311.

[16] P. Berthelot, *Cohomologie Cristalline des Schémas de Caractéristique* $p > 0$, Lecture Notes in Math. 407, Springer, 1974.

[17] V. Drinfeld, Langlands' conjecture for GL(2) over functional fields, in: *Proc. ICM Helsinki*, 1978, 565–574.

[18] P. Deligne, Cohomologie étale (SGA $4\frac{1}{2}$), Lecture Notes in Math. 569, Springer, 1977.

[19] K. Kato, p-adic Hodge theory and values of zeta functions of modular forms, *Astérisque* 295 (2004), 117–290.

[20] M. Kisin, Moduli of finite flat group schemes, and modularity, *Ann. of Math.* 170 (2009), 1085–1180.

[21] M. Harris and R. Taylor, *The Geometry and Cohomology of Some Simple Shimura Varieties*, Annals of Math. Studies 151, Princeton, 2001.

[22] Y. Choquet-Bruhat, Théorème d'existence pour certains systèmes d'équations aux dérivées partielles non linéaires, *Acta Math.* 88 (1952), 141–225.

[23] The Mathlib Community, *Mathlib: the Lean mathematical library*, 2024. DOI: 10.5281/zenodo.10886692.

[24] P. C.-K. Lee, Standard conjecture D is necessary for morphism decidability (Paper 73, CRM Series format guide), Zenodo, 2026. DOI: 10.5281/zenodo.18765700.

[25] P. C.-K. Lee, The omniscience cost of the Hodge conjecture: a CRM analysis (Paper 87), Zenodo, 2026. DOI: 10.5281/zenodo.18809903.

[26] P. C.-K. Lee, The BSD squeeze: CRMLint decomposition of the Gross–Zagier–Kolyvagin proof (Paper 95), Zenodo, 2026. DOI: 10.5281/zenodo.18821019.

[27] P. C.-K. Lee, The root number bifurcation: CRM cost of BSD detection controlled by the functional equation (Paper 96), Zenodo, 2026. DOI: 10.5281/zenodo.18820969.