

Breed Predispositions to Disease in Dogs and Cats

Third Edition

Alex Gough · Alison Thomas · Dan O'Neill



WILEY Blackwell

Breed Predispositions to Disease in Dogs and Cats

Breed Predispositions to Disease in Dogs and Cats

THIRD EDITION

Alex Gough

Bath Veterinary Referrals, Bath, UK

Alison Thomas

Blue Cross Animal Hospital Victoria, London, UK

Dan O'Neill

Royal Veterinary College, London, UK

WILEY Blackwell

This edition first published 2018
© 2018 John Wiley & Sons Ltd

Edition History

John Wiley & Sons (1e, 2004)
John Wiley & Sons (2e, 2010)

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of Alex Gough, Alison Thomas and Dan O'Neill to be identified as the authors of this work has been asserted in accordance with law.

Registered Offices

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA
John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Office

9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Gough, Alex, author. | Thomas, Alison, 1964– author. | O'Neill, Dan (Dan G.), author.
Title: Breed predispositions to disease in dogs and cats / Alex Gough, Alison Thomas, Dan O'Neill.
Description: Third edition. | Hoboken, NJ : Wiley, [2018] | Includes bibliographical references. |
Identifiers: LCCN 2017043483 (print) | LCCN 2017055418 (ebook) | ISBN 9781119225553 (pdf) |
ISBN 9781119225577 (epub) | ISBN 9781119225546 (pbk.)
Subjects: LCSH: Dogs–Diseases. | Cats–Diseases. | MESH: Dog Diseases–genetics | Cat Diseases–genetics |
Genetic Predisposition to Disease | Pedigree
Classification: LCC SF991 (ebook) | LCC SF991 .G67 2018 (print) | NLM SF 991 | DDC 636.7/089607–dc23
LC record available at <https://lccn.loc.gov/2017043483>

Cover Design: Wiley

Cover Image: (Center right) Labrador retriever © Alison Thomas; (Top left) Cat: Seal point Ragdoll, (Bottom left) Dog: French Bulldog © Dan O'Neill

Set in 9/11pt Minion by SPi Global, Pondicherry, India

Alex Gough

To loved ones, friends and colleagues, and of course to my wife Naomi and daughter Abigail for bearing with me through another huge project.

Alison Thomas

To all the wonderful colleagues I have had the pleasure of working with over the years. Most of all to my partner Richard, and children Tom and Harry, for making my life so much fun.

Dan O'Neill

To my wife, best friend and inspiration, Joanne. And to my three children and lights of my life, Alistair, Megan and Clodagh. Thank you each for being you.

This book is also dedicated to all those vets suffering from mental illness, and particularly to those who have lost their lives due to this common condition affecting our profession.

Contents

<i>Author Biographies</i>	<i>xi</i>
---------------------------	-----------

<i>Foreword</i>	<i>xiii</i>
-----------------	-------------

<i>Preface</i>	<i>xv</i>
----------------	-----------

<i>Abbreviations</i>	<i>xvii</i>
----------------------	-------------

Introduction	1	Bolognese	36
Basic and Clinical Genetics	1	Bolonka Zwetna	36
Epidemiology	7	Border Collie	36
Longevity	10	Border Terrier	38
Methods	14	Borzoi	38
		Boston Terrier	39
		Bouvier Des Flandres	40
Part I: Dog Breeds	17	Boxer	41
Affenpinscher	19	Boykin Spaniel	46
Afghan Hound	19	Briard	46
African Boerboel	20	British Bulldog	46
Airedale Terrier	20	Brittany Spaniel	46
Akbash	21	Brussels Griffon	47
Akita Inu	21	Buhund	47
Alaskan Husky	21	Bull Terrier	47
Alaskan Klee Kai	21	Bull Terrier – Miniature	48
Alaskan Malamute	21	Bulldog – American	49
Alsatian	22	Bulldog – English	49
American Bulldog	22	Bullmastiff	51
American Cocker Spaniel	22	Cairn Terrier	52
American Eskimo	22	Cane Corso	53
American Pit Bull Terrier	22	Catahoula Leopard Dog	53
American Staffordshire Terrier	23	Cavalier King Charles Spaniel	54
Anatolian Shepherd Dog	24	Chesapeake Bay Retriever	56
Australian Cattle Dog	24	Chihuahua	56
Australian Kelpie	25	Chin	57
Australian Shepherd Dog	25	Chinese Crested Dog	58
Australian Silky Terrier	26	Chinese Shar Pei	58
Australian Terrier	27	Chow Chow	58
Basenji	27	Clumber Spaniel	59
Basset Hound	28	Cocker Spaniel	60
Beagle	29	Collies – Rough and Smooth	64
Bearded Collie	31	Coonhound	66
Beauceron	32	Corgi	66
Bedlington Terrier	32	Coton De Tulear	66
Belgian Shepherd Dog	32	Curly-Coated Retriever	67
Berger Blanc Suisse	32	Dachshund	67
Bernese Mountain Dog	33	Dalmatian	70
Bichon Frise	34	Dandie Dinmont Terrier	72
Bloodhound	35	Danish Chicken Dog	72
		Danish/Swedish Farmdog	72

Deerhound	72	Irish Red and White Setter	111
Deutscher Wachtelhund	72	Irish Setter	111
Dobermann Pinscher	72	Irish Terrier	113
Dogo Argentino	76	Irish Water Spaniel	114
Dogue De Bordeaux	77	Irish Wolfhound	114
Dutch Partridge Dog	77	Italian Greyhound	116
Dutch Sheepdog	77	Italian Mastiff	116
Dutch Spaniel	77	Italian Spinone	116
Elkhound	77	Jack Russell Terrier	117
English Bulldog	77	Jämthund	119
English Cocker Spaniel	78	Japanese Akita Inu	119
English Setter	78	Japanese Chin	121
English Shepherd	79	Japanese Shiba Inu	121
English Springer Spaniel	79	Japanese Tosa	121
English Toy Spaniel	79	Jindo	122
Entlebucher Mountain Dog	79	Keeshond	122
Eskimo Dog	79	Kelpie	123
Estrela Mountain Dog	80	Kerry Blue Terrier	123
Finnish Hound	80	King Charles Spaniel	123
Finnish Lapphund	80	Klee Kai	124
Finnish Spitz	80	Kooiker Dog	124
Flat-Coated Retriever	80	Korean Jindo	124
Fox Terrier	81	Kromfohrländer	124
Foxhound	83	Kuvasz	124
French Alpine Mastiff	83	Labrador Retriever	124
French Bulldog	83	Lakeland Terrier	131
French Mastiff	85	Lancashire Heeler	131
French Spaniel	85	Lapland Reindeer Dog	131
Frisian Water Dog	85	Lapphund	131
Galgos Español	85	Leonberger	131
German Pinscher	85	Lhasa Apso	132
German Shepherd Dog	85	Lowchen	134
German Spaniel	93	Mah Thai	134
Glen of Imaal Terrier	93	Malamute	134
Golden Retriever	93	Maltese	134
Gordon Setter	99	Manchester Terrier	136
Grand Bleu De Gascogne Hound	100	Mastiff	136
Great Dane	100	Mastiff (Unspecified)	136
Great Pyrenees	104	Mcnab Shepherd	137
Greater Swiss Mountain Dog	104	Mexican Hairless	138
Greyhound	104	Miniature Bull Terrier	138
Greyhound – Spanish	107	Miniature Pinscher	138
Griffon Bruxellois	107	Mountain Dog	139
Hamilton Hound	107	Münsterländer – Pointer	139
Havanese	108	Neapolitan Mastiff	139
Hokkaido	108	New Zealand Huntaway	140
Hovawart	108	Newfoundland	140
Hungarian Kuvasz	109	Norfolk Terrier	142
Hungarian Puli	110	Norwegian Buhund	143
Hungarian Vizsla	110	Norwegian Elkhound	143
Hungarian Water Dog	111	Norwegian Lundehund	144
Husky	111	Norwegian Sheepdog	144

Norwich Terrier	145	Shiba Inu	195
Nova Scotia Duck Tolling Retriever	145	Shih Tzu	195
Ogar Polski	146	Siberian Husky	198
Old English Mastiff	146	Silken Windhound	199
Old English Sheepdog	146	Silky Terrier	199
Otterhound	147	Skye Terrier	199
Papillon	147	Sloughi	200
Parson Russell Terrier	148	Smooth Collie	200
Pekingese	149	Soft-Coated Wheaten Terrier	200
Petit Basset Griffon Vendeen	150	Spanish Greyhound	201
Phu Quoc Ridgeback	151	Spanish Mastiff	201
Pinscher – Miniature	151	Spanish Water Dog	201
Pointer	151	Spinone Italiano	201
Polish Hound	154	Springer Spaniel	201
Polish Lowland Sheepdog	154	Staffordshire Bull Terrier	204
Pomeranian	154	Staffordshire Bull Terrier – American	206
Poodle (Unspecified Variant)	156	Sussex Spaniel	206
Poodle – Miniature	158	Swedish Cattle Dog	206
Poodle – Standard	161	Swedish Elkhound	206
Poodle – Toy	162	Swedish Farmdog	207
Portuguese Podengo Pequeno	164	Swedish Lapphund	207
Portuguese Water Dog	164	Swedish Vallhund	207
Presa Canario	165	Swiss Mountain Dog	207
Pug	165	Tenterfield Terrier	208
Puli	168	Thai Ridgeback	208
Pyrenean Mountain Dog	168	Tibetan Mastiff	208
Pyrenean Shepherd	169	Tibetan Spaniel	209
Rat Terrier	170	Tibetan Terrier	209
Red and White Setter	170	Tosa	210
Red Kelpie	170	Treeing Walker Coonhound	210
Red Setter	170	Turkish Shepherd Dog	210
Retriever	170	Vallhund	211
Rhodesian Ridgeback	170	Vizsla	211
Rottweiler	172	Vorstehhund	211
Rough Collie	177	Wachtelhund	211
Russian Black Terrier	177	Weimaraner	211
St Bernard	177	Welsh Corgi	213
Saluki	179	Welsh Springer Spaniel	214
Samoyed	179	Welsh Terrier	214
Savoy Sheepdog	180	West Highland White Terrier	215
Schapendoes	180	Wetterhoun	218
Schipperke	181	Whippet	218
Schnauzer (Unspecified Variant)	181	White Swiss Shepherd Dog	219
Schnauzer – Giant	181	Wolfhound – Irish	220
Schnauzer – Miniature	183	Wolfspitz	220
Schnauzer – Standard	187	Yorkshire Terrier	220
Scottish Deerhound	187		
Scottish Terrier	188	Part II: Cat Breeds	225
Sealyham Terrier	190		
Sennenhund	190	Abyssinian	227
Shar Pei	190	American Shorthair	229
Shetland Sheepdog	192	Asian	229

Australian Mist	229	Maine Coon	241
Balinese	229	Manx	242
Bengal	230	Norwegian Forest Cat	243
Birman	231	Ocicat	244
British Shorthair	232	Oriental Shorthair	244
Burmese	233	Persian	245
Chartreux	236	Ragdoll	247
Chinchilla	236	Russian Blue	249
Colourpoint Persian	236	Savannah	249
Cornish Rex	236	Scottish Fold	249
Devon Rex	236	Seychellois	249
Egyptian Mau	237	Siamese	249
European Shorthair	237	Siberian Forest Cat	253
Exotic Shorthair	238	Singapura	253
Foreign Shorthair	238	Snowshoe	253
Foreign White	238	Somali	254
Havana Brown	238	Sphynx	254
Himalayan	239	Tonkinese	255
Japanese Bobtail	240	Turkish Angora	255
Javanese	240	Turkish Van	255
Korat	240		
La Perm	241	Part III: Disease Descriptions	257

Author Biographies

Alex Gough MA VetMB CertSAM CertVC PGCert MRCVS

Head of Medicine Referrals, Bath Veterinary Referrals

Alex graduated from Cambridge University Vet School in 1996 and achieved RCVS certificates in Small Animal Medicine and Veterinary Cardiology, as well as a human Postgraduate Certificate in Neuroimaging for Research from Edinburgh University. He worked in mixed, mainly small animal practice for six years, then in referral practice, seeing referrals in medicine, cardiology and neurology, with a particular interest in medical neurology. He is the author of *Differential Diagnosis in Small Animal Medicine* (2007), has written a column summarizing the latest research for the *Veterinary Times* since 2003 as well as chapters on neurology and clinical genetics in two BSAVA manuals, and sits on the advisory clinical board of a large veterinary group. In his spare time, he writes historical fiction novels and plays guitar.

Alison Thomas BVSc, Cert SAM, MRCVS

Senior Veterinary Surgeon, Victoria Hospital, Blue Cross

Clinical Leadership Team, Blue Cross

Alison graduated from Liverpool University in 1987 and has spent most of her career working in charity small animal practice. After two years in private practice in Aylesbury, she moved to Asia, spending two years in private practice in Singapore, followed by seven years at the SPCA in Hong Kong. Since 1998 she has worked at Blue Cross in Victoria, London, becoming Senior Veterinary Surgeon in 2007 and joining the newly formed Clinical Leadership Team of Blue Cross in 2016. She gained the RCVS Certificate in Small Animal Medicine in 2001, and was awarded Advanced Practitioner Status in Small Animal Medicine in 2017. In her spare time she enjoys hiking, travelling and reading.

Dan O'Neill MVB BSc(hons) GPCert(SAP) GPCert(FelP) GPCert(Derm) GPCert(B&PS) MSc(VetEpi) PhD MRCVS

Senior Lecturer, Companion Animal Epidemiology, Royal Veterinary College

After graduating from Dublin Vet School in 1987, Dan worked in industry and general practice for 22 years, latterly running his own companion animal practice in Petts Wood, Kent, for 12 years. During these years, he was awarded additional qualifications in pharmacology, general practice, dermatology, feline practice and business management. In 2009 he undertook an MSc supported by BBSRC and then a PhD supported by the RSPCA in veterinary epidemiology at the RVC to develop the VetCompass programme of primary-care veterinary clinical research. After postdoctoral posts supported by Dogs Trust and Kennel Club Charitable Trust, Dan was appointed as Senior Lecturer in Companion Animal Epidemiology at the RVC in 2017. As well as teaching and publishing, Dan focuses on expanding VetCompass internationally, with particular emphasis on breed-related health. In his spare time, he is a keen ITF taekwondo enthusiast and currently holds a 2nd Dan black belt.

Foreword

For anyone interested in the health and welfare of dogs and cats, there are few topics that engender such emotional energy, diverse opinion and heated debate as the potential negative impacts of selective breeding on canine and feline disease occurrence, and the optimal ways to manage and/or eliminate such impacts. But energy, opinion and debate can only bring true positive change when they are based on good evidence. In this respect, the third edition of *Breed Predispositions to Disease in Dogs and Cats* is hugely timely. The book comes not only at a time of increasing awareness of the impacts that breed characteristics may have on health, but also when there is growing appreciation of the glaring underuse of objective data to support traditional perceptions and opinions which have become accepted as ‘fact’ in breeding folklore and veterinary science. This book aims to remove the roles of speculation, opinion and anecdote from the discussion on breed health issues and instead to refocus and underpin these discussions based on solid evidence-based principles.

There is no doubt that the Bateson Report on pedigree dog health and its far-reaching recommendations (Bateson, 2010), the creation of the Royal College of Veterinary Surgeons’ RCVS Knowledge initiative to promote the generation and application of veterinary clinical evidence (RCVS Knowledge, 2017) and a general increase in commitment to evidence based veterinary medicine over the past decade have resulted in a greater appreciation of the need for reliable evidence on the health impacts of breed characteristics in dogs and cats. Fortunately, this awareness of the need for valid evidence has coincided with the development of exciting new tools which allow us to collect and interpret large volumes of relevant data from primary and referral veterinary practices and to analyse these in robust and less biased ways. As a result, for the first time, we are increasingly able to provide some reliable real-world context to the likely impact of breed characteristics on animal health and welfare. The development and international adoption of standardized systems of nomenclature such as the VeNom initiative (VeNom Coding Group, 2017) and ground-breaking research tools such as the Royal Veterinary College’s VetCompass Programme (VetCompass, 2017) allow researchers to explore vast amounts of clinical data from first-opinion and referral veterinary practices. These developments have transformed how we can investigate companion animal diseases and their impact on animals, their owners and their breeders. The era of ‘Big Data’ for companion animals and its impact on animal health and welfare is now truly upon us.

Realization of the powers from developments such as these means that this new edition of *Breed Predispositions to Disease in Dogs and Cats* really does herald a new and more valid perspective in our understanding of the types of disorders and their likely impact on different dog and cat breeds. The new edition has been completely rewritten using ‘evidence-based veterinary medicine’ criteria that are applied to international data and consequently provides an accurate reference resource on disease predispositions that is relevant to breeds from all corners of the world. The information is comprehensive and detailed but presented in a readily understandable and searchable format. Prevalence, odds and risk ratio values, as well as study design details, are provided so that the more motivated reader can go beyond an awareness that a predisposition has been reported and start to examine the context and strength of the reported associations. This book truly is a cornucopia of breed health information.

This third edition will become an invaluable and constant resource for students, vets, breeders, owners, scientists and indeed anyone interested in companion animal welfare. We are privileged to live at a pivotal tipping point in the generation and application of evidence for better decision-making in companion animal health. This book will play a key role in centralizing our current knowledge into a single resource, and is thereby a torch-bearer that will finally enable us to move beyond endless circular discussion to positive action that will benefit the welfare of our cats and dogs.

*Professor David Church BVSc PhD MACVSc FHEA MRCVS
Deputy Principal
and Professor of Small Animal Studies,
The Royal Veterinary College*

Preface

It is widely accepted that almost all dog and cat breeds have specific diseases to which they are particularly prone (i.e. predisposed). Indeed, many textbooks and published research papers include lists of breed predispositions as a standard feature when describing specific disease conditions. To extend this focus, the first edition of *Breed Predispositions to Disease in Dogs and Cats*, published in 2004, aimed to provide a single reference resource for breed predispositions that would better illuminate our understanding of breed health (Gough & Thomas, 2004). The concept for the first edition was born during discussions between the two original authors (Alex Gough and Alison Thomas) while preparing for their RCVS Certificate in Small Animal Medicine exams in 2001. This book was the first of its kind to focus purely on breed-specific predispositions and was widely welcomed by academics, veterinarians, breeders and owners. That original edition was compiled mainly from secondary sources of evidence such as textbooks, reviews and conference proceedings and did not provide detailed reference citations for all the breed–disease combinations reported. The second edition, published in 2010, redressed many of these shortcomings and was updated with more recent publications while also ensuring that every cited disease had at least one supporting reference. However, much of the disease information still came from secondary sources such as textbooks, review articles and conference proceedings. The implication of this was that the second edition was substantially reliant on expert opinion. At that time, almost a decade ago, this approach may have been acceptable, but as we progress into the modern age of evidence-based veterinary medicine (EBVM), expert opinion is now generally considered to be weak evidence, and reliance should instead be placed on the results of original research (Holmes & Ramey, 2007).

In consequence, preparations began for a third edition that would have a strong emphasis on improved academic rigour, better compliance with the modern principles of EBVM and a sound epidemiological infrastructure. To meet these lofty aspirations, Alex and Alison enlisted Dr Dan O'Neill to join as a third co-author, to ensure high epidemiological standards and also to reduce the individual workload for each author. Dan is an epidemiologist working on the VetCompass Programme at the Royal Veterinary College (VetCompass, 2017), but with that rare academic attribute of essentially still being a general veterinary practitioner. We are confident that this third edition has achieved our academic goals – but we must also sadly report that we failed in our aspiration to reduce the workload. Indeed, it was quite the reverse, as the new perspectives introduced by Dan entailed a complete rewrite. Ah well, you can't win them all – but we do hope that you enjoy the end result of our combined labours.

For this new edition, we have consulted and referenced primary sources of evidence almost exclusively (i.e. the original published papers that reported the primary research) and have restricted inclusion to just those diseases where primary research identified sufficient evidence for the existence of a breed predisposition. As might be expected, this new approach led to some challenges and several discussions between the authors on the optimal threshold of evidence for disease inclusion. During our rewrite, we became painfully aware of just how little evidence actually exists for much of what we may 'believe' to be true about companion animal health. When we examined the literature closely, many of the predispositions commonly reported as 'knowns' in textbooks and introductions to peer-reviewed publications had very little, if any, reliable supporting evidence. This realization reinforced our determination that the new edition should follow rigorous evidence-based principles, but it also meant that the new book would entail a total rewrite, with substantial work required to freshly identify those diseases with and without a solid evidence base. We found ourselves painfully deleting many conditions that had been included in the previous book based on expert opinion but which lacked adequate evidence. 'Believing in a predisposition' and 'having evidence for a predisposition' are not always the same thing. The positive side of our new EBVM approach, however, was a refreshing discovery that our new detailed trawl of the primary literature led us to identify many new breed–disease combinations that had not made it into the previous two editions. This may have been because the predisposition was first reported after the second edition was published or because our previous reliance on expert opinion had failed to uncover the association.

Our new search methods and inclusion criteria are described more fully in the *Methods* section. We hope that these changes support a more robust and defensible evidential and scientific body of information in this third edition, which also includes additional supporting information for breed predispositions where possible. Such additional information may describe the population studied in the original papers – such as geographical location, referral or general practice population (many academic papers are based on referral populations, and their results do not necessarily generalize well to the populations of patients commonly seen in general practice) – while the date of the referenced papers may assist with a perspective on the temporal relevance of the results. Information is also provided on the comparator populations used in the studies (e.g. crossbreds or all study dogs) and the numerical results, which show the strength of the reported predisposition (e.g. odds ratio or prevalence). Taken together, these new segments of information should help the reader to piece together the likelihood of the reported predispositions being real and relevant in relation to his or her own personal animals and interests. In the first and second editions, the research was divided between the authors by body system. For this third edition, we have instead divided the research alphabetically by breed. Consequently, some differences in writing style and emphasis may be apparent between the authors' sections. However, by sticking to pre-agreed methods, we hope to have maintained a satisfactory level of consistency across the work. We have also updated the genetics section as well as adding new explanatory sections on methods, longevity and epidemiology. We hope that these will give the interested reader some useful background on these topics as well as suggestions on where to find further information as required.

Companion animals are often bred according to the whims and needs of mankind rather than following the harsh survival rules of natural selection, and therefore breed-related disease has become an important anthropogenic welfare issue. In consequence, it behoves everyone with an interest in companion animals to strive to reduce these animal welfare costs. A critical first step in this process is the need to define which breed–disease combinations (i.e. predispositions) have strong supporting evidence. We hope that the third edition of our book meets this need and provides a solid evidence base from which other companion animal stakeholders can develop effective strategies to improve animal welfare. Breeders and breeding organizations can use this book to identify priorities when considering the genetic health of their breeds. The show community, both those showing and those judging, may use this book to refine their opinions on optimal conformations and temperaments within individual breeds. The strong evidence-based approach of the book can help veterinary students and veterinarians with diagnosis and when advising prospective and current owners on breed-specific disease proclivities. Owners may find the book useful when deciding on breed selection or considering on how best to care for their current or prospective dog or cat. Ultimately, this book aims to enhance the welfare of current and future generations of cats and dogs by increasing our awareness of those diseases which commonly affect individual breeds and which may therefore be prevented or diagnosed earlier. Good evidence on breed predispositions empowers us all to combat disease occurrence and should lead to improvements in the lives of our dogs and cats.

We have thoroughly enjoyed writing this book: it became a labour of love for the three of us and consumed our lives for over a year, but we are very proud of the final product. There will obviously be some parts that we will re-read later and decide we could have done better, and we welcome the reader letting us know about these. There will also be some opportunities that were missed in this edition, and we will be glad to receive suggestions. However, we hope you will forgive these shortcomings for now and simply accept this third edition for what it is: an evidence-based blueprint for the current state of knowledge on breed predispositions to disease in dogs and cats. We hope you enjoy reading this book.

Alex Gough, Alison Thomas and Dan O'Neill

Abbreviations

95% CI	95% confidence interval
AKC	American Kennel Club
ANA	antinuclear antibodies
aPTT	activated partial thromboplastin time
AV	atrioventricular
CI	confidence interval
CYAR	cat years at risk
DYAR	dog years at risk
<i>E. coli</i>	<i>Escherichia coli</i>
EBVM	evidence-based veterinary medicine
ECG	electrocardiography
GSD	German Shepherd Dog
Ig	immunoglobulin, including isotypes IgA, IgG and IgM
IR	incidence rate
IRR	incidence rate ratio
KC	Kennel Club
MHC	major histocompatibility complex
MRI	magnetic resonance imaging
OR	odds ratio
PCR	polymerase chain reaction
PR	prevalence ratio
PT	prothrombin time
RR	relative risk, <i>or</i> risk ratio
SBT	Staffordshire Bull Terrier
T ₄	thyroxine
TRH	thyrotropin-releasing hormone
TSH	thyroid-stimulating hormone
VMDB	Veterinary Medical Database
WHWT	West Highland White Terrier

Introduction

BASIC AND CLINICAL GENETICS

Inherited diseases and breed predisposition

It has long been recognized that many traits, desirable and undesirable, can be passed along family lines. Darwin noted in 1868 that there is a 'unanimity of ... belief by veterinaries of all nations in the transmission of various morbid tendencies'. Inherited diseases in dogs and cats can be categorized as those associated with adherence to breed standards and those unrelated to breed standards. The brachycephalic head shape is particularly associated with a number of diseases such as brachycephalic obstructive airway syndrome, dystocia and corneal ulceration (Packer *et al.*, 2015; O'Neill *et al.*, 2017b, 2017c). Diseases not directly related to breed standards include many intraocular diseases, haematological and immune-mediated diseases, and endocrine diseases (although the creation of small gene pools for a breed because of adherence to breed standards may have contributed to the prevalence of these diseases). For ease of use, the accounts in this book have been arranged

by body system rather than in relation to breed standards.

It should be noted that while most conditions with breed predispositions are likely to be truly hereditary, this is not always the case. (Note that in our text, we use *hereditary*, *genetic* and *inherited* as synonyms.) Some conditions may arise because of the use to which the animal is commonly put, such as racing injuries in greyhounds, or to their behaviour, such as the searching behaviour of spaniels making them prone to grass awn (grass seed) foreign bodies. Nevertheless, even diseases such as these will have a genetic component, for example in influencing the behaviour of the spaniel, or giving the greyhound the athletic ability that means it is used for racing, and therefore can still be considered to have some inheritability.

Domestication and the canine and feline genome

Dogs are thought to be descended from a common ancestor with wolves, with estimates for the timing of the divergence ranging from 15 000 to 100 000 years ago. Domestication may have occurred more than once, and there may have

been further interbreeding with wolves subsequently. At least two bottlenecks have occurred in canine genetic history, one when they diverged from wolves, and another more recently when modern dog breeds were created. Nevertheless, the dog has an enormous variation in phenotype, as shown by characteristics such as size, colour, coat type and behaviour.

It has been shown that there is more variation in functional genes in domestic dogs than in wolves (Cruz *et al.*, 2008), and alterations in functional genes are often deleterious to welfare. Population bottlenecks and selective breeding may have exacerbated this, and natural selection against these deleterious conditions is less likely in domestic animals than in their wild counterparts. Domestic dogs may therefore be more prone to inherited diseases than wolves.

Dogs were originally bred to fulfil many different purposes, such as hunting, fighting, guarding, herding and companionship. Sight-hound type hunting dogs have been noted in archaeological records dating back 4000 years, and in Ancient Roman times, Columella described the division of dog breeds into working and hunting types. However, most modern dog breeds originate in the last 150 years, with the development of dog breeding as a hobby of the middle- and upper-class Victorian.

In a study (Parker *et al.*, 2007) that examined the DNA of a large number of dogs representing 161 breeds, the authors were able to divide the breeds into 'clades', that is, breeds with common ancestors. This paper shows how complex the genetic history of the dog is, but certain interesting points stand out. One is that single mutations can cause recognizable changes across multiple breeds within a clade. This has the consequence that dogs in a single clade may be prone to similar inherited diseases. Since most dog breeds are young in evolutionary terms, there has been little time for new mutations to occur, and so most disease-causing genetic mutations are thought to have occurred before the breeds were founded. It is also notable that related breeds came out of certain times and geographical locations. For example, dog fighting was popular in Ireland in the 1800s, and many mastiffs and bull terrier crosses from this location and period later developed into recognized breeds. The introduction of dogs into North America by European settlers and later Asian migrations largely replaced the indigenous domesticated canine

population which had been introduced by the first American settlers over 10 000 years previously. However, this study showed that breeds related to animals brought by European settlers likely had some interbreeding with the more ancient American breeds, and so American breeds of European origin retain some of the genetic material of the previous indigenous breeds.

Phenotypic variation (e.g. conformation and behaviour) is much smaller in cats than in dogs. Cats are thought to have been domesticated later than dogs, but are probably of less direct use to humans as working animals than dogs, since they are harder to train. Deliberate breeding was therefore more limited, and domesticated cats show as much genetic diversity as the wildcat.

The canine and feline genomes have both been sequenced, and the body of research into the genome and into genetic diseases in these species is rapidly growing.

Basic genetics

All mammalian life is based on the genetic code stored within the nucleus of a cell. This genetic code is stored in a long molecule called *deoxy-ribonucleic acid* (DNA). Each DNA molecule is composed of a string of units, called bases. There are four different bases, and they attract each other in pairs – *guanine* to *cytosine* and *adenine* to *thymine*. When attached together, they form the famous *double helix*. The order in which these bases (or base pairs, since they always match together) occur along the molecule provides the code for the synthesis of proteins. Proteins are then responsible for most of the functions of the body, from the structure of tissues, to the biological catalysts called enzymes, to the hormones which regulate the body's metabolic processes.

A length of DNA which codes for a particular protein is called a *gene*. Long strings of genes, interspersed with areas of DNA which do not code for proteins, make up *chromosomes*. Each nucleus of a mammalian cell contains a set number of chromosomes, except the sex cells (*gametes*) – sperm and ovum. For dogs this number is 78, and for cats it is 38.

When a somatic (body) cell divides, the chromosomes shorten and thicken within the nucleus, so they become visible under a microscope. They then replicate, and one copy of each chromosome separates into a new nucleus before the cell splits. This process is called *mitosis*.

However, in the production of the gametes (the process of *meiosis*), the chromosomes line themselves up in the middle of the cell with a companion. This companion is always the same, and two chromosomes that associate together are called *homologous pairs*. These homologous pairs separate, so the gametes have half the number of chromosomes as normal cells. This means that when a sperm and ovum combine at fertilization, the newly formed cell (the *zygote*) has the correct number of chromosomes.

Homologous pairs code for related genes, but are not identical. The two genes, one on each chromosome, interact in different ways. Sometimes one gene is *dominant* to the other, the less dominant gene being termed *recessive*, and the expression of the gene, that is, the protein that is produced, will be determined by the dominant gene. In other cases both genes will play a role in the production of the protein, a situation called *co-dominance*.

The exception to the homologous pairs are two chromosomes called the sex chromosomes (all the other chromosomes are called the *autosomes*). These chromosomes determine the sex of an animal. In most mammals, including dogs and cats (and humans), a female's somatic cells contain two X chromosomes while the male's somatic cells contain an X and a Y. At meiosis, the ova acquire a single X chromosome from the mother, whereas the sperm inherit either an X or a Y from the father. This has significance for the inheritance of conditions carried on the X chromosome, and means that some inherited diseases can be more prevalent in one sex than another.

Although any one animal will carry only up to two versions of a gene, many more can exist within a population because of mutation and natural selection. These different versions of the gene are called *alleles*.

In conditions and characteristics that are inherited in a simple way, that is, the conditions are autosomal dominant or recessive, a system of genetics devised by the monk Gregor Mendel (hence Mendelian genetics) can be used to predict the likely offspring of two parents, if the parents' genetic make-up is known. For example, the gene that codes for Labrador coat colours is dominant for black and recessive for brown. A Labrador with two alleles for black colour (call the allele B) is described as BB and hence the coat will be black. If it has one allele for black and one

for brown (call the allele b), it will be described as Bb but the coat colour will still be black since this colour is dominant. However, if the dog possesses two alleles for brown (bb), it will be brown. The genetic make-up is called the *genotype*, whereas the physical expression of the genes is called the *phenotype*.

The situation is slightly more complex when looking at matings, and a matrix can be used to aid prediction of offspring types. Take the example of a BB black male crossed with a bb brown female. The BB male will produce sperm each carrying a single B gene, and the female will produce ova each carrying a single b gene. These are then recombined at random to produce offspring. The matrix would therefore look like this:

		Male	
		B	b
Female	b	Bb	Bb
	b	Bb	Bb

This means that all the offspring would be Bb. They all carry the b gene for brown coat, but because this allele is recessive, the coat colour is black. An animal with two identical alleles (e.g. BB or bb) is called a *homozygote*, while an animal with two different alleles (e.g. Bb) is called a *heterozygote*. If a black Bb female were then crossed with a black Bb male a different pattern would emerge:

		Male	
		B	b
Female	B	BB	Bb
	b	Bb	bb

On average three of the offspring would be black: one a homozygote (BB) and two heterozygotes (Bb). The fourth would be a homozygote for brown coat colour (bb), and, since this b allele is not now being suppressed by the dominant B allele, the brown coat colour phenotype is expressed.

In fact, since the fertilization process is random, a litter of four pups may not be born in the exact 1:2:1 ratio, but if this were repeated enough times, the proportions of pups of the various genotypes would approximate to this.

Generally, the alleles separate randomly from each other. If a parent has two genetic conditions, just because one condition is expressed in an

offspring, that does not mean that the other will be. However, some alleles that are closely positioned on a chromosome tend to be passed on together. Thus, two traits controlled by different genes may often be found together in the same individual, and the presence of one of these traits may act as a marker for the other. This process is known as linkage.

When one allele is not dominant over another co-dominance exists. For example, certain flowers that have alleles for red flowers (R) and white flowers (W) will be coloured red if homozygous for red (RR), white if homozygous for white (WW) but pink if heterozygous (RW).

Some genes, even if dominant, do not always produce a physical effect in the host. For example, the condition polycystic kidney disease in cats is inherited as an autosomal dominant trait, but not all cats with the genes have cysts in their kidneys. This situation is called *incomplete penetrance*. Penetrance is the proportion of individuals with a particular genotype that demonstrate the characteristics normally expected with that genotype.

Some characteristics are carried on the X chromosome, and this can lead to the phenomenon of *sex linkage*. For example, Golden Retrievers are predisposed to a condition called X-linked muscular dystrophy. The allele for muscular dystrophy (call it M) is carried on the X chromosome, as is the allele for a normal dog not suffering from the condition (call it N). M is recessive to N. Therefore a female carrying a single affected X chromosome (genetic make-up $X^M X^N$) would not show the effects of the disease. If this female were mated with a normal male ($X^N Y$), then the matrix for their offspring would be as follows:

		Male	
		X^N	Y
Female	X^M	$X^M X^N$	$X^M Y$
	X^N	$X^N X^N$	$X^N Y$

All of the females born to this cross will be clinically unaffected by the disease, but 50% of the females will be carriers of the disease. These will not show the disease, since they have a normal gene on the other X chromosome which suppresses the abnormal, recessive gene. However, the males only possess a single X chromosome, so the 50% of males born $X^M Y$ will show the disease (since they do not possess another X chromosome

with a normal gene). The 50% of males born $X^N Y$ will not show the disease and will not carry it.

Because of this process, sex-linked diseases usually affect only males, and males cannot normally be asymptomatic carriers. Females are often carriers, but the only way they can express the disease is if their mother was a carrier and their father was affected. This situation is rare in nature, especially for uncommon genes, but can occur in domestic animals due to inbreeding.

Some disease inheritances are more complex still, because more than one gene may determine the expression of a disease, or the interaction of genes and environment can determine the outcome in an individual. For example, more than one gene is considered to be responsible for hip dysplasia but the dog's nutrition, exercise and other factors can also influence the severity of the disease.

Finally, some diseases are not inherited through the DNA of the cell nucleus at all, but through the DNA present within the *mitochondria* (which are intracellular organelles responsible for energy production). Mitochondria are entirely inherited from the mother, hence characteristics and diseases caused by mitochondrial DNA can only be passed down from the mother. Although conditions caused by mitochondrial DNA are rare, some canine myopathies are thought to be inherited this way.

In summary, an autosomal dominant trait is transmitted from generation to generation without skipping. Each affected offspring has at least one affected parent, unless the disease has arisen because of mutation. If the disease is lethal, then it will be very rare. An autosomal recessive disease may skip generations. If the two parents are affected, then all the offspring are affected. With an X-linked dominant condition, affected males mated to normal females transmit the gene to their daughters, who are all affected, but not their sons. Affected females then pass the condition on to approximately half of their sons and half of their daughters. In the overall population, the incidence in females tends to be twice that of males. With an X-linked recessive disease, the condition may skip generations. The incidence is more common in males. Affected males do not transmit the disease when mated to a normal female, but all female offspring will be carriers. Females showing the disease who are mated with normal males will pass the condition on to all their sons, and all their daughters will be carriers.

Clinical genetics

As noted above, genetic diseases may be more frequently encountered in domestic animals than in most wild populations. The process of domestication involves selecting animals for desirable traits from a human point of view. Initially, these traits would have been practical: speed in a horse, fertility and milk production in a cow, herding instincts in a sheepdog and so on. Over time, for animals such as dogs and cats that came to be kept for their companionship and aesthetic appeal, selection pressures switched towards features that made the animals fit in well to the human environment or made them look 'cute' – for example, miniaturization or achondroplasia – but which may have reduced adaptation to survive in the wild. As breeding practices were refined and the science of genetics was developed, inbreeding was used to create breeds that bred true with respect to certain desired characteristics (i.e. offspring greatly resembled their parents).

Unfortunately, inbreeding reduces the genetic variation within a breed, and tends to accentuate the expression of diseases that are due to recessive genes. Population bottlenecks occur through the importation of a small number of founder animals to a new country or because of regenerations of previously extinct breeds, and they are complicated by the *popular sire effect* whereby desirable individuals such as a show champion are overused (particularly males, which can produce many more offspring than a female). Most of the characterized genetic diseases of the dog are inherited as autosomal recessive traits. This may be because of inbreeding, but it is also due to the difficulty in identifying and eliminating recessive traits in breeding programmes.

It should be noted that inbreeding of itself does not cause genetic disease, and some degree of inbreeding can be of benefit for the concentration of desirable genes. In fact, some inbred strains of mice and rats are entirely homozygous and yet are quite healthy (Beck *et al.*, 2000). Inbreeding promotes homozygosity, and thus deleterious recessive genes are exposed by increasing the probability of their expression. However, by exposing these genes, it is possible to eliminate them by further selective breeding.

Data are currently sparse regarding the prevalence of disease caused by the spontaneous

appearance of new mutations. It seems likely that most of the genotypic variability of the domestic dog was present in its common ancestor with the wolf. However, it has been suggested that the Canidae family have elevated genome-wide basal slippage rates, meaning an increased rate of creation of new mutations due to errors in replication, compared to humans and cats (Shearin & Ostrander 2010). In most of those limited cases studied, the mutation seems to be uniform within a breed. This suggests that a *founder effect* applies, that is, a single initial mutation was propagated throughout the breed. In some cases, closely related breeds may have the same mutation causing a disease – for example, phosphoructokinase deficiency in English Springers and American Cocker Spaniels (and presumably Sprockers) – suggesting that a common ancestor was responsible for the original mutation. Some diseases, however, have more than one mutation in the same gene (*allelic heterogeneity*) or mutations in different genes which can lead to the same outcome. For example, oculoskeletal dysplasia is caused by mutations of different genes in the Labrador and the Samoyed, while multifocal retinopathy is caused by two different mutations of the same gene (Guziewicz *et al.*, 2007; Goldstein *et al.*, 2010).

When determining whether a disease is heritable, certain typical characteristics increase suspicion of a genetic predisposition. Often the first thing to suggest that a disease is inherited is that the disease occurs with a higher frequency in a group of related animals than in the general population. This can help distinguish an inherited disease from a breed predisposition (although it can be argued that in most cases breed predispositions are related to genetics in some sense, and a breed predisposition is suggestive of a genetic cause). For example, St Bernards are predisposed to osteosarcomas (Egenvall *et al.*, 2007), but it is possible this is merely a reflection of their large size, the faster growth rate leading to more mistakes being made in DNA replication, leading to cancer. However, analysis of pedigrees shows that there is a familial clustering pattern to cases of the disease, which suggests a specific gene or group of genes being responsible.

A hereditary defect often involves the same anatomic site in a group of related animals. This is often seen in congenital heart disease in dogs. Also, a hereditary disease is often seen to increase

in frequency with inbreeding. Hereditary diseases often have an early onset, and those that do not often have a consistent age of onset. Hereditary diseases usually affect a few individuals within a litter, as opposed to intoxications and infectious diseases, which frequently affect higher proportions. Some genetic diseases will cause abortion or resorption, and these are often difficult to recognize clinically. Similarly, some hereditary diseases will cause a failure to thrive, the 'fading kitten (or puppy) syndrome', and again it can be hard to determine the cause in these cases.

There is an extremely wide range of severity of hereditary diseases, from the relatively benign to the invariably fatal. Diagnosis of a hereditary disease is usually based on history, clinical signs, history of disease in related individuals, test matings, specific imaging or clinicopathological tests for diseases and genetic testing.

Test matings are often suggested in order to identify autosomal recessive diseases, but this does have problems. With late-onset defects, the results of the mating will be known too late to be useful in selecting which individuals to use for breeding. Test matings can be more useful for early-onset diseases, but the ethics of keeping a known affected animal purely for test purposes, and what to do with affected offspring, can be problematic. Furthermore, the results of test matings may be unreliable. For example, in the case of a recessive disease in which the N allele is normal and n is abnormal, a mating of a suspected carrier (N?) to a known carrier (Nn) which produced six normal puppies would give only an 82.2% certainty that the N? was not a carrier (NN). However, a single abnormal pup would confirm carrier status.

The results of random matings, if performed often enough and with respect to a sufficiently prevalent gene, can provide useful information without the need to maintain a carrier or affected animal, and with less likelihood of breeding unwanted affected individuals.

Specific tests for diseases include ultrasonography and histopathology for polycystic kidney disease, MRI for syringomyelia, and von Willebrand factor assay for von Willebrand's disease. Some laboratories will test samples using enzyme and immunological assays to detect diseases, and the results may indicate whether an individual is a homozygote or heterozygote. An example of this is testing for haemophilia B.

A defect in an affected protein's size, function or amount allows the identification of carriers of a disease in some cases, although there may be an overlap with normal values. Also, compensatory rises in other proteins, such as an isoenzyme related to pyruvate kinase in pyruvate kinase deficiency, may reduce the accuracy of this sort of test.

Causal molecular defects have been identified for some inherited diseases. Examples identified on the X chromosome include haemophilia B, severe combined X-linked immunodeficiency and hereditary nephropathy. Some autosomal recessive traits for which the mutation has been identified include copper toxicosis in Bedlington Terriers, progressive retinal atrophy in Irish Setters, von Willebrand's disease in Scottish Terriers and pyruvate kinase deficiency in Basenjis.

Many specific DNA tests are now commercially available to identify genetic diseases, and the number of tests is increasing rapidly, to include tests for such diverse diseases as degenerative myelopathy, von Willebrand's disease, copper toxicosis and anal furunculosis.

Specific DNA test results for diseases should be interpreted with caution, and what may seem a clear-cut result may often be misleading. For example, a positive genetic test result for degenerative myelopathy means the individual is at some risk for developing the disease. However, it does not mean that developing the disease is inevitable, nor does it mean that a clinically affected individual does not have another cause of its clinical signs, such as disc disease or neoplasia. Another pitfall is that there may be more than one type of mutation responsible for a disease, particularly between breeds. For example, the mutation causing muscular dystrophy in the Cavalier King Charles Spaniel is different from the one causing muscular dystrophy in the Golden Retriever.

DNA tests are either linkage-based or mutation-based. Linkage-based tests look for a marker gene that is physically near the gene of interest. Mutation-based tests look for the specific mutation causing a disease. Linkage tests may be inaccurate in a small number of cases where chromosomal recombination has occurred in the region between the marker and the mutation.

DNA testing shows great promise for the identification and elimination of genetic diseases

in dogs and cats. The inherited diseases can be identified before an animal is bred, and affected animals can either be removed from the breeding pool or, in the case of recessive traits, bred only to normal individuals, to preserve desirable characteristics. This allows the genetic diversity of breeds to be retained while inherited diseases are eliminated.

The limitations of DNA testing, such as the limited availability of tests, and the fact that its utility is largely restricted to single-gene diseases, mean that there is still a vital role for screening programmes to eliminate inherited diseases. Screening programmes currently in operation in the UK include the British Veterinary Association/Kennel Club programmes for hip and elbow dysplasia and eye diseases, and the International Cat Care scheme for polycystic kidney disease.

EPIDEMIOLOGY

The first and second editions of this book applied relatively loose evidence-based approaches to report on predispositions to diseases in dogs and cats. These evidence-based methods were not explicitly defined and were heavily reliant on expert opinion from textbooks, review articles and conference proceedings. Expert opinion is sometimes called 'eminence-based' veterinary medicine; it represents the personal view of recognized experts or self-appointed commentators without explicit external critical appraisal applied to its quality as evidence. Expert opinion is commonly promulgated as 'evidence' in veterinary medicine, particularly at conferences, in editorials and during undergraduate teaching. However, it is widely considered to be weak evidence at best, unless underpinned by a solid and stated evidential platform (Holmes, 2007). This is because many cognitive biases are inevitably inherent within the belief systems of any individual expert, which also explains why experts so often vehemently disagree on specific issues.

This third edition of *Breed Predispositions to Disease in Dogs and Cats* aims to be more explicit about how the epidemiological results for each breed-disease combination were chosen for inclusion. The book also aimed to cite those references that related to the highest available quality of study designs for each disease. Most of the references in this new book describe original research

that has been published in high-quality peer-reviewed journals. To support our new emphasis on reporting quantitative results from primary research, we have penned this introduction to epidemiology, to explain the epidemiologic metrics (e.g. prevalence, odds) that are reported throughout the book. Further information on this fascinating science of epidemiology is available in several useful texts (Thrusfield, 2007; Dohoo *et al.*, 2009; Pfeiffer, 2010).

Epidemiology is the study of disease levels in populations and of factors that determine the occurrence of these diseases. Veterinary epidemiology is a structured scientific approach towards collecting, integrating, analysing and interpreting data on health and demographics at a population level. Epidemiology aims to describe quantitatively the population under investigation. For example, we can define the proportion of German Shepherd Dogs among a known population of dogs [demography] or the proportion of these German Shepherd Dogs with aggression [prevalence] or whether male German Shepherd Dogs have a higher risk of aggression than females [risk factor analysis] (O'Neill *et al.*, 2017a). By identifying key relationships in biological systems, we can develop options for prevention or better diagnosis of future disease cases. New epidemiological evidence is interpreted within the wider body of basic scientific knowledge to contribute towards understanding and solving problems. For example, if we know that Pugs are predisposed to corneal ulceration, then we can recommend that owners are diligent with ocular care and more alert to eye problems in this breed, and that veterinarians are more vigilant during ophthalmological examinations, especially to assess for keratoconjunctivitis sicca (dry eye) which predisposes to corneal ulceration (O'Neill *et al.*, 2016a, 2017b). This evidence-based approach is very different to the traditional anecdotal approach where opinion (no matter how expert) and personal experience are the dominant forces. In the new epidemiological paradigm, data reign supreme.

At its most basic level, there are two main types of epidemiology: descriptive and analytic. Descriptive epidemiology describes the world that is defined by the data under examination in order to understand its demographic, disease or risk-factor features. In the context of the current book, descriptive epidemiology is used to report the frequency of the occurrence of disease within specific breeds. One measure of disease occurrence

is prevalence, which is commonly reported as the proportion or percentage of animals in a group that are affected by the disease at any one point in time (*point prevalence*) or during any specified period (*period prevalence*). So, for example, a statement that the ‘St Bernard had 19.4% prevalence for elbow dysplasia in the UK’ means that 19.4% of the St Bernard dogs in that study group of dogs in the UK had a diagnosis of elbow dysplasia during that particular study. Prevalence does not draw any distinction between long-standing cases that pre-existed the study period and new cases that were first diagnosed during the study period: these all count equally towards the prevalence total. Cases that are newly diagnosed during a specified period are called *incident cases*. These are reported as either the *incidence risk* (the proportion of animals that were not affected at the start of the study and that are newly diagnosed during the study period) or the *incidence rate* (number of new cases diagnosed divided by the sum of the length of time at risk for each animal in the study overall). Incidence rate is a useful measure to assess the rapidity with which animals develop disease over time.

Measures that describe deaths associated with specific diseases are called mortality. These are very useful pieces of data that offer information on the severity and impact of the condition. The *mortality rate* is derived similarly to the incidence rate and reports the number of deaths during a specific period diagnosed divided by the sum of the length of time at risk of death for each animal in the study overall. *Case fatality* describes the probability of death in affected animals. This is generally reported as a proportion (from 0.0 to 1.0) or percentage (0% to 100%) that describes the number of deaths divided by the total number of animals diseased.

Whereas descriptive epidemiology describes patterns of disease and can report absolute values for disease (e.g. prevalence tells us what percentage of a group of animals have a disease of interest), analytic epidemiology explores risk factors for diseases. *Risk factors* are attributes of an animal that affect its probability of developing a specific disease. For example, important risk factors for patellar luxation in dogs include body-weight, breed, age, sex and neutering (O'Neill *et al.*, 2016b). In the context of this book, breed is the most important attribute that we explore as a risk factor for disease. A key feature of analytic

epidemiology is that it requires a comparison group. If we aim to identify whether some category within a risk factor increases the probability of disease, we need to compare this probability to some other category and then report the relative results. For example, in relation to sex as a risk factor, we might choose to compare disease levels in males versus females to assess whether being male is associated with increased probability of a disease such as epilepsy (Kearsley-Fleet *et al.*, 2013). These comparative results can be reported using metrics such as *risk ratio* or *relative risk* (RR), *odds ratio* (OR) or *incidence risk/rate ratio* (IRR). The reader will see these terms used throughout this book. These metrics report the relative value for the risk-factor category of interest compared with the baseline (comparator) category. These ratios can be broadly interpreted in an equivalent way: a value above 1.0 suggests an increased probability of disease whereas a value below 1.0 suggests that the category is protective and may be associated with reduced probability of disease.

When exploring breed as a risk factor for disease, it is very important to select a logical comparator group to assist meaningful inference from the results. Options for the breed comparator group include ‘all dogs in the study’, ‘all remaining dogs in the study’, ‘all crossbred dogs’, or even another specified ‘single breed’. Swapping the comparator category from a group with a low risk of disease to a group with a high risk of disease can cause an odds ratio to change from above 1.0 to below 1.0 and give an illusion of reversal of risk. From this, it is clear that we must interpret results with great care. As Mark Twain aptly observed, ‘There are three kinds of lies: lies, damned lies, and statistics.’

A *census* is an epidemiological study that examines every animal in a population and can give an exact (true) value for the overall population, provided that all other aspects of the study design are perfect. However, owing to financial and logistical constraints, most epidemiological studies rely on just subsets (*samples*) of the overall population and are therefore restricted to reporting values that can then be extrapolated from the sample to the overall population. Statistical methods allow studies to report the exact value for the sample and then also to provide a spread of lower and higher values between which the study is 95% confident the true value in the wider general population lies.

This spread is called the 95% confidence interval (often abbreviated to 95% CI) and defines the statistical uncertainty associated with the reported measure of disease occurrence. The *power of a study* to confidently report precise results increases as the study sample size increases. This means that larger and possibly more reliable studies can report narrower spreads for the 95% CI; in other words, they have greater *precision*. When interpreting results, it is very important to examine not just the central exact value for the sample but also the width of the 95% CI that describes the inference for the wider population. A wide spread for the 95% CI suggests that the study was low-powered (i.e. a small sample size) and that precise conclusions may be difficult or unsafe to accept (Poole, 2001). The 95% CI for a prevalence value can be interpreted loosely as the range of values within which we are 95% confident that the true prevalence in the wider target population exists. For an odds ratio or risk ratio, if the lower limit of the 95% CI is above 1.0, then we can be highly confident ($p < 0.05$) of an increased odds/risk compared with the comparator group.

The *p-value* is another tool that helps to infer the strength of evidence for statistical results. Ideally, all analytic test results should report an associated *p-value*. The *p-value* defines the probability (from 0.0 to 1.0) of obtaining a result equal to or more extreme than what was observed, generally assuming that there is no true difference between the groups under comparison. A very low *p-value* describes a very low probability that the current result would have been found if there truly was no difference between the groups, and therefore we interpret this as suggesting that the groups are likely to truly differ. Previously, a simplistic approach was taken to the interpretation of the *p-value*, whereby any value less than 0.05 was taken as being 'statistically significant' and many older papers just reported whether *p-values* were above or below this cut-off. While many sources continue to use this heuristic, it is preferable to report the actual *p-value* for fuller interpretation by the reader in conjunction with other aspects of the result such as the size of the effect, the width of the confidence interval and the nature of the comparator groups chosen (Jeffery, 2015).

Causality (causation) deals with interpretation of possible causal relationships between a risk factor and a disease. A true causal relationship (i.e. the risk factor can be stated as an absolute

cause of the disease) is often very difficult to establish, even in the face of large volumes of supporting evidence. Most diseases have a complicated web of genetic, epigenetic, environmental and temporal factors that interact to promote disease occurrence. Even then, there may be random elements in play that determine which individuals from an apparently similar group get diseased. The relative effects and directions of causal factors can be problematic to unravel and quantify. For this reason, it is generally wise to avoid ascribing causality to risk factors, but instead to report just what the evidence usually suggests, which is an association. For example, rather than saying that 'being a Yorkshire Terrier causes patellar luxation', it is safer to say that 'there is an association between being a Yorkshire Terrier and having patellar luxation' (McGwin, 2010; O'Neill *et al.*, 2016b).

Many of the diseases investigated in the early days of epidemiological analyses had relatively simple and direct causal pathways. Exposure to a risk factor such as canine distemper virus regularly resulted in a very clear disease outcome called distemper. In such cases, analyses that only took account of a single risk factor (variable) were often adequate to answer the research question about probable causality. Such analyses are called *univariable* and explore associations between just one risk factor and a disease outcome. Nowadays, however, most of these 'simple' questions have already been answered and we are left with the more complex questions where multiple causal factors may be implicated in disease: the web of causation. To answer these multifactorial questions, we need more complex statistics called *multivariable* analyses that account for several variables in a single analysis. Thankfully, modern computing power now enables most researchers to carry out multivariable analysis easily and the cautious reader should seek out multivariable results rather than just to accept the more simplistic and possibly misleading findings from univariable analyses.

Confounding is a critically important concept when trying to understand the web of factors associated with disease occurrence. Confounding (meaning 'mixing up' or 'confusing') occurs when the effects of the risk factor of interest (e.g. breed) are mixed up with some other associated factor that is also associated with the disease outcome. Pet insurance is a good example of possible confounding. It is now well established that insured

pets are more likely to have disease diagnosed than uninsured pets, especially for those diseases that are more expensive or complicated to diagnose. It is also the case that certain dog breeds are more likely to be insured than others because of breed-related owner attitudes or differential pricing of policies by insurers, among other reasons. Consequently, a simple (univariable) analysis that directly reports the odds of diagnosis of a specific disease between two breeds of dog that have differing levels of pet insurance may appear to show the more insured breed as apparently more diseased. However, a multivariable analysis that also takes insurance into account can remove the effects of the differential uptake of insurance across the breeds and give a truer comparison of inherent predisposition between the breeds. Confounders, both known and unknown, should always be considered when planning studies, and especially when interpreting the results from any study.

Epidemiological studies are population-based analyses. This means that they are essentially reporting the cases (the *numerator*) that are identified from some underlying group of animals (the *denominator*). We often place inordinate focus on the numerator animals, because these are the ones with the disease of interest, but we ignore the denominator population at our own peril. It is critical to learn as much as possible about the denominator population so that we can extrapolate study results safely to wider or different populations. We especially need to know where these denominator animals lived (e.g. the UK or the USA), the dates for the study (e.g. 1990 or 2010), and some basic demography on these animals (e.g. ages, breeds, insurance). These key pieces of information are needed to assess the representativeness of the sample animals for the target population within that study. Additionally, this information allows comparison with our own population of interest so that we can evaluate the generalizability of the results, for example, in the same breed but in a different country and 10 years after the original study. The astute epidemiologist recognizes that time and location/setting are associated with many other ‘hidden’ changes such as economics or DNA testing that may affect the propensity for true or apparent disease diagnosis/occurrence.

Epidemiological studies rarely examine true disease status (i.e. whether an animal is truly diseased or not in the real world) but instead gener-

ally apply some belief or knowledge about the disease status. Hence, the same individual animal at the same time point may have differing disease status recorded depending upon how the data are collected. For example, a puppy with a diagnosis of congestive heart failure from a primary-care veterinary practice may be recorded as a congenital ventricular septal defect case by a referral specialist but may also be recorded as having no cardiac disease by an insurance database if the heart disease preceded the inception of the insurance policy or if the policy did not cover congenital diseases. Meanwhile, the owner of this same animal may record the disease as coughing or collapsing. The fact that none of these disease status reports is incorrect in its own context highlights the importance of carefully stating the case definition for the disease of interest and exploring how the disease data were derived. In this book, we try to help the reader by describing where possible how the data were collected – for example, from an owner survey, from an insurance database, or from referral or primary-care veterinary data.

In summary, epidemiology is the best method yet devised to understand the demography and health of dogs and cats. It can unlock secrets of disease that are otherwise impossible to discover. It can tell us which breeds are predisposed to which diseases. But with great power comes great responsibility. It behoves the user of epidemiology to understand the basics of this science so that we do not abuse or misuse its power.

LONGEVITY: UNDERSTANDING AND INTERPRETING THE DATA

Longevity (lifespan) and mortality (causes of death) statistics offer tantalizing prospects for unique insights into health and welfare variation in domestic dogs. However, interpretation of published statistics is fraught with pitfalls for the unwary, who may rush to draw conclusions without deeper consideration about such data drawn from various sources. Longevity (lifespan) is hugely uncertain for any individual animal and can be heavily influenced by unexpected disease, environmental effects or accidents. However, estimating average longevity for general populations of animals (e.g. a specific breed) can

be much more accurate, especially for comparison across breeds within restricted geographical and temporal limits taken from the same dataset.

We considered adding data on longevity for each breed into this edition to provide another perspective on comparative breed health that might act as a proxy value for the summative effects of all diseases within each breed. After all, surely the average duration that a breed lives should be an excellent and trustworthy measure of the health of that breed. But could it really be so simple? Following prolonged consideration, we rejected the option of adding longevity because of current limitations on the availability of reliable comparative population-based data, and because there are many caveats to the use of such data that often go unrecognized or ignored. However, to provide some information on breed longevity, and to highlight some of the pitfalls to the safe interpretation of these data, Table 1 shows some longevity results from the VetCompass Programme in the UK that illustrate apparently wide lifespan variation across breeds. Many other reports have shown similar results (Michell, 1999; Proschowsky *et al.*, 2003b; Adams *et al.*, 2010; Fleming *et al.*, 2011).

Reviewing the data shown in Table 1, the reader may feel very comfortable to accept these longevity values across a range of breeds as incontrovertible evidence that can rank breeds based on the summative effects of their general health, robustness and proclivity to disease. There are several other sources of good published evidence on longevity in dogs and cats that appear to tell a similar story (Proschowsky *et al.*, 2003a, 2003b; Fleming *et al.*, 2011; O'Neill *et al.*, 2013a). However, a good conceptual grasp of longevity is needed before we can safely move from our current position of data access to a position of data understanding and thence to a desired position of new beliefs (Proschowsky *et al.*, 2003a, 2003b; Fleming *et al.*, 2011; O'Neill *et al.*, 2013a). 'Having data' and 'understanding data' are not always synonymous.

Firstly, it is worth emphasizing that longevity in dogs is influenced by many factors other than just breed effects. The domestic dog (*Canis lupus familiaris*) exhibits unparalleled morphological diversity, from the 1 kg Chihuahua to the 85 kg Mastiff (Alderton & Morgan, 1993; Neff & Rine, 2006). There is now substantial evidence that average longevity reduces as breeds increase in average body size (Patronek *et al.*, 1997; Michell, 1999; Galis *et al.*, 2007; Greer *et al.*, 2007; Adams *et al.*, 2010; O'Neill *et al.*, 2013a). Lifespan reduc-

tion in larger dogs has been attributed to a range of genetic differences and pathological conditions induced by artificial selection and accelerated growth (Galis *et al.*, 2007; Urfer *et al.*, 2007; Fleming *et al.*, 2011; Salvin *et al.*, 2012; Kraus *et al.*, 2013). Consequently, perhaps we should restrict comparison to breeds of similar body-weight if we wish to compare reliably true breed-related health as opposed to longevity effects that are related to bodysize irrespective of breed. In other words, is it fair to directly compare the lifespans of the Cairn Terrier and the Great Dane from Table 1 and to assume that we can draw safe conclusions about their relative health from these values alone, or are we really just seeing differences that come mainly from comparing small and large breeds?

Euthanasia is another phenomenon in companion animals that needs to be considered when evaluating longevity as an indicator of health. Most humans undergo unassisted (so-called 'natural') deaths, but the converse is generally true for domestic pet species in developed countries. Reported euthanasia rates for dogs vary from 52% to 86% (Gobar, 1998; Michell, 1999; O'Neill *et al.*, 2013a), while 86% of deaths of UK cats involve euthanasia (O'Neill *et al.*, 2015a). By definition, euthanasia means that these animals have died prematurely before reaching the end of their so-called natural lifespan. The average ages at euthanasia may therefore be a highly reliable indicator of the inflection point at which quality of life dips below an acceptable threshold, and therefore may be a better measure of lifetime health than the maximum achievable lifespan up to a 'natural' death (McCutcheon & Fleming, 2001/2002). However, high euthanasia rates in dogs and cats also mean that longevity is influenced heavily by the opinions and decision-making patterns of owners, which may take welfare and suffering into account but may be additionally influenced by economic, performance and social factors. Varying decision-making on the acceptability and timing of euthanasia across breeds, diseases, countries and time could therefore impact differentially on subsequent longevity results. For example, owners may perceive the need for euthanasia differently between larger and smaller breeds for issues such as canine aggression, incontinence, mobility or even surgery, because larger breeds may cost more to treat, pose more risk to owners or offer shorter potential future lives than smaller breeds.

Table 1

Longevity for common dog breeds attending primary veterinary practices in England ranked by median age at death. The interquartile range (IQR), range and number of study dogs are also shown ($n=5095$) (O'Neill *et al.*, 2013a).

Breed	Median (years)	IQR	Range	No. of dogs
Miniature Poodle	14.2	11.1–15.6	2.0–19.4	20
Bearded Collie	13.7	12.2–14.3	4.0–17.0	25
Border Collie	13.5	11.5–15.0	0.1–19.1	184
Miniature Dachshund	13.5	9.2–14.3	2.0–19.5	25
West Highland White Terrier	13.5	10.4–14.9	0.2–21.0	128
Cairn Terrier	13.4	10.6–15.4	0.2–21.6	27
Jack Russell Terrier	13.4	9.3–15.7	0.0–24.0	298
Shih Tzu	13.3	9.2–15.6	0.0–18.6	79
English Springer Spaniel	13.3	10.4–14.8	0.3–19.4	111
Dalmatian	13.3	11.5–14.0	0.9–17.2	27
Crossbreed	13.1	10.1–15.0	0.0–22.0	1120
Yorkshire Terrier	13.0	10.0–15.1	0.01–20.6	217
Lhasa Apso	13.0	7.7–15.3	0.0–16.7	32
Bichon Frise	12.7	9.5–14.8	0.1–18.5	56
Weimaraner	12.6	11.1–13.5	6.5–17.0	36
Labrador Retriever	12.5	10.6–14.0	0.0–18.0	418
Golden Retriever	12.5	11.0–14.1	0.1–17.6	114
Shetland Sheepdog	12.5	11.7–13.8	8.5–14.6	20
Rough Collie	12.0	9.4–13.8	1.0–17.1	28
Border Terrier	12.0	8.9–13.1	1.2–21.2	31
King Charles Spaniel	12.0	10.0–14.2	0.0–15.3	26
Scottish Terrier	12.0	9.1–12.7	0.3–15.9	21
Cocker Spaniel	11.5	7.5–13.7	0.0–18.0	145
Bull Terrier	11.2	7.3–13.0	1.4–16.3	36
German Shepherd Dog	11.0	9.2–12.9	0.0–18.0	312
Greyhound	10.8	8.1–12.0	2.5–16.3	88
Staffordshire Bull Terrier	10.7	4.7–14.0	0.0–18.1	300
Boxer	10.0	7.7–11.6	0.0–16.5	91
Cavalier King Charles Spaniel	9.9	8.1–12.3	0.0–17.2	124
Dobermann	9.2	6.2–11.0	2.1–13.0	37
Bulldog	8.4	3.2–11.3	0.4–15.2	26
Rottweiler	8.0	5.5–10.2	0.0–16.6	105
Chihuahua	7.1	1.0–11.9	0.0–19.9	36
Mastiff	7.1	2.0–9.0	0.0–13.8	35
Great Dane	6.0	4.0–9.0	0.0–11.0	23
Dogue de Bordeaux	5.5	3.3–6.1	0.0–8.8	21

In the current context of dog and cat breeds, longevity defines the average lifespan for each breed. These longevity values are generally identified using death data from large populations such as referral or primary-care veterinary records, pet insurance data or owner/breeder surveys (Proschowsky *et al.*, 2003b; Bonnett *et al.*, 2005; Adams *et al.*, 2010; Fleming *et al.*, 2011; O'Neill *et al.*, 2013a, 2015a). A steady state of breed popularity over time is a key assumption when using these data for comparison between breeds, but this assumption is rarely true. If the

relative proportion of all puppies born each year is constant over a prolonged period for each breed, then the ages at death for each breed should be a reliable indicator of average breed longevity. However, breeds often rise and fall markedly in popularity over time, resulting in waves of young and old individuals for these breeds that complicate the interpretation of subsequent death data. For example, the popularity of Pugs in the UK has increased sharply in recent years, rising from less than 1% of all puppies born before 2008 to 2.8% of puppies born in




Figure 1

Annual proportional birth rates (2003–2014) for Pugs among all dogs ($n = 263\,456$) attending VetCompass primary-care veterinary clinics in England. The annual birth count of Pugs is shown in each bar (O'Neill *et al.*, 2016a).

2013 (Figure 1). Given that a high proportion of the Pugs that existed in 2014 were therefore young, it follows that younger Pugs have a greater probability of inclusion in mortality statistics compared with breeds that have not recently increased in population counts. Consequently, results from mortality data during 2014 that report the average longevity of Pugs will bias the true age at death for the Pug downwards and give misleading results showing that the breed dies younger than it truly does. The converse effect applies for breeds that are in numerical decline, which have relatively higher proportions of older dogs available to die during any one year. This effect is known as cohort bias, and it renders direct comparisons between average breed longevity highly problematic (Urfer, 2008).

In addition to breed popularity, several other factors that vary across breeds can also influence breed longevity results and complicate comparisons that aim to use longevity as a measure of breed health. Levels of neutering uptake can vary between countries. For example, some Scandinavian countries have historically restricted neutering in dogs to direct health-related purposes, whereas the procedure is routinely promoted in other countries such as the UK for population control and health prophylaxis, with the outcome that much higher

proportions of UK dogs are neutered (Anfinssen *et al.*, 2011; O'Neill *et al.*, 2014b). Likewise, pet insurance uptake in dogs varies widely: 0.3–3.0% in America, 4% in Canada, 34.0–40.3% in the UK, 68.4% in Sweden (O'Neill *et al.*, 2014a). Both neutering and pet insurance status are differentially associated with the health status, diagnostic rates and lifespans of dogs, and therefore variable uptake across breeds can influence the longevity achieved by these breeds, independently of inherent breed health characteristics (Egenvall *et al.*, 1998; Hart *et al.*, 2016; O'Neill *et al.*, 2016c; Belanger *et al.*, 2017).

This *Longevity* section started with an unchallenged thesis that longevity, if it could be reliably interpreted, could be a useful metric to evaluate and compare breed health and welfare. But, as we have seen, this is perhaps too simple a perspective, for many reasons. For example, can we even believe that longevity and welfare are linearly related, and that welfare continues to score higher as longevity increases to its maximum? It is not necessarily true that a long life is an indicator of high animal welfare, and perhaps therefore the focus of welfare studies should be more on the quality of the overall life lived, rather than on the quantity. In human medicine, scientists are now turning their attention to the concept of healthspan

(healthy longevity) and towards quantifying both the length and the proportion of lifespan that qualifies as healthspan (Waters, 2011). This is yet another example of how a simple comparative analysis of breed longevity could direct the unwary towards unsafe conclusions.

METHODS

This third edition of *Breed Predispositions to Disease in Dogs and Cats* has substantially bolstered the scientific methods used in the two earlier editions in order to place stronger emphasis on compliance with modern principles of evidence-based veterinary medicine (EBVM) (Holmes, 2007). In line with this progression, this new *Methods* section explains the processes followed during the literature searching and reporting in the current edition, while the *Epidemiology* section (above) describes the epidemiological approaches used.

Information on breed predispositions is available from a wide variety of sources. *Primary sources* of information describe original studies and show the information that was first published. For instance, for a scientific study that describes the common diseases of dogs in England, the primary source is the paper originally published by the scientists who performed the research, 'Prevalence of disorders recorded in dogs attending primary-care veterinary practices in England' (O'Neill *et al.*, 2014b). *Secondary sources* are documents such as websites, the press, books, editorials and review articles that may include information taken from primary sources. In relation to the paper describing the common diseases of dogs in England, a secondary source might be a newspaper article entitled 'Pedigree dogs "as healthy as mongrels", say vets' (Copping, 2014). Such sources often add further discussion or interpretation that extends or does not necessarily reflect the true intent of the original primary research, that may cherry-pick certain aspects of the original research, or that may not be completely accurate. While secondary sources can make for interesting general reading, they may not always tell an accurate story, or the full story, or an unbiased story. Consequently, it is wise to validate secondary reports by following the trail back to the original primary research before accepting the veracity of any conclusions.

Human medicine moved towards applying more rigorous standards for defining information sources that were 'good evidence' in the 1980s, and the Cochrane Collaboration aimed to provide clinicians with valid publications and guidelines to assist improved decision-making in public health from 1993 (Cochrane, 2017). Later, in the 1990s, evidence-based medicine (EBM) was recognized in human medicine as a distinct discipline that should be founded on the best available clinically relevant research (Sackett *et al.*, 1996). Since then, evidence-based veterinary medicine (EBVM) has become increasingly accepted in the veterinary field and is similarly reliant on understanding and using the most reliable sources of evidence when making decisions or developing beliefs (Cockcroft & Holmes, 2003).

A critical aspect of EBVM is to identify the most reliable sources of evidence from the ever-increasing deluge of information that is available in the modern era of electronic publication and data dissemination. The hierarchy of evidence quality is stylized as a pyramid that narrows progressively from the wider volume of lower reliability material at the base to a smaller volume of higher reliability material at the tip (Figure 2). The higher quality evidence towards the top of the pyramid tends to be individual or amalgamated analyses based on well-designed original pieces of research (studies) that have been through the peer-review process. These high-quality studies are designed to reduce selection or information biases, to be large enough to reduce random error, and to have appropriate statistical analytic methods (Vandeweerd *et al.*, 2012). Although there is some debate about whether the study design or the quality of the execution are more important for validating the reliability of the results, it is generally accepted that the pyramid of evidence is a useful model for the quality of evidence (Rosner, 2012). This EBVM aspiration was at the forefront of our minds when we designed the research and reporting methods for the current edition. Where possible, we aimed to reference only original peer-reviewed scientific publications and to avoid the inclusion of conference proceedings, review articles, editorials, websites or veterinary textbooks.

The research and writing process that we used was as follows. Each of the three authors was allocated a random subset of breeds for which




Figure 2
Hierarchy of evidence quality.

we aimed to identify all disease predispositions with sufficient supporting evidence. A breed was considered predisposed to a disease if some available evidence reported an increased incidence, prevalence or risk compared with an appropriate comparator group, preferably within a peer-reviewed primary publication. Specifically, where an odds ratio, a risk ratio or an incidence ratio was the reported metric of comparison, this ratio value would be greater than 1.0 and be supported by a p-value of <0.05 or by a 95% confidence interval that spanned values entirely greater than 1.0.

Each author followed the same general literature search strategies to ensure that the probability of disease discovery was similar across the breeds. The literature search covered a spectrum of electronic bibliographic databases including *CAB Direct*, *Google Scholar*, *IVIS* (International Veterinary Information Service), *PubMed*, *Science Direct*, *Veterinary Information Network* and *Web of Knowledge*. The precise search strategies included various keyword search combinations from relevant categories, including [BREED NAME], [DISEASE NAME], INHERIT*, HERED*, CONGEN*, GENETIC* and PREDISPOS*.

Relevant findings were merged from multiple searches for each breed. Analytic studies reporting the results of comparative studies that

reported increased incidence or prevalence in the breed of interest compared with some other meaningful comparator group (e.g. crossbreds) were prioritized. Where this level of evidence was not available, high-quality descriptive studies were accepted. We also accepted results from genetic studies that identified the mutations for specific diseases or evidence of inheritance within the specified breeds in combination with some evidence of increased incidence or prevalence. Case reports, studies conducted on laboratory research animals, and literature not published in the English language were generally excluded. The constraints of working within the available literature meant that a variety of comparator groups were accepted; these included 'all study dogs', 'all remaining study dogs', 'all crossbred dogs', or single or a combination of other breeds. If an author was in doubt about a source, then all three authors independently reviewed the original article to reach a majority consensus. From the final list of accepted publications, available information was extracted that described (1) the predisposed breed and disease, (2) the strength of the predisposition and the comparator group, (3) the geographic location, (4) the authors and date of the original publication, and (5) any other information of potential relevance such as inheritance and

signalment (e.g. sex or age) associations. These metadata were summarized and reproduced in this book.

For breed predispositions with several supporting publications, we applied some criteria to decide which publications to include in the book. Reports that were larger, more recent and had stronger study designs according to the pyramid of evidence (Figure 2), or those papers that provided evidence on the genetic mechanisms, were favoured for inclusion. Preference was also given to references based on studies with larger underlying denominator populations or those that were deemed more representative of the wider populations. Priority was additionally given to studies based on multivariable statistical analyses rather than univariable results. Multivariable statistical methods take account of multiple risk factors when reporting the breed effects and therefore account for confounding effects from other factors such as insurance status, age and neutering in order to provide less biased inference (O'Neill *et al.*, 2013b).

The lists of breeds included in the current edition have been extended from those presented in previous editions. The current edition includes predispositions to over 650 diseases across 204 breeds of dog and 45 breeds of cat. In line with moves towards international standardization of veterinary language, the breed names and synonyms used in this edition were based on the breed lists available within the VeNom coding system (VeNom Coding Group, 2017) along with additional breed terms identified from the VetCompass Programme database (VetCompass, 2017). This third edition defined breed and purebred as any dog types that were achieved through the process of selective breeding and that would breed true. *Breeding true* was taken to mean that when any two individuals from the same breed are mated, their progeny show consistent, replicable and predictable characteristics typical of the parents. *Pedigreed* animals were defined as that subset of individual breeds with known parentage for several generations. The terms *crossbred* and *mixed breed* were taken as synonyms to describe any dog types that

were not a purebred, regardless of whether their parentage was known or not.

In this edition, we have aimed to include only information on disease predisposition. A contrarian approach that we contemplated as a useful adjunct towards improved understanding of breed health was to also include evidence for diseases against which specific breeds are protected (i.e. they are less likely to get this disease than a comparator group of animals). Ultimately, however, we did not tackle this task in the current text because there is little information published on disease protection within breeds, but it may be included in a future edition.

Usage of the term 'inherited' was downplayed in the current edition, because it is now recognized that the majority of diseases in dogs and cats have both inherited and environmental components to their causality. For example, information on the inheritance of hip dysplasia in dogs has been widely published, and so hip dysplasia is widely considered as an inherited disorder (Lewis *et al.*, 2011b; Wilson *et al.*, 2013). However, age and sex are also known to be associated with hip dysplasia (Witsberger *et al.*, 2008). The current edition therefore uses the term *breed predisposition* to cover the combined effects from all factors (including genetic, epigenetic, environmental and owner-related) associated with increased probability of disease in a given breed.

It is also worth noting that the current edition is restricted to breeds and their diseases with published supporting evidence. It is clear that breed-based research is not carried out at random but may be biased towards common or popular breeds, human translational research, working breeds, laboratory breeds or perceptions of priority topics where funding for research may be more readily available. This means that, while this edition may accurately identify the evidenced predispositions within breeds, this does not necessarily mean that these are the only or even a representative selection of the true predispositions for each breed. As always, absence of evidence is not evidence of absence.

PART I

DOG BREEDS

AFFENPINSCHER

Dermatological conditions

Canine follicular dysplasia (seasonal flank alopecia)

- Reported in a small case series
- In this breed, low plasma levels of sex hormones were not considered the cause of the condition (Waldman, 1995)

AFGHAN HOUND

Cardiovascular conditions

Heart block

- This breed reported to be predisposed to high-grade second-degree or third-degree heart block in a US case series
- Heavier, older and sexually intact female dogs over-represented (Schrope & Kelch, 2006)

Musculoskeletal conditions

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.9 compared to mixed breeds (LaFond *et al.*, 2002)

Neurological conditions

Afghan myelopathy

- Reported in two case series
- Considered to be inherited in an autosomal recessive fashion
- Onset in young adolescents (Averill & Bronson, 1977; Cummings & de Lahunta, 1978)

Ocular conditions

Cataract

- Prevalence of primary cataract 2.36%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Prevalence declined over the years 1964–2003
- Highest prevalence at age 1–2 years in this breed (Gelatt & MacKay, 2005)

Corneal oedema (due to infection or vaccination with canine adenovirus type 1)

- Increased susceptibility (less commonly seen with the development of canine adenovirus type 2 vaccines)
- Afghans showed a more profound clinical response than Beagles experimentally (Curtis & Barnett, 1981)

Respiratory conditions

Chylothorax

- Usually idiopathic
- Afghan hounds comprised 37.5% of dogs with idiopathic chylothorax and 26.5% of all dogs with chylothorax
- No sex predisposition noted
(Fossum *et al.*, 1986)

Laryngeal paralysis–polyneuropathy syndrome

- Afghans reported to be predisposed
- May be inherited by an autosomal dominant mode
(Burbidge, 1995)

Lung lobe torsion

- Afghans reported to be over-represented compared to hospital population, with 4/22 cases
(Johnson & Feeney, 1984; Neath *et al.*, 2000)

AFRICAN BOERBOEL

Musculoskeletal conditions

Elbow dysplasia

- Common in this breed in South Africa
- > 38% incidence
- Males predisposed
(Kirberger & Stander, 2007)

Neurological conditions

Cervical spondylomyelopathy (cervical vertebral malformation, wobblers syndrome)

- Seen in first 2 years of life in this breed
- Reported in a South African case series
(Gray *et al.*, 2003)

AIREDALE TERRIER

Cardiovascular conditions

Dilated cardiomyopathy (DCM)

- Increased prevalence with age
- Approximately twice as common in males as in females
- Thought to be familial or genetic
(Tidholm & Jonsson, 1997)

Electrocardiographic abnormalities

- All 42 dogs of this breed investigated in a screening survey had ECG abnormalities
- Abnormalities included mean electrical axis deviations, low-voltage QRS complexes and first-degree AV block
(Amberger *et al.*, 1996)

Dermatological conditions

Grass awn migration

- Increased prevalence in this breed compared to hospital population
- Common in the summer months
(Brennan & Ihrke, 1983)

Canine follicular dysplasia (seasonal flank alopecia)

- Neutered females predisposed
- A marked predilection in this breed implies a genetic basis for this group of diseases
- Hair loss begins at 2–4 years of age and occurs mainly on the flank
(Miller & Dunstan, 1993)

Endocrine conditions

Hypothyroidism

- Breed at increased risk ($p < 0.01$)
- Genetic component suspected
- May occur at a younger age in breeds at risk (2–3 years)
- Ratio of affected males:females higher in at-risk breeds compared to non-high-risk breeds
(Milne & Hayes, 1981; Larsson, 1986)

Haematological/immunological conditions

Haemophilia B

- Severe factor IX deficiency in this breed
- Familial in this breed
(Brooks, 1999)

von Willebrand's disease (vWD)

- Type I seen in this breed
(Brooks, 1999)

Musculoskeletal conditions

Congenital umbilical hernia

- This breed reported to be significantly over-represented
- Females reported to be at excess risk
(Hayes, 1974a)

Hip dysplasia

- OR 3.9 compared to mixed breeds
- Neutered male dogs predisposed
(LaFond *et al.*, 2002)

Neoplastic conditions

Bladder and urethral tumours

- Airedales significantly over-represented compared to a hospital population
- Male:female ratio 1.95:1, but this was not statistically significant ($p > 0.05$)
(Norris *et al.*, 1992)

Nasal cavity tumours

- Breed at increased risk in a US teaching hospital case series
- Relative risk (RR) 4.6 (95% CI 2.24–9.25)
- Median age 9 years
- Males over-represented in most studies (Hayes *et al.*, 1982)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at moderately increased risk in a Finnish population (Niskanen & Thrusfield, 1998)

AKBASH

See *Turkish Shepherd Dog*

AKITA INU

See *Japanese Akita Inu*

ALASKAN HUSKY**Endocrine conditions**

Hypothyroidism (lymphocytic thyroiditis)

- Breed with a higher prevalence of thyroid hormone autoantibodies (THAA)
- In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, Huskies had an OR of 1.45 ($p=0.001$) of being affected compared to dogs of all other breeds
- Across the study, females were over-represented, and the highest prevalence was in dogs 2–4 years old

(Nachreiner *et al.*, 2002)

Neurological conditions

Lysosomal storage disease – GM₁ gangliosidosis

- Autosomal recessive inheritance; mutation identified
- Symptoms include proportional dwarfism and neurological deficits (ataxia and dysmetria) from 5–7 months of age

(Kreutzer *et al.*, 2005)

Mitochondrial encephalopathy

- Inherited, mutation identified
- Genetic defect at the level of the thiamine transporter

(Vernau *et al.*, 2015)

ALASKAN KLEE KAI**Haematological/immunological conditions**

Factor VII deficiency

- 6/18 client-owned dogs of this breed had this deficiency in an American study
- Inherited condition

(Kaae *et al.*, 2007)

ALASKAN MALAMUTE**Endocrine conditions**

Hypothyroidism

- More than 30% of Malamutes had a low T₄ in a sample of 2033 dogs of various breeds, compared to 10% for Dachshunds and Schnauzers
- Median TSH concentration significantly lower in this breed in a series of 693 dogs from 7 different breeds
- American populations studied

(Blake & Lapinski, 1980; Hegstad-Davies *et al.*, 2015)

Gastrointestinal conditions

Pancreatitis

- This breed reported to be predisposed in a Hungarian series of 80 cases

(Pápa *et al.*, 2011)

Haematological/immunological conditions

Stomatocytosis

- Reported in a few cases of Malamutes with chondrodysplasia in a Canadian study
- May be associated with anaemia

(Fletcher & Pinkerton, 1972)

Musculoskeletal conditions

Alaskan Malamute chondrodysplasia

- Autosomal recessive inheritance with complete penetrance and variable expression
- American population studied

(Sande *et al.*, 1982; Bingel *et al.*, 1985)

Cranial cruciate ligament (CCL) disease

- Prevalence in this breed 3.25% (OR 1.29, 95% CI 1.10–1.50; $p=0.018$)
- Population studied was from 27 teaching hospitals in the USA
- Neutered female dogs predisposed (Witsberger *et al.*, 2008)

Hip dysplasia

- 7.8% prevalence (OR 2.33, 95% CI 2.10–2.58; $p<0.001$)
- Population studied was from 27 teaching hospitals in the USA
- Neutered male dogs predisposed (Witsberger *et al.*, 2008)

Neoplastic conditions**Tracheal and laryngeal tumours**

- 5/26 dogs in a series and literature review were Alaskan Malamutes
- 10 of the 26 cases in the study were a Spanish population, the rest were a worldwide literature review (Ramírez *et al.*, 2015)

Sebaceous gland tumours

- Breed at risk of sebaceous adenoma and epithelioma in case series
- American population (Scott & Anderson, 1990)

Neurological conditions**Idiopathic polyneuropathy in Alaskan Malamutes**

- Affects mature young adults
- Previously considered eliminated by breeding programmes, but more cases have arisen recently in the USA and northern Europe
- Autosomal recessive inheritance due to a single gene mutation (Braund *et al.*, 1997; Bruun *et al.*, 2013)

Ocular conditions**Cone degeneration (hemeralopia or day blindness)**

- Autosomal recessive inheritance
- Different underlying mutations reported in American versus Australian populations (Seddon *et al.*, 2006; Sidjanin *et al.*, 2002)

ALSATIAN

See *German Shepherd Dog*

AMERICAN BULLDOG

See *Bulldog – American*

AMERICAN COCKER SPANIEL

See *Cocker Spaniel*

AMERICAN ESKIMO

See *Eskimo Dog*

AMERICAN PIT BULL TERRIER**Gastrointestinal conditions****Parvovirus enteritis**

See under *Infectious conditions*

Infectious conditions**Babesiosis**

- High incidence reported in this breed in a number of countries, including USA, Australia and Romania
- In Romania, significantly associated with fighting-dog breeds, especially American Pit Bulls

(Birkenheuer *et al.*, 2005;

Jefferies *et al.*, 2007; Imre *et al.*, 2013)

Parvovirus enteritis

- Breed at increased risk in cases series
- Age 6 weeks to 6 months at higher risk (Houston *et al.*, 1996)

Ocular conditions**Retinal dysplasia**

- Reported in one purpose-bred colony from a single affected founder dog in Brazil
- Authors extrapolate that this condition is inherited in this breed
- Autosomal dominant inheritance (Rodarte-Almeida *et al.*, 2016)

Renal and urinary conditions**Urolithiasis – cystine**

- Breed at significantly increased risk in case series

(Case *et al.*, 1992)

AMERICAN STAFFORDSHIRE TERRIER

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- American Staffordshire Terrier was reported with high aggressivity in Italy
- 5/16 cases which involved repeated biting or shaking of the victim were this breed (all male dogs)
(Notari & Goodwin, 2007; Zapata *et al.*, 2016; Wright, 1985)

Cardiovascular conditions

Arrhythmia

- American Staffordshire Terrier had the highest prevalence (28.6%) of supraventricular arrhythmias among referred cardiology cases in Poland
(Noszczyk-Nowak *et al.*, 2017)

Congenital heart disease

- Includes a range of congenital heart disorders
- OR 4.2 compared with all referral dogs in Italy
(Oliveira *et al.*, 2011)

Dermatological conditions

Atopic dermatitis (atopy)

- In this breed, symptoms started between 1 and 2 years of age more often than at other ages
- Incidence rate (IR) 7.6 per 1000 DYAR (95% CI 4.2–11.0)
- Hungarian and Swedish populations studied
- This breed frequently has adverse reactions to food

(Nødtvedt *et al.*, 2006; Tarpataki *et al.*, 2006)

Demodicosis

- An inheritance pathway has been described
- This breed had the highest odds of juvenile-onset generalized demodicosis: OR 35.6 (95% CI 4.6–277.0) compared with an overall US first-opinion population
(It *et al.*, 2010; Plant *et al.*, 2011)

Endocrine conditions

Hypothyroidism

- Females and younger dogs are predisposed to having serum thyroid hormone autoantibodies (THAA) that are associated with hypothyroidism

- American Staffordshire Terrier had OR 1.78 ($p=0.001$) for THAA compared with all other breeds

(Nachreiner *et al.*, 2002)

Infectious conditions

Babesiosis

- Purebreds and dogs living in rural areas predisposed, especially in the autumn and spring
- American Staffordshire Terrier was significantly over-represented, with a prevalence of 33.3% among dogs in Poland showing typical symptoms of babesiosis

(Adaszek *et al.*, 2011)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Estimated heritability of 0.48 and is a highly polygenic complex trait
- Neutered and older individuals are predisposed
- OR 1.62 (95% CI 1.42–1.82) compared with an overall referral population in the USA
- American Staffordshire Terrier had 6.5% prevalence, compared with 1.58% overall in referred dogs in the Czech Republic

(Nečas *et al.*, 2000; Witsberger *et al.*, 2008; Baker *et al.*, 2017)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.0 compared to mixed breeds
(LaFond *et al.*, 2002)

Neoplastic conditions

Canine cutaneous histiocytoma

- OR 2.60 (95% CI 1.45–4.68) compared with crossbreeds in the USA
(Goldschmidt & Mcmanus, 2000; Fulmer & Mauldin, 2007)

Lymphoma

- German population
- Two comparison populations: own clinic ($n=52$ 142) and insured population ($n=123$ 423)
- Odds ratios 3.3 and 4.6 respectively for the two populations
(Ernst *et al.*, 2016)

Mast cell tumour (MCT)

- Mean age at presentation between 7.5 and 9 years, but can occur at any age
- OR 8.57 (95% CI 5.14–8.13) compared with crossbreeds in Austria

- American Staffordshire Terrier had the third-highest breed RR (7.33) compared with all breeds in laboratory records in the USA
- OR 2.07 (95% CI 1.44–2.99) compared with crossbreeds in the USA

(Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011; Leidinger *et al.*, 2014; Mochizuki *et al.*, 2016)

Neurological conditions

Cerebellar degeneration

- Autosomal recessive inheritance suspected
- Prevalence of 1 in 400 estimated in an American population
- Onset 18 months to 9 years
- Described in American Staffordshire Terriers from several countries, and characterized using MRI in Germany

(Olby *et al.*, 2004; Henke *et al.*, 2008)

Severe subacute necrotizing encephalopathy (Leigh-like syndrome)

- Reported in a case series of 17 dogs from seven closely related litters
- Age of onset 6–8 weeks
- Suspected to be inherited

(Collins *et al.*, 2013)

Ocular conditions

Retinal degeneration

- Causative mutations identified in American Staffordshire Terrier in the USA: *PDE6B* deletion mutation and *IQCB1* insertional mutation

(Goldstein *et al.*, 2013)

Renal and urinary conditions

Hyperuricosuria

- American Staffordshire Terrier had 3.17% prevalence of carriers for the mutation in the USA

(Karmi *et al.*, 2010a)

Reproductive conditions

Prostate disorders

- Survey of 72 300 male dogs in a French population
- Prostate disorders included: benign prostatic hyperplasia (45.9%), prostatitis (38.5%), abscesses (7.7%), cysts (5.0%), neoplasia (2.6%)
- OR in this breed 3.8 (95% CI 2.5–5.8)

(Polisca *et al.*, 2016)

ANATOLIAN SHEPHERD DOG

Behavioural conditions

Tail-chasing

- This breed over-represented in a Turkish population

(Yalcin, 2010)

Musculoskeletal conditions

Carpal laxity syndrome

- Turkish population
- 9/43 dogs in a case series were this breed
- 6–8 weeks usual age of onset

(Cetinkava *et al.*, 2007)

AUSTRALIAN CATTLE DOG

Behavioural conditions

Aggression

- 9/33 dog attacks on postal delivery officers in Queensland where breed was reported were Australian Cattle Dogs
- This breed showed increased aggression to strangers in Australia compared to seven other common breeds

(Podberscek & Blackshaw, 1991;

Duffy *et al.*, 2008)

Gastrointestinal conditions

Congenital portosystemic shunt

- 13/62 dogs were this breed in an Australian case series
- Large intrahepatic shunts often involve the right liver lobe in this breed (OR 5.6 right vs. left)

(Tisdall *et al.*, 1994; Hunt, 2004;

Krotscheck *et al.*, 2007)

Musculoskeletal conditions

Patellar luxation

- This breed reported to be over-represented in a Korean case series of 134 dogs

(Alam *et al.*, 2007)

Neoplastic conditions

Mast cell tumour (MCT)

- Breed at increased risk in an Australian case series of 70 mast cell tumours

(Baker-Gabb *et al.*, 2003)

Neurological conditions

Congenital deafness

- 10.8% prevalence in 899 dogs tested
- Australian population

- Females predisposed, with an OR of 1.7 (95% CI 1.0–2.8)
- Autosomal recessive inheritance with incomplete penetrance
- Heritability of 0.21
(Strain, 2004; Sommerlad *et al.*, 2012)

Hereditary polioencephalomyelopathy of the Australian Cattle dog

- Affects young dogs
- Thought to be due to an inherited biochemical defect
(Brenner *et al.*, 1997)

Neuronal ceroid lipofuscinosis

- Suspected to be autosomal recessive inheritance
- Reported in case reports and a case series forming a US genetic study
(Kolichski *et al.*, 2016)

Ocular conditions

Cataract

- Suspected to be inherited
- Prevalence of primary cataract 2.32%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Gelatt & MacKay, 2005)

Glaucoma – primary

- 1.51% prevalence in this breed in a North American population 1994–2002
- Males predisposed in this breed
(Gelatt & MacKay, 2004a)

Glaucoma – secondary

- More diagnoses than expected in this breed compared to an American hospital population
(Johnsen *et al.*, 2006)

Renal and urinary conditions

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population
(Case *et al.*, 1992)

AUSTRALIAN KELPIE

Musculoskeletal conditions

Perineal herniation

- Intact male dogs predisposed
- Mean age of onset: 9.4 years

- This breed reported to be over-represented in an Australian population
(Bellenger, 1980)

Neurological conditions

Cerebellar degeneration

- Autosomal recessive inheritance
- Signs seen at 6–12 weeks
- A 3 Mb region on CFA 3 containing 29 genes identified as the location for the cerebellar abiotrophy mutation in Australian Kelpie in Australia
(Fletcher *et al.*, 2010; Shearman *et al.*, 2011)

Physiological conditions

Pain perception and fear memory resilience

- Selective sweep spanning three megabases on chromosome 3 identified in Australian Working Kelpie
- May be associated with the *HOMER1* gene
- Selection allows Australian Working Kelpies to work effectively in harsh environmental conditions
(Arnott *et al.*, 2015)

AUSTRALIAN SHEPHERD DOG

Dermatological conditions

Sterile nodular panniculitis

- This breed significantly over-represented in a series of 39 dogs in an American population
(Contreary *et al.*, 2015)

Drug reactions

Ivermectin and milbemycin

- Associated with the mutant *MDR1* allele
- 16.6% of dogs of this breed tested had the mutation in an American study from volunteer breeders and owners
- Mutation occurred with a frequency of 46% in this breed in a UK population
(Neff *et al.*, 2004; Mealey *et al.*, 2005; Tappin *et al.*, 2012)

Endocrine conditions

Hyperadrenocorticism (Cushing's syndrome)

- Reported to be common in this breed in a case series of 153 dogs in an American population
(Wood *et al.*, 2007)

Haematological/immunological conditions

Haemophilia A

- Moderate factor VIII deficiency in this breed
- Familial in this breed

(Brooks, 1999)

Selective malabsorption of cobalamin (vitamin B₁₂)

- Genetic mutation identified

(He *et al.*, 2005)

Musculoskeletal conditions

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- Uncommon condition
- OR 1.91 in one study (compared to mixed breeds)

(LaFond *et al.*, 2002)

Neoplastic conditions

Pituitary tumour resulting in hyperadrenocorticism

See under *Endocrine conditions*

Neurological conditions

Epilepsy

- 50 epileptic Australian Shepherds with epilepsy compared to 50 non-epileptic Australian Shepherds in a German population
- 56% had poor seizure control
- Non-merle coat phenotype was associated with poorer seizure control
- Genetic basis suspected

(Weissl *et al.*, 2012)

Ocular conditions

Cataract

- Autosomal dominant inheritance suspected
- Prevalence of primary cataract 1.71%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Mellersh *et al.*, 2006; Gelatt & MacKay, 2005)

Chronic superficial keratitis (pannus)

- Breed at risk in case series, suspected to be inherited
- Age of onset: 4–7 years
- Prevalence and severity increase at higher altitude

(Chavkin *et al.*, 1994)

Collie eye anomaly

- Congenital condition, suspected to be inherited

(Munyard *et al.*, 2007)

Multiple ocular defects

- Autosomal recessive inheritance with incomplete penetrance
- Defects seem to be associated with merles with predominantly white coats
- Defects may include microphthalmia, microcornea, cataract, persistent pupillary membranes, equatorial staphylomas, colobomas and retinal dysplasia

(Gelatt *et al.*, 1981)

Persistent hyaloid remnants

- Congenital, suspected to be inherited
- 13/223 dogs of this breed in a case series were affected by this condition

(Munyard *et al.*, 2007)

Renal and urinary conditions

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population

(Case *et al.*, 1992)

AUSTRALIAN SILKY TERRIER

Dermatological conditions

Atopic dermatitis (atopy)

- OR 6.8 (95% CI 2.1–22.2) compared with an overall referral population in Australia

(Jaeger *et al.*, 2010)

Gastrointestinal conditions

Congenital portosystemic shunt

- Usually presents in young dogs
- Referred Australian Silky Terriers (8.5% incidence) had an OR of 9.3 (95% CI 5.5–15.6) compared with an overall Australian referral population

(Hunt, 2004)

Sialocoele

- Australian Silky Terrier was over-represented ($p < 0.01$), comprising 6.6% of cases compared with 1.8% of a referral population in Australia

(Bellenger & Simpson, 1992)

Musculoskeletal conditions

Patellar luxation

- Mainly medial luxation observed, often bilateral

- OR 16.0 (95% CI 3.5–73.7) compared to crossbreeds in the USA, and over-represented in Korea
(LaFond *et al.*, 2002; Alam *et al.*, 2007)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 10.29%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Silky Terrier had 22.76% prevalence, compared with 4.04% for crossbreeds in a US referral study

(Gelatt & MacKay, 2005;
Bellumori *et al.*, 2013; Donzel *et al.*, 2016)

Toxicity

Cane toad toxicity

- Smaller breeds, and especially terriers, predisposed
- Australian Silky Terrier was the second most affected dog, comprising 14.4% of a referral caseload in Australia

(Reeves, 2004)

AUSTRALIAN TERRIER

Endocrine conditions

Diabetes mellitus

- Swedish population
- Incidence rate (IR) was 183 per 10 000 DYAR (mixed-breed IR 15)

(Fall *et al.*, 2007)

Musculoskeletal conditions

Patellar luxation

- OR 8.0 (compared to mixed breeds) in a US population

(LaFond *et al.*, 2002)

Ocular conditions

Cataract

- Prevalence of primary cataract 4.10%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

BASENJI

Gastrointestinal conditions

Immunoproliferative enteropathy (of Basenjis)

- Familial, likely inherited
- Usually presents before 3 years
(Ochoa *et al.*, 1984; MacLachlan *et al.*, 1988)

Haematological/immunological conditions

Pyruvate kinase deficiency

- Inherited as an autosomal recessive trait
- Prevalence of 20 carriers for the disease out of 186 dogs tested in an Australian population in 1978
(Hogg *et al.*, 1978; Whitney & Lothrop, 1995)

Musculoskeletal conditions

Pyruvate kinase deficiency

See under *Haematological/immunological conditions*

Ocular conditions

Persistent pupillary membranes

- Suspected to be genetic
- 75/105 Basenjis affected in one Australian study

(James, 1991; Mason, 1976)

Physiological conditions

Reduced thyroxine levels

- Basenjis have lower T₄ reference range than mixed-breed dogs
- Australian population studied
(Seavers *et al.*, 2008)

Renal and urinary conditions

Fanconi syndrome

- Familial, autosomal recessive inheritance suspected
- 10% Basenjis in the USA affected in one study
- 37% 'probably carriers' and 6% 'probably affected' in a Czech and German population
- Females outnumbered males 3:1
(Bovee *et al.*, 1978; Noonan & Kay, 1990; Načeradská, 2009)

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population
(Case *et al.*, 1992)

BASSET HOUND

Dermatological conditions

Malassezia dermatitis

- This breed significantly over-represented in American and UK case series
- Predisposition in this breed to overcolonization with yeast thought to be due to a primary keratinization defect and/or the presence of deep skin folds
(Bond *et al.*, 1996; Mauldin *et al.*, 1997; Guillot *et al.*, 2003)

Primary seborrhoea

- Probably inherited as an autosomal recessive trait
- Signs first appear at an early age and get worse
- In this breed, otitis, greasy seborrhoea and dermatitis are seen, often in the body folds
(Bond *et al.*, 1998)

Skin tumours

See under *Neoplastic conditions* (*Trichoepithelioma*)

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed ranked seventh-highest at risk in a case series
- Increased weight generally increased risk, but Basset belonged to the lowest weight class of all the purebred dogs in this study
(Glickman *et al.*, 1994)

Mycobacterium avium complex infection

See under *Infectious conditions*

Haematological/immunological conditions

Hereditary thrombopathy

- Also known as Basset Hound thrombopathia
- Familial, autosomal recessive inheritance
- Reported in Canadian and American case series
- Signal transduction defect in this breed
(Patterson *et al.*, 1989; Brooks, 1999)

Severe combined immunodeficiency

- Inherited as an X-linked recessive trait
- Thymic hypoplasia and lymphopenia seen
(Perryman, 2004)

Infectious conditions

Malassezia dermatitis

See under *Dermatological conditions*

Mycobacterium avium complex infection

- Reported in a small American case series
- Possibly due to an inherited defect in cell-mediated immunity

(Carpenter *et al.*, 1988)

Musculoskeletal conditions

Elbow dysplasia

- Ununited anconeal process (OR 2.7 compared to mixed breeds) and fragmented coronoid process (OR 19.5) seen in this breed
- In Basset Hounds, ununited anconeal process is secondary to non-traumatic premature closure of the distal ulnar growth plate
- Also reported to be at increased risk in an American case series of 284 dogs with this condition

(Hayes *et al.*, 1979; LaFond *et al.*, 2002)

Inguinal/scrotal herniation

- Significantly increased risk in an American case series

(Hayes, 1974a)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 3.5 compared to mixed breeds

(LaFond *et al.*, 2002)

Patellar luxation

- OR 2.0 compared to mixed breeds

(LaFond *et al.*, 2002)

Neoplastic conditions

Nasal cavity tumours

- Breed at increased risk in a US teaching hospital case series
- RR 3.3 (95% CI 1.89–5.83)
- Median age 9 years
- Males over-represented in most studies
(Hayes *et al.*, 1982)

Trichoepithelioma

- Breed at risk in an American case series
(Scott & Anderson, 1991)

Neurological conditions

Cervical spondylomyelopathy (cervical vertebral malformation, wobbler syndrome)

- 12/224 cases were Bassets in a British series
(Lewis, 1989)

Intervertebral disc disease (IVDD)

- 10/229 dogs were Bassets in an American study
- This breed typically has its first episode of this disease at a greater age than other breeds
- Ratio of back length to height at withers is positively associated with increased risk (Mayhew *et al.*, 2004; Packer *et al.*, 2013)

Lafora's disease

- Reported in this breed in several case reports and a German histological study
- Hereditary and progressive (Kaiser *et al.*, 1991)

Ocular conditions**Glaucoma – primary**

- Inheritance suspected
- Prevalence increases with age (mean age of onset 6.3 years)
- Females predisposed in one study
- 11.36% prevalence in this breed in an American study
- Pectinate ligament dysplasia had a prevalence of 38.4% in a UK study
- Autosomal recessive inheritance
- A genetic study of 223 UK Bassets found a mutation frequency of 0.081, with a predicted frequency of affected dogs in the population of 0.007 (Gelatt & MacKay, 2004a; Dees *et al.*, 2014; Oliver *et al.*, 2015, 2016a)

Lens luxation – secondary

- Significant predisposition in this breed, in a Swiss population (Betschart *et al.*, 2014)

Physiological conditions**Osteochondrodysplasia**

- Accepted as breed standard
- Histopathologically confirmed that this breed is chondrodysplastic (Martinez *et al.*, 2007)

Renal and urinary conditions**Urolithiasis – cystine**

- Breed at risk in case series in American, Irish and New Zealand populations
- Young dogs affected (2–5 years)
- Almost all cases are male (Brown *et al.*, 1977; Jones *et al.*, 1998, 2001)

BEAGLE

Note: Because of the Beagle's extensive use in research settings, a large number of conditions are described in the breed that are not described in detail in other breeds. However, these may not represent true breed predispositions. It may be because the beagle is the only breed in which a condition has been studied in detail, the study being carried out to assist in human medical research. The conditions listed below are likely to represent true breed predispositions in the Beagle.

Behavioural conditions**Aggression**

- Reported to be one of the breeds most likely to direct aggression towards owners (Duffy *et al.*, 2008)

Withdrawal

- Beagles more likely to withdraw from stimuli compared to three other breeds (Plutchik, 1971)

Cardiovascular conditions**Mitral valve disease**

- 25.3% of healthy Beagles in a series with an age range 1.4–11.7 years had a murmur consistent with mitral valve disease, confirmed by colour-flow Doppler (Vörös *et al.*, 2015)

Dermatological conditions**Familial vasculopathy**

- Early onset of signs (4–10 months)
- No sex predilection noted (Scott-Moncrieff *et al.*, 1992)

Skin tumours

See under *Neoplastic conditions (Palpebral neoplasia)*

Endocrine conditions**Diabetes mellitus**

- Swedish population
- Incidence rate (IR) 24 cases per 10 000 DYAR, compared with 15 for mixed breeds (Fall *et al.*, 2007)

Hyperadrenocorticism (Cushing's syndrome)

- Reported to be common in this breed in a case series of 153 dogs in an American population (Wood *et al.*, 2007)

Hypoadrenocorticism (Addison's disease)

- Breed reported to commonly affected in one Brazilian case series

(Romão, 2011)

Hypothyroidism

- Reported in American and Czech Beagle colonies
- 8% incidence in Czech colony

(Fritz *et al.*, 1970; Vajner *et al.*, 1997)**Thyroid neoplasia**

- Breed at increased risk in Swiss and US case series

(Harari *et al.*, 1986; Wenger *et al.*, 2005)**Gastrointestinal conditions****Selective malabsorption of cobalamin (vitamin B₁₂)**

- Autosomal recessive inheritance suspected
- Genetic mutation described in a US series

(Fyfe *et al.*, 1991)**Haematological/immunological conditions****Factor VII deficiency**

- Inherited as an autosomal dominant trait
- Reported in multiple Beagle colonies in USA and UK

(Spurling *et al.*, 1972, 1974; Brooks, 1999)**Haemophilia A**

- Severe factor VIII deficiency in this breed
- Familial in this breed

(Brooks, 1999)

Meningitis and polyarthritisSee under *Neurological conditions***Non-spherocytic haemolytic anaemia**

- Inherited condition
- Causes increased osmotic fragility of red cells

(Maggio-Price *et al.*, 1988; Pekow *et al.*, 1992)**Pyruvate kinase deficiency**

- In a US case series of 39 anaemic and 29 non-anaemic Beagles, 35% were PK-deficient

(Inal Gultekin *et al.*, 2012a)**Neoplastic conditions****Extraskeletal soft-tissue osteosarcomas**

- Breed reported to be at increased risk in a case series

- No sex predilection
- Older animals predisposed

(Langenbach *et al.*, 1998)**Palpebral neoplasia**

- Increased risk compared to mixed-breed dogs
- Median age 9.6 years

(Roberts *et al.*, 1986)**Perianal (hepatoid) gland adenomas**

- Breed at risk in US case series
- Average age was 10.5 years
- RR 2.4 (95% CI 1.73–3.73) in males and 1.9 (not statistically significant) in females (VMDB)

(Hayes & Wilson, 1977)

Pituitary tumour resulting in hyperadrenocorticismSee under *Endocrine conditions***Thyroid neoplasia**See under *Endocrine conditions***Urinary tract neoplasia**

- Beagles significantly over-represented ($p < 0.05$) in a Canadian study, compared to hospital population, and in a Japanese study

(Norris *et al.*, 1992; Aoki *et al.*, 2012)**Neurological conditions****Idiopathic epilepsy**

- Reported in a colony of Beagles and an American case series

(Bielfelt *et al.*, 1971; Ekenstedt *et al.*, 2011)**Intervertebral disc disease (IVDD)**

- 29.4% were Beagles in a US retrospective case series of cervical intervertebral discs
- 43.9% of herniated cervical discs in this breed were at C2–C3
- 11/229 dogs with thoracolumbar intervertebral disc disease were Beagles

(Mayhew *et al.*, 2004; Hakozaiki *et al.*, 2015)**Meningitis and polyarthritis**

- Also known as Beagle pain syndrome, steroid-responsive meningitis
- OR 11.51 (95% CI 2.56–51.72; $p = 0.01$) compared to selected controls in a UK population
- Suspected to be genetic

(Rose *et al.*, 2014)

Ocular conditions**Cataract**

- 24/1314 Beagles in a 1974 case study had punctate anterior opacities
- In a Korean population, 7.5% of beagles had cataracts

(Hirth *et al.*, 1974;
BongKyeong *et al.*, 2001)

Corneal dystrophy

- Inheritance suspected
- One report found that 15% of Beagles examined were affected
- Progressive but rarely affects vision

(Roth *et al.*, 1981)

Glaucoma – primary, open angle

- Autosomal recessive inheritance
- Beagles at higher risk than mixed-breed dogs in an American population

(Slater & Erb, 1986)

Ocular neoplasia

See under *Neoplastic conditions* (*Palpebral neoplasia*)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk
- 10/71 cases were Beagles in a Brazilian case series
- Usually presents in the first 1–2 years of life

(Merlini *et al.*, 2014)

Tapetal degeneration

- Autosomal recessive inheritance
- Reported in US laboratory dogs

(Burns *et al.*, 1988)

Renal and urinary conditions**Renal amyloidosis**

- Breed at increased risk in American case series
- Most cases > 6 years at diagnosis
- Females predisposed in one study

(DiBartola *et al.*, 1989)

Renal dysplasia

- May result in unilateral agenesis
- High prevalence reported in some families of Beagles

(Robbins, 1965; Bruder *et al.*, 2010)

Urinary tract neoplasia

See under *Neoplastic conditions*

Urolithiasis

- Breed at significantly increased risk ($p < 0.05$) in a case series

(Brown *et al.*, 1977)

BEARDED COLLIE**Dermatological conditions****Pemphigus foliaceus**

- Increased risk compared to the hospital population and the dermatology caseload
- Mean age of onset 4.2 years

(Ihrke *et al.*, 1985)

Endocrine conditions**Hypoadrenocorticism (Addison's disease)**

- 9.4% of an American population of 635 Bearded Collies were affected
- Heritability estimated to be 0.76
- Breed at increased risk in Swedish population (RR 7.43, 95% CI 5.11–10.5; $p < 2.2 \times 10^{-16}$)
- Females at increased risk in Swedish population (RR 1.85, 95% CI 1.55–2.22; $p < 0.001$)
- Prevalence in this breed in Swedish population of 0.322% (95% CI 0.088–0.825%)

(Oberbauer *et al.*, 2002; Hanson *et al.*, 2016)

Musculoskeletal conditions**Hip dysplasia**

- Neutered male dogs predisposed in an American population
- OR 3.1 (95% CI 1.1–8.7) compared to mixed breeds

(Lafond *et al.*, 2002)

Ocular conditions**Cataract**

- Prevalence of primary cataract 1.94%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Lens luxation – secondary

- Significant predisposition in this breed in a Swiss population

(Betschart *et al.*, 2014)

BEAUCERON

Neoplastic conditions

Mast cell tumour (MCT)

- Significantly increased risk for this breed in an Austrian population

(Leidinger *et al.*, 2014)

Squamous cell carcinoma – digit

- Over-represented in a French population
- Mean age 10.2 years
- No sex predisposition

(Belluco *et al.*, 2013)

BEDLINGTON TERRIER

Gastrointestinal conditions

Chronic hepatitis (copper storage hepatopathy, copper toxicosis)

- Inherited as an autosomal recessive trait
- Clinical onset seen in young to middle-aged dogs
- High prevalence worldwide
- 68/90 dogs of this breed affected in a 1979 American study

(Twedt *et al.*, 1979; Johnson *et al.*, 1980; Hultgren *et al.*, 1986)

Ocular conditions

Cataract

- Prevalence of primary cataract 8.49%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Congenital alacrima (congenital keratoconjunctivitis sicca)

- Significantly over-represented in an American study ($p = 0.04$)

(Westermeyer *et al.*, 2009)

BELGIAN SHEPHERD DOG (INCLUDING GROENENDAEL, LAEKENOIS, Tervuren, and Malinois)

Behavioural conditions

Aggression

- This breed reported to have higher odds of inter-dog dominance aggression

(Rugbjerg *et al.*, 2003)

Dermatological conditions

Vitiligo

- Presumed to be hereditary
- Antimelanocyte antibodies found in all 17 affected Belgian Tervurens tested and none of the 11 normal Belgian Tervurens tested in one study

(Mahaffey *et al.*, 1978)

Gastrointestinal conditions

Gastric carcinoma

See under *Neoplastic conditions*

Neoplastic conditions

Gastric carcinoma

- 8/11 dogs in one case series were of this breed
- Male dogs more commonly affected in case series
- Mean age of occurrence: 8–10 years

(Fonda *et al.*, 1989; Scanziani *et al.*, 1991)

Neurological conditions

Idiopathic epilepsy

- Common in the Danish population of this breed
- Late onset in this breed increases prevalence, owing to dogs already being used for breeding by the time they first show signs
- In a Finnish survey, 9.6% of Belgian Shepherds had had a seizure. These were all Groenendael and Tervuren variants, and none Malinois or Laekenois

(Teikari, 1996; Berendt *et al.*, 2008)

Ocular conditions

Chronic superficial keratitis (pannus)

- Breed at risk in case series, suspected to be inherited
- Age of onset: 4–7 years
- Prevalence and severity increase at higher altitude

(Chavkin *et al.*, 1994)

Physiological conditions

Leucopenia

- 6/9 healthy Belgian Tervurens sampled had white blood cell counts in the range $2.4\text{--}5.4 \times 10^9/\text{l}$

(Greenfield *et al.*, 1999)

BERGER BLANC SUISSE

See *White Swiss Shepherd Dog*

BERNESE MOUNTAIN DOG

Dermatological conditions

Skin tumours

See under *Neoplastic conditions*

Leptospirosis

- This breed significantly over-represented in a German study

(Geisen *et al.*, 2008)

Gastrointestinal conditions

Gastric or mesenteric dilatation/volvulus

- 1.8% mortality rate from these conditions in this breed in a Swiss population

(Klopfenstein *et al.*, 2016)

Hepatocerebellar degeneration

See under *Neurological conditions*

Oesophageal foreign bodies

- This breed over-represented in a Swiss hospital population

(Gianella *et al.*, 2009)

Haematological/immunological conditions

Factor I deficiency

- Autosomal inheritance

(Brooks, 1999)

Factor VII deficiency

- Autosomal inheritance

(Brooks, 1999)

Infectious conditions

Borreliosis (Lyme disease)

- Seroprevalence of 58% in this breed, compared to 15% in control dogs in a Swiss study ($p < 0.001$)

(Gerber *et al.*, 2007)

Musculoskeletal conditions

Elbow dysplasia

- Ununited anconeal process (OR 50.5, 95% CI 25.9–98.6 compared to mixed breeds) and fragmented coronoid process (OR 140.1, 95% CI 62.2–301.4) seen in this breed
- Fragmented coronoid process thought to be inherited

(Ubbink *et al.*, 1999; LaFond *et al.*, 2002; Temwichitr *et al.*, 2010)

Hip dysplasia

- Neutered male dogs predisposed
- OR 7.2 (95% CI 4.2–12.3) in one study (compared to mixed breeds)
- Decreasing prevalence seen in this breed in a French study

(LaFond *et al.*, 2002; Genevois *et al.*, 2008)

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 47.1 (95% CI 26.4–84.0) compared to mixed breeds

(Rudd *et al.*, 1990; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.8 (95% CI 1.5–6.0) compared to mixed breeds

(LaFond *et al.*, 2002)

Neoplastic conditions

Histiocytic sarcoma complex – disseminated histiocytoma

- Polygenic mode of inheritance
- Affects older dogs (7–8 years)
- More common in males
- In a Danish population, 13/812 dogs of this breed were diagnosed with this condition, and 11 of these were related
- OR 17.92 in a Japanese population

(Moore & Rosin, 1986; Padgett *et al.*, 1995; Nielsen *et al.*, 2010; Shiokawa *et al.*, 2013)

Lymphoma

- German population
- Two comparison populations: own clinic ($n = 52$ 142) and insured population ($n = 123$ 423)
- Odds ratios 2.4 and 2.0 respectively for the two populations

(Ernst *et al.*, 2016)

Mast cell tumour (MCT)

- This breed reported to be over-represented in a German population and an Austrian population

(Kessler *et al.*, 1997; Leidinger *et al.*, 2014)

Systemic histiocytosis

- Benign, technically non-neoplastic condition
- Polygenic inheritance suspected
- Mainly affects young to middle-aged males

(Moore, 1984; Padgett *et al.*, 1995)

Neurological conditions

Alexander disease

- Out of a total of 11 reports of this disease in dogs, 5 were Bernese Mountain Dogs and 1 was a Bernese cross

(Weissenböck *et al.*, 1996;
Wrzosek *et al.*, 2015)

Cervical spondylomyelopathy (cervical vertebral malformation, wobbler syndrome)

- Reported in a US case series of 7 Bernese (Eagleson *et al.*, 2009)

Degenerative myelopathy

- *SOD1:c.118A* has a frequency of 38% in this breed
- The *SOD1:c.52 T* allele appears to be restricted to this breed, with a frequency of 3.5%

(Zeng *et al.*, 2014)

Hepatocerebellar degeneration

- Reported in three related litters from two kennels in the USA
- Age of onset 4–6 weeks
- Autosomal recessive inheritance

(Carmichael *et al.*, 1996)

Idiopathic epilepsy

- Polygenic mode of inheritance in this breed

(Kathmann *et al.*, 1999)

Ocular conditions

Progressive retinal atrophy (PRA)

- Familial, inheritance suspected
- Early-onset retinopathy has been described in this breed in France

(Chaudieu & Molon-Noblot, 2004)

Physiological conditions

Prolonged activated partial thromboplastin time (aPTT)

- Higher levels of lupus anticoagulants and anticardiolipin in this breed

(Nielsen *et al.*, 2011)

Renal and urinary conditions

Familial renal disease (membranoproliferative glomerulonephritis and interstitial nephritis)

- Autosomal recessive with expression affected by a second sex-linked locus
- Affected dogs present at 2–5 years with renal failure and marked proteinuria

- Most of the dogs in one study had a high titre to *Borrelia burgdorferi*, suggesting that this organism may have had a role in the development of the condition (see also *Infectious conditions*)

- Incidence of renal disease in this breed was 51 cases per 10 000 DYAR (95% CI 41–61), compared with a mean incidence for all breeds of 15.8 per 10 000 DYAR, in a Swiss population

(Minkus *et al.*, 1994;

Reusch *et al.*, 1994; Pelander *et al.*, 2015)

Urolithiasis – struvite

- This breed reported to be at statistically increased risk of struvite in a Hungarian population

(Bende *et al.*, 2015)

Reproductive conditions

Prostate disease

- Over-represented in a French population (OR 2.5, 95% CI 1.3–4.7)

(Polisca *et al.*, 2016)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk in Swedish case series

(Egenvall *et al.*, 2001)

Respiratory conditions

Histiocytic sarcoma complex – disseminated histiocytoma

See under *Neoplastic conditions*

BICHON FRISE

Dermatological conditions

Atopic dermatitis (atopy)

- Increased risk in an Australian hospital population (OR 3.3, 95% CI 1.81–5.87)

- Males may be predisposed in this breed ($p=0.05$)

(Mazrier *et al.*, 2016)

Sarcoptic mange

- Reported to be at risk in a Chinese population
- Prevalence 3.19% (95% CI 0–6.75%) compared to average prevalence of pet dogs of 1.18% (95% CI 0.85–1.52%)

(YiZhou *et al.*, 2014)

Endocrine conditions

Diabetes mellitus

- Incidence rate (IR) 17 per 10 000 DYAR in a Swedish population (mixed-breed IR was 15) (Fall *et al.*, 2007)

Hyperadrenocorticism

- OR 6.5 (95% CI 3.5–12.1; $p < 0.001$) in a UK general practice population
- Dogs aged 12 years and above had an OR of 5.7 (95% CI 3.7–8.7; $p < 0.001$) of hyperadrenocorticism compared with dogs aged 6–8.9 years (O'Neill *et al.*, 2016d)

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed significantly over-represented in an Australian case series
- More common in females than males in this breed (12:2; $p < 0.001$)
- Usually extrahepatic (Hunt, 2004)

Haematological/immunological conditions

Immune-mediated haemolytic anaemia (IMHA)

- Usually affects young adult and middle-aged animals
- May be more common in bitches
- May be seasonal variations
- OR 5.3 (95% CI 1.2–22.5; $p = 0.024$) compared to hospital population in a US study (Miller *et al.*, 2004)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- This breed over-represented (11/55) in a case series of dogs <15 kg presenting for cranial cruciate ligament surgery (US population) (Campbell *et al.*, 2016)

Patellar luxation

- OR 4.8 (95% CI 2.5–9.3) compared to mixed breeds in an American population (LaFond *et al.*, 2002)

Ocular conditions

Cataract

- Autosomal recessive inheritance suspected
- High prevalence in this breed: 11.45% in a US population, compared to 1.61% in mixed-breed dogs, in a retrospective report based on VMDB data, 1964–2003

- Age of onset 2–8 years (Gelatt & MacKay, 2005; Adkins & Hendrix, 2005; Wallace *et al.*, 2005)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in large Canadian case series (Lulich *et al.*, 1999; Houston *et al.*, 2004)

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population (Case *et al.*, 1982)

Urolithiasis – struvite

- Breed at increased risk in large Canadian case series
- Females predisposed (16:1 in one large study) (Houston *et al.*, 2004)

Respiratory conditions

Primary ciliary dyskinesia

- Inherited defect
- Several case series affecting related dogs in this breed (Maddux *et al.*, 1991; Vaden *et al.*, 1991; Jamne *et al.*, 1998)

BLOODHOUND

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in case series
- 30.5% of deaths in this breed were from this condition in one study
- Prevalence ratio (PR) of 13.2 (95% CI 9.4–18.6; $p < 0.0001$) GDV deaths compared to 23 breeds with three or more deaths due to GDV (Evans & Adams, 2010b)

Musculoskeletal conditions

Hip dysplasia

- Neutered male dogs predisposed
- OR 4.5 (95% CI 2.0–9.9) compared to mixed breeds (LaFond *et al.*, 2002)

Neurological conditions

Degenerative myelopathy

- High frequency of mutation for this disease reported in a Belgian population (Beckers *et al.*, 2016)

BOLOGNESE

Renal and urinary conditions

Urolithiasis – purine

- Reported to be at increased risk in a Hungarian population

(Bende *et al.*, 2015)

BOLONKA ZWETNA

Ocular conditions

Lens luxation – secondary

- Significantly increased incidence in a German population in this breed

(Betschart *et al.*, 2014)

BORDER COLLIE

Cardiovascular conditions

Patent ductus arteriosus

- Breed association noted in one Australian study

(Aherne & Beijerink, 2013)

Drug reactions

Vincristine-associated myelosuppression

- 3/5 dogs of this breed with the *ABCB1* wild-type gene developed vincristine-associated myelosuppression, compared to 0/21 dogs without this genotype (Border Collies and other breeds) – this was significantly different

(Lind *et al.*, 2013)

Endocrine conditions

Diabetes mellitus

- Incidence rate (IR) 36 per 10 000 DYAR in a Swedish population, compared with a mixed-breed IR of 15

(Fall *et al.*, 2007)

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed at risk in case series
- Clinical signs usually seen in young dogs < 1 year

(Hunt, 2004)

Gastrointestinal foreign bodies

- Breed at increased risk in case series

(Hayes, 2009)

Inflammatory bowel disease

- Significantly increased risk in a UK population
- OR 1.99 (95% CI 1.17–3.41; $p=0.012$)

(Kathrani *et al.*, 2011)

Selective malabsorption of cobalamin (vitamin B₁₂)

- Autosomal recessive inheritance suspected
- Median age 11.5 months

(Battersby *et al.*, 2005; Lutz *et al.*, 2013)

Haematological/immunological conditions

Selective malabsorption of cobalamin (vitamin B₁₂)

See under *Gastrointestinal conditions*

Trapped neutrophil syndrome

- Autosomal recessive inheritance suspected
- Reported in numerous countries
- Carrier frequency 11.1% in a Japanese study

(Allan *et al.*, 1996;

Sharman & Wilton, 2007;

Mizukami *et al.*, 2013)

Infectious conditions

Cryptococcosis

- Breed at increased risk in an Australian case series
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors

(O'Brien *et al.*, 2004)

Musculoskeletal conditions

Achilles tendon rupture

- 7/45 dogs in a UK study with this condition were Border Collies or Border Collie crosses

(Corr *et al.*, 2010)

Central tarsal bone fracture

- Reported in case series of 6 UK border collies
- May have the radiographic appearance of a luxation

(Guilliard, 2007)

Gastrocnemius musculotendinopathy

- 8/9 dogs with this condition were Border Collies in a Swiss case series
- May be related to biomechanical forces or motion pattern

(Stahl *et al.*, 2010)

Hip dysplasia

- Neutered male dogs predisposed
- OR 2.1 compared to mixed breeds (95% CI 1.4–3.1)

(LaFond *et al.*, 2002)

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 15.0 compared to mixed breeds (95% CI 9.6–23.3)
- Increased risk of shoulder osteoarthritis in this breed

(LaFond *et al.*, 2002;
Maddox *et al.*, 2013)

Trauma from agility work

- This breed at increased risk of injury compared to its exposure

(Levy *et al.*, 2009)

Neoplastic conditions**Lingual haemangiosarcoma**

- Reported to be at increased risk in a large case series

(Dennis *et al.*, 2006)

Testicular neoplasia

See under *Reproductive conditions*

Neurological conditions**Acute non-compressive nucleus pulposus extrusion**

- Border collies over-represented in a UK case series

(Fenn *et al.*, 2016)

Adult-onset deafness

- Onset between 3 and 7 years
- Autosomal dominant pattern of inheritance suspected
- 4/6 dogs aged 12 years or older were deaf in one or both ears

(Schmutz, 2014)

Border Collie collapse

- Reported in a US case series of 165 Border Collies
- Median age of onset 2 years
- Genetic basis suspected

(Taylor *et al.*, 2016a, 2016b)

Congenital deafness

- Prevalence of 2.8% estimated in a UK population of this breed in one UK study
- In another UK study, 2.0% were unilaterally deaf and 0.4% bilaterally deaf
- Higher rates of merle coat pigmentation, blue iris pigment, and excess white on the head compared to Border Collies with normal hearing ($p < 0.001$ for all of these)
- Suspected to be inherited

(Platt *et al.*, 2006; De Risio *et al.*, 2011)

Epilepsy

- Often juvenile onset in this breed
- Reduced mean survival time compared to other breeds ($p = 0.01$)
- 13/136 dogs with juvenile-onset epilepsy were this breed in a UK population
- 11/13 Border Collies experienced cluster seizures in this study
- In a German study of Border Collies with idiopathic epilepsy, 49% had a severe course of the disease, and drug resistance was found in 71% of dogs treated with two or more antiepileptic drugs
- Mutations in the *ABCB1* gene were associated with antiepileptic drug resistance in this breed

(Hulsmeyer *et al.*, 2010;

Alves *et al.*, 2011; Arrol *et al.*, 2012)

Neuronal ceroid lipofuscinosis

- Autosomal recessive inheritance
- Signs seen at 1–2 years
- Gene responsible identified in a Australian study
- Carrier frequency estimated at 8.1% in a Japanese study

(Melville *et al.*, 2005;

Mizukami *et al.*, 2011)

Steroid-responsive meningitis–arteritis (SRMA)

- At increased risk in a UK study: multivariable analysis showed OR 6.91 (95% CI 2.20–21.66; $p = 0.001$)
- All cases < 2 years of age at presentation

(Rose *et al.*, 2014)

Ocular conditions**Chronic superficial keratitis (pannus)**

- Breed at risk in case series, suspected to be inherited
- Age of onset: 4–7 years
- Prevalence and severity increase at higher altitude

(Chavkin *et al.*, 1994)

Collie eye anomaly

- Congenital disorder; autosomal recessive inheritance suspected
- 6% prevalence in this breed in a 1982 UK survey

(Bedford, 1982a)

Lens luxation – primary

- Simple autosomal recessive inheritance suggested
- Age of onset: 3–5 years (mean 4.7 years)

(Foster *et al.*, 1986)**Neuronal ceroid lipofuscinosis**See under *Neurological conditions***Progressive retinal atrophy (PRA)**

- X-linked, generalized and central forms of PRA have been identified in this breed
- 33/161 dogs had retinal lesions in a French study
- Frequency decreased from 20% in 2001 to 8% in 2012 in a French study
- 91/326 dogs of this breed in a Norwegian study were affected by multifocal retinal degeneration

(Chaudieu, 2001; Vilboux *et al.*, 2008; Kjær *et al.*, 2010; Chaudieu *et al.*, 2014)**Physiological conditions****Methylmalonic aciduria**

- Found as an incidental finding in healthy eucobalaminaemic Border Collies and Border Collies with clinical signs of cobalamin deficiency

(Lutz *et al.*, 2012)**Reproductive conditions****Testicular neoplasia**

- Breed at increased risk of Sertoli cell tumour in case series
- Common tumour in male dog
- Median age 9.5 years

(Weaver, 1983)

BORDER TERRIER**Dermatological conditions****Generalized sebaceous gland hyperplasia**

- Possible genetic predisposition

(Dedola *et al.*, 2010)**Neurological conditions****Canine epileptoid cramping syndrome**

- First episode usually occurs before 3 years of age

(Black *et al.*, 2014)**Epilepsy**

- OR 2.70 (95% CI 1.57–4.62; $p < 0.001$) in a UK general practice population
- As these cases were not from a referral population, some of the cases seen may represent a misdiagnosis of canine epileptoid cramping syndrome
- Also reported in a German survey of breeders and owners, where incidence was estimated at 13.1%

(Kloene *et al.*, 2008;Kearsley-Fleet *et al.*, 2013)**Ocular conditions****Corneal ulceration (ulcerative keratitis)**

- OR 2.21 (95% CI 1.18–4.14; $p = 0.014$) compared to crossbreeds in a UK general practice population

(O'Neill *et al.*, 2017b)**Renal and urinary conditions****Ectopic ureter**

- Significantly increased risk in this breed in a UK population

(Holt *et al.*, 2000)**Reproductive conditions****Dystocia**

- Odds ratio in a UK first-opinion practice study of 4.2 (95% CI 2.0–8.5; $p < 0.001$)

(O'Neill *et al.*, 2017c)**BORZOI****Gastrointestinal conditions****Gastric dilatation/volvulus (bloat, GDV)**

- Breed at increased risk

(Burrows & Ignaszewski, 1990)

Neurological conditions**Cervical spondylomyelopathy (cervical vertebral malformation, wobblers syndrome)**

- Reported to be inherited as an autosomal recessive trait in a small group of related Borzois

(Jaggy *et al.*, 1988)

Ocular conditions

Inherited focal retinal degeneration

- Inherited as a simple autosomal trait
- Reported in a French population (Chaudieu, 1995)

Lens luxation – secondary

- Significant predisposition in this breed
- Swiss population (Betschart *et al.*, 2014)

Multifocal chorioretinitis (Borzoi chorioretinopathy)

- Inheritance suspected, but not by simple autosomal or sex-linked pattern
- Reported in dogs aged 7 months to 7 years
- Some reports suggest a male predisposition (Storey *et al.*, 2005)

BOSTON TERRIER**Dermatological conditions**

Atopic dermatitis (atopy)

- OR 2.0 compared to an American hospital population
- Age of onset usually 4–6 years (Scott, 1981)

Calcinosis circumscripta

- This breed is predisposed to lesions on the cheek and at the base of the pinna (Scott & Buerger, 1988)

Localized parakeratotic hyperkeratosis

- Reported in a US case series of 16 dogs of this breed (Lee *et al.*, 2016)

Endocrine conditions

Hyperadrenocorticism (Cushing's syndrome)

- Reported to be common in this breed in a case series of 153 dogs in an American population (Wood *et al.*, 2007)

Gastrointestinal conditions

Atresia ani

- This breed showed an increased incidence of this rare disease in an American population
- Females predisposed (Vianna & Tobias, 2005)

Musculoskeletal conditions

Patellar luxation

- Common condition
- OR 4.2 (95% CI 2.9–6.0) compared to mixed breeds (LaFond *et al.*, 2002)

Perineal herniation

- Intact males predisposed (Hosgood *et al.*, 1995)

Neoplastic conditions

Chemodectomas (aortic body and carotid body)

- Breed at significantly increased risk
- Older male dogs predisposed
- 5/16 cases in a recent US study were of this breed (Hayes & Sass, 1988; Mai *et al.*, 2015)

Mammary neoplasia

- Increased incidence in this breed in a US population (MacVean *et al.*, 1978)

Mast cell tumour (MCT)

- Relative risk (RR) > 8 in a 1969 US study (Peters, 1969)

Pituitary tumour resulting in hyperadrenocorticism

See under *Endocrine conditions*

Primary brain tumour

See under *Neurological conditions*

Neurological conditions

Primary brain tumour

- 8/41 post-mortem examinations of this breed had this condition ($p=0.0001$)
- Increased risk of glial neoplasms in this breed (astrocytomas and oligodendrogliomas) (Song *et al.*, 2013)

Ocular conditions

Cataract

- Autosomal recessive inheritance suspected
- Significantly increased odds ratio for the condition in this breed
- Prevalence of primary cataract 11.11%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

- Early- and late-onset inherited cataracts in this breed are thought to be genetically distinct conditions
(Adkins & Hendrix, 2005; Gelatt & MacKay, 2005; Mellersh *et al.*, 2006, 2007)

Glaucoma – primary

- Suspected to be inherited
- 2.88% prevalence in this breed in a North American population
(Gelatt & MacKay, 2004a)

Glaucoma – secondary

- This breed at increased risk compared to a US hospital population
- Increased risk of glaucoma and other complications after phacoemulsification for cataract removal in this breed. Odds ratio of 290.44 of postoperative blindness in this breed compared to mixed breeds (95% CI 4.77–17.673; $p = 0.007$)
- 10.5% of cases in this breed eviscerated or enucleated after cataract surgery
(Lannek & Miller, 2001; Johnsen *et al.*, 2006; Klein *et al.*, 2011; Scott *et al.*, 2013)

Lens luxation – secondary

- Significant predisposition in this breed
- Swiss population
(Betschart *et al.*, 2014)

Traumatic proptosis

- One of the most commonly affected breeds in a Brazilian population
(Brandão *et al.*, 2005)

Uveal cysts

- Breed at increased risk
- Mean age 6.8 years in one study
(Corcoran & Koch, 1993)

Renal and urinary conditions

Hypospadias

- Congenital defect with a higher incidence in this breed, suggesting a possible genetic basis
- Predominantly affects male dogs
(Hayes & Wilson, 1986)

Reproductive conditions

Dystocia

- Breed predisposition to obstructive dystocia due to dorsoventrally flattened maternal pelvic canal and large fetuses with large heads

- 92.3% of this breed required caesarean sections
- OR 17.1 (95% CI 7.8–37.6; $p < 0.001$) in a UK first-opinion series
(Eneroth *et al.*, 1999; Evans & Adams, 2010a; Martins-Bessa *et al.*, 2015; O'Neill *et al.*, 2017c)

Hypospadias

See under *Renal and urinary conditions*

Respiratory conditions

Brachycephalic obstructive airway syndrome (BOAS)

- Most commonly present with clinical signs aged 1–4 years
- 9% of cases were this breed in one US study
(Riecks *et al.*, 2007; Fasanella *et al.*, 2010)

Tracheal hypoplasia

- This breed represented 15% of cases in an American case series
- Age at diagnosis from 2 days to 12 years
(Coyné & Fingland, 1992)

BOUVIER DES FLANDRES

Gastrointestinal conditions

Muscular dystrophy of pharyngeal and oesophageal muscles causing dysphagia

- Hereditary
(Peeters *et al.*, 1991; Peeters & Ubbink, 1994)

Musculoskeletal conditions

Elbow dysplasia

- Fragmented coronoid process: OR 19.5 (95% CI 8.1–46.9) compared to mixed breeds
(LaFond *et al.*, 2002)

Hip dysplasia

- Neutered male dogs predisposed
- OR 4.1 (95% CI 2.8–6.1) compared to mixed breeds
(LaFond *et al.*, 2002)

Muscular dystrophy

See under *Gastrointestinal conditions*

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 12.1 (95% CI 5.8–25.4) compared to mixed breeds
(LaFond *et al.*, 2002)

Neoplastic conditions

Gastric carcinoma

- Males predisposed
- Median age 10 years
- Proportional mortality ratio 36.5 in a Norwegian population

(Seim-Wikse *et al.*, 2013)

Prostatic carcinoma

- OR 8.44 (95% CI 4.38–16.1)
- Older dogs affected
- Some studies suggest an increased risk in castrated dogs

(Teske *et al.*, 2002)

Ocular conditions

Glaucoma – primary

- Inheritance suspected
- Goniodysgenesis common

(van der Linde-Sipman, 1987)

Reproductive conditions

Prostatic carcinoma

See under *Neoplastic conditions*

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Age of onset 4–6 months
- Neurogenic and hereditary in this breed

(Venker-van Haagen *et al.*, 1978)

BOXER

Behavioural conditions

Aggression

- Increased risk of being involved in human and animal attacks in a German study
- In an Australian study, Boxers were only reported to attack animals

(Blackshaw, 1991; Unshelm *et al.*, 1993)

Cardiovascular conditions

Aortic stenosis – subaortic stenosis (SAS)

- High prevalence of murmurs consistent with aortic stenosis in this breed in various populations, for example 77% in a Norwegian and Swedish population
- Male predisposition

(Heiene *et al.*, 2000; Chetboul *et al.*, 2006b; Bussadori *et al.*, 2009)

Arrhythmogenic right ventricular cardiomyopathy

- Familial and thought to be inherited
- Prognosis better for younger dogs without syncope
- All cases in the UK thought to come from a small number of Boxers imported from the USA

(Basso *et al.*, 2004; Caro-Vadillo *et al.*, 2013; Cattanaach *et al.*, 2015)

Atrial septal defect

- 31.9% of cases were this breed in a French cardiology population
- OR 15.28 (95% CI 10.24–22.84; $p < 0.05$) compared to hospital population
- No sex predisposition
- Mean age of diagnosis 4.1 years

(Chetboul *et al.*, 2006a, 2006b)

Dilated cardiomyopathy (DCM)

- At least one gene has been identified that is associated with the DCM phenotype in this breed
- This breed may exhibit the ‘fatty infiltration–degenerative’ type, or the ‘attenuated wavy fibre’ type of DCM

(Tidholm & Jonnson, 1997; Meurs *et al.*, 2013)

Mitral valve dysplasia

- Recognized as common in this breed in a French population

(Chetboul *et al.*, 2006b)

Pulmonic stenosis

- Congenital
- 3.3% prevalence in this breed in a Swiss study

(Matic, 1988; Chetboul *et al.*, 2006b; Höpfner *et al.*, 2010)

Dermatological conditions

Atopic dermatitis (atopy)

- Above average risk for this condition in this breed in a Swedish population
- OR 1.9 (95% CI 1.06–2.67) compared to Australian hospital population

(Nødtvedt *et al.*, 2006; Mazrier *et al.*, 2016)

Calcinosis circumscripta

- No sex predisposition
- This breed is predisposed to lesions on the cheek and at the base of the pinna
- Commonly reported breed in a US population

(Scott & Buerger, 1988; Doerr *et al.*, 2013)

Canine follicular dysplasia (seasonal flank alopecia)

- Presumed genetic basis
- Seen concurrently with interface dermatitis in one case series in a US population
- Alopecia starts at 2–4 years of age and is restricted to the flank in this breed (Miller & Dunstan, 1993; Rachid *et al.*, 2003)

Canine leproid granuloma

See under *Infectious conditions*

Demodicosis

- Predisposed in an Argentine population (Barrientos *et al.*, 2013)

Idiopathic sterile granuloma and pyogranuloma

- This breed over-represented in one US study (Panich *et al.*, 1991)

Otitis externa

- Breed over-represented in a Polish study (Sapierzynski, 2009)

Skin tumours

See under *Neoplastic conditions*

Endocrine conditions**Hyperadrenocorticism (Cushing's syndrome)**

- Breed at increased risk in case series
- Median age 10 years (Ling *et al.*, 1979)

Hypothyroidism

- Breed at increased risk
- May occur at a younger age in breeds at risk (2–3 years)
- Females and neutered males at increased risk (Nesbitt *et al.*, 1980)

Thyroid neoplasia

- Breed at increased risk in a Swiss study (Wenger *et al.*, 2005)

Gastrointestinal conditions**Cleft palate**

- Congenital
- Probable autosomal recessive inheritance
- Prevalence of 2.3% in a Dutch population (Nielen *et al.*, 2001)

Folate – low serum level

- Reported to be significantly at risk in a UK clinical pathology laboratory study (Dandrieux *et al.*, 2013)

Gastric dilatation/volvulus (bloat, GDV)

- 5.4% of cases in a Brazilian post-mortem population were Boxers (Castro *et al.*, 2013)

Gingival and oropharyngeal neoplasia

See under *Neoplastic conditions*

Inflammatory bowel disease

- Increased risk reported in UK population (OR 1.70, 95% CI 1.04–2.76; $p = 0.0328$) (Kathrani *et al.*, 2011)

Pancreatitis

- Breed at risk of chronic pancreatitis in case series of post-mortem dogs
- Relative risk 3.0 (95% CI 2.4–3.8) (Watson *et al.*, 2007)

Splenic disease

- Boxers significantly over-represented in an Australian histopathological case series ($p = 0.0148$) compared to Australian National Kennel Council population data (Christensen *et al.*, 2009)

Ulcerative colitis (histiocytic ulcerative colitis)

- Occurs predominantly in this breed (Churcher & Watson, 1997; German *et al.*, 2000a)

Haematological/immunological conditions**Eccentrocytosis**

- Boxers reported to be over-represented in an Italian case series (Caldin *et al.*, 2005)

Factor I deficiency

- Autosomal inheritance (Brooks, 1999)

Factor II deficiency

- Autosomal inheritance (Brooks, 1999)

Haemophilia A

- Severe factor VIII deficiency in this breed
- Familial in this breed (Brooks, 1999)

Inflammatory myopathy

- This breed over-represented in a US case series (Evans *et al.*, 2004)

Infectious conditions**Babesiosis**

- This breed over-represented in a Brazilian population (Ungar de Sá *et al.*, 2007)

Canine leproid granuloma

- Boxers most commonly affected in a Brazilian population (Conceição *et al.*, 2011)

Cryptococcosis

- Breed at increased risk in an Australian case series
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors (O'Brien *et al.*, 2004)

Leishmaniasis

- Breed significantly over-represented in a Spanish population (Miranda *et al.*, 2008)

Protothecosis

- 7/17 cases in an Australian population were Boxers or Boxer crosses (Stenner *et al.*, 2007)

Musculoskeletal conditions**Cranial cruciate ligament (CCL) disease**

- Neutered individuals are predisposed
- Older animals are predisposed
- OR in this breed 2.14 (95% CI 2.00–2.30; $p < 0.01$)
- In a UK referral population, 9% with this condition were Boxers (Witsberger *et al.*, 2008; Guthrie *et al.*, 2012)

Hip dysplasia

- Heritability of 0.24 in an Italian population (Sturaro *et al.*, 2006)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males possibly predisposed
- OR in this breed 18.4 (95% CI 7.1–47.5) (LaFond *et al.*, 2002)

Osteochondrosis – lumbosacral

- This breed heavily represented in a UK study
- Mean age at diagnosis 6.3 years (Hanna, 2001)

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 2.2 (95% CI 1.1–4.5) compared to mixed breeds (LaFond *et al.*, 2002)

Osteochondrosis – stifle

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 56.3 (95% CI 24.8–127.8) compared to mixed breeds (LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.8 (95% CI 1.4–2.3) compared to mixed breeds (LaFond *et al.*, 2002)

Perineal herniation

- Intact males predisposed (Robertson, 1984)

Radial carpal bone fracture

- 11/15 dogs were Boxers in a UK case series (Li *et al.*, 2000)

Spondylosis deformans

- Usually clinically insignificant
- 84% of boxers had spondylosis lesions in one Italian study (Canier *et al.*, 2004)

Neoplastic conditions**Canine cutaneous histiocytoma**

- Breed at increased risk in Australian case series (9/80 cases were this breed) (Er & Sutton, 1989)

Chemodectomas (aortic body and carotid body)

- Breed at significantly increased risk
- Older male dogs predisposed (Hayes & Sass, 1988)

Chondrosarcoma (skeletal)

- 24% of cases in an American case series were Boxers (Brodey *et al.*, 1974)

Fibroadnexal hamartoma

- Increased risk in a Brazilian population
- Average age of onset 6.3 years
- No sex predisposition
(Loures & Conceição, 2009)

Gingival and oropharyngeal neoplasia

- Breed at increased risk in case series
- Particularly prone to epuli
(Dorn & Priester, 1976; Delverdier *et al.*, 1991)

Haemangioendothelioma

- Significant predisposition in this breed in a German post-mortem study
(Giesel *et al.*, 1986)

Lymphoma

- Breed at increased risk
- Most cases are seen in middle-aged dogs (mean 6–7 years)
- OR 3.26 (95% CI 1.57–6.76; $p=0.002$) in a UK population, compared to other breeds
(Edwards *et al.*, 2003; Lurie *et al.*, 2004)

Mammary neoplasia

- Lifetime prevalence of 0.38 in a Norwegian survey
- Incidence rate (IR) of 35.47 malignant tumours per 1000 bitches per year in a Norwegian population
(Moe, 2001; Dahl *et al.*, 2002)

Mast cell tumour (MCT)

- 12/53 dogs with mast cell tumours were Boxers in an Australian case series
- Boxers presented significantly younger than other breeds
- OR 10.7 (95% CI 3.7–30.4) in a UK study
(Baker-Gabb *et al.*, 2003; Warland & Dobson, 2013; Shoop *et al.*, 2015)

Osteosarcoma

- Breed at risk in a US case series
- Males over-represented
- Seen in older dogs
(Misdrop & Hart, 1979)

Pituitary tumour resulting in hyperadrenocorticism
See under *Endocrine conditions***Primary brain tumour**See under *Neurological conditions***Testicular neoplasia**See under *Reproductive conditions***Thyroid neoplasia**See under *Endocrine conditions***Transmissible venereal tumour**

- Breed significantly predisposed in a Brazilian population
(Brandão *et al.*, 2002)

Vaginal/vulval neoplasia

- This breed reported to be predisposed in a Polish study
(Sapierzyński, 2007)

Vascular tumours

- Boxers were reported to be at increased risk of primary vascular neoplasia in a Norwegian population
- Annual incidence rate (IR) of primary malignant vascular tumours in this breed of 1.5 per 1000 dogs
(Moe *et al.*, 2008)

Neurological conditions**Central vestibular disease**

- Reported to be more common in Boxers than in mixed-breed dogs in a Brazilian population
(Chaves *et al.*, 2014)

Degenerative myelopathy

- Reported to have a high incidence in a Belgian population
- 14/52 Boxers in a South African study were carriers and one was affected
(Zeiler *et al.*, 2013; Beckers *et al.*, 2016)

Discoispondylitis

- Young/middle-aged dogs affected
- Males twice as likely to be affected as females
- Odds ratio in this breed 3.5 (95% CI 1.8–6.9) in a US study
(Burkert *et al.*, 2005)

Epilepsy

- Prevalence of 2.4% in a Dutch population
- More likely to suffer cluster seizures compared to Labradors ($p=0.01$)
(Nielen *et al.*, 2001; Monteiro *et al.*, 2012)

Idiopathic head tremor syndrome

- 13% of cases were this breed in one multinational survey of vets

(Shell *et al.*, 2015)

Primary brain tumour

- Older dogs affected
- 16/18 dogs of this breed with brain tumours had gliomas in one study
- Boxers significantly over-represented in a US post-mortem case series ($p = .0001$, 28/212 cases)
- Boxers also over-represented in a histological case series of meningiomas

(Snyder *et al.*, 2006; Sturges *et al.*, 2008;

Song *et al.*, 2013)

Progressive axonopathy

- Autosomal recessive inheritance
- Reported in a UK case series

(Griffiths *et al.*, 1985)

Steroid-responsive meningitis–arteritis (SRMA)

- Multivariable analysis of a UK population showed an OR of 4.39 (95% CI 1.14–16.98; $p = 0.032$)
- Prognosis better in this breed than in other breeds

(Behr & Cauzinille, 2006; Rose *et al.*, 2014)

Ocular conditions**Distichiasis**

- Breed at increased risk

(Lawson, 1973)

Spontaneous chronic corneal epithelial defects (refractory corneal ulceration, indolent ulcers)

- Breed at increased risk
- Age of onset 7–9 years
- Odds ratio for corneal ulceration in this breed 13.84 (95% CI 10.05–19.06; $p < 0.001$) compared to crossbreeds in a UK general practice population

(Bentley, 2005; Chandler *et al.*, 2010; O'Neill *et al.*, 2017b)

Physiological conditions**Blood group**

- <20% of Boxers were DEA 1.1-positive in a South African study
- All Boxers were DEA 1.1-negative in one Portuguese and one Swiss study

(van der Merwe *et al.*, 2002;

Ferreira *et al.*, 2011; Riond *et al.*, 2011)

Polyodontia

- 16/47 dogs in a German case series with supernumerary teeth were Boxers

(Kuiper *et al.*, 1982)

Renal and urinary conditions**Renal disease**

- Incidence rate (IR) in a large Swedish case series of 36 cases per 10 000 DYAR (95% CI 27–44), mean age of diagnosis 5.3 years
- Juvenile nephropathy reported in a UK case series

(Chandler *et al.*, 2007; Pelander *et al.*, 2015)

Urethral sphincter mechanism incompetence

- Breed at risk in case series (65% of Boxers reported to be affected in a German study)
- Spayed females predisposed

(Blendinger *et al.*, 1995; Arnold, 1997)

Reproductive conditions**Cryptorchidism**

- Congenital defect believed to be inherited as a sex-limited autosomal recessive trait
- Breed at risk in case series
- Prevalence 10.7% in a Dutch population

(Nielen *et al.*, 2001)

Dystocia

- Dystocia occurred in 27.7% of all whelpings in a Swedish population
- Caesarean sections were performed in 22.8% of all the whelpings and in 80.1% of cases of dystocia
- 60% of cases of dystocia were due to uterine inertia
- In a UK first-opinion study, OR was 2.5 (95 CI 1.3–4.8; $p = 0.006$)

(Linde Forsberg & Persson, 2007; O'Neill *et al.*, 2017c)

Testicular neoplasia

- Breed at increased risk
- Seen at earlier age than in other breeds (mean age 7.2 years in one study)
- One study suggests increased risk of Sertoli cell tumours

(Hayes & Pendergrass, 1976; Weaver, 1983)

Vaginal hyperplasia/vaginal prolapse

- This breed most often reported to be affected

(Post *et al.*, 1991)

BOYKIN SPANIEL

Cardiovascular conditions

Pulmonic stenosis

- Reported in 4 dogs with a common ancestry
- May be familial in this breed
- May be a polygenic mode of inheritance (Jacobs *et al.*, 1990)

Haematological/immunological conditions

Haemophilia A

- Severe factor VIII deficiency in this breed
- Familial in this breed (Brooks, 1999)

Musculoskeletal conditions

Hip dysplasia

- 40% prevalence in a North American screening population (Tsai & Murphy, 2006)

Neurological conditions

Exercise-induced collapse

- In a North American study, 36% were heterozygous and 10% homozygous for the *DNM1* mutation causing this condition (Minor *et al.*, 2011)

BRIARD

Neoplastic conditions

Lymphoma

- German population
- Two comparison populations: own clinic ($n=52$ 142) and insured population ($n=123$ 423)
- Odds ratios 5.6 and 9.5 respectively for the two populations (Ernst *et al.*, 2016)

Squamous cell carcinoma – digit

- Over-represented in a French population
- Mean age 10.2 years
- No sex predisposition (Belluco *et al.*, 2013)

Ocular conditions

Hereditary retinal dystrophy of Briards (congenital stationary night blindness)

- Autosomal recessive inheritance with variable expression (Wrigstad *et al.*, 1994; Narfström, 1999a)

Retinal pigment epithelial dystrophy (RPED, central progressive retinal atrophy)

- Autosomal recessive inheritance has been suggested
- Age of onset varies
- A 1984 study showed that 31% of Briards of 18 months or older were affected
- Becoming much less prevalent following the introduction of control schemes (Bedford, 1984, 2009; Lightfoot *et al.*, 1996)

Physiological conditions

Hypercholesterolaemia

- Possible primary abnormality in cholesterol metabolism (Watson *et al.*, 1993)

Renal and urinary conditions

Ectopic ureter

- Significant increased risk in this breed in a UK population (Holt *et al.*, 2000)

BRITISH BULLDOG

See *Bulldog – English*

BRITTANY SPANIEL

Dermatological conditions

Grass awn migration

- Predisposition due to behaviour
- Increased prevalence in this breed compared to hospital population (Brennan & Ihrke, 1983)

Otitis externa

- This breed over-represented in a Greek case series (Saridomichelakis *et al.*, 2007)

Sterile nodular panniculitis

- This breed significantly over-represented in a series of 39 dogs in an American population (Contreary *et al.*, 2015)

Gastrointestinal conditions

Cleft palate

- Autosomal recessive inheritance
- 26.9% incidence in a breeding colony (Richtsmeier *et al.*, 1994)

Haematological/immunological conditions

Deficiency of third component of complement

- Discovered in a colony of Brittany Spaniels with inherited spinal muscular atrophy
- Inherited as an autosomal recessive
- Inherited separately from the spinal atrophy gene (Blum *et al.*, 1985)

Infectious conditions

Histoplasmosis

- Breed at increased risk
- Most dogs are <2 years, males predisposed
- Mostly seen in the Americas, India and Southeast Asia

(Selby *et al.*, 1981;

Clinkenbeard *et al.*, 1989)

Musculoskeletal conditions

Hip dysplasia

- Neutered male dogs predisposed
- OR 1.7 (95% CI 1.3–2.3) compared to mixed breeds

(LaFond *et al.*, 2002)

Spinal muscular atrophy

See under *Neurological conditions*

Neoplastic conditions

Thyroid neoplasia

- Breed over-represented in a French case series (Cohn-Bendit, 1995)

Neurological conditions

Spinal muscular atrophy

- Autosomal dominant inheritance
- Age of clinical onset <1 year (Cork *et al.*, 1982, 1990)

Spinocerebellar degeneration (late onset)

- Signs seen at 7–13 years
- Familial
- Reported in a small American case series (Higgins *et al.*, 1998)

Ocular conditions

Lens luxation – primary

- Reported in a French population
- *ADAMTS17* mutation responsible for this condition in some breeds not identified in cases of primary lens luxation in this breed in a UK population (Chaudieu *et al.*, 1993; Gould *et al.*, 2011)

Sudden acquired retinal degeneration (SARD, amaurosis)

- 7/17 dogs in a French case series with this condition were Brittany Spaniels (Goulle, 2010)

Renal and urinary conditions

Renal amyloidosis

- Breed over-represented in a French case series
- 54% of cases were aged 5–8 years (Pagès, 1988)

BRUSSELS GRIFFON

See *Griffon Bruxellois*

BUHUND

See *Norwegian Buhund*.

BULL TERRIER

Cardiovascular conditions

Aortic stenosis – subaortic stenosis (SAS)

- Inheritance possibly autosomal dominant with modifying genes, or polygenic
- Mean age of onset for acquired disease 7.6 years
- Associated with polycystic kidney disease in this breed

(O'Leary *et al.*, 2005;

O'Leary and Wilkie, 2009)

Mitral dysplasia

- Congenital
- Genetic basis suspected
- Associated with polycystic kidney disease in this breed (Malik & Church, 1988; O'Leary *et al.*, 2005; O'Leary & Wilkie, 2009)

Dermatological conditions

Acrodermatitis

- Inherited as an autosomal recessive trait
- Low IgA levels may predispose to infection
- May be associated with zinc deficiency (McEwan *et al.*, 2000, 2003)

Atopic dermatitis (atopy)

- Dogs between 1 and 2 years of age have highest probability of an insurance claim for atopy

- This breed had a risk factor of 21 cases per 1000 DYAR (if 1000 dogs were followed for 1 year, 21 would have an insurance claim for atopy)
- Some studies show no sex predilection, others show females predisposed
- Increased risk of adverse food reaction in this breed according to a Hungarian study (Nødtvedt *et al.*, 2006; Tarpataki *et al.*, 2006)

Otitis externa

- 4% of cases were this breed in a Romanian case series (Mircean *et al.*, 2008)

Haematological/immunological conditions

Acrodermatitis

See under *Dermatological conditions*

Musculoskeletal conditions

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.6 (95% CI 1.0–2.4) compared to mixed breeds (LaFond *et al.*, 2002)

Patellar luxation

- Mainly medial luxation observed
- Breed predisposed in a Korean case series (Alam *et al.*, 2007)

Neoplastic conditions

Lymphoma

- Breed over-represented in an Australian case series (Wyatt & Robertson, 1998)

Mast cell tumour (MCT)

- Breed at increased risk in an Australian case series
- May be seen at any age (from 4 months onwards), but usually seen in older animals (Er & Sutton, 1989)

Squamous cell carcinoma

- Breed over-represented in a Mexican case series (Silva-Hidalgo *et al.*, 2015)

Neurological conditions

Congenital deafness

- 96.29% of deaf puppies of this breed had a white coat (Strain, 2004; De Risio *et al.*, 2016b)

Tail-chasing

- 14/32 dogs with this condition were this breed in an Australian population
- Other abnormal behaviours may also be seen in this breed, possibly due to complex partial seizures

(Blackshaw *et al.*, 1994;

Dodman *et al.*, 1996;

Moon-Fanelli & Dodman, 1998)

Trance-like syndrome

- Reported in a UK case series
- Possibly a compulsive disorder (Lowrie *et al.*, 2015a)

Renal and urinary conditions

Familial renal disease (glomerular basement membrane disorder)

- Autosomal dominant inheritance
- Condition believed to be similar to Alport syndrome in humans
- Proteinuria may be an early indicator, with cases progressing to renal failure at 1–8 years of age
- Most reports of the condition come from Australia
- May be seen concurrently with polycystic kidney disease

(Hood *et al.*, 2002a;

O'Leary & Atwell, 2003)

Polycystic kidney disease (PKD)

- Autosomal dominant inheritance
- Bilateral cysts (1–25 mm) found in cortex and medulla
- May be seen concurrently with hereditary nephritis (see *Familial renal disease*)
- Dogs with PKD had an increased risk of mitral valve disease and left ventricular out-flow obstruction (O'Leary *et al.*, 1999, 2002, 2005)

Respiratory conditions

Tracheal collapse

- Thought to be primary in this breed
- Reported in a UK case series (Spaull & Friend, 2014)

BULL TERRIER – MINIATURE

See *Miniature Bull Terrier*

BULLDOG – AMERICAN**Dermatological conditions**

Ichthyosis

- 32/545 dogs of this breed homozygous for the suspected mutation
- All affected dogs homozygous and their parents heterozygous
- Autosomal recessive inheritance
- US population

(Mauldin *et al.*, 2015)**Neurological conditions**

Neuronal ceroid lipofuscinosis

- Autosomal recessive mode of inheritance
- Affected American Bulldogs were homozygous for the A allele of a G to A transition in the cathepsin D gene (*CTSD*)
- A allele not detected in DNA samples from 131 randomly selected dogs of 108 breeds other than American Bulldog
- A allele had a frequency of 0.28 in 123 American Bulldogs tested

(Evans *et al.*, 2005; Awano *et al.*, 2006)**Ocular conditions**

Iridociliary cysts

- Reported in a case series of 7 dogs of this breed
- Brazilian and Spanish population
- Some dogs closely related, so possibly inherited

(Pereira *et al.*, 2014)

Neuronal ceroid lipofuscinosis

See under *Neurological conditions***BULLDOG – ENGLISH
(BRITISH BULLDOG)****Behavioural conditions**

General behavioural traits

- In a questionnaire survey, this breed reported to jump up at people frequently, and to be active and hard to train

(Ricciarelli, 2016)

Cardiovascular conditions

Pulmonic stenosis

- May be polygenic mode of inheritance
- This breed accounted for 15.3% of cases of this condition in a Canadian referral population

- May be associated with anomalous coronary artery development in this breed
(Darke, 1989; Ramos *et al.*, 2014)

Tetralogy of Fallot

- Congenital

(Darke, 1989)

Ventricular tachycardia

- Reported in a small Italian case series
- Possibly associated with segmental arrhythmogenic right ventricular cardiomyopathy
- Arrhythmogenic right ventricular cardiomyopathy may be an emerging disease in this breed

(Santilli *et al.*, 2011, 2014)**Dermatological conditions**

Atopic dermatitis (atopy)

- Breed at increased risk in an Australian case series

(Mazrier *et al.*, 2016)

Demodicosis

- Younger dogs predisposed
- Susceptibility to generalized demodicosis may be inherited
- Increased incidence in this breed reported in a Russian population

(Kuznetsova *et al.*, 2012)

Idiopathic nasodigital hyperkeratosis

- Breed possibly predisposed in an American case series

(Scott & Miller, 2012a)

Idiopathic sterile granuloma and pyogranuloma

- This breed over-represented in one US study
(Panich *et al.*, 1991)

Gastrointestinal conditions

Cleft palate

- Probable autosomal recessive inheritance
(Mulvihill *et al.*, 1980)

Hiatal hernia

- Breed predisposed in a US case series
(Lorinson & Bright, 1998)

Primary splenic torsion

- English Bulldogs accounted for 11.2% of cases in a US case series
(DeGroot *et al.*, 2016)

Haematological/immunological conditions

Haemophilia A

- Severe factor VIII deficiency in this breed
- Familial in this breed

(Brooks, 1999)

Metabolic conditions

Hypomagnesaemia

- Period of prevalence of 15% in this breed
- Risk ratio (RR) 1.8 (95% CI 1.3–2.7) compared to Boxers
- Possibly related to sleep apnoea or arterial hypertension

(Mellema & Hoareau, 2014)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Neutered individuals are predisposed
- Older animals are predisposed
- OR in this breed 2.14 (95% CI 1.99–2.37; $p < 0.001$)

(Witsberger *et al.*, 2008)

Hip dysplasia

- Torsional deformity common in this breed
- Neutered male dogs predisposed
- OR in this breed 1.27 (95% CI 1.15–1.40; $p < 0.001$)

(Witsberger *et al.*, 2008; Gnudi *et al.*, 2009)

Osteochondrosis – stifle

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 44.2 (95% CI 17.4–112.6) compared to mixed breeds

(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.9 (95% CI 1.5–2.5) compared to mixed breeds

(LaFond *et al.*, 2002)

Patellar luxation

- OR 6.1 (95% CI 4.2–9.0) compared to mixed breeds

(LaFond *et al.*, 2002)

Swimmer puppy syndrome

- 8.33% of cases in a Thai population were English Bulldogs

(Nganvongpanit & Yano, 2013)

Vertebral anomalies

See under *Neurological conditions*

Neoplastic conditions

Chemodectomas (aortic body and carotid body)

- Breed at increased risk
- Older male dogs predisposed

(Hayes & Sass, 1988; Owen *et al.*, 1996)

Granulosa-theca cell ovarian tumour

- This breed reported to be predisposed in Polish and US studies

(Hayes & Young, 1978; Sapierzyński, 2007)

Lymphoma

- Breed at increased risk
- Most cases are seen in middle-aged dogs (mean 6–7 years)
- OR 4.73 (95% CI 1.42–15.83; $p = 0.012$) in a UK population, compared to other breeds

(Edwards *et al.*, 2003)

Perianal (hepatoid) gland adenomas

- Breed at risk in US case series
- Average age was 10.5 years
- Relative risk (RR) in males 4.6 (95% CI 2.11–9.89) compared to VMDB data
- Not seen in females in this breed

(Hayes & Wilson, 1977)

Neurological conditions

Discospondylitis

- Young/middle-aged dogs affected
- Males twice as likely to be affected as females
- OR in this breed 3.0 (95% CI 1.2–7.5) in a US study

(Burkert *et al.*, 2005)

Idiopathic head tremor syndrome

- 37% of cases were this breed in one multinational survey of vets

(Shell *et al.*, 2015)

Otitis media

- This breed has increased risk of having material in middle ear on MRI, compared to the total population undergoing MRI

(Owen *et al.*, 2004)

Ventriculomegaly

- This breed has larger cerebral lateral ventricles than other breeds

(Vite *et al.*, 1997; Ryan *et al.*, 2014)

Vertebral anomalies

- Congenital
- Thoracic vertebral anomalies very common in this breed
(Schlensker & Distl, 2012; Ryan *et al.*, 2017)

Ocular conditions**Corneal ulceration (ulcerative keratitis)**

- OR 6.53 (95% CI 3.96–10.78; $p < 0.001$) compared to crossbreeds in a UK general practice population
(O'Neill *et al.*, 2017b)

Prolapse of the gland of the nictitating membrane ("cherry eye")

- Breed at increased risk
- Usually presents in the first 1–2 years of life
(Morgan *et al.*, 1993; Mazzucchelli *et al.*, 2012; Prémont *et al.*, 2012; Multari *et al.*, 2016)

Renal and urinary conditions**Ectopic ureter**

- Significant increased risk in this breed in UK and US populations
(Hayes, 1984; Holt *et al.*, 2000)

Urethral prolapse

- Generally seen in male dogs at 4 months to 5 years of age
- Odds ratio in a US population 366.99 (95% CI 265.83–506.65) compared to all breeds
(Kirsch *et al.*, 2002; Carr *et al.*, 2014)

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population
- Also reported in a New Zealand population
- Odds of a Bulldog being affected with cystine was 154.1 times greater than for other breeds
(Case *et al.*, 1992; Bartges *et al.*, 1994; Jones *et al.*, 1998)

Urolithiasis – urate

- Breed at risk in case series
- Average age at diagnosis in one large study: males 5.5 years, females 4.6 years
- Males are predisposed
- Odds of a Bulldog being affected with urate was 43.0 times greater than for other breeds
(Houston *et al.*, 2004; Bartges *et al.*, 2014)

Reproductive conditions**Cryptorchidism**

- Congenital defect believed to be inherited as a sex-limited, autosomal recessive trait
- Breed at risk in case series
(Hayes *et al.*, 1985)

Dystocia

- This breed at increased risk of both emergency and elective caesarean section in a North American case series
- Odds ratio in a UK first-opinion practice study of 5.7 (95% CI 3.1–10.5; $p < 0.001$)
(Moon *et al.*, 1998; O'Neill *et al.*, 2017c)

Urethral prolapse

See under *Renal and urinary conditions*

Vaginal hyperplasia/vaginal prolapse

- This breed one of the most often reported to be affected
(Post *et al.*, 1991)

Respiratory conditions**Brachycephalic obstructive airway syndrome (BOAS)**

- Most commonly present with clinical signs aged 1–4 years
- 61% of cases were this breed in one US study
- 19% of this breed in a large UK primary-care study were suffering from upper respiratory tract disorders
- Bronchial disorders also common with this disorder

(Riecks *et al.*, 2007; De Lorenzi *et al.*, 2009; Fasanella *et al.*, 2010; O'Neill *et al.*, 2015b)

Tracheal hypoplasia

- This breed represented 55% of cases in an American case series
- Age at diagnosis from 2 days to 12 years
(Coyne & Fingland, 1992)

BULLMASTIFF**Musculoskeletal conditions****Elbow dysplasia**

- Fragmented coronoid process OR 38.9 (95% CI 19.8–76.5) compared to mixed breeds
(LaFond *et al.*, 2002)

Hip dysplasia

- Neutered male dogs predisposed
- OR 3.5 (95% CI 2.2–5.6) compared to mixed breeds

(LaFond *et al.*, 2002)**Osteochondrosis – hock**

- Affects dogs 5–9 months of age
- OR 85.9 (95% CI 35.8–206.4) compared to mixed breeds

(LaFond *et al.*, 2002)**Osteochondrosis – shoulder**

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 6.7 (95% CI 2.9–15.5) compared to mixed breeds

(LaFond *et al.*, 2002)**Neoplastic conditions****Lymphoma**

- Two German comparison populations – own clinic ($n=52\ 142$) and insured population ($n=123\ 423$)
- Odds ratio 7.8 and 5.0 respectively for the two populations
- Familial in a 1984 study, with estimated incidence of 5000 cases per 100 000 dogs
- High incidence also reported in this breed in a UK study: OR 6.76 (95% CI 2.03–22.51; $p=0.002$)

(Onions, 1984; Edwards *et al.*, 2003; Ernst *et al.*, 2016)**Neurological conditions****Cerebellar degeneration**

- Autosomal recessive inheritance suggested
- Signs seen at 4–9 weeks
- May be seen with hydrocephalus

(Johnson *et al.*, 2001)**Ocular conditions****Entropion**

- Polygenic inheritance suspected
- Associated with macroblepharon

(Read & Broun, 2007)

Progressive retinal atrophy (PRA)

- Autosomal dominant inheritance suspected (may be more than one genetic form)
- Ophthalmoscopic signs at 4 months to 3 years

(Kijas *et al.*, 2003)**Renal and urinary conditions****Urolithiasis – cystine**

- Breed at risk in case series
- Young dogs affected (2–5 years)
- Almost all cases are male

(Case *et al.*, 1992)**Reproductive conditions****Pyometra (cystic endometrial hyperplasia–pyometra complex)**

- Breed over-represented in a UK case series ($p<0.0001$)

(Gibson *et al.*, 2013)**Vaginal hyperplasia/vaginal prolapse**

- This breed one of the most often reported to be affected

(Post *et al.*, 1991)**CAIRN TERRIER****Cardiovascular conditions****Mitral valve disease**

- Odds ratio in a UK study 2.11 (95% CI 1.34–3.32) in males

(Thrusfield *et al.*, 1985)**Dermatological conditions****Atopic dermatitis (atopy)**

- Some studies show no sex predilection, others show females predisposed
- Breed predisposed in a US study
- This breed 5.7 times more frequently diagnosed than the general hospital population

(Schick & Fadok, 1986)

Endocrine conditions**Diabetes mellitus**

- Swedish population
- Incidence rate (IR) 183 per 10 000 DYAR, compared with a mixed-breed IR of 15

(Catchpole *et al.*, 2005; Fall *et al.*, 2007)**Hypoadrenocorticism (Addison's disease)**

- Increased risk in an insured Swedish population
- Relative risk 3.39 (95% CI 2.17–5.06)

(Hanson *et al.*, 2016)**Thyroid neoplasia**See under *Neoplastic conditions*

Gastrointestinal conditions

Chronic hepatitis

- Reported to be predisposed in a UK case series
- Mean age of diagnosis 8 years
(Bexfield *et al.*, 2012a)

Congenital portosystemic shunt

- Breed at risk in case series
- 58/6372 dogs screened in a Dutch population had portosystemic shunt
- Clinical signs usually seen in young dogs < 1 year
- Usually extrahepatic in this breed
- Inheritance autosomal, monogenic or polygenic
(van Straten *et al.*, 2005)

Microvascular portal dysplasia

- Complex inheritance in this breed
- Reported in a US case series
(Schermerhorn *et al.*, 1996;
van Steenbeek *et al.*, 2012)

Musculoskeletal conditions

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- OR 17.9 (95% CI 6.5–49.2) compared to mixed breeds in a US study
(LaFond *et al.*, 2002)

Cranio-mandibular osteopathy (lion jaw)

- This breed reported to be at high risk in a US study
(Munjar *et al.*, 1998)

Patellar luxation

- Odds ratio 1.9 (95% CI 1.1–3.2) compared to mixed breeds in a US study
(LaFond *et al.*, 2002)

Neoplastic conditions

Testicular neoplasia

See under *Reproductive conditions*

Thyroid neoplasia

- Breed significantly over-represented in a UK case series
(Sullivan *et al.*, 1987a)

Ocular conditions

Cataract

- Prevalence of primary cataract 3.89%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Gelatt & MacKay, 2005)

Glaucoma

- May be related to ocular melanosis in this breed (see below)
- Prevalence of 1.82% in this breed in a US case series
(Gelatt & MacKay, 2004a)

Lens luxation – secondary

- Significant predisposition in this breed
- Swiss population
(Betschart *et al.*, 2014)

Ocular melanosis (abnormal pigment deposition)

- Familial, autosomal dominant inheritance suspected
- Predisposes to glaucoma
- Reported in US case series
(Petersen-Jones *et al.*, 2007)

Renal and urinary conditions*E. coli* urinary tract infection

- Commonly reported in males of this breed in a large study of urolithiasis
(Ling *et al.*, 1998a)

Ectopic ureter

- Significant increased risk in this breed in a UK population
(Holt *et al.*, 2000)

Renal dysplasia

- Reported in a US case series of related animals
(Seiler *et al.*, 2010)

Reproductive conditions

Testicular neoplasia

- Breed at increased risk of Sertoli cell tumour in case series
- Mean age 9–11 years
(Weaver, 1983)

CANE CORSO

See *Italian Mastiff*

CATAHOULA LEOPARD DOG**Cardiovascular conditions**

Heart block

- This breed reported to be predisposed to high-grade second-degree or third-degree heart block in a US case series

- Heavier, older and sexually intact female dogs over-represented
(Schrope & Kelch, 2006)

CAVALIER KING CHARLES SPANIEL

Behavioural conditions

Fear behaviour in puppies

- 53% of puppies in this breed exhibited fear behaviour in a UK and US study
(Morrow *et al.*, 2015)

Cardiovascular conditions

Femoral artery occlusion

- 2.3% of dogs of this breed had an undetectable right or left femoral pulse in one study
- A further 4.2% had a weak right or left femoral pulse
- Probably clinically insignificant, due to collateral circulation
- Possible genetic predisposition
(Buchanan *et al.*, 1997)

Mitral valve disease

- Higher hazard of death from this condition in this breed in a UK general practice study
- Heritability for grade of murmur for 4–5-year-old Cavaliers estimated at 0.67
- Inheritance not thought to be due to a single major gene effect
- Odds ratio of cardiovascular disease in this breed in an insured Japanese population of 16.2 (95% CI 14.4–18.2) compared to the Miniature Dachshund
(Lewis *et al.*, 2011a; French *et al.*, 2012; Mattin *et al.*, 2015a, 2015b; Inoue *et al.*, 2016)

Dermatological conditions

Ichthyosis

- Congenital and probably hereditary
- Often associated with keratoconjunctivitis sicca in this breed
(Barnett, 2006; Hartley *et al.*, 2012)

Persistent scratching

See under *Neurological conditions* (Chiari malformation/syringomyelia)

Gastrointestinal conditions

Exocrine pancreatic insufficiency (EPI)

- Breed at risk in case series
(Batchelor *et al.*, 2007)

Hepatic disease

- 11.1% of dogs of this breed had primary hepatic disease in a post-mortem population
(Kent *et al.*, 2016)

Pancreatitis

- Breed at risk of chronic pancreatitis in case series of post-mortem dogs
- Relative risk (RR) 3.2 (95% CI 2.5–4.1)
(Watson *et al.*, 2007)

Haematological/immunological conditions

Immunoglobulin deficiency

- Associated with a protozoal pneumonia (*Pneumocystis carinii*)
 - Inheritance uncertain
- See also under *Infectious conditions*
(Watson *et al.*, 2006)

Infectious conditions

Angiostrongylosis

- Increased risk of infection in this breed in a UK study
(Blehaut *et al.*, 2014)

Pneumocystis carinii infection

- Increased susceptibility in this breed due to immunodeficiency
- See also under *Haematological/immunological conditions* (Immunoglobulin deficiency)
(Watson *et al.*, 2006)

Musculoskeletal conditions

Humeral condylar fractures

- Breed over-represented in a UK case series
(Cockett & Jones, 1985)

Inguinal/scrotal herniation

- Breed over-represented in an Australian population compared to hospital population
(Bellenger, 1996)

Patellar luxation

- May be inherited as an autosomal recessive trait
- Medial luxation more common in this breed
- OR 9.1 (95% CI 3.3–25.1) compared to mixed breeds
(LaFond *et al.*, 2002)

Temporomandibular joint dysplasia

- Usually asymptomatic
(Dickie *et al.*, 2002)

Neoplastic conditions**Anal sac adenocarcinoma**

- OR 3.36 (95% CI 2.17–5.20) in the UK (Polton *et al.*, 2006)

Neurological conditions**Chiari malformation/syringomyelia**

- Also known as caudal occipital malformation syndrome; persistent scratching
- Heritability of syringomyelia 0.37 ± 0.15 SE in a UK population
- Prevalence of asymptomatic syringomyelia in this breed in the UK estimated at 25% by age 12 months and 70% at age 72 months or more
- In a UK general practice population, signs suggestive of this condition had a period prevalence in this breed of 1.6% (95% CI 1.2–2.06%) compared to 0.05% (95% CI 0.04–0.06%) for all breeds
- In a study of all Danish Kennel Club registered Cavaliers, prevalence of 15.4% was found, with heritability of symptomatic syringomyelia estimated at 0.81 (Lewis *et al.*, 2010; Parker *et al.*, 2011; Thøfner *et al.*, 2015; Sanchis-Mora *et al.*, 2016)

Episodic falling

- Reported in UK and USA
- Age of clinical onset: 3–4 months
- Mutation associated with the condition found in a US population
- 12.9% of US population were carriers for this mutation (Gill *et al.*, 2012)

Idiopathic epilepsy

- Inherited in this breed
- Age of onset 6 months to 6 years
- All coat colours affected, but more common in lines originating from whole-colour ancestors from the late 1960s (Rusbridge & Knowler, 2004; Rusbridge, 2005; Driver *et al.*, 2013)

Primary secretory otitis media

- Reported in Swedish and US case series (Stern-Bertholtz, 2003; Cole *et al.*, 2015)

Ocular conditions**Cataract**

- Prevalence of primary cataract 3.90%, compared to 1.61% in mixed-breed dogs, in a retrospective

study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005)

Corneal ulceration (ulcerative keratitis)

- OR 6.74 (95% CI 4.82–9.41; $p < 0.001$) compared to crossbreeds in a UK general practice population (O'Neill *et al.*, 2017b)

Keratoconjunctivitis sicca

- 14/87 cases were this breed in a Danish case series
- One study suggested a predisposition for males
- Associated with ichthyosis in this breed (Nicolet, 2006; Sanchez *et al.*, 2007; Hartley *et al.*, 2012)

Physiological conditions**Giant platelets and thrombocytopenia**

- Affects around 50% of Cavaliers
- Giant platelets may lead to reduced count of platelets if using automated methods
- Plateletcrit gives a more accurate picture of the platelet mass than other automated techniques
- Inherited as an autosomal recessive trait (Pedersen *et al.*, 2002; Cowan *et al.*, 2004; Kelley *et al.*, 2014)

Renal and urinary conditions**Chronic kidney disease**

- Breed predisposed in a UK general practice population
- 52.2% of this breed had renal lesions at post-mortem examination. 16.7% in this series had an ante-mortem diagnosis of chronic kidney disease (O'Neill *et al.*, 2013b; Kent *et al.*, 2016)

Reproductive conditions**Dystocia**

- Odds ratio in a UK first-opinion practice survey of 2.0 (95% CI 1.1–3.9; $p = 0.033$) (O'Neill *et al.*, 2017c)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk in Swedish case series (Egenvall *et al.*, 2001)

Respiratory conditions**Angiostrongylosis**

See under *Infectious conditions*

Brachycephalic obstructive airway syndrome (BOAS)

- 20.5% of cases in an Australian referral population were this breed
(Torrez & Hunt, 2006)

Pneumonia due to *Pneumocystis carinii* infection

See under *Haematological/immunological conditions* and *Infectious conditions*

CHESAPEAKE BAY RETRIEVER

Dermatological conditions

Adult-onset hair loss

- Reported in a US case series
(Cerundolo *et al.*, 2005)

Haematological/immunological conditions

von Willebrand's disease

- This breed is affected by type III disease
- Inherited as an autosomal recessive trait
(Brooks, 1999)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- This breed predisposed in a US case series
- Neutered individuals may be predisposed
(Duval *et al.*, 1999)

Hip dysplasia

- Neutered male dogs predisposed
- OR 4.4 (95% CI 3.0–6.5) compared to mixed breeds in a US study
(LaFond *et al.*, 2002)

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 7.7 (95% CI 3.3–17.8) compared to mixed breeds in a US study
(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.8 (95% CI 1.1–3.0) compared to mixed breeds in a US study
(LaFond *et al.*, 2002)

Ocular conditions

Cataract

- Dominant mode of inheritance with incomplete penetration has been suggested
- Age of onset: young adult (6 months to 2 years), may progress
- 12.5% prevalence in a Norwegian population
- Prevalence of primary cataract 2.79%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Bjerkås, 1991; Gelatt & MacKay, 2005)

Progressive rod–cone degeneration

- Frequency of the disease-causing mutation found to be 0.14 in this breed in a Czech study
(Dostal *et al.*, 2011)

Neurological conditions

Exercise-induced collapse

- 56/320 dogs of this breed were heterozygous (carriers) and 8/320 were homozygous (affected) for the *DNM1* gene mutation causing this condition in a North American study
(Minor *et al.*, 2011)

CHIHUAHUA

Cardiovascular conditions

Mitral valve disease

- Odds ratios in a UK study 4.63 (95% CI 2.37–9.01) in males and 7.62 (3.41–17.01) in females
(Thrusfield *et al.*, 1985)

Patent ductus arteriosus

- Breed over-represented in a US case series
(Henrich *et al.*, 2011)

Gastrointestinal conditions

Gallbladder mucocoele

- OR 2.18 (95% CI 1.58–5.94; $p = 0.002$) in a US study, compared to hospital population
(Kutsunai *et al.*, 2014)

Infectious conditions

Parvovirus

- Breed over-represented in a US case series
(Lefebvre, 2013)

Musculoskeletal conditions

Atlantoaxial subluxation/instability

See under *Neurological conditions*

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- OR 26.8 (95% CI 11.7–61.7) in one US study, compared to mixed breeds (LaFond *et al.*, 2002)

Inguinal/scrotal herniation

- Breed over-represented in an Australian population compared to hospital population (Bellenger, 1996)

Patellar luxation

- Mainly medial luxation observed
- OR 8.9 (95% CI 6.1–12.8) in a US study, compared to mixed breeds
- OR 5.9 (95% CI 4.4–7.9, $p < 0.001$) in a UK general practice case series (LaFond *et al.*, 2002; Alam *et al.*, 2007; O'Neill *et al.*, 2016c)

Neurological conditions

Atlantoaxial subluxation/instability

- Congenital
- Mean age of presentation 2.7 years
- Over-represented in a UK case series (Denny *et al.*, 1988)

Hydrocephalus

- Congenital
- Over-represented in a US case series (Biel *et al.*, 2013)

Necrotizing meningoencephalitis

- Reported in a small US case series (Higgins *et al.*, 2008)

Chiari malformation/syringomyelia

- Increased risk in a UK general practice epidemiological study (OR 7.4, 95% CI 1.24–44.71; $p = 0.028$) (Sanchis-Mora *et al.*, 2016)

Ocular conditions

Cataract

- Prevalence of primary cataract 1.84%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005)

Corneal ulceration (ulcerative keratitis)

- OR 2.04 (95% CI 1.21–3.43; $p = 0.008$) compared to crossbreeds in a UK general practice population (O'Neill *et al.*, 2017b)

Progressive rod cone degeneration

- Mutant allele detected with a frequency of 0.02 in this breed in a Japanese genetic study (Kohyama *et al.*, 2016)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed significantly over-represented in a UK study (Roe *et al.*, 2012)

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population (Case *et al.*, 1992)

Reproductive conditions

Cryptorchidism

- Congenital defect believed to be inherited as a sex-limited, autosomal recessive trait
- Breed at increased risk in US case series (Pendergrass & Hayes, 1975)

Dystocia

- Increased risk in a US case series
- Incidence rates (IR) per 1000 DYAR in a Swedish population were 31.9 (short-hair) and 31.4 (long-hair)
- Odds ratio in a UK first-opinion study was 12.3 (95% CI 8.2–18.2; $p < 0.001$) (Gaudet, 1985; Bergström *et al.*, 2006; O'Neill *et al.*, 2017c)

Eclampsia (puerperal tetany)

- Breed over-represented compared to hospital population in US study (Drobatz & Casey, 2000)

Respiratory conditions

Tracheal collapse

- 4/100 cases were this breed in a UK case series (fourth most common breed affected)
- 8% of cases were this breed in a Korean case series
- Usually acquired in older dogs but can be congenital (White & Williams, 1994; KiChang *et al.*, 2004)

CHIN

See *Japanese Chin*

CHINESE CRESTED DOG

Dermatological conditions

Canine ectodermal dysplasia (hairlessness)

- Chinese Crested is a hairless breed
- The breed is produced by a dominant gene for hypotrichosis being combined with the gene for long hair
- Homozygotes (HH) for hypotrichosis die prenatally
- All Cresteds are Hh if hairless
- hh are coated (called 'powder puffs')
- Three sub-phenotypes have marked differences histologically

(Robinson, 1985;
Drögemüller *et al.*, 2008b;
Wiener *et al.*, 2013)

Neurological conditions

Canine multiple system degeneration

- Presents as a movement disorder
 - Autosomal recessive inheritance
- (O'Brien *et al.*, 2005)

Ocular conditions

Cataract

- Prevalence of primary cataract 5.66%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Keratoconjunctivitis sicca

- Predisposed in a Norwegian population compared to Papillons and Toy/Miniature Poodles
- (Ulfeng *et al.*, 2009)

Lens luxation – primary

- This breed over-represented in a Swiss referral population

(Betschart *et al.*, 2014)

Progressive rod–cone degeneration (PRCD)

- Mutant allele detected with a frequency of 0.02 in a Czech genetic study

(Dostal *et al.*, 2011)

CHINESE SHAR PEI

See *Shar Pei*

CHOW CHOW

Cardiovascular conditions

Heart block

- This breed reported to be predisposed to high-grade second-degree or third-degree heart block in a US case series
- Heavier, older and sexually intact female dogs over-represented

(Schrope & Kelch, 2006)

Dermatological conditions

Alopecia X (castration-responsive alopecia)

- Relative risk 23.5 in a Canadian referral population

(Scott & Paradis, 1990)

Flea-bite hypersensitivity

- Most studies show no breed predisposition, but one French study showed this breed was predisposed

(Prélaud & Guaguere, 1998)

Pemphigus foliaceus

- No age or sex predispositions noted
- OR 12.3 for this breed in a US pathological case series

(Kuhl *et al.*, 1994)

Gastrointestinal conditions

Exocrine pancreatic insufficiency (EPI)

- Breed at risk in a UK clinicopathological case series
- Prevalence significantly higher in this breed than in a control population of insured dogs
- Females over-represented in this breed

(Batchelor *et al.*, 2007)

Gastric carcinoma

See under *Neoplastic conditions*

Melanoma – oral

See under *Neoplastic conditions*

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Neutered individuals are predisposed
- Older animals are predisposed
- OR 1.73 (95% CI 1.55–1.93) compared to a US referral population

(Witsberger *et al.*, 2008)

Elbow dysplasia

- Fragmented coronoid process OR 16.6 (95% CI 8.0–34.7) compared to mixed breeds
- Ununited anconeal process OR 13.3 (95% CI 7.8–22.6)

(LaFond *et al.*, 2002)**Hip dysplasia**

- Neutered male dogs predisposed
- OR in this breed 1.90 (95% CI 1.73–2.07) in one US study
- In another US study odds ratio compared to mixed breeds was 5.4 (95% CI 4.2–7.0)

(LaFond *et al.*, 2002;
Witsberger *et al.*, 2008)**Myotonia**

- Inherited condition
- Onset of signs at around 2 months of age
- Reported in one small case series from New Zealand and one from Australia

(Jones *et al.*, 1977; Farrow & Malik, 1981)**Panosteitis (enostosis, eosinophilic panosteitis)**

- Young males predisposed
- OR 1.7 (95% CI 1.3–2.2) compared to mixed breeds

(LaFond *et al.*, 2002)**Patellar luxation**

- OR 6.1 (95% CI 3.9–9.4) compared to mixed breeds

(LaFond *et al.*, 2002)**Pituitary dwarfism**

- Reported predisposed in a US population

(Scott & Walton, 1986)

Neoplastic conditions**Gastric carcinoma**

- OR 23.53 ($p < 0.01$) in an Austrian case series
- Also reported as one of the most frequently presented breeds in a German case series

(Bilek & Hirt, 2007;
Babo *et al.*, 2012)**Melanoma – oral**

- Breed at increased risk in case series (13/113 cases of oral melanoma at a US diagnostic investigation laboratory)
- Average age 11.4 years

(Ramos-Vara *et al.*, 2000;
Schultheiss, 2006)**Neurological conditions****Dysmyelination of the central nervous system**

- Tremors present from birth
- Reported in five Chow Chows from three different litters
- Not reported recently

(Vandevelde *et al.*, 1978)**Myotonia**See under *Musculoskeletal conditions***Ocular conditions****Cataract**

- Familial, congenital
- May be seen with other ocular anomalies but relationship unclear

(Collins *et al.*, 1992)**Glaucoma**

- Prevalence of 4.7% in a US case series

(Gelatt & MacKay, 2004a)

Reproductive conditions**Pyometra (cystic endometrial hyperplasia–pyometra complex)**

- This breed represented 6.4% of cases of pyometra in a Belgian population despite the breed only comprising 1.6% of the population
- Pyometra also more severe in this breed, with anaemia and endotoxaemia

(Troyer & Schepper, 1989)

CLUMBER SPANIEL**Musculoskeletal conditions****Mitochondrial myopathy**

- Primary defect is in mitochondrial function
- Reported in USA and UK

(Herrtage & Houlton, 1979; Shelton, 1999)

Pyruvate dehydrogenase phosphatase 1 deficiency

- 20% of current Clumber and Sussex Spaniel population are carriers for the affected gene

(Cameron *et al.*, 2007)**Ocular conditions****Cataract**

- Prevalence of primary cataract 5.32%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Reproductive conditions

Dystocia

- One of the top 10 breeds reported to need caesarean sections in a UK study
- Uterine inertia more common problem than physical blockage in this breed (Evans & Adams, 2010a)

COCKER SPANIEL

Behavioural conditions

Aggression

- Reported to be at increased risk of aggression towards the owner, aggression towards strangers and indoor urination/defecation in a Danish study
- In a Spanish study, owner-directed aggression was more common in this breed than in other breeds, and the golden coat colour was more common in aggressive than in non-aggressive Cocker Spaniels (Lund *et al.*, 1996; Amat *et al.*, 2009)

Cardiovascular conditions

Bradyarrhythmia

- This breed reported to be predisposed to high-grade second-degree or third-degree heart block in a US case series
- Heavier, older and sexually intact female dogs over-represented
- Also reported to be predisposed to sinus node dysfunction/sick sinus syndrome in US case series (Schrope & Kelch, 2006; Wess *et al.*, 2006; Ward *et al.*, 2016)

Dilated cardiomyopathy (DCM)

- DCM in this breed is often related to taurine deficiency
- May respond to taurine and L-carnitine supplementation
- Increased prevalence with age
- Approximately three times as common in males as females in this breed
- Thought to be familial or genetic
- Reported common in this breed in a UK case series (Kittleson *et al.*, 1997; Martin *et al.*, 2009)

Dermatological conditions

Atopic dermatitis (atopy)

- Reported to be strongly predisposed in a Greek population (Saridomichelakis *et al.*, 1999)

Grass awn migration

- This breed over-represented in a Czech case series (Crha *et al.*, 2003)

Idiopathic nasodigital hyperkeratosis

- Breed possibly predisposed in an American case series (Scott & Miller, 2012a)

Malassezia dermatitis

- This breed significantly over-represented in American and UK case series (Bond *et al.*, 1996; Mauldin *et al.*, 1997)

Otitis externa

- 60% of cases of severe otitis externa in one US case series were Cocker spaniels
- Significantly over-represented in a Greek study in which grass awns and *Malassezia* were a factor in this breed
- Cockers significantly over-represented in a UK case series (Angus *et al.*, 2002; Saridomichelakis *et al.*, 2007; Zur *et al.*, 2011)

Tail injuries

- 56.6% of undocked working spaniels received a tail injury in one shooting season in a UK study
- Breed also predisposed in a UK general practice case-control study (Diesel *et al.*, 2010; Lederer *et al.*, 2014)

Vitamin-A-responsive dermatosis

- Reported in a small case series in this breed (Scott, 1986)

Endocrine conditions

Hyperadrenocorticism (Cushing's syndrome)

- Reported to be common in this breed in a case series of 153 dogs in an American population (Wood *et al.*, 2007)

Hypoadrenocorticism (Addison's disease)

- Increased risk in an insured Swedish population
- Relative risk (RR) 2.94 (95% CI 1.87–4.43) (Hanson *et al.*, 2016)

Hypothyroidism

- Breed at increased risk in a US case series from the 1960s and 1970s

- May occur at a younger age in breeds at risk (2–3 years)
- Females and neutered males at increased risk (Milne & Hayes, 1981)

Chronic hepatitis

- American and English Cocker Spaniels
- Breed at increased risk
- Males predisposed
- Some cases may be associated with hepatic accumulation of alpha-1 antitrypsin in this breed
- Odds ratios in a 2008 UK population 21.6 (95% CI 9.7–47.9; $p < 0.001$) (American cockers) and 2.4 (1.8–3.2; $p < 0.001$) (English cockers) (Sevelius *et al.*, 1994; Watson, 2004; Bexfield *et al.*, 2012a)

Gallbladder mucocoele

- OR 6.52 (95% CI 3.13–25.53; $p = 0.002$) in a US study, compared to hospital population (Kutsunai *et al.*, 2014)

Oropharyngeal neoplasia

- Breed at risk in a US 1960s and 1970s case series (Dorn & Priester, 1976)

Pancreatitis

- Breed at risk of acute and chronic pancreatitis in a case series of post-mortem dogs
- RR 2.8 (95% CI 2.3–2.5) for acute and chronic pancreatitis combined (Watson *et al.*, 2007)

Haematological/immunological conditions

Factor II deficiency

- Probably inherited as an autosomal recessive trait
- Severe signs in this breed (Brooks, 1999)

Factor X deficiency

- Familial
- May be inherited as an autosomal recessive trait
- Severe signs in this breed (Brooks, 1999)

Haemophilia A

- Severe factor VIII deficiency in this breed
- Familial in this breed (Brooks, 1999)

Immune-mediated haemolytic anaemia (IMHA)

- Usually affects young adult and middle-aged animals
- May be more common in bitches
- May be seasonal variations
- OR 12.2 (95% CI 4.5–33.1; $p < 0.001$) in a US study, compared to hospital population
- Increased risk in a Canadian study ($p = 0.012$) (Carr *et al.*, 2002; Miller *et al.*, 2004)

Immune-mediated thrombocytopenia (IMTP)

- Breed over-represented compared to a US hospital population (O'Marra *et al.*, 2011)

Platelet disorder

- Delta storage pool disorder in this breed (Brooks, 1999)

Infectious conditions

Angiostrongylosis

- Increased risk of infection in this breed in a UK study (Blehaut *et al.*, 2014)

Babesiosis

- Breed over-represented in a Brazilian case series
- 41.93% frequency of detection in suspicious clinical cases in this breed (Ungar de Sá *et al.*, 2007)

Cryptococcosis

- American Cocker Spaniel
- Breed at increased risk in two US case series
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors (Berthelin *et al.*, 1994; Trivedi *et al.*, 2011)

Rhipicephalus sanguineus

- Increased susceptibility to this tick compared to Beagles in experimental studies (Louly *et al.*, 2009, 2010)

Metabolic conditions

Overweight/obesity

- Breed predisposed in a North American private practice population
- OR 1.9 (95% CI 1.6–2.2; $p < 0.0001$) (Lund *et al.*, 2006)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- This breed reported to be at increased risk in a Czech study

(Nečas *et al.*, 2000)

Incomplete ossification of the humeral condyle

- Polygenic recessive mode of inheritance
- Males predisposed

(Marcellin-Little *et al.*, 1994)

Inguinal/scrotal herniation

- Breed over-represented in an Australian population compared to hospital population

(Bellenger, 1996)

Patellar luxation

- American Cocker Spaniel
- May be inherited as an autosomal recessive trait
- Lateral luxation more common in this breed
- OR 2.1 (95% CI 1.7–2.7) compared to mixed breeds in a US population

(LaFond *et al.*, 2002)

Temporomandibular joint dysplasia

- American Cocker Spaniels
- Congenital
- Usually affects dogs from 6 months of age

(Hoppe & Svalastoga, 1980)

Neoplastic conditions

Anal sac adenocarcinoma

- English Cocker Spaniel had an OR of 11.00 (95% CI 8.27–14.64) in the UK

(Polton *et al.*, 2006)

Canine acanthomatous ameloblastoma

- Breed over-represented in a US case series

(Fiani *et al.*, 2011)

Canine cutaneous histiocytoma

- Breed at increased risk in an Australian case series (9/80 cases were this breed)

(Er & Sutton, 1989)

Lingual plasma cell tumour

- Increased risk in a US case series

(Dennis *et al.*, 2006)

Lymphoma

- Relative risk (RR) 1.15 ($p < 0.001$) in a French study

(Pastor *et al.*, 2009)

Mammary neoplasia

- Significantly higher risk in this breed in a Czech case study ($p < 0.01$)
- In one US study of male dogs with mammary tumours, 7/18 dogs were Cocker Spaniels
- This breed consistently affected according to one Mexican study

(Zatloukal *et al.*, 2005; Bearss *et al.*, 2012;

Salas *et al.*, 2015)

Melanoma

- Breed at increased risk in a US histopathological case series
- Mean age 8.9 years
- Predilection site of the lip in this breed

(Schultheiss, 2006)

Oropharyngeal neoplasia

See under *Gastrointestinal conditions*

Perianal (hepatoid) gland adenomas

- Breed at risk in US case series
- Average age was 10.5 years
- Relative risk compared to US veterinary medical database of 2.0 (95% CI 1.41–3.03) in males and 3.3 (95% CI 2.03–7.30) in females

(Hayes & Wilson, 1977)

Sebaceous gland carcinoma

- Incidence of 15.6% in a US case series

(Strafuss, 1976)

Neurological conditions

Intervertebral disc disease (IVDD)

- Reported to be common in this breed in a US case series
- English Cocker more likely to have caudal and midlumbar intervertebral disc extrusions than Dachshunds

(Mayhew *et al.*, 2004; Cardy *et al.*, 2016)

Multisystem neuronal degeneration

- Familial and suspected inherited
- Age of clinical onset 10–14 months

(Jaggy & Vandeveld, 1988)

Ocular conditions

Cataract

- Autosomal recessive inheritance suspected
- Relatively common condition in this breed, possibly higher incidence in females
- Prevalence of primary cataract 8.77% in American Cocker Spaniel and 8.23% in English

Cocker Spaniel, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

- Lens-induced uveitis common in this breed (Olesen *et al.*, 1974; van der Woerd *et al.*, 1992; Adkins & Hendrix, 2005; Gelatt & MacKay, 2005)

Corneal ulceration (ulcerative keratitis)

- OR 2.33 (95% CI 1.58–3.42 < 0.001) compared to crossbreeds in a UK general practice population (O'Neill *et al.*, 2017b)

Distichiasis

- Extremely high prevalence in a Danish study (49.3%)
- Females predisposed
- Heritability moderate to high (0.22–0.51) (Petersen *et al.*, 2015)

Glaucoma

- Inheritance suspected
- Predisposition for females suggested
- May be associated with goniodysgenesis
- Glaucoma secondary to cataracts also seen
- In a North American study, prevalence among American Cocker Spaniels was 5.52%, and 1.35% in English Cocker (Gelatt & MacKay, 2004a, 2004b; Johnsen *et al.*, 2006)

Keratoconjunctivitis sicca

- Breed at increased risk in a UK study
- One study suggested a female predisposition in English Cocker Spaniels (Sanchez *et al.*, 2007)

Progressive rod–cone degeneration (PRCD)

- Frequency of the disease-causing mutation found to be 0.34 (English Cocker) and 0.08 (American Cocker) in a Czech study (Dostal *et al.*, 2011)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- American Cocker Spaniel
- Increased risk in this breed in a US case series (Morgan *et al.*, 1993)

Retinal detachment

- One of the most common breeds affected in a US case series (Hendrix *et al.*, 1993)

Retinal dysplasia – multifocal

- American Cocker Spaniels
- Congenital, autosomal recessive inheritance suspected (MacMillan & Lipton, 1978)

Retinal pigment epithelial dystrophy (RPED, central progressive retinal atrophy)

- English Cocker Spaniels
- Inheritance suspected
- More prevalent in the UK than in the USA. Becoming less prevalent following the introduction of control schemes
- May be related to vitamin E deficiency due to an underlying metabolic abnormality in this breed (McLellan *et al.*, 2002)

Traumatic proptosis

- One of the most commonly affected breeds in a Brazilian population (Brandão *et al.*, 2005)

Renal and urinary conditions

Chronic kidney disease

- Breed predisposed in a UK general practice population (O'Neill *et al.*, 2013b)

Familial renal disease (glomerular basement membrane disorder)

- English Cocker Spaniels
- Autosomal recessive inheritance suggested
- The condition is believed to be similar to Alport syndrome in humans
- Cases present at between 6 months and 2 years of age with proteinuria and chronic renal failure (Lees *et al.*, 1998; Davidson *et al.*, 2007)

Urolithiasis – silica

- Breed at risk in a US series
- Mean age at diagnosis 5.8 years
- Males predisposed (Osborne *et al.*, 1999)

Urolithiasis – struvite

- Breed at risk in case series
- Average age at diagnosis in one large study: males 6.0 years, females 5.7 years
- Females predisposed (16:1 in one large study) (Osborne *et al.*, 1999; Houston *et al.*, 2004)

Urolithiasis – urate

- Breed at risk in a Czech case series
- Male dogs predisposed (Kučera, 2007)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- English Cocker Spaniels
- Breed at risk in a Swedish case series (Egenvall *et al.*, 2001)

XX sex reversal

- American Cocker Spaniels
- Autosomal recessive inheritance (Meyers-Wallen & Patterson, 1988; Pujar *et al.*, 2005)

Respiratory conditions

Bronchiectasis

- American Cocker Spaniels had increased risk in a US case series
- Usually affects middle-aged to older dogs
- Usually occurs secondary to chronic pulmonary disease (Hawkins *et al.*, 2003)

COLLIES – ROUGH AND SMOOTH

Behavioural conditions

Aggression and fearfulness

- Reported to be at increased risk of aggression to strangers and general fearfulness (Arvelius *et al.*, 2014; Lund *et al.*, 1996)

Dermatological conditions

Bacterial folliculitis/furunculosis

- Relative risk (RR) 2.0 compared to Canadian hospital population (Scott & Paradis, 1990)

Cutaneous lupus erythematosus

- Reported in two US case series (Jackson, 2004; Jackson *et al.*, 2004)

Demodicosis

- Breed over-represented in a case series (Scott & Miller, 2012b)

Dermatophytosis (ringworm)

- Collies were commonly represented in a Croatian case series (Pinter *et al.*, 1999)

Familial canine dermatomyositis

- Inherited as an autosomal dominant trait with incomplete penetrance
- No predispositions for sex, coat colour or coat length (Haupt *et al.*, 1985; Evans *et al.*, 2017)

Idiopathic sterile granuloma and pyogranuloma

- This breed over-represented in one study
- No age or sex predisposition (Panich *et al.*, 1991)

Idiopathic ulcerative dermatosis in Shetland Sheepdogs and Collies

- No sex predisposition
- Affects middle-aged to older dogs (Jackson & Olivry, 2001; Moriello, 2004)

Pemphigus foliaceus

- OR 3.9 for this breed in a US pathology case series (Kuhl *et al.*, 1994)

Drug reactions

Ivermectin and milbemycin

- Associated with the mutant *MDR1* allele
- 22% of this breed were homozygous and 46.8% heterozygous for the mutation in US and UK populations (Neff *et al.*, 2004; Mealey *et al.*, 2005)

Endocrine conditions

Diabetes mellitus

- One of the most common breeds affected in a UK population (Davison *et al.*, 2005)

Insulinoma

See under *Neoplastic conditions*

Gastrointestinal conditions

Colorectal polyps

- This breed the most common in a US case series (4/17 cases were Collies) (Seiler, 1979)

Exocrine pancreatic insufficiency (EPI)

- Rough Collies
- Typical age of onset: 36 months
- Prevalence in a clinicopathological case series (of animals with suggestive clinical signs) of 10.4% (95% CI 5.9–16.6%) in this breed (Wiberg *et al.*, 1999; Batchelor *et al.*, 2007)

Gastric carcinoma

See under *Neoplastic conditions*

Gastric dilatation/volvulus (bloat, GDV)

- Incidence rate (IR) 21 cases per 1000 DYAR (95% CI 8–34) in a US study (Glickman *et al.*, 2000a)

Pancreatitis

- Breed at risk of chronic pancreatitis in a case series of post-mortem dogs
- RR 2.0 (95% CI 1.2–3.2) (Watson *et al.*, 2007)

Haematological/immunological conditions**Factor I deficiency**

- Autosomal inheritance (Brooks, 1999)

Platelet disorder

- Storage pool and signal transduction defect (Brooks, 1999)

Immune-mediated haemolytic anaemia (IMHA)

- Increased risk in this breed in a UK case–control study (Miller *et al.*, 2004)

Haemophilia A

- Familial and moderate severity in the rough collie (Brooks, 1999)

Metabolic conditions**Hypomagnesaemia**

- OR 3.9 (95% CI 1.3–6.8; $p < 0.02$) in a referral hospital population (Khanna *et al.*, 1998)

Musculoskeletal conditions**Perineal herniation**

- Older, intact males predisposed in a case series (Robertson, 1984)

Neoplastic conditions**Gastric carcinoma**

- Rough Collies
- Male dogs more commonly affected in case series
- Mean age of occurrence 8–10 years
- Proportional morbidity ratio of 26.1 (95% CI 6.4–106.5) in a Norwegian case series (Sullivan *et al.*, 1987b; Seim-Wikse *et al.*, 2013)

Insulinoma

- Reported to be common in this breed in a US case series (Kruth *et al.*, 1982)

Nasal cavity tumours

- Breed at increased risk in a US teaching hospital case series
- RR 2.9 (95% CI 2.05–4.12)
- Median age 9 years
- Males over-represented in most studies (Hayes *et al.*, 1982)

Testicular neoplasia

- Rough Collies and Shetland Sheepdogs five times more likely to have testicular tumours than the rest of a Norwegian population
- Over-represented in a Slovenian study (Nødtvedt *et al.*, 2011; Švara *et al.*, 2014)

Neurological conditions**Cerebellar degeneration**

- Reported in an Australian case series
- Autosomal recessive inheritance (Hartley *et al.*, 1978)

Idiopathic epilepsy

- *ABCB1* mutation, which has a >50% prevalence in Collies, is associated with outcome (Muñana *et al.*, 2012)

Ocular conditions**Collie eye anomaly**

- Rough and Smooth Collies
- Congenital; autosomal recessive inheritance suspected. Some authors suggest polygenic inheritance
- High incidence in this breed throughout the world (50–90% has been reported), now reduced through selective breeding (Bedford, 1982b; Wallin-Håkanson *et al.*, 2000)

Corneal dystrophy

- Rough Collies
- Inheritance suspected
- Reported in a UK case series
- Crystalline lipid stromal dystrophy
- Age of onset: 1–4 years
- May or may not be related to Collie eye anomaly (Crispin, 1988)

Myopia (short-sightedness)

- Breed at increased risk in a large US study (Kubai *et al.*, 2008)

Nodular granulomatous episclerokeratitis

- Young to middle-aged Collies predisposed in a case series (Paulsen *et al.*, 1987)

Retinopathy (rod–cone dystrophy)

- Reported to be inherited in a recessive mode in this breed (Wolf *et al.*, 1978)

Renal and urinary conditions**Renal amyloidosis**

- Breed at increased risk in a US case series
- Most cases > 6 years at diagnosis
- Females predisposed
- Renal amyloidosis may be seen secondary to dermatomyositis in this breed (DiBartola *et al.*, 1989)

Reproductive conditions**Pyometra (cystic endometrial hyperplasia–pyometra complex)**

- Rough Collie
- Breed at increased risk in a Swedish insurance database
- This breed may have increased risk at an earlier age compared to other breeds
- OR 2.05 (95% CI 1.21–3.47) in a UK population (Niskanen & Thrusfield, 1998; Egenvall *et al.*, 2001)

COONHOUND**Dermatological conditions****Blastomycosis**

See under *Infectious conditions*

Infectious conditions**Blastomycosis**

- Breed at increased risk in case series
- Young (2–4 years) entire males predisposed
- Geographic distribution: mainly North America (Mississippi, Missouri, Ohio River valleys, mid-Atlantic States, Quebec, Manitoba and Ontario)
- Proximity to a body of water is a risk factor in endemic areas

(Rudmann *et al.*, 1992; Arceneaux *et al.*, 1998)

Histoplasmosis

- Reported to be common in this breed in a US case series (Burk *et al.*, 1978)

Musculoskeletal conditions**Hip dysplasia**

- Treeing Walker Coonhound predisposed
- Neutered male dogs predisposed
- OR 3.1 (95% CI 1.3–7.4) compared to mixed breeds in a US study (LaFond *et al.*, 2002)

Renal and urinary conditions**Renal amyloidosis**

- Treeing Walker Coonhounds at increased risk in case series
- Most cases > 6 years at diagnosis
- Females predisposed in one study (DiBartola *et al.*, 1989)

CORGI

See *Welsh Corgi*

COTON DE TULEAR**Neurological conditions****Neonatal cerebellar ataxia**

- Also known as Bandera's neonatal ataxia
- Found in 7/32 pups from five litters in one US study
- In a genetic study of 112 dogs of this breed, 15 were homozygous for the mutant gene and were affected, 43 were heterozygotes, and 54 were homozygous for the ancestral gene
- Ataxia started at 2 weeks of age
- Autosomal recessive inheritance

(Coates *et al.*, 2002; Zeng *et al.*, 2011)

Ocular conditions**Multifocal retinopathy**

- Autosomal recessive inheritance
- Develops from 11 weeks of age
- Reported in 10 affected dogs of this breed which were compared to 20 unaffected dogs of this breed, in a Canadian study (Grahn *et al.*, 2006, 2008)

Renal and urinary conditions

Primary hyperoxaluria

- Autosomal recessive inheritance
- Onset from 3–4 weeks
- Reported in 7 dogs from four apparently unrelated litters in a Finnish study (Vidgren *et al.*, 2012)

CURLY-COATED RETRIEVER

Musculoskeletal conditions

Glycogen storage disease type III

- Also known as Cori's disease
- Females predisposed
- Signs first seen at 2 months of age
- Carriers reported in the USA, Australia, New Zealand and Finland (Gregory *et al.*, 2007)

Ocular conditions

Cataract

- Hereditary
- 12.1% prevalence in a Norwegian study (Bjerkås, 1991)

Neurological conditions

Exercise-induced collapse

- 84/251 dogs of this breed were heterozygous (carriers) and 49/251 were homozygous (affected) for the *DNM1* gene mutation causing this condition in a North American study (Minor *et al.*, 2011)

DACHSHUND

Behavioural conditions

Aggression

- This breed predisposed to increased inter-dog aggression in a Danish study
- In a Dutch study, breed predisposed to aggression against both humans and other dogs (Rugbjerg *et al.*, 2003; Duffy *et al.*, 2008)

Cardiovascular conditions

Mitral valve disease

- Odds ratios in a UK study 1.97 (95% CI 1.07–3.62) in males and 2.31 (1.09–4.89) in females
- Mitral valve prolapse is common in this breed
- Polygenic mode of inheritance suspected

- Dachshunds frequently affected by mitral type murmurs in a French study of six breeds (Thrusfield *et al.*, 1985; Olsen *et al.*, 1999; Serfass *et al.*, 2006)

Dermatological conditions

Atopic dermatitis (atopy)

- 6% prevalence in this breed in an Italian study (Tognetti *et al.*, 1987)

Colour dilution alopecia

- Coat-colour genes are involved in the pathogenesis
- Autosomal recessive inheritance (Beco *et al.*, 1996)

Sterile nodular panniculitis

- Miniature Dachshunds predisposed in a Japanese case series
- Neutered and younger dogs predisposed (Yamagishi *et al.*, 2007)

Endocrine conditions

Diabetes mellitus

- More likely to develop anti-insulin antibodies than crossbreeds in a UK study
- 14/77 cases in a Swedish case series were this breed (Forsberg, 1986; Holder *et al.*, 2015)

Hyperadrenocorticism (Cushing's syndrome)

- Reported to be common in this breed in a case series of 153 dogs in an American population (Reusch & Feldman, 1991; Wood *et al.*, 2007)

Hypothyroidism

- Breed at increased risk in a US 1960s and 1970s case series
- May occur at a younger age in breeds at risk (2–3 years)
- Females and neutered males at increased risk (Milne & Hayes, 1981)

Thyroid neoplasia

- Breed at increased risk in Swiss study (Wenger *et al.*, 2005)

Gastrointestinal conditions

Bacterial cholecystitis and bactibilia

- 5/10 cases in a retrospective case-control series were Dachshunds (Lawrence *et al.*, 2015)

Colorectal polyps

- Reported in a Japanese case series
- Reported to be associated with dysbiosis in a Japanese population
- More commonly found ventrally
(Ohmi *et al.*, 2012; Igarashi *et al.*, 2015; Uchida *et al.*, 2016)

Pancreatitis

- Breed at increased risk in a Hungarian case series
(Pápa *et al.*, 2011)

Sialocoele

- Dachshunds significantly over-represented in an Australian study compared to hospital population
(Bellenger & Simpson, 1992)

Haematological/immunological conditions**Haemophilia A**

- Miniature Dachshunds
- Moderate factor VIII deficiency in this breed
- Familial in this breed
(Brooks, 1999)

Immunoglobulin deficiency

- Causes a protozoal pneumonia (*Pneumocystis carinii*)
 - Associated with low IgG levels
 - Inheritance uncertain
- See also under *Infectious conditions*
(Lobetti *et al.*, 1996; Lobetti, 2000)

von Willebrand's disease (vWD)

- Type I seen in this breed
(Brooks, 1999)

Infectious conditions***Pneumocystis carinii* infection**

- Age affected: < 1 year
- See also under *Haematological/immunological conditions* (*Immunoglobulin deficiency*)
(Lobetti *et al.*, 1996; Lobetti, 2000)

Metabolic conditions**Overweight/obesity**

- Breed predisposed in a North American private practice population
- OR 1.6 (95% CI 1.2–2.2; $p = 0.0005$)
(Lund *et al.*, 2006)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- OR 4.8 (95% CI 2.0–11.2) in one US study, compared to mixed breeds
(LaFond *et al.*, 2002)

Inguinal/scrotal herniation

- Breed over-represented in an Australian population compared to hospital population
(Bellenger, 1996)

Olecranon fracture

- 43/67 dogs in a German case series were Dachshunds or Dachshund crosses
(Brunnberg *et al.*, 1983)

Osteogenesis imperfecta

- Reported in two related litters of Dachshunds
- Associated with a recessive mutation which had a prevalence of 12.9% in this breed in a European study
(Seeliger *et al.*, 2003; Eckardt *et al.*, 2013)

Patellar luxation

- 7.9% of cases in a German series were this breed
(Schafer *et al.*, 1982)

Pes varus

- Reported in two case series
- Seen from 5–6 months of age
- Possible autosomal recessive inheritance
(Radasch *et al.*, 2008; Petazzoni *et al.*, 2012)

Tail anomalies

- Congenital tail anomalies found in 0.93% of short-haired, 0.56% of rough-haired and 2.21% of long-haired Dachshunds in a German population in the 1970s
- Thought to be inherited as an autosomal recessive trait
(Fritsch & Ost, 1983)

Neoplastic conditions**Adrenocortical tumour resulting in hyperadrenocorticism**

- Breed at risk
 - Females slightly over-represented
- See also under *Endocrine conditions*
(Reusch & Feldman, 1991)

Haemangiosarcoma – cardiac

- 7/51 cases with this condition in a Japanese case series were Miniature Dachshunds (Yamamoto *et al.*, 2013)

Mammary neoplasia

- This breed accounted for 37.3% of cases in a German case series
- Significantly increased risk in this breed in a Czech series (Bomhard & Drejack, 1977; Zatloukal *et al.*, 2005)

Mast cell tumour (MCT)

- Breed over-represented in a German case series (Kessler *et al.*, 1997)

Melanoma

- Dachshunds reported to be predisposed to amelanotic melanoma in a Brazilian case series
- Miniature Dachshunds predisposed to oral melanoma in a Thai case series (Rolim *et al.*, 2012; Choisunirachon *et al.*, 2014)

Oral fibrosarcoma

- Reported to be at increased risk compared to mixed-breed dogs in a Polish case series (Sapierzyński *et al.*, 2007b)

Pituitary tumour resulting in hyperadrenocorticism

See under *Endocrine conditions*

Squamous cell carcinoma – digit

- Breed at increased risk in case series
- Middle-aged to older dogs (Henry *et al.*, 2005)

Neurological conditions**Idiopathic epilepsy**

- Thought to be familial in this breed
- Age of onset 6 months to 6 years (Cunningham & Farnbach, 1988)

Intervertebral disc disease (IVDD)

- Prevalence of 15.7 % (95 % CI 14.1–17.3) in a UK epidemiological study
- Standard short-haired Dachshunds had the highest risk compared to other varieties of Dachshund
- Older and neutered dogs at increased risk (Mayhew *et al.*, 2004; Packer *et al.*, 2016a)

Vestibular disease

- Breed over-represented in a Brazilian case series
- 2/3 of cases in this series involved central vestibular disease (Chaves *et al.*, 2014)

Ocular conditions**Cataract**

- Prevalence of primary cataract in long-haired Dachshunds 2.10%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005)

Progressive retinal atrophy (PRA)

- Miniature long-haired Dachshunds
- Autosomal recessive inheritance
- Rod–cone dystrophy
- Onset and progression variable
- Reported in a UK case series and a Norwegian breeding study (Turney *et al.*, 2007; Ropstad *et al.*, 2008)

Keratoconjunctivitis sicca

- 12/200 cases were this breed in a US case series
- Females predisposed (Sansom & Barnett, 1985)

Sudden acquired retinal degeneration (SARD, amaurosis)

- This breed reported to be frequently diagnosed in a US case series (Montgomery *et al.*, 2008)

Physiological conditions**Chondrodysplasia**

- Accepted as breed standard
- Short, bowed legs, but normal skulls seen (Martinez *et al.*, 2000)

Haematological differences

- Haematocrit (PCV, packed cell volume), haemoglobin and red cell count were higher in this breed than in mixed-breed dogs (Torres *et al.*, 2014)

Heart rate

- Lower mean heart rate in this breed under anaesthesia compared to other breeds (Harrison *et al.*, 2012)

Renal and urinary conditions

Urolithiasis – cystine

- Almost all cases reported were male
- Significantly increased risk of cystine calculus formation in an American population
- Reported to be common in this breed in a Swedish study and a large American study (Case *et al.*, 1992; Wallerström *et al.*, 1992; Ling *et al.*, 1998b)

Reproductive conditions

Dystocia

- Breed predisposition to obstructive dystocia due to a narrow maternal pelvic canal, and also dystocia due to uterine inertia
- 5.5% of cases of dystocia were this breed in a US case series
- Also over-represented in a German case series
- Odds ratio in a UK first-opinion practice (Miniature Dachshund) 7.9 (95% CI 4.0–15.4; $p < 0.001$) (Gaudet, 1985; Trautmann & Nolte, 2003; O'Neill *et al.*, 2017c)

Respiratory conditions

Nasopharyngeal stenosis

- Reported in a South African case series or seven smooth-haired Dachshunds
- Mean age at presentation 3 years (Kirberger *et al.*, 2006)

Pneumonia due to *Pneumocystis carinii* infection

See under *Infectious conditions*

DALMATIAN

Dermatological conditions

Actinic (solar) keratosis

- Breed at increased risk
- Areas of light-coloured/sparse haircoat affected. Exposure to sun increases the risk
- May undergo neoplastic transformation (Power & Prélaud, 2011)

Atopic dermatitis (atopy)

- Dogs between 1 and 2 years of age have the highest probability of an insurance claim for atopy
- Breed at increased risk in a US series of 264 cases (1981–1984)
- Some studies show no sex predilection, others show females predisposed

- This breed had a risk factor of 5.4 cases per 1000 DYAR, compared to 1.7 for the general population (Schick & Fadok, 1986; Nødtvedt *et al.*, 2006)

Cutaneous neoplasia

See under *Neoplastic conditions*

Sterile nodular panniculitis

- Breed identified at high risk in a hospital review when compared to all other dogs seen during the study period at University of California, Davis, USA (Contreary *et al.*, 2015)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Breed with a higher prevalence of thyroid hormone autoantibodies (THAA)
- In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, Dalmatians had an OR of 1.74 ($p = 0.001$) of being affected compared to dogs of all other breeds
- Across the study, females were over-represented and the highest prevalence was in dogs 2–4 years old (Nachreiner *et al.*, 2002)

Gastrointestinal conditions

Chronic hepatitis

- Breed at increased risk in a UK-based study of 551 cases (OR 4.1, 95% CI 2.2–7.7; $p < 0.001$) compared to a control population. Across the whole study, median age was 4 years 7 months and females were over-represented
- Some studies suggest Dalmatians are at risk of copper-associated hepatitis (Webb *et al.*, 2002; Watson, 2004; Bexfield *et al.*, 2012a)

Infectious conditions

Cryptococcosis

- Breed at increased risk in a case series in Australia
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors (O'Brien *et al.*, 2004)

Musculoskeletal conditions

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older

- OR 3.1 (95% CI 1.6–6.2) compared to mixed breeds in a large retrospective case series based on cases recorded in the VMDB, 1986–1995 (Nečas, 1999; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.5 (95% CI 2.0–3.1) compared to mixed breeds in a large retrospective case series based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Conjunctival haemangiosarcoma

- Uncommon tumour, representing 1% of all ocular tumours
- Dalmatians were over-represented (compared to VMDB population) in a retrospective case series in North America
- Exposure to ultraviolet (UV) light may be a risk factor

(Pirie *et al.*, 2006)

Primary brain tumour

See under *Neurological conditions*

Squamous cell carcinoma – cutaneous

- Breed at increased risk of cutaneous squamous cell carcinoma in a retrospective study of VMDB data, 1964–2002: OR 6.94 (95% CI 4.26–11.32) compared to all other dogs
- Increased frequency noted in Australian Dalmatians

(Rothwell *et al.*, 1987; Villamil *et al.*, 2011)

Neurological conditions

Congenital deafness

- Up to 30% of the breed reported to be affected in the USA. Lower percentages reported in Europe
- Mode of inheritance unclear
- Dogs with blue eyes more commonly affected
- More commonly unilateral than bilateral (Juraschko *et al.*, 2003; Cargill *et al.*, 2004; Strain, 2004)

Idiopathic epilepsy

- Breed at increased risk in case series (Licht *et al.*, 2002; Short *et al.*, 2011)

Laryngeal paralysis–polyneuropathy syndrome

See under *Respiratory conditions*

Neuronal ceroid lipofuscinosis

- Rare
- Inheritance suspected
- Clinical onset 6 months. Can be diagnosed at 4–5 months by skin biopsy
- Signs include tremors, dysmetria, ataxia and visual deficits

(Goebel *et al.*, 1988)

Primary brain tumour – choroid plexus neoplasm

- Breed at significantly increased risk compared to the general population, in a study of post-mortem results of 435 dogs with intracranial neoplasia

(Song *et al.*, 2013)

Ocular conditions

Neuronal ceroid lipofuscinosis

See under *Neurological conditions*

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Autosomal dominant inheritance
- Dalmatians suffer from a glomerular basement membrane disorder. Renal failure is seen at a median age of 18 months
- Condition is believed to be similar to Alport syndrome in humans

(Hood *et al.*, 2002b)

Hyperuricosuria

- Hyperuricosuria (associated with an *SLC2A9* mutation) is inherited recessively and predisposes to urate urolithiasis

(Bannasch *et al.*, 2008)

Urolithiasis – urate

- Common in this breed. Prevalence was 34% in male Dalmatians in one 2004 study
- Average age at diagnosis in one large study: males 5.5 years, females 4.6 years
- In a retrospective case series of 25 499 laboratory stone submissions in the USA, 1985–2006, OR was 32 (95% CI 29–35.3) compared to mixed-breed dogs. Dogs <7 years old at increased risk ($p < 0.001$) across all breeds
- In a retrospective study of 14 008 uroliths from dogs in the UK, 1997–2006, OR was 130.20 (95% CI 113.43–149.44; $p < 0.0001$) compared to the national insurance database
- Males predisposed (OR 14.0 in one study)

(Bannasch *et al.*, 2004;

Albasan *et al.*, 2005; McCue *et al.*, 2009;

Low *et al.*, 2010; Roe *et al.*, 2012)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Autosomal recessive inheritance suspected
- Affects young dogs (9 months or younger)
- In this condition, laryngeal paralysis is associated with a more generalized polyneuropathy (Braund *et al.*, 1994a)

Respiratory distress syndrome

- Rare condition; two separate reports of cases seen in young related dogs
- Possibly inherited as an autosomal recessive trait (Järvinen *et al.*, 1995; Syrjä *et al.*, 2009)

DANDIE DINMONT TERRIER

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed at increased risk in case series: OR 31.7 compared to mixed-breed dogs
- Clinical signs usually seen in young dogs < 1 year (Tobias & Rohrbach, 2003)

Ocular conditions

Glaucoma – primary

- Breed at increased risk of pectinate ligament dysplasia. 22.1% were moderately or severely affected in a survey of 95 Dandie Dinmont Terriers in the UK
- Age of onset 7 years and older (Oliver *et al.*, 2016a)

Reproductive conditions

Delivery by caesarean section

- In the top 10 breeds in a cross-sectional study conducted by the Kennel Club in the UK
- The study found a caesarean rate of 41.4% in litters of Dandie Dinmont Terriers born over a 10-year period (Evans & Adams, 2010a)

DANISH CHICKEN DOG (GAMMEL DANSK HØNSEHUND)

Neurological conditions

Congenital myasthenia gravis

- Autosomal recessive inheritance
- This form of myasthenia gravis is not improved by anticholinesterase (Flagstad *et al.*, 1989)

DANISH/SWEDISH FARMDOG

Dermatological conditions

Atopic dermatitis (atopy)

- In a study of Swedish insurance data, Danish/Swedish Farmdogs had a risk factor of 3.6 cases per 1000 DYAR, compared to 1.7 cases per 1000 DYAR for all breeds combined
- Birth in autumn or summer is a predisposing factor
- Dogs between 1 and 2 years of age have highest probability of an insurance claim for atopy
- Some studies show no sex predilection, others show females predisposed (Nødtvedt *et al.*, 2006)

Reproductive conditions

Dystocia

- Breed at increased risk in a retrospective study of Swedish insurance data (195 931 bitches 1995–2002)
- Incidence in the Danish/Swedish Farmdog was 11.1 cases per 1000 DYAR, compared to 5.7 cases per 1000 DYAR for the general population (Bergström *et al.*, 2006)

DEERHOUND

See *Scottish Deerhound*

DEUTSCHER WACHTELHUND

See *Wachtelhund*

DOBERMANN PINSCHER (DOBERMANN)

Behavioural conditions

Flank-sucking

- High incidence reported in Dobermanns
- Dogs with flank-sucking have a higher incidence of pica (the eating of non-food substances) than the normal population (Moon-Fanelli *et al.*, 2007; Ogata *et al.*, 2013)

Cardiovascular conditions

Atrial fibrillation (AF)

- Breed at increased risk in case series
- In a study of 109 cases of AF from two North American veterinary teaching hospitals,

Dobermanns were over-represented ($p < 0.0001$) in the group of cases where structural heart disease and overt congestive failure was present

- Males were found to be significantly over-represented
- OR 5.66 in a separate large study of 3542 cases in the VMDB
(Menaut *et al.*, 2005; Westling *et al.*, 2008)

Atrial septal defect

- Uncommon disease
- Congenital – inheritance not known
(Lee *et al.*, 2007; Meurs, 2010)

Dilated cardiomyopathy (DCM)

- Very common in Dobermann Pinschers. One study suggests a cumulative prevalence of 58.2% in Dobermanns > 7 years old in Europe. Prevalence in North America has been reported as 45–63%
- Autosomal dominant inheritance suggested
- Prevalence increases with age
- Earlier reports suggested DCM is twice as common in males as in females. A recent study has suggested that females are equally affected but tend to have a disease that progresses more slowly, with only ECG abnormalities rather than the echocardiographic changes more commonly seen in males. They are therefore less likely to be diagnosed and may also develop congestive heart failure later in life
- Screening has been recommended by Holter ECG and echocardiography from 2 years of age and repeated annually
(Tidholm & Jonsson, 1997; Tidholm *et al.*, 2001; Borgarelli *et al.*, 2006; Wess *et al.*, 2010)

Patent ductus arteriosus

- Breed at increased risk (OR 2.8; $p = 0.0007$) compared to the general hospital population in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- This study found a female predisposition (OR 2.7)
(Oliveira *et al.*, 2011)

Dermatological conditions

Blastomycosis

See under *Infectious conditions*

Colour dilution alopecia

- Reported in blue or fawn Dobermanns and Miniature Pinschers
- Frequency of the disease may be as high as 93% in blues and 75% in fawns
- Coat-colour genes play a role in the inheritance of this condition
(Miller, 1990a)

Cryptococcosis

See under *Infectious conditions*

Cutaneous neoplasia

See under *Neoplastic conditions*

Demodicosis

- Breed at increased risk in a number of case series
- Most common in dogs < 1 year of age
(Nayak *et al.*, 1997; Solanki *et al.*, 2007)

Follicular dysplasia

- Affects black or red Dobermanns
- Age of onset: 1–4 years
- Affects caudodorsum and flanks
(Miller, 1990b)

Oculocutaneous albinism

- Affected dogs have white coats, pale irises, and increased risk of melanoma
(Winkler *et al.*, 2014)

Drug reactions

Sulfonamide-associated hypersensitivity

- Dobermanns are at increased risk of cutaneous reactions and polyarthropathy following the use of this drug
(Trepanier, 1999)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Breed at increased risk in case series. OR 6.06 ($p = 0.001$) compared to a large university hospital population. Association with a rare MHC class 2 haplotype documented in Dobermanns. May occur at a younger age in at-risk breeds (2–3 years of age). Neutered animals (both sexes) were at increased risk
- Breed with higher prevalence of thyroid hormone autoantibodies (THAA). In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, Dobermanns had an OR of 1.24 ($p = 0.001$) of being affected compared to dogs of all other breeds

- Across the study, females were over-represented, and the highest prevalence was in dogs 2–4 years old
(Panciera, 1994; Nachreiner *et al.*, 2002; Kennedy *et al.*, 2006a)

Gastrointestinal conditions

Chronic hepatitis

- Breed at increased risk in a UK-based study of 551 cases: OR 11.5 (95% CI 7.6–17.3; $p < 0.001$) compared to a control population. Median age was 5 years 4 months, and females were over-represented across the whole study
- A very aggressive form of hepatitis, sometimes referred to as chronic active hepatitis, is seen in Dobermanns
- Increased hepatocellular copper is seen and may be secondary to cholestasis rather than a primary defect in hepatic copper metabolism, but this is debated
- An autoimmune pathogenesis has been proposed
(Speeti *et al.*, 2003; Mandigers *et al.*, 2004; Bexfield *et al.*, 2012a)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in case series
- One of the four most common breeds represented in an internet-based survey of dog owners (2551 submissions)
(Mackenzie *et al.*, 2010; Pipan *et al.*, 2012)

Parvovirus enteritis

See under *Infectious conditions*

Haematological/immunological conditions

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance
(Brooks, 1999)

von Willebrand's disease (vWD)

- High incidence in Dobermanns: 73% had abnormal (<50%) von Willebrand factor (vWF) in a study of 5554 Dobermanns in 1988
- Autosomal dominant inheritance with incomplete penetration suspected
- Mean age of affected Dobermanns was 4.6 years in one study
- Not all at-risk dogs will bleed
(Brooks *et al.*, 1992; Riehl *et al.*, 2000; Gentilini & Turba, 2013)

Infectious conditions

Blastomycosis

- Breed at increased risk in case series
- Young dogs (2–4 years) predisposed
- Geographic distribution: mainly North America (Mississippi, Missouri, Ohio River valleys, mid-Atlantic States, Quebec, Manitoba and Ontario)
- Proximity to a body of water is a risk factor in endemic areas
(Kerl, 2003)

Cryptococcosis

- Breed at increased risk in a case series of 195 cases (1981–2001) in Australia
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors
(Malik *et al.*, 1995; O'Brien *et al.*, 2004)

Demodicosis

See under *Dermatological conditions*

Parvovirus enteritis

- Breed at increased risk in a number of case series
- OR 3.1 compared to the general hospital population in a University of Pennsylvania study
- Age 6 weeks to 6 months at higher risk
- One study has suggested increased risk is associated with a stronger pro-inflammatory cytokine response
(Glickman *et al.*, 1985; Houston *et al.*, 1996; Nemzek *et al.*, 2007)

Musculoskeletal conditions

Carpal laxity syndrome

- Breed at increased risk in case series
- Puppies affected at 6–12 weeks of age. Males over-represented
- Usually self-limiting
(Cetinkava *et al.*, 2007)

Osteosarcoma

See under *Neoplastic conditions*

Neoplastic conditions

Cutaneous neoplasia

- In a retrospective study of 104 histopathological samples examined at a Bulgarian university clinical pathology department (1991–2000), the Dobermann was the breed with the third-highest prevalence

- Average age across the study was 9.58 years (range 6–11.7)
- Males were significantly more affected (Dinev, 2002)

Lipoma

- Breed at increased risk
- Most common in middle-aged, obese female dogs
- Dobermanns may be predisposed to infiltrative lipomas (Kim *et al.*, 2005)

Mammary neoplasia

- 42% of insured Swedish Dobermanns had developed mammary tumours at 10 years of age (ranked second), compared to 13% for the general population
- Increased risk is reflected in case studies in other parts of the world
- Risk increases with age
- Risk significantly decreased with early neutering (Jitpean *et al.*, 2012)

Melanoma

- Breed at increased risk of cutaneous melanoma in a retrospective study of VMDB data, 1964–2002: OR 3.02 (95% CI 2.32–3.93) compared to all other dogs (Villamil *et al.*, 2011)

Neoplasia – overall

- Several studies have suggested a higher risk (Michalska & Michalski, 1997; Srivastava *et al.*, 2009)

Osteosarcoma

- Breed at increased risk in case series
- OR 2.3 (95% CI 1.8–3.0) compared to German Shepherd Dogs in a retrospective study of VMDB data, 1980–1994
- Across all breeds there was an increased risk with increasing age, plateauing at 10 years
- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes
- Breed also at high risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Dobermanns had an overall incidence of 24 cases (95% CI 13–35) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR

- Males over-represented
- Median age was 7.5 years in Dobermanns; risk increases with increasing age (Ru *et al.*, 1998; Egenvall *et al.*, 2007)

Synovial myxoma

- 4/6 cases in a USA-based series were Dobermanns or their crosses (66.7%; $p < 0.0001$)
- Mean age 9.2 (6–13) years
- The stifle was the most commonly affected joint (Craig *et al.*, 2002)

Neurological conditions

Cervical spondylomyelopathy (cervical vertebral malformation, wobblers syndrome)

- Breed at high risk
- Defects are stenosis of the vertebral canal and malformation of cervical vertebrae (C5–C7, most commonly C6–C7 in the Dobermann)
- Mean age 6.8 years in the Dobermann (da Costa, 2010)

Dancing Dobermann disease

- Uncommon
- Age of clinical onset 6 months to 7 years
- Poorly understood but thought to be a peripheral neuropathy (Chrisman, 1990)

Deafness and vestibular disease

- Autosomal recessive inheritance suspected
- Affected puppies are deaf and demonstrate a head tilt and circling behaviour from 3 weeks (Wilkes & Palmer, 1990)

Discospondylitis

- Dobermanns were at higher risk in a study of 513 cases in North America: OR 2.3 (95% CI 1.3–4.0) compared to mixed-breed dogs
- Higher risk with increasing age
- Males twice as likely to be affected as females: OR 2.0 (95% CI 1.7–2.4) (Burkert *et al.*, 2005)

Idiopathic head tremor syndrome

- Breed at increased risk in a case series (retrospective study of 291 cases)
- Average age of onset 29 months
- Possible trigger events found in 21% of cases
- Mentation was normal in 93% cases (Wolf *et al.*, 2011; Shell *et al.*, 2015)

Intervertebral disc disease (IVDD)

- Breed at increased risk in a review of Swedish insurance data
- Dobermanns had 88.6 occurrences per 10 000 DYAR (third-highest), compared to 27.8 occurrences per 10 000 DYAR for the general population
- Incidence higher in males than females (Bergknut *et al.*, 2012)

Narcolepsy–cataplexy

- Autosomal recessive inheritance
- Symptoms appear around 4 weeks, increase with age and plateau around 10–32 weeks (John *et al.*, 2004; Tonokura *et al.*, 2007)

Ocular conditions**Multiple ocular defects**

- Congenital; autosomal recessive inheritance suspected
- May include microphthalmia, anterior segment dysgenesis, congenital cataract and retinal dysplasia (Bergsjø *et al.*, 1984; Lewis *et al.*, 1986)

Persistent hyperplastic tunica vasculosa lentis and persistent hyperplastic primary vitreous (PHPV)

- Congenital; dominant inheritance with incomplete penetration suggested
- Reported widely in Europe, but rare in the USA (Boevé, 1988; Leppänen & Saloniemi, 1998)

Physiological conditions**Blood group**

- In a study of 274 dogs in Portugal in which 56.9% were DEA 1.1-positive and 43.1% were DEA 1.1-negative, all 12 Dobermanns in the study were negative (Ferreira *et al.*, 2011)

Renal and urinary conditions**Familial renal disease (familial nephropathy)**

- Mode of inheritance unknown
- Dobermanns are believed to suffer from membranoproliferative glomerulonephritis, a glomerular basement membrane disorder which may progress to glomerulonephritis
- The disease presents at 1–6 years with proteinuria and chronic renal failure (Wilcock & Patterson, 1979; Chew *et al.*, 1983)

Urethral sphincter mechanism incompetence

- Breed at increased risk in case series
- Spayed females predisposed (Holt & Thrusfield, 1993; Arnold, 1997)

Urolithiasis – calcium oxalate

- Breed at increased risk in a Canadian case series of urolith submissions, 1998–2001: OR 5.60 (95% CI 1.83–17.18) compared to mixed-breed dogs (Ling *et al.*, 2003)

Reproductive conditions**Mammary neoplasia**

See under *Neoplastic conditions*

Prostate disease

- Dobermann was the breed at greatest risk of prostate disease in a study of 177 cases over 5.5 years
- Mean age of onset 8.9 years
- The study included all types of prostate disease; prostatic adenocarcinoma was the most common type in neutered male dogs across the study (Krawiec & Heflin, 1992)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 43% of insured Swedish Dobermanns had developed pyometra at 10 years of age (ranked 17th) compared to 19% for the general population (Jitpean *et al.*, 2012)

DOGO ARGENTINO**Neoplastic conditions****Mast cell tumour (MCT)**

- Breed at increased risk in case series (Leidinger *et al.*, 2014)

Neurological conditions**Congenital deafness**

- Prevalence of hearing disorders in a study of 299 Dogo Argentino dogs was 27%
- The Dogo Argentino seems to be affected by complete cochleosaccular degeneration, which can be bilateral or unilateral (Pellegrino *et al.*, 2009; Blum & Distl, 2013)

DOGUE DE BORDEAUX (FRENCH MASTIFF)

Cardiovascular conditions

Aortic stenosis – subaortic stenosis (SAS)

- Autosomal recessive inheritance proposed
- OR 11.2 ($p < 0.0001$) compared to the hospital population in a retrospective series of 976 cases of congenital heart disease in Italy
- Both healthy and affected dogs had a smaller aortic annulus than other breeds
(Höllermer *et al.*, 2008; Oliveira *et al.*, 2011; Ohad *et al.*, 2013)

Tricuspid valve dysplasia

- Autosomal recessive inheritance proposed
- In Israel, Dogue de Bordeaux had the highest prevalence of all dogs in the country
(Oliveira *et al.*, 2011; Ohad *et al.*, 2013)

Dermatological conditions

Footpad hyperkeratosis

- Familial
- Lesions appear by 6 months of age
(Lachaume *et al.*, 1998)

Endocrine conditions

Hypothyroidism

- In a study of 44 Dogue de Bordeaux, prevalence of hypothyroidism was 4.5%
- 7 of the 10 male dogs in the study had low T_4 levels (although 6 of these had normal TSH levels)
(Segalini *et al.*, 2009)

Neurological conditions

Thoracic spinal stenosis

- Breed at increased risk in case series
- 8/23 dogs with thoracic spinal stenosis were Dogue de Bordeaux
- 21/23 dogs were male
- Mean age was 23 months (range 4–105 months)
- Most common sites were T2–3 and T3–4
- Symptoms included proprioceptive deficits in the hindlimbs
(Johnson *et al.*, 2012)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Autosomal recessive inheritance suspected
- Progressive disease, mean age at death 20 months

- In a study of 102 adult Dogue de Bordeaux dogs, 33% had abnormal proteinuria likely to be related to a glomerulopathy
(Lavoué *et al.*, 2010, 2015)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk in a case series based on the populations of five RSPCA hospitals in the UK between 2006 and 2011
- Dogue de Bordeaux had a prevalence of 4.6%, compared to 2.2% for all dogs over the study period
- Mean age was 3.3 years for the Dogue de Bordeaux, compared to a mean age of 7.7 years for the overall population
(Gibson *et al.*, 2013)

DUTCH PARTRIDGE DOG (DRENTSE PATRIJSHOND)

Gastrointestinal conditions

Stomatocytosis–hypertrophic gastritis

- Autosomal recessive inheritance suspected
- Rare
- Mean age at presentation 9.5 months old
- Affected dogs had diarrhoea, haemolytic anaemia and ataxia
(Slappendel *et al.*, 1991)

Haematological/immunological conditions

Stomatocytosis–hypertrophic gastritis

See under *Gastrointestinal conditions*

DUTCH SHEEPDOG

See *Schapendoes*

DUTCH SPANIEL

See *Wetterhoun*

ELKHOUND

See *Norwegian Elkhound*; *Swedish Elkhound*

ENGLISH BULLDOG

See *Bulldog – English*

ENGLISH COCKER SPANIEL

See *Cocker Spaniel*

ENGLISH SETTER

Dermatological conditions

Atopic dermatitis (atopy)

- Breed at increased risk in a US series of 264 cases (1981–1984)
- Birth in autumn or summer is a predisposing factor
- One study showed dogs between 1 and 2 years of age have the highest probability of an insurance claim for atopy
- Some studies show no sex predilection, others show females predisposed

(Schick & Fadok, 1986;
Nødtvedt *et al.*, 2006)

Cutaneous neoplasia

See under *Neoplastic conditions*

Malassezia dermatitis

- Breed at increased risk: OR 7.7 (compared to the hospital population) in a retrospective study of 86 cases at a US university pathology department
- Age of incidence 6 months to 15 years
- Spayed females (RR 4.51) and castrated males (RR 4.19) were at increased risk compared to the hospital population
- May be seasonal

(Mauldin *et al.*, 1997)

Symmetric lupoid onychodystrophy

- Rare condition
- Breed at increased risk
- Affects dogs at 3–8 years of age

(Dahlgren *et al.*, 2016)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- English Setters may be predisposed to lymphocytic thyroiditis

(Graham *et al.*, 2007)

Gastrointestinal conditions

Intestinal adenoma

See under *Neoplastic conditions*

Musculoskeletal conditions

Elbow dysplasia

- Common cause of forelimb lameness
- Increased risk of unilateral anconeal process: OR 3.7 (95% CI 1.6–8.3) compared to mixed breeds in a study of 847 cases from the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Hip dysplasia

- Breed at increased risk in case series
- OR 1.49 (95% CI 1.35–1.65; $p < 0.001$) compared to all other dogs in a large retrospective study of VMDB cases, 1964–2003. Male neutered dogs and dogs < 4 years of age (of all breeds) combined found to be at increased risk

(Witsberger *et al.*, 2008)

Osteochondrosis – shoulder

- OR 10.1 (95% CI 6.5–15.9) compared to mixed breeds in a large study of 1242 cases from the VMDB, 1986–1995
- Males predisposed
- Age of onset usually 4–7 months, but can be older

(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- Breed at increased risk in case series
- OR 1.9 (95% CI 1.3–2.8) compared to mixed breeds in a large study of 5633 cases from the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Intestinal adenoma

- Breed at increased risk ($p < 0.01$) in a case series of 3827 gastrointestinal biopsies in the Czech Republic

(Frgelecová *et al.*, 2013)

Mast cell tumour (MCT)

- Breed at increased risk of cutaneous MCT in a retrospective study of VMDB data, 1964–2002: OR 1.88 (95% CI 1.34–2.64) compared to all other dogs

(Villamil *et al.*, 2011)

Primary brain tumour

See under *Neurological conditions*

Neurological conditions**Congenital deafness**

- 7.9% prevalence in one study of 3656 English Setters
- Signs seen from birth

(Strain, 2004)

Neuronal ceroid lipofuscinosis

- Rare
- Autosomal recessive inheritance. Mutation identified
- Signs seen at 1–2 years

(Lingaas *et al.*, 1998; Katz *et al.*, 2005)**Primary brain tumour – choroid plexus neoplasm**

- Breed at significantly increased risk compared to the general population in a study of post-mortem results of 435 dogs with intracranial neoplasia

(Song *et al.*, 2013)**Ocular conditions****Neuronal ceroid lipofuscinosis**See under *Neurological conditions***ENGLISH SHEPHERD****Drug reactions****Ivermectin and milbemycin**

- High doses can cause tremors, ataxia, coma and death
- Associated with the mutant *MDR1* allele

(Dowling, 2006)

ENGLISH SPRINGER SPANIELSee *Springer Spaniel***ENGLISH TOY SPANIEL**See *King Charles Spaniel***ENTLEBUCHER MOUNTAIN DOG****Ocular conditions****Cataract**

- Autosomal recessive inheritance
- In a Swiss study of 285 of this breed over 10 years, 20.4% had cataracts

- Age at diagnosis was 5.34 ± 2.71 years
- Localization: posterior pole. Non-progressive (Speiss, 1994; Kuster *et al.*, 2011)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance suspected
- In a Swiss study of 285 of this breed over 10 years, 6.3% had PRA
- Age at diagnosis was 4.93 ± 1.32 years

(Heitmann *et al.*, 2005; Kuster *et al.*, 2011)**Renal and urinary conditions****Ectopic ureter**

- High incidence
- In a study of 565 dogs, 67.1% had at least one ectopic ureter
- Males more often affected than females
- Mode of inheritance not determined

(Fritsche *et al.*, 2014)**ESKIMO DOG****Haematological/immunological conditions****Bone marrow necrosis**

- Significantly over-represented in a retrospective study of 34 cases

(Weiss, 2005)

Immune-mediated polyarthritisSee under *Musculoskeletal conditions***Musculoskeletal conditions****Immune-mediated polyarthritis**

- Breed at increased risk in case series
- Over-represented for both reactive immune-mediated non-erosive polyarthritis (11% of 18 cases) and idiopathic non-erosive immune-mediated polyarthritis (7% of 43 cases)
- Median ages respectively 6.7 years and 4.8 years

(Stull *et al.*, 2008)**Ocular conditions****Cataract**

- Prevalence of primary cataract 1.84%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Progressive retinal atrophy (PRA)

- American Eskimo Dog
- Autosomal recessive inheritance
- Progressive rod–cone degeneration
- Age of onset: 3–5 years

(Moody *et al.*, 2005)**ESTRELA MOUNTAIN DOG****Musculoskeletal conditions****Hip dysplasia**

- In a study of 313 Estrela Mountain Dogs, 66% were found to be affected

(Ginja *et al.*, 2009)**FINNISH HOUND****Endocrine conditions****Diabetes mellitus**

- Breed at increased risk in a retrospective review of Swedish insurance data, 1995–2004
- Incidence in the Finnish Hound was 36 cases per 10 000 DYAR, compared to an overall incidence in all breeds of 13 cases per 10 000 DYAR
- Females at significantly increased risk overall

(Fall *et al.*, 2007)**Neurological conditions****Cerebellar ataxia**

- Inherited, mutation identified
- Early onset (clinical signs from 3 months) and rapidly progressive

(Kyöstilä *et al.*, 2012)**FINNISH LAPPHUND****Neurological conditions****Lysosomal storage disease – glycogen storage disease type II (Pompe disease)**

- Autosomal recessive inheritance
- In one study 5% (5/95) dogs carried the genetic mutation for this disease

(Seppälä *et al.*, 2013)**Ocular conditions****Progressive retinal atrophy (PRA)**

- Autosomal recessive inheritance suspected
- Progressive rod–cone degeneration
- Late onset: clinical signs at 3–6 years

(Aguirre-Hernández *et al.*, 2007)**FINNISH SPITZ****Endocrine conditions****Diabetes mellitus**

- OR 2.32 (95% CI 1.22–4.41) compared to mixed-breed dogs in a retrospective report of 6860 dogs with diabetes mellitus from the VMDB, 1970–1999
- Females predisposed (OR 1.37)

(Guptill *et al.*, 2003)**Gastrointestinal conditions****Atresia ani**

- Breed at increased risk of this rare disease: OR 19.401 compared to mixed-breed dogs
- Females predisposed: OR 1.796
- Symptoms seen at weaning

(Vianna & Tobias, 2005)

Haematological/immunological conditions**Immune-mediated haemolytic anaemia (IMHA)**

- Breed at increased risk in case series
- Usually affects young adult and middle-aged animals
- Females predisposed

(Miller *et al.*, 2004)**Neurological conditions****Idiopathic epilepsy**

- Inherited as a polygenic trait
- In a study of 2141 Finnish Spitz in Finland, over 10 years, the prevalence was 5.36%
- Males predisposed
- Median age of onset was 3 years
- 85% of seizures had a focal onset, 54% were secondary generalized

(Viitmaa *et al.*, 2007, 2013)**Ocular conditions****Cataract**

- Prevalence of primary cataract 2.47%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

FLAT-COATED RETRIEVER**Musculoskeletal conditions****Bone tumour**See under *Neoplastic conditions*

Hip dysplasia

- Breed at increased risk described in several countries
- High heritability, with a greater degree in dams than sires, has been described (Leppänen & Saloneiemi, 1999; Wood *et al.*, 2000, 2004)

Patellar luxation

- Breed at increased risk in a retrospective report of cases from the VMDB, 1986–1995: OR 2.9 (95% CI 1.0–8.2) compared to mixed-breed dogs
- Males predisposed (OR 1.8) in a separate retrospective report of patellar luxation in large-breed dogs (>15 kg) (LaFond *et al.*, 2002; Gibbons *et al.*, 2006)

Spondylosis deformans and diffuse idiopathic skeletal hyperostosis (DISH)

- Breed at increased risk in a retrospective radiographic study at a Netherlands veterinary teaching hospital (2003–2008)
- OR 2.834 ($p < 0.001$) for spondylosis and OR 7.671 ($p < 0.001$) for DISH compared to all dogs having radiographs taken during the study period
- Prevalence increases with age (Kranenburg *et al.*, 2011)

Neoplastic conditions**Bone tumour**

- Breed at high risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Flat-coated Retrievers had an overall incidence of 35 cases (95% CI 27–43) per 10 000 DYAR compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Males over-represented
- Median age was 8.6 years in Flat-coated Retrievers; risk increases with increasing age (Egenvall *et al.*, 2007)

Sarcoma – soft tissue

- Breed at increased risk of histiocytic sarcoma
- May affect muscle/fascia of limbs or visceral organs (primarily spleen)
- Average age of onset for soft-tissue sarcomas is 8–9 years, but it may be seen earlier in this breed

(Morris *et al.*, 2002; Constantino-Casas *et al.*, 2011)

Ocular conditions**Glaucoma – primary**

- Breed at increased risk of glaucoma associated with pectinate ligament dysplasia
- In a UK study, 34.7% of 389 randomly selected Flat-coated Retrievers examined between 1994 and 1996 had a degree of pectinate ligament dysplasia

(Read *et al.*, 1998; Wood *et al.*, 1998)

FOX TERRIER**Cardiovascular conditions****Pulmonic stenosis**

- Described as a predisposed breed in a number of papers

(Darke, 1989)

Ventricular septal defect

- Breed at increased risk in case series
- Fox Terriers represented 4 of the 56 cases (7%)
- Mean age at diagnosis was 9 months (range 2.2 months to 11.8 years)

(Bomassi *et al.*, 2015)

Dermatological conditions**Atopic dermatitis (atopy)**

- Reported as a breed at increased risk (OR 2.9 compared to the general hospital population) in a study in the USA of 383 cases
- Some studies show no sex predilection, others show females are predisposed (Scott, 1981; Nødtvedt *et al.*, 2006; Nicorescu & Crivineanu, 2007)

Endocrine conditions**Diabetes mellitus**

- OR 3.02 (95% CI 1.94–4.70) compared to mixed-breed dogs in a retrospective report of 6860 dogs with diabetes mellitus from the VMDB, 1970–1999
- Females predisposed: OR 1.37

(Guptill *et al.*, 2003)

Hypothyroidism – congenital

- Toy Fox Terriers
- Autosomal recessive inheritance
- Affected pups had delayed eye-opening, abnormal haircoat, inactivity and goitre (Fyfe *et al.*, 2003)

Gastrointestinal conditions

Congenital megaesophagus

- Wire-haired and Smooth-haired Fox Terriers
- Autosomal recessive inheritance
- Presumed secondary to myasthenia gravis (see under *Neurological conditions*) (Jenkins *et al.*, 1976; Miller *et al.*, 1983)

Pancreatitis

- Breed at increased risk in a series of 80 cases (Pápa *et al.*, 2011)

Sialorrhoea with spirocercosis

- Breed at increased risk
- In a retrospective study of 298 spirocercosis cases, sialorrhoea was present in 33. Fox Terriers comprised 36% of cases with sialorrhoea but only 5% of the whole group (van der Merwe *et al.*, 2012)

Musculoskeletal conditions

Congenital myasthenia gravis

See under *Neurological conditions*

Patellar luxation

- Breed at increased risk in case series
- In a large retrospective case series based on cases recorded in the VMDB, 1986–1995, odds ratios were 12.8 (95% CI 4.1–40.1) (Toy Fox Terrier) and 2.2 (95% CI 1.1–4.5) (Wire-haired Fox Terrier) compared to mixed breeds (LaFond *et al.*, 2002)

Neurological conditions

Congenital myasthenia gravis

- Smooth-haired Fox Terrier
- Autosomal recessive inheritance
- Rare
- Age of clinical onset: 6–8 weeks (Miller *et al.*, 1983, 1984)

Hereditary ataxia

- Smooth-haired and Toy Fox Terriers
- Autosomal recessive inheritance suspected
- Originally reported in Sweden
- Rare
- Age of clinical onset: 2–6 months (Rohdin *et al.*, 2015)

Ocular conditions

Cataract

- Breed at increased risk compared to mixed-breed dogs in a retrospective report of dogs

with primary cataracts from the VMDB, 1964–2003

- Prevalence of primary cataract was 11.7% in the Smooth-haired Fox Terrier, 4.88% in the Wire-haired Fox Terrier, and 3.78% in the Toy Fox Terrier, compared to 1.61% in mixed-breed dogs

(Gelatt & MacKay, 2005)

Glaucoma – primary

- Toy and Wire-haired Fox Terriers
- Breed at increased risk compared to the overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
- Prevalence was 2.28% in Wire-haired Fox Terriers for the years 1994–2002, compared to an overall prevalence in all dogs of 0.89% in the same period
- Prevalence was 2.64% in Toy Fox Terriers and 1.97% in Wire-haired Fox Terriers for the years 1984–1993, compared to an overall prevalence in all dogs of 0.76% in the same period

(Gelatt & MacKay, 2004a)

Glaucoma – secondary

- Toy and Wire-haired Fox Terriers
- Breed at increased risk compared to the overall population of dogs in a retrospective study of 9695 dogs with secondary glaucoma from the VMDB, 1964–2003
- Wire-haired Fox Terriers were at increased risk of glaucoma secondary to cataracts, lens luxation, cataract surgery and uveitis, compared to mixed-breed dogs
- Toy Fox Terriers were at increased risk of glaucoma secondary to lens luxation, and cataract surgery, compared to mixed-breed dogs

(Gelatt & MacKay, 2004b)

Lens luxation – primary

- Toy and Wire-haired Fox Terriers
- Autosomal recessive inheritance
- Age of onset between 3 and 8 years (Curtis & Barnett, 1980; Gould *et al.*, 2011)

Renal and urinary conditions

Ectopic ureter

- Congenital anomaly
- Breed at increased risk in a series of 217 cases in female dogs recorded in the VMDB (1964–1981)

- OR 5.5 compared to all female dogs seen in the same period
- Usually presents < 1 year of age
- More commonly diagnosed in females (Hayes, 1984)

FOXHOUND

Haematological/immunological conditions

Pelger–Huet anomaly

- Inherited leucocyte anomaly which is often asymptomatic (Bowles *et al.*, 1979)

Infectious conditions

Leishmaniasis

- Visceral form is endemic in Foxhounds in North America
- In the absence of the insect vector, the disease may be transmitted directly or vertically in kennels (Schantz *et al.*, 2005)

FRENCH ALPINE MASTIFF

See *Savoy Sheepdog*

FRENCH BULLDOG

Cardiovascular conditions

Pulmonic stenosis

- Breed at greatly increased risk (OR 19.1; $p < 0.0001$ compared to the general hospital population) in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- This study found a male predisposition across all breeds (OR 1.5) (Oliveira *et al.*, 2011)

Ventricular septal defect

- Breed at greatly increased risk (OR 7.2; $p = 0.0003$) compared to the general hospital population) in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy

- 48% of the cases of ventricular septal defect in this study were seen with another cardiac defect, most commonly pulmonic stenosis (Oliveira *et al.*, 2011)

Dermatological conditions

Atopic dermatitis (atopy)

- Breed at increased risk in a Swiss case series
- Breed at increased risk in a study of Hamburg cases: OR 46.9 (95% CI 16.7–131.3) compared to the general hospital population
- Prevalence was found to vary with geographical location (Picco *et al.*, 2008; Jaeger *et al.*, 2010)

Demodicosis

- French Bulldogs are at increased risk of juvenile-onset, generalized demodicosis, based on a large retrospective case–control study of a US practice database: OR 5.0 (95% CI 1.4–18.0) compared to the hospital population
- Juvenile onset was defined as < 18 months old
- Other risk factors associated with demodicosis included pyoderma, short hair and concurrent coccidiosis (Plant *et al.*, 2011)

Gastrointestinal conditions

Gastrointestinal changes seen with BOAS

- French Bulldogs have a high incidence of brachycephalic obstructive airway syndrome (BOAS: see under *Respiratory conditions*)
- Gastrointestinal changes (inflammatory disease of the oesophagus, stomach and duodenum) are commonly seen with this condition (Garcia-Sancho *et al.*, 2013)

Ulcerative colitis (histiocytic ulcerative colitis)

- Predisposed to a condition similar to that seen in Boxers
- Believed to be associated with intramucosal *E. coli*, and has been shown to be responsive to fluoroquinolones (Tanaka *et al.*, 2003; Manchester *et al.*, 2013)

Musculoskeletal conditions

Fracture of the thoracic limb – humeral condyle

- Breed at increased risk: OR 59.0 (95% CI 18.44–166.89) in a retrospective series of 189 cases seen at the Norwegian College of Veterinary Medicine, 1971–1991 (control group was the Norwegian Kennel club population over the same period)

- Lateral condyle more commonly affected than medial (OR 90.3)
- Most fractures occurred at 3–4 months old (Rørvik, 1993)

Hemivertebrae (wedge-shaped vertebrae)

See under *Neurological conditions*

Patellar luxation

- Breed at increased risk: OR 5.4 (95% CI 3.1–9.3; $p < 0.001$) compared to mixed-breed dogs, using the data of 210 824 dogs attending 119 clinics across England
- The study found that females (OR 1.3) and neutered dogs (OR 2.4) were at increased risk (O'Neill *et al.*, 2016c)

Neoplastic conditions

Primary brain tumour

See under *Neurological conditions*

Neurological conditions

Arachnoid cyst – spinal

- Rare condition
- Breed at increased risk in a retrospective case series of 122 cases
- French Bulldogs were the third most common breed with 13 of the 122 cases
- All cases in French Bulldogs were in the thoracolumbar spine
- Across the study, males were over-represented (95/122 cases were male)
- In a separate study of 343 French Bulldogs presenting with neurological disease, 2002–2016, 25 were affected by spinal arachnoid cysts, of which 88% were thoracic and 12% were cervical (Mauler *et al.*, 2014; Mayousse *et al.*, 2017)

Hemivertebrae (wedge-shaped vertebrae)

- Selection for 'screw tail' (hemivertebrae in the tail) in French Bulldogs predisposes to hemivertebrae in other areas of the spine
- May be associated with neurological signs including weakness of the hindlimbs, faecal and urinary incontinence (Schlensker & Distl, 2016)

Intervertebral disc disease (IVDD)

- Breed at increased risk
- In common with other chondrodystrophoid breeds, the disease can occur at a younger age in French Bulldogs and peaks at 3–7 years

- Males were predisposed (57.4%) in French Bulldogs in one study
- French Bulldogs were in the top five breeds affected in a large USA-based retrospective study of 90 004 dogs presented to a university teaching hospital. 27.06% were affected, compared to 4.43% of mixed breeds
- In a separate study of 343 French Bulldogs presenting with neurological disease, 2002–2016, 156 were affected by IVDD. 39.8% of cases were cervical and 60.2% were non-cervical. C3–C4 was the most commonly affected site (Bellumori *et al.*, 2013; Aikawa *et al.*, 2014; Mayousse *et al.*, 2017)

Primary brain tumour – oligodendroglioma

- Breed at significantly increased risk compared to the general population in a study of post-mortem results of 435 dogs with intracranial neoplasia
- In a separate study of 343 French Bulldogs presenting with neurological disease, 2002–2016, 25 were affected by brain tumours, of which 17 were gliomas (Song *et al.*, 2013; Mayousse *et al.*, 2017)

Ocular conditions

Corneal ulceration (ulcerative keratitis)

- In a UK-based study of first-opinion practice, the prevalence in the French Bulldog was 1.87% (95% CI 0.97–3.24), compared to an overall prevalence of 0.8% (95% CI 0.75–0.86) (O'Neill *et al.*, 2017b)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk in a retrospective study of 114 cases
- Males over-represented (74/114)
- 75.4% cases occurred in dogs < 1 year
- Often bilateral in French Bulldogs (Mazzucchelli *et al.*, 2012)

Renal and urinary conditions

Urolithiasis – cystine

- Breed at increased risk in a retrospective case series of 2543 uroliths examined at the Budapest Urolith Centre in Hungary (2001–2012)
- Mutation in the *SLC3A1* and *SLC7A9* genes suspected (Harnevik *et al.*, 2006; Bende *et al.*, 2015)

Reproductive conditions

Delivery by caesarean section

- French Bulldogs were in the top 10 breeds in a cross-sectional study conducted by the Kennel Club in the UK. The study found a caesarean rate of 84% in 80 litters of French Bulldogs born over a 10-year period
- Dystocia is usually a result of fetopelvic disproportion in French Bulldogs
- French Bulldogs are excluded from insurance cover for caesarean section by Swedish pet insurance companies (Bergström *et al.*, 2006; Evans & Adams, 2010a)

Respiratory conditions

Brachycephalic obstructive airway syndrome (BOAS)

- Breed at increased risk due to conformation
- Risk increases with shorter muzzle length, increased neck girth and obesity
- French Bulldog kennel club registrations in the UK have grown from 350 in 2004 to 6990 in 2013 making this a condition of enormous welfare concern
- Concurrent gastrointestinal lesions may occur in this breed

(Poncet *et al.*, 2005; Emmerson, 2014; Liu *et al.*, 2015)

FRENCH MASTIFF

See *Dogue de Bordeaux*

FRENCH SPANIEL**Dermatological conditions**

Acral mutilation and analgesia

- Rare condition. 13 cases identified in French Spaniels in Canada
- Signs noted between 3.5 and 12 months of age (Paradis *et al.*, 2005)

FRISIAN WATER DOG

See *Wetterhoun*

GALGOS ESPAÑOL

See *Spanish Greyhound*

GERMAN PINSCHER**Cardiovascular conditions**

Vascular ring anomaly – persistent right aortic arch (PRAA)

See under *Gastrointestinal conditions*

Gastrointestinal conditions

Vascular ring anomaly – persistent right aortic arch (PRAA)

- High prevalence noted in the German Pinscher
- Symptoms of regurgitation are seen at weaning (Menzel & Distl, 2011)

Ocular conditions

Cataract

- Autosomal recessive inheritance suspected
- In a retrospective review of ophthalmological examinations of 122 German Pinschers in Finland, 6.5% were found to have cataracts
- In a retrospective review of examinations of 409 German Pinschers in Germany (1997–2013) by certified veterinary ophthalmologists, 15.6% were diagnosed with primary cataracts (Leppänen *et al.*, 2001; Pfahler *et al.*, 2015)

Persistent hyperplastic tunica vasculosa lentis

- Congenital, autosomal recessive inheritance suspected
- In a retrospective review of ophthalmological examinations carried out in 122 German Pinschers in Finland, 8.4% were affected
- In a retrospective review of examinations of 409 German Pinschers in Germany (1997–2013) by certified veterinary ophthalmologists, 3.2% were affected (Leppänen *et al.*, 2001; Pfahler *et al.*, 2015)

GERMAN SHEPHERD DOG (GSD or ALSATIAN)**Cardiovascular conditions**

Aortic stenosis – subaortic stenosis (SAS)

- Breed at mildly increased risk (OR 1.8; $p=0.0166$) compared to the general hospital population in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- This study found a male predisposition (OR 1.7). Previous studies have found no gender difference

- Inheritance possibly autosomal dominant with variable penetrance

(Oliveira *et al.*, 2011)

Haemangiosarcoma – right atrial

See *Pericardial effusion*, below, and under *Neoplastic conditions (Haemangiosarcoma)*

Patent ductus arteriosus

- Breed at markedly increased risk (OR 5.2; $p < 0.0001$) compared to the general hospital population) in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- This study found a female predisposition (OR 2.7)

(Oliveira *et al.*, 2011)

Pericardial effusion

- Breed at increased risk in a number of studies
- Average age at diagnosis 9.7 years
- Haemangiosarcoma of the right atrium is a common cause in GSDs

(Mellanby & Herrtage, 2005; MacDonald *et al.*, 2009)

Vascular ring anomaly – persistent right aortic arch (PRAA)

See under *Gastrointestinal conditions*

Ventricular arrhythmia and sudden death

- Polygenic inheritance likely
- Affects young dogs, with a window of vulnerability for sudden death at 3–18 months (peak 6–7 months)
- Sudden death occurs more frequently during rapid-eye-movement (REM) sleep and after exercise

(Cruickshank *et al.*, 2009)

Ventricular septal defect

- Breed at markedly increased risk (OR 3.7; $p = 0.001$) compared to the general hospital population) in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- Almost half of the ventricular septal defect cases in this study were seen with another cardiac defect, most commonly pulmonic stenosis

(Oliveira *et al.*, 2011)

Dermatological conditions

Anal furunculosis (perianal fistula)

- GSDs represent >80% of reported cases
- Most common in middle-aged to older dogs
- Some similarities to perianal Crohn's disease in humans
- Immune-mediated pathogenesis suspected (Kennedy *et al.*, 2008; Massey *et al.*, 2014)

Atopic dermatitis (atopy)

- Breed at increased risk in a study of Hamburg cases: OR 6.6 (95% CI 3.9–11.2) compared to the general hospital population
- The same study showed an increased risk in Munich cases: OR 5.0 (95% CI 2.9–8.6) compared to the general hospital population
- The same study showed an increased risk in Melbourne cases: OR 2.4 (95% CI 1.4–4.1) compared to the general hospital population
- This study suggests that the overall breed risk profile varies substantially depending on geographical location (Nødtvedt *et al.*, 2006; Jaeger *et al.*, 2010; Tengvall *et al.*, 2013)

Cutaneous neoplasia

See under *Neoplastic conditions*

Discoid lupus erythematosus

- Uncommon disease of the nasal planum
- GSDs and their crosses were over-represented in one study (44% of 27 cases) (Wiemelt *et al.*, 2004)

Erythema multiforme

- Rare condition
- Breed at increased risk ($p = 0.0039$) in a small study of 16 cases seen at a dermatology referral centre at Cornell University, US (Scott & Miller, 1999)

Familial vasculopathy

- Autosomal recessive inheritance
- Seen in Canada, plus one case report from the UK
- Occurs at 4–7 weeks of age
- Possible association with vaccination (Weir *et al.*, 1994; Rest *et al.*, 1996)

Focal metatarsal fistulation (sterile idiopathic pedal panniculitis)

- Rare disease, seen almost exclusively in GSDs
- Age of onset usually 2.5–4 years (Kunkle *et al.*, 1993; Paterson, 1995)

German Shepherd Dog pyoderma

- Described as a distinct clinical entity in the GSD. Also known as folliculitis–furunculosis and cellulitis syndrome (FFCS)
- In a prospective study in France, 80/277 cases of pyoderma (superficial and deep) were GSDs
- 23 of the GSDs in this study had no underlying disease
- Of these, middle-aged dogs were predisposed (mean age 7.5 years)
- Males were over-represented (20 of the 23)
- Abnormalities of cell-mediated immunity and low IgA levels have been implicated in the pathogenesis

(Denerolle *et al.*, 1998)**Leishmaniasis**See under *Infectious conditions***Mucocutaneous pyoderma**

- Uncommon condition; may be difficult to differentiate from discoid lupus erythematosus
- The GSD and its crosses appear to be predisposed

(Bassett *et al.*, 2004; Wiemelt *et al.*, 2004)**Multiple collagenous naevi (nodular dermatofibrosis)**

- Autosomal dominant inheritance
- Occurs with renal cystadenocarcinomas and uterine leiomyomas

(Castellano & Idiart, 2005; Zanatta *et al.*, 2013)**Otitis externa**

- Breed at increased risk ($p < 0.001$) when compared with the hospital's normal population in a study of 430 cases at a veterinary teaching hospital in Jerusalem

(Zur *et al.*, 2011)**Symmetric lupoid onychodystrophy**

- GSD at increased risk in a number of studies
- GSDs represented 22.2% cases but only 4% of the hospital population in one retrospective study (RR 5.6)

(Scott *et al.*, 1995; Harvey & Maxwell, 1996)**Systemic lupus erythematosus**See under *Haematological/immunological conditions***Drug reactions****Ivermectin and milbemycin**

- Breed at increased risk of ivermectin and milbemycin toxicosis
- High doses can cause tremors, ataxia, coma and death

(Merola *et al.*, 2009)**Endocrine conditions****Acromegaly**

- Rare condition, seen mainly in intact female dogs
- GSDs were found to be at increased risk in a retrospective study of the medical records of 34 380 dogs attending two referral veterinary hospitals in Italy between 2004 and 2008. The incidence in the intact female GSD population was 1.65%, significantly higher ($p < 0.0001$) than the incidence of 0.15% in the overall entire female population

(Fracassi *et al.*, 2014)**Pituitary dwarfism**

- Rare
- Autosomal recessive mode of inheritance suggested
- Clinical signs (failure to grow) seen at 2–3 months

(Voorbij *et al.*, 2011; Tsai *et al.*, 2012)**Gastrointestinal conditions****Acquired megaesophagus**

- Breed at increased risk in case series
- Megaesophagus can be idiopathic, or secondary to peripheral neuropathies, myasthenia gravis, oesophagitis and chronic or recurrent gastric dilatation

(Gaynor *et al.*, 1997)**Anal furunculosis (perianal fistula)**See under *Dermatological conditions***Antibiotic-responsive diarrhoea (small intestinal bacterial overgrowth)**

- Breed at increased risk in case series
- May be associated with IgA deficiency in GSDs (see under *Haematological/immunological conditions*)

(Batt *et al.*, 1991; German *et al.*, 2000b)**Congenital megaesophagus**

- Autosomal dominant inheritance with incomplete penetrance has been proposed

- Varies in severity, with mild cases resolving spontaneously after several months
(Tsai *et al.*, 2012)

Exocrine pancreatic insufficiency (EPI)

- Very high prevalence in the GSD (approximately 60% of all cases)
- The underlying cause is pancreatic acinar atrophy (PAA)
- Typical age of onset 36 months
- Higher prevalence in females
(Batchelor *et al.*, 2007; Tsai *et al.*, 2012, 2013)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in case series
- 28.6% of dogs were GSDs in a series of 112 cases
(Brockman *et al.*, 1995; Buber *et al.*, 2007)

Haemangiosarcoma – splenic

See under *Neoplastic conditions*

Hypocobalaminaemia

- Breed at increased risk of decreased cobalamin (vitamin B₁₂) (<100 ng/l)
- In a UK-based study of 9960 dogs, GSDs were found to have an OR of 1.5 (95% CI 1.7–3.9; $p < 0.004$) compared to all dogs in the study
- This study found a lower risk of decreased folate (<3.5 µg/dl): OR 0.7 (95% CI 0.6–0.8; $p = 0.004$) for GSDs

(Dandrieux *et al.*, 2013)

Inflammatory bowel disease

- Breed at increased risk in a case series (18/80 cases were GSDs)
- Mean age 4.9 years
- May be associated with antibiotic responsive diarrhoea and IgA deficiency
(Batt *et al.*, 1991; Craven *et al.*, 2004; Lee *et al.*, 2015)

Intussusception – gastro-oesophageal

- Rare condition seen relatively more frequently in this breed
- Usually seen in dogs < 3 months old
- Higher incidence in males
- May be associated with congenital megaoesophagus
(Roach & Hecht, 2007; Shilby *et al.*, 2014)

Parvovirus enteritis

See under *Infectious conditions*

Small intestinal volvulus

- Rare condition
- Few cases are reported in the literature, but a high proportion of those cases are in GSDs, suggesting a predisposition
- Usually associated with chronic intestinal disorders
(Cairo *et al.*, 1999)

Splenic masses

- Breed at increased risk in case series
- In a retrospective study, 25/249 cases were in GSDs
- Median age at diagnosis was 10 years
- Male:female ratio was 2:1
- 53% of cases were malignant (73.5% of those were haemangiosarcoma)
(Eberle *et al.*, 2012)

Splenic torsion

- Rare condition
- Breed at increased risk in a retrospective review of case records at seven North American referral hospitals. GSDs represented 24/102 cases (23.5%)
- Males were at increased risk
(DeGroot *et al.*, 2016)

Vascular ring anomaly – persistent right aortic arch (PRAA)

- High prevalence noted in the GSD (12/52 cases in one study, and 16/44 cases in another, were GSDs)
- Symptoms of regurgitation are seen at weaning
(Buchanan, 2004; Krebs *et al.*, 2014)

Haematological/immunological conditions

Babesiosis

See under *Infectious conditions*

Ehrlichiosis

See under *Infectious conditions*

Haemophilia A

- Factor VIII deficiency
- X-linked recessive inheritance
- Male dogs predominantly affected
- A moderately severe form of the disease in this breed allows survival to adulthood
(Parry *et al.*, 1988; Aslanian *et al.*, 2014)

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance (Brooks, 1999)

Hypocobalaminaemia

See under *Gastrointestinal conditions*

Selective IgA deficiency

- Breed at increased risk in case series
- 14% of 319 GSDs were found to have low serum IgA in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease
- In GSDs, low IgA levels may be associated with antibiotic-responsive diarrhoea, inflammatory bowel disease, atopy and pancreatic acinar atrophy (Batt *et al.*, 1991; Olsson *et al.*, 2014)

Systemic lupus erythematosus

- Uncommon (incidence is approximately 0.03% of the general canine population)
- Affects mainly males in the GSD
- Average age of 5 years
- Symptoms may include skin lesions, pyrexia and polyarthrititis (Fournel *et al.*, 1992)

Infectious conditions**Babesiosis**

- Breed at increased risk of testing seropositive to *Babesia* organisms in a study of 651 blood samples from dogs in Hungary
- A genetic predisposition towards carrier status is proposed (Hornok *et al.*, 2006)

Cryptococcosis

- Breed at increased risk in a case series in Australia (where cryptococcosis is the most common systemic mycosis)
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors (O'Brien *et al.*, 2004)

Disseminated aspergillosis

- Breed at increased risk in case series
- In a retrospective study which reviewed the medical records of the University of California, Davis, teaching hospital, GSDs had an OR of

43.0 (95% CI 20.0–91.0; $p < 0.0001$) compared to the general hospital population

- Females were over-represented: OR 2.9 (95% CI 1.2–6.7; $p = 0.02$)
- Median age 4.5 years (range 2–8 years) (Schultz *et al.*, 2008)

Ehrlichiosis

- Breed considered to be at high risk in many areas of the world
- In a Kenyan study of 514 dogs with ehrlichiosis, GSDs represented 42.6% cases
- In a retrospective case-control study in Peru, the odds ratio was 12.2 ($p < 0.01$) compared to the general population (Contreras *et al.*, 2009; Kitaa *et al.*, 2014)

Leishmaniasis

- Breed at increased risk in a study of 390 cases seen at a university veterinary clinic in Barcelona
- GSDs represented 13.6% of dogs with leishmaniasis, compared to 7.5% of the general population ($p < 0.001$)
- Bimodal age distribution, with peaks at 2–4 years and 7+ years
- Male dogs are predisposed (Miranda *et al.*, 2008)

Parvovirus enteritis

- Breed at increased risk in several case series from a variety of continents
- Dogs aged between 6 weeks and 6 months are at highest risk
- Absence of, or inadequate vaccination is a risk factor (Houston *et al.*, 1996)

Musculoskeletal conditions**Calcinosis circumscripta**

- GSDs < 2 years old account for > 50% of reported cases
- May affect periarticular soft tissues resulting in lameness, soft tissues surrounding the spinal cord resulting in spinal cord compression or lingual soft tissue resulting in lesions on the tongue (Roudebush *et al.*, 1988; Dennis *et al.*, 2006)

Elbow dysplasia

- Common cause of forelimb lameness in young dogs
- Breed at increased risk in a number of case series

- In a large retrospective case series based on cases recorded in the VMDB, 1986–1995, GSDs were recorded with the following risks: fragmentation of the medial coronoid process, OR 43.7 (95% CI 30.1–63.2) compared to mixed breeds; ununited anconeal process, OR 8.2 (95% CI 6.2–10.9) compared to mixed breeds
- Polygenic inheritance. Primary lesions of elbow dysplasia may be inherited separately (LaFond *et al.*, 2002; Stock *et al.*, 2011)

Gracilis contracture (fibrotic myopathy)

- Uncommon condition which is mainly seen in young male GSDs
- Commonly occurs in athletic individuals
- May affect other muscles including sartorius, semimembranosus, semitendinosus, and biceps femoris (Lewis *et al.*, 1997; Steiss, 2002; Spadari *et al.*, 2008)

Hip dysplasia

- Breed at increased risk in case series
- OR 3.61 (95% CI 3.52–3.71; $p < 0.001$) compared to all other dogs in a large retrospective study of VMDB cases, 1964–2003. Male neutered dogs and dogs <4 years of age of all breeds combined found to be at increased risk
- OR 5.7 (95% CI 5.0–6.3) compared to mixed-breed dogs in a separate retrospective study of VMDB data, 1986–1995
- Prevalence in GSDs has been estimated as 35% (LaFond *et al.*, 2002; Witsberger *et al.*, 2008; Fels & Distl, 2014)

Lumbosacral disease

See under *Neurological conditions*

Lumbosacral transitional vertebrae

- Significantly higher prevalence in GSDs (5.7%) compared to all other breeds combined (3.5%; $p < 0.001$), in a Swiss study reviewing radiographs of 4000 dogs of medium-large breeds
- Predisposes to lumbosacral disease (see under *Neurological conditions*) (Damur-Djuric *et al.*, 2006)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males may be predisposed

- OR 9.6 (95% CI 4.3–21.0) compared to mixed breeds in a retrospective study of VMDB data, 1986–1995 (LaFond *et al.*, 2002; Safra *et al.*, 2013)

Osteochondrosis – sacrum

See under *Neurological conditions (Lumbosacral disease)*

Osteochondrosis – shoulder

- Breed at increased risk in case series. OR 2.1 (95% CI 1.4–3.1) compared to mixed-breed dogs in a retrospective study of VMDB data, 1986–1995
- Age at presentation 4–14 months
- Males over-represented (25/36 cases in one study in Poland) (LaFond *et al.*, 2002; Biezyński *et al.*, 2012)

Osteochondrosis – stifle

- Breed at increased risk in case series. OR 17.5 (95% CI 7.5–40.7) compared to mixed-breed dogs in a retrospective study of VMDB data, 1986–1995
- Mean age at presentation 5.9 months
- Males over-represented (103/135 cases of multiple breeds in a review of cases in Europe and the USA) (Montgomery *et al.*, 1989; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- Breed at increased risk
- OR 3.3 (95% CI 2.9–3.7) compared to mixed breeds in a retrospective study of VMDB data, 1986–1995 (LaFond *et al.*, 2002)

Spondylosis deformans

- Common condition in ageing dogs
- Breed at increased risk in a retrospective study of radiographic records (OR 3.327; $p < 0.001$)
- Usually clinically insignificant but may be associated with type II intervertebral disc disease (Levine *et al.*, 2006; Kranenburg *et al.*, 2011)

Systemic lupus erythematosus

See under *Haematological/immunological conditions*

Neoplastic conditions**Anal sac adenocarcinoma**

- Breed at increased risk in a number of small studies (7/42 cases in one study)
- Mean age 10.2–10.8 years
- Some surveys suggest a female predisposition (Bennett *et al.*, 2002; Potanas *et al.*, 2015)

Haemangiosarcoma

- Breed at high risk of haemangiosarcoma in all locations
- In one retrospective study of 104 cases, 34 were GSDs
- Most common location was the spleen, then the skin/subcutis, liver, heart and lung
- Mean age was reported as 10 years
- In another study of 92 dogs with splenic haemangiosarcoma, GSDs had an OR of 4.7 (95% CI 2.7–7.8) compared with all other purebred dogs
- Haemangiosarcoma is the most common tumour of the heart (Brown *et al.*, 1985; Ng & Mills, 1985; Prymak *et al.*, 1988)

Mammary neoplasia

- Most common tumour in entire female dogs
- Breed at increased risk compared to the general population of insured dogs in a study based on Swedish insurance data: OR 3.2 (99% CI 2–4.9) (Egenvall *et al.*, 2005)

Melanoma

- Breed at increased risk of limbal melanoma
- In a retrospective histopathological study of 244 cases submitted to the Comparative Ocular Pathology Laboratory of Wisconsin, 1988–1998, 41% of limbal melanomas were from GSDs
- In another retrospective report of 30 cases seen at the Animal Health Trust in England, the GSD was not found to be at increased risk compared to the general population (Giuliano *et al.*, 1999; Donaldson *et al.*, 2006)

Perianal adenocarcinomas

- Breed at increased risk (compared to the hospital population at the time) in a series of 41 cases in male dogs at Colorado State University
- Entire males at significantly greater risk than neutered males (Vail *et al.*, 1990)

Renal cystadenocarcinomas

See under *Renal and urinary conditions*

Sinonasal neoplasia

- In a retrospective review of cases seen at the Animal Medical Center in New York, GSDs represented 27.2% (9/33) of chondrosarcomas and 12.5% (2/16) of osteosarcomas
- Average age was 7 years (chondrosarcomas) and 10 years (osteosarcomas)
- Males were over-represented (Patnaik *et al.*, 1984b)

Testicular neoplasia

See under *Reproductive conditions*

Thymoma

- Uncommon tumour
- In one retrospective study, GSDs represented 5/18 cases, suggesting a possible predisposition
- In a more recent retrospective study, GSDs represented only 4/116 cases
- Middle-aged and older dogs affected
- May be associated with myasthenia gravis, hypercalcaemia or immune-mediated disease (Aronsohn, 1985; Day, 1997b; Robat *et al.*, 2013)

Trichoepithelioma

- Breed at increased risk in case series
- Mean age 8.6 years
- Predilection sites: limbs, back and trunk (Scott & Anderson, 1991)

Uterine leiomyoma

Occurs with renal cystadenocarcinomas (see under *Renal and urinary conditions*)

Neurological conditions**Degenerative myelopathy**

- Common in GSDs
- Caused by a mutation in the superoxide dismutase 1 (*SOD1*) gene which has a high degree of penetrance
- Age of clinical onset > 5 years (Tsai *et al.*, 2012; Holder *et al.*, 2014)

Discospondylitis

- GSDs were at higher risk in a study of 513 cases in North America: OR 2.6 (95% CI 1.8–3.9) compared to mixed-breed dogs
- Risk increases with increasing age

- Males twice as likely to be affected as females: OR 2.0 (95% CI 1.7–2.4)

(Burkert *et al.*, 2005)

Giant axonal neuropathy

- Rare condition
 - Autosomal recessive inheritance suspected
 - Age of clinical onset 14–15 months
- (Duncan & Griffiths, 1981)

Idiopathic epilepsy

- GSDs had an increased risk compared to crossbreeds, with an OR of 1.9 (95% CI 1.28–2.80; $p=0.001$) in a large retrospective study of data from 92 UK-based primary veterinary practices
- Males over 1.5 times more likely to be affected than females (95% CI 1.44–2.06; $p<0.001$)
- Age of onset 6 months to 6 years
- GSDs are also at increased risk of cluster seizures

(Kearsley-Fleet *et al.*, 2013; Packer *et al.*, 2016b)

Intervertebral disc disease (IVDD)

- GSDs were found to be at higher risk when compared to other dogs in a number of studies
- GSDs were the third most commonly affected breed after mixed-breed dogs and Dachshunds in one retrospective study of 946 cases seen at a referral hospital in Switzerland (representing 7% of cases)
- GSDs accounted for 52.4% of 21 cases in a retrospective study of dogs with T1–T9 lesions diagnosed by MRI at a New York referral hospital
- GSDs were found to be the breed at highest risk of L7–S1 disc lesions amongst non-chondrodystrophic breeds
- Older dogs are more likely to be affected; GSDs often had lesions at multiple sites

(Fluehmann *et al.*, 2006; Suwankong *et al.*, 2008; Hearon *et al.*, 2014)

Lumbosacral disease

- GSDs are frequently reported as the breed at highest risk
- Factors include early degeneration of the lumbosacral (LS) intervertebral disc, instability of the LS joint, sacral osteochondrosis, LS transitional vertebrae and LS step formations, which are higher in the GSD than in other large-breed dogs

- Congenital, with developmental and acquired factors believed to be involved

(Suwankong *et al.*, 2008; Amort *et al.*, 2012; Ondreka *et al.*, 2012)

Lumbosacral transitional vertebrae

- Predisposes to lumbosacral disease. See above and under *Musculoskeletal conditions*

(Damur-Djuric *et al.*, 2006)

Ocular conditions

Chronic superficial keratitis (pannus)

- GSDs have the highest incidence of all breeds
- Immune-mediated basis, linked to MHC class 2 genes
- Median age 4 years
- When seen in younger dogs it is rapidly progressive and more severe
- Prevalence and severity increase at higher altitude
- May be associated with plasma cell infiltration of the nictitating membrane

(Jokinen *et al.*, 2011)

Melanoma

See under *Neoplastic conditions*

Plasma cell infiltration of the nictitating membrane (plasmoma)

- Breed at increased risk
 - May be associated with pannus
- (Jokinen *et al.*, 2011)

Renal and urinary conditions

Renal cystadenocarcinomas

- Autosomal dominant inheritance suggested
 - Mean age at diagnosis: 8.2 years
 - Multiple, bilateral tumours are seen, with generalized nodular dermatofibrosis and (in female dogs) multiple uterine leiomyomas
- (Lium & Moe, 1985; Castellano *et al.*, 2000; Castellano & Idriat, 2005)

Urolithiasis – silica

- Breed at increased risk in several case series
- However, a more recent large retrospective study of 1697 cases over 20 years failed to show a predisposition in the GSD
- Mean age at diagnosis 5.8 years
- Males predisposed

(Osborne *et al.*, 1981, 1999; Aldrich *et al.*, 1997)

Reproductive conditions**Cryptorchidism**

- Congenital defect believed to be inherited as a sex-limited, autosomal recessive trait
- Breed at increased risk in case series (Romagnoli, 1991; Yates *et al.*, 2003)

Mammary neoplasia

See under *Neoplastic conditions*

Prostate disease

- Breed at increased risk: OR 2.1 (95% CI 1.5–2.9; $p < 0.001$) compared to the general hospital population a large retrospective study conducted at Alfort Veterinary College, France (2002–2009)
- Study included dogs with benign prostatic hypertrophy, prostatitis, cysts, abscesses, neoplasia and squamous metaplasia
- Mean age was 8.6 ± 3.2 years
- Entire males were at significantly increased risk: OR 3.096 (95% CI 1.466–6.541) except for prostatic neoplasia (Polisca *et al.*, 2016)

Testicular neoplasia

- In a study of 232 routine necropsy cases, 16/59 dogs found to have testicular tumours were GSDs
- Mean age 9–11 years
- Retained testicles at increased risk
- One study suggests GSDs have an increased risk of seminoma (Hayes & Pendergrass, 1976; Sapierzyński *et al.*, 2007a; Grieco *et al.*, 2008)

Uterine leiomyoma

Seen with renal cystadenocarcinomas; see under *Renal and urinary conditions*

GERMAN SPANIEL

See *Wachtelhund*

GLEN OF IMAAL TERRIER**Ocular conditions****Progressive retinal atrophy (PRA)**

- Autosomal recessive inheritance
- Mutation in the ADAM 9 gene identified
- Late-onset rod–cone dystrophy (Kropatsch *et al.*, 2010)

GOLDEN RETRIEVER**Behavioural conditions****Aggression**

- Breed identified with significantly lower levels of aggression towards both humans and other dogs in a study based on Canine Behavioural Assessment and Research Questionnaires (C-BARQ) completed by owners in the USA (Duffy *et al.*, 2008)

Cardiovascular conditions**Aortic stenosis – subaortic stenosis (SAS)**

- OR 9.63 (95% CI 4.21–22.01) compared to the general hospital population, in an Australian retrospective study of 40 dogs with SAS seen between 2001 and 2012
- In an Italian study of 241 cases OR was 3.6 ($p < 0.0001$) (Oliveira *et al.*, 2011; Aherne & Beijerink, 2013)

Cardiomyopathy – X-linked muscular dystrophy

- Also known as Golden Retriever muscular dystrophy
- Similar to Duchenne muscular dystrophy in humans
- Rare
- Most clinically affected cases are male (Jeanson-Leh *et al.*, 2014)

Haemangiosarcoma – right atrial

- A cause of pericardial effusion (see below)
 - Breed at increased risk
 - Golden Retrievers represented 8/23 cases in a retrospective report of cases confirmed by histopathology as haemangiosarcoma
 - Mean age was 9.1 years
- See also under *Neoplastic conditions* (Weisse *et al.*, 2005)

Mitral valve dysplasia

- OR 25.36 (95% CI 6.04–106.49) compared to the general hospital population, in an Australian retrospective study of 8 dogs with mitral valve dysplasia seen between 2001 and 2012
- Golden retrievers represented 3/13 dogs with mitral valve dysplasia in a retrospective study of 151 cases of congenital heart disease seen in a Swedish referral hospital (1989–1996) (Tidholm, 1997; Aherne & Beijerink, 2013)

Pericardial effusion

- Breed at increased risk
- May be idiopathic or a result of a cardiac mass
- Golden Retrievers represented 20/107 pericardial effusion cases in a retrospective study carried out at University of California, Davis, USA, 1985–2006
- Golden Retrievers represented 47/143 cases in a retrospective study at a referral centre in the UK. 40 of these cases were negative for a cardiac mass on echocardiogram (mean age of these dogs was 8.5 years). 7 cases had a cardiac mass on echocardiogram (mean age of these dogs was 9.5 years)

(Stafford Johnson *et al.*, 2004;
MacDonald *et al.*, 2009)

Tricuspid valve dysplasia

- In an Italian retrospective case series of 976 dogs with congenital heart defects, Golden Retrievers had an OR of 6.6 ($p = 0.0022$)

(Oliveira *et al.*, 2011)

Dermatological conditions**Atopic dermatitis (atopy)**

- Breed at increased risk in a study of Munich cases (OR 8.6; 95% CI 4.7–15.6) compared to the general hospital population
- Breed at increased risk in a study of Campbell, USA cases. OR 5.1 (95% CI 3.1–8.4) compared to the general hospital population
- This study suggested that overall breed risk profile varies substantially depending on geographical location

(Scott & Paradis, 1990;
Jaeger *et al.*, 2010)

Cutaneous neoplasia

See under *Neoplastic conditions*

Grass awn migration

- Breed at increased risk in case series. May be due to behaviour/lifestyle
- Common in the summer months

(Brennan & Ihrke, 1983)

Ichthyosis

- Autosomal recessive inheritance, mutation identified
- Usually diagnosed in young dogs median age 2.29 years

(Mauldin *et al.*, 2008; Grall *et al.*, 2012)

Pyotraumatic folliculitis (acute moist dermatitis, hot spot, wet eczema)

- Breed at increased risk in case series. Relative risk 1.31 compared to the general hospital population

- Dogs < 4 years old at increased risk ($p < 0.0001$)
(Holm *et al.*, 2004)

Endocrine conditions**Hypothyroidism**

- Breed at increased risk in a case series. OR 2.86 ($p = 0.036$) compared to a large university hospital population. Neutered animals (both sexes) were at increased risk
- Breed with higher prevalence of thyroid hormone autoantibodies (THAA). In a cohort study of 287 948 serum samples from dogs in the US with clinical signs of hypothyroidism, Golden Retrievers had an OR 1.9 ($p = 0.001$) of being affected compared to dogs of all other breeds. Across the study, females were over-represented and the highest prevalence was in dogs 2–4 years old
- Hypothyroidism seems to occur at a younger age (2–3 years) in at risk breeds

(Panciera, 1994; Nachreiner *et al.*, 2002)

Thyroid neoplasia

May be associated with hyper- or hypothyroidism, but most are euthyroid. See under *Neoplastic conditions*

Gastrointestinal conditions**Acquired megaesophagus**

- Breed at increased risk in case series
- Megaesophagus can be idiopathic or secondary to peripheral neuropathies, myasthenia gravis, oesophagitis and chronic or recurrent gastric dilatation

(Gaynor *et al.*, 1997)

Congenital portosystemic shunt

- Breed at increased risk in case series
- Golden Retrievers represented 11/100 cases in a retrospective case series of intrahepatic shunts at the University of Pennsylvania
- Golden Retrievers represented 4/11 cases of intrahepatic shunts in a retrospective case series at Tufts University
- Clinical signs usually seen in young dogs < 1 year
- Golden Retrievers have intrahepatic shunts more commonly than extrahepatic shunts

(D'Anjou *et al.*, 2004; Weisse *et al.*, 2014)

Cricopharyngeal dysfunction

- Inheritance has been demonstrated; a single recessive allele of large effect contributes to the disease in Golden Retrievers (Davidson *et al.*, 2004)

Folate – low serum level

- Breed at increased risk of low serum folate level in a retrospective study of 9960 samples over 12 years at Liverpool University, UK
- OR 1.9 (95% CI 1.6–2.3)
- This indicates that either the reference interval is lower in Golden Retrievers or that they are at increased risk of proximal small intestinal disease

(Dandrieux *et al.*, 2013)

Haemangiosarcoma – splenic

- Often metastasizes to the liver, omentum or lungs
- Breed at increased risk. Golden Retrievers represented 13/59 cases in a retrospective report of cases at the University of Guelph, Canada

See also under *Neoplastic conditions*

(Kim *et al.*, 2007)

Melanoma

See under *Neoplastic conditions*

Haematological/immunological conditions**Haemophilia A**

- Golden Retrievers experience a mild factor VIII deficiency
- X-linked recessive inheritance

(Brooks *et al.*, 2005)

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance

(Brooks, 1999)

Hereditary spectrin deficiency

- Spectrin deficiency has been estimated to be present in 11–24.6% of Dutch Golden Retrievers
- Autosomal dominant inheritance
- Causes spherocytosis and fragility of red blood cells

(Slappendel *et al.*, 2005)

Selective IgA deficiency

- Breed at increased risk in case series
- 13% of 168 Golden Retrievers were found to have low serum IgA (<0.07 g/l) in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease (Olsson *et al.*, 2014)

Infectious conditions**Blastomycosis**

- Breed at increased risk in case series
- Golden Retrievers represented 15.2% of 125 cases in a retrospective series of dogs with pulmonary blastomycosis seen at the University of Minnesota, 1989–2006. Mean age was 4 years
- Entire males were predisposed
- Geographic distribution: mainly North America (Mississippi, Missouri, Ohio River valleys, mid-Atlantic States, Quebec, Manitoba and Ontario)
- Proximity to a body of water is a risk factor in endemic areas (Rudmann *et al.*, 1992; Crews *et al.*, 2008)

Borreliosis (Lyme disease)

- Breed at increased risk of nephritis (protein-losing nephropathy) (Dambach *et al.*, 1997)

Metabolic conditions**Overweight/obesity**

- 51.9% of Golden Retrievers attending veterinary clinics in China were classified as obese (Courcier *et al.*, 2010; Mao *et al.*, 2013; Raffan *et al.*, 2016)

Musculoskeletal conditions**Cranial cruciate ligament (CCL) disease**

- Breed at increased risk in a number of studies
- OR 1.9 (95% CI 1.1–3.3) compared to mixed-breed dogs in one retrospective study of data from UK first-opinion practices (VetCompass data)
- In this study dogs of all breeds which had high body weight within their breed were at increased risk (OR 3.4; $p < 0.001$) compared to dogs of low body weight within their breed; dogs 9–11.9 years old were at increased risk (OR 4.4; $p < 0.001$) compared to dogs <3 years old; neutered female dogs were at increased risk (OR 2.1; $p < 0.001$) compared to entire females

- In a large retrospective case series based on cases recorded in the VMDB, 1964–2003, OR 1.11 (95% CI 1.05–1.17; $p=0.007$) compared to all dogs. Neutered males, neutered females and dogs >4 years of age of all breeds combined were at increased risk

(Witsberger *et al.*, 2008; Guthrie *et al.*, 2012; Torres de la Riva *et al.*, 2013; Taylor-Brown *et al.*, 2015)

Elbow dysplasia

- Developmental joint disease. Common cause of forelimb lameness in young dogs
- Breed at increased risk in a number of studies
- In a large retrospective case series based on cases recorded in the VMDB, 1986–1995 Golden Retrievers were recorded with the following risks: fragmentation of the medial coronoid process, OR 5.5 (95% CI 3.3–9.2) compared to mixed breeds; ununited anconeal process, OR 4.9 (95% CI 3.6–6.7) compared to mixed breeds; osteochondrosis, OR 42.2 (95% CI 18.7–95.3)
- Polygenic inheritance. Primary lesions of elbow dysplasia may be inherited separately (LaFond *et al.*, 2002; Lavrijsen *et al.*, 2012)

Hip dysplasia

- Breed at increased risk in a number of studies
- OR 3.3 (95% CI 3.0–3.8) compared to mixed breeds in a large retrospective case series based on cases recorded in the VMDB, 1986–1995
- In a large retrospective case series based on cases recorded in the VMDB, 1964–2003, OR 2.70 (95% CI 2.61–2.8; $p<0.001$) compared to all dogs. Neutered males and dogs <4 years of age of all breeds combined were at increased risk (LaFond *et al.*, 2002; Witsberger *et al.*, 2008)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males may be predisposed
- OR 5.4 (95% CI 2.5–11.7) compared to mixed breeds in a large retrospective case series based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Muscular dystrophy – X-linked

- X-linked recessive inheritance
- Similar to Duchenne muscular dystrophy in humans

- Degenerative muscular condition resulting in death usually from cardiac or respiratory failure

See also under *Cardiovascular conditions* and *Respiratory conditions*

(Cooper *et al.*, 1988; Jeanson-Leh *et al.*, 2014)

Osteochondrosis – shoulder

- Breed at increased risk (9/36 cases were Golden Retrievers in one Polish study). Males predisposed across all breeds (25 of the 36 cases were male)
- Age of onset usually 4–7 months but can be older
- OR 12.6 (95% CI 10.00–15.9) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002; Biezyński *et al.*, 2012)

Osteochondrosis – stifle

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 3.9 (95% CI 1.2–12.8) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Osteosarcoma

See under *Neoplastic conditions*

Neoplastic conditions

Canine acanthomatous ameloblastoma

- Increased risk based on retrospective case series at University of California, Davis, USA
- Mean age 8.8 years
- Commonly located in the rostral mandible
- Neutered dogs of both sexes were over-represented

(Fiani *et al.*, 2011)

Fibrosarcoma – lingual

- Breed at increased risk
- OR 3.64 (95% CI 1.56–8.47) compared to all other breeds in a retrospective case series in Colorado, 1995–2004

(Dennis *et al.*, 2006)

Fibrosarcoma – oral

- Breed at increased risk
- 13/25 dogs were Golden Retrievers in a retrospective case series in Colorado, USA

- Males may be predisposed
- Mean age of onset is 7.5 years, but up to 25% of cases have been seen at < 5 years of age (Ciekot *et al.*, 1994)

Haemangiosarcoma

- 20% of Golden Retrievers in the USA develop haemangiosarcomas
- Most common sites are the spleen (28–50%), right atrium/auricle (3–30%), skin and subcutaneous tissue (13%)
- Breed at increased risk of cutaneous haemangiosarcoma in a retrospective study of VMDB data, 1964–2002: OR 2.80 (95% CI 2.25–3.49) compared to all other dogs
- Neutered females are at 4 times higher risk than entire females
- Shared predisposing genetic loci with lymphosarcoma have been identified (Villamil *et al.*, 2011; Kahn *et al.*, 2013; Torres de la Riva *et al.*, 2013; Thomas *et al.*, 2014; Tonomura *et al.*, 2015)

Histiocytic sarcoma

- Breed at increased risk in case series
- OR 5.0 in a series of 73 cases in Japan (Affolter & Moore, 2002; Takahashi *et al.*, 2014)

Lymphoma

- Most common haematological malignancy
- Golden Retrievers are predisposed to B- and T-cell lymphoma
- 13% of Golden Retrievers in the USA develop lymphoma
- Cutaneous lymphoma – Golden Retrievers have an OR of 2.35 (95% CI 1.76–3.13) compared to all other dogs, based on data from the VMDB, 1964–2002
- Neutered male Golden Retrievers are 3 times more at risk than entire males
- Most cases are seen in middle-aged dogs (mean 6–7 years)
- Shared predisposing genetic loci with haemangiosarcoma have been identified (Villamil *et al.*, 2011; Torres de la Riva *et al.*, 2013; Boerkamp *et al.*, 2014; Elvers *et al.*, 2015; Tonomura *et al.*, 2015)

Mast cell tumour (MCT)

- Breed at increased risk
- In a large UK study based on VetCompass data, Golden Retrievers had a prevalence of

1.39%, compared to an overall prevalence of 0.27% in all dogs

- Golden Retrievers are at increased risk of multiple tumours compared to all breeds (OR 3.8)
- Neutered female Golden Retrievers are at higher risk than entire females
- May be seen at any age (from 4 months onwards), but usually seen in older animals (Murphy *et al.*, 2006; Torres de la Riva *et al.*, 2013; Shoop *et al.*, 2015)

Melanoma – cutaneous

- Breed at increased risk of cutaneous melanoma in a retrospective study of VMDB data, 1964–2002: OR 2.59 (95% CI 2.01–3.35) compared to all other dogs (Villamil *et al.*, 2011)

Melanoma – ocular

- Breed at increased risk of limbal and canine anterior uveal melanoma in a retrospective case series at the Animal Health Trust, UK
- Mean age was 6.2 years (with a bimodal distribution of 3–4 years and 7–10 years) (Donaldson *et al.*, 2006)

Melanoma – oral

- Breed at increased risk of oral melanoma in a retrospective study of the University of Missouri, USA, database
- OR 1.84 ($p < 0.001$) compared to a database population
- Average age 11.4 years (Ramos-Vara *et al.*, 2000)

Osteosarcoma

- Breed at increased risk in case series
- OR 2.1 (95% CI 1.6–2.8) compared to German Shepherd Dogs in a retrospective study of VMDB data, 1980–1994
- Across all breeds there was an increased incidence with age, plateauing at 10 years
- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes (Ru *et al.*, 1998)

Primary brain tumour

See under *Neurological conditions*

Sarcoma – soft tissue

- Breed at increased risk of cutaneous soft-tissue sarcoma in a retrospective study of VMDB data, 1964–2002: OR 2.88 (95% CI 2.46–3.36) compared to all other dogs (Villamil *et al.*, 2011)

Sweat gland tumour

- Breed at increased risk of apocrine ductal adenoma in a case series
- Mean age was 9.1 years (Kalahar *et al.*, 1990)

Thyroid neoplasia

- Represents 1.1% of all neoplasms
- Breed at increased risk in case series
- OR 2.2 (95% CI 1.75–2.84) compared to all dogs in a retrospective study of data from the VMDB, 1995–2006
- Older animals at increased risk: 57% were between 10 and 15 years of age (Wucherer & Wilke, 2010)

Trichoepithelioma

- Breed at increased risk in case series
- Mean age 8.6 years (Scott & Anderson, 1991)

Neurological conditions**Horner's syndrome**

- Golden Retrievers represented 110/155 dogs with Horner's syndrome referred to a UK veterinary ophthalmologist over a 10-year period
- 95 of the 110 Golden Retrievers were male (Boydell, 2000)

Idiopathic epilepsy

- Breed at increased risk
- In a retrospective review of 394 cases seen at a Munich Veterinary teaching hospital, 2002–2008, Golden Retrievers had an OR of 2.04 ($p < 0.001$) compared to the hospital population
- Age of onset: 6 months to 6 years (Zimmermann *et al.*, 2009)

Primary brain tumour

- Breed at increased risk in a number of case series
- Approximately 50% of primary brain tumours in the Golden Retriever are meningiomas
- Golden Retrievers are also at increased risk of choroid plexus tumours

- Older dogs affected (Snyder *et al.*, 2006; Westworth *et al.*, 2008; Song *et al.*, 2013)

Sensory ataxic neuropathy

- Reported in Swedish Golden Retrievers
- Mutation identified
- Insidious onset between 2 and 8 months of age (Jaderlund *et al.*, 2007; Baranowska *et al.*, 2009)

Trigeminal neuropathy

- Breed at increased risk in case series (Mayhew *et al.*, 2002)

Ocular conditions**Cataract**

- Dominant inheritance with incomplete penetration suggested
- In a UK study of eye examinations of 2251 Golden Retrievers the incidence of cataracts was 7.4%
- Prevalence of primary cataract 2.20%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Localization: posterior subcapsular cortex
- Age of onset 6–18 months; slow-growing or non-progressive
- Usually bilateral
- Perinuclear cataracts are seen less frequently. These usually progress to generalized opacity
- Cataracts may also be seen secondary to pigmented uveitis/iridociliary cysts (Curtis & Barnett, 1989; Gelatt & MacKay, 2005)

Glaucoma – secondary

- Associated with pigmented uveitis and iridociliary cysts
- Autosomal dominant inheritance with partial penetrance suspected
- Usually bilateral
- May be associated with secondary cataracts (Deehr, 1998; Sapienza *et al.*, 2000; Townsend & Gornik, 2013; Holly *et al.*, 2016)

Melanoma

See under *Neoplastic conditions*

Pigmentary uveitis

See *Glaucoma – secondary*

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance suspected
- A number of different forms exist in Golden Retrievers, each having a different causal mutation. Three forms with identified mutations exist at the time of writing, and these have been termed prcd-PRA, PRA_{GR1} and PRA2

(Downs *et al.*, 2014b)**Retinal dysplasia – geographic**

- Inheritance suspected

(Holle *et al.*, 1999)**Retinal dysplasia – multifocal**

- Congenital, autosomal recessive inheritance suspected

(Long & Crispin, 1999)

Retinal pigment epithelial dystrophy (RPED, central progressive retinal atrophy)

- More prevalent in the UK than in the USA. Becoming less prevalent following the introduction of control schemes

(Curtis, 1988)

Spontaneous chronic corneal epithelial defects (refractory corneal ulceration, indolent ulcers)

- Breed at increased risk in case study
- Mean age 9.25 years

(Murphy *et al.*, 2001)**Uveal cysts**

- Often associated with pigmentary uveitis and secondary glaucoma in Golden Retrievers

See also *Glaucoma – secondary*(Deehr, 1998; Sapienza *et al.*, 2000)**Renal and urinary conditions****Ectopic ureter**

- Uncommon congenital anomaly
- Breed at increased risk in case series
- Golden Retrievers represented 16.6% of 24 cases in one study at Ohio State University veterinary teaching hospital, USA
- Usually presents < 1 year of age
- More commonly diagnosed in females

(Holt & Moore, 1995; Cannizzo *et al.*, 2003)**Familial renal disease (familial nephropathy)**

- Reported in young dogs < 3 years old
- Cases present in renal failure

(Kerlin & Van Winkle, 1995;

de Moraes *et al.*, 1996)**Lyme nephritis**

- Breed at increased risk to developing protein-losing nephropathy as a result of *Borrelia burgdorferi* infection

(Dambach *et al.*, 1997)**Reproductive conditions****Pyometra (cystic endometrial hyperplasia–pyometra complex)**

- Breed at increased risk in case series
- In a study of data from five RSPCA hospital populations in the UK between 2006 and 2011. The prevalence in Golden Retrievers was 5.4%, compared to 2.2% in all dogs over the study period
- Common disease of older entire bitches (mean age 7.25 years)
- Most cases present within 12 weeks of oestrus

(Egenvall *et al.*, 2001; Gibson *et al.*, 2013)**Respiratory conditions****Muscular dystrophy – X-linked**

- May affect the diaphragm muscle, trachea and alveoli resulting in respiratory failure

See also under *Musculoskeletal conditions*(Lessa *et al.*, 2014)**GORDON SETTER****Dermatological conditions****Black hair follicular dysplasia**

- Reported in Gordon Setters in Norway
- Males over-represented
- Antinuclear antibodies were found in some cases, leading to a suspicion that there may be an autoimmune basis
- May be related to symmetric lupoid onychodystrophy, suggesting a common genetic predisposition

(Øvrebø Bohnhorst *et al.*, 2001)**Canine juvenile cellulitis**

- Uncommon condition
- One report suggested that 24.4% of Gordon Setter litters were affected, 1985–1988
- Age of onset usually 1–4 months

(Mason & Jones, 1989)

Symmetric lupoid onychodystrophy

- Reported in Gordon Setters in Norway
- One report suggested 12.6% prevalence
- Males over-represented

- Immune-mediated disease, autoimmune basis suspected
- May be related to black hair follicular dysplasia, suggesting a common genetic predisposition

(Øvrebø Bohnhorst *et al.*, 2001; Wilbe *et al.*, 2010b; Dahlgren *et al.*, 2016)

Endocrine conditions

Hypothyroidism

- Breed at increased risk in a case series at the University of Montreal, Canada
- In a Norwegian study of 291 8-year-old Gordon Setters, 2.7% had hypothyroidism
- Mean age 6.4 years
(Scott & Paradis, 1990; Bianchi *et al.*, 2015; Ziener *et al.*, 2015)

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a case series based on data from the VMDB, 1980–1989
- OR 4.1 (95% CI 1.8–9.3)
(Glickman *et al.*, 1994)

Musculoskeletal conditions

Elbow dysplasia

- Fragmentation of the medial coronoid process is common in Gordon Setters
- OR 19.8 (95% CI 9.2–42.8) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002)

Hip dysplasia

- OR 2.0 (95% CI 1.4–3.0) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Decreasing prevalence seen in Gordon Setters in a French study
- One study in Norway suggested Gordon Setters that were puppies in spring and summer had a lower incidence than those that were puppies in autumn and winter
(Hanssen, 1991; LaFond *et al.*, 2002; Genevois *et al.*, 2008)

Neurological conditions

Cerebellar degeneration

- Rare
- Autosomal recessive inheritance suspected
- Signs seen at 6 months to 4 years of age

- Affected dogs develop a slowly progressive ataxia
(de Lahunta *et al.*, 1980; Agler *et al.*, 2014)

Lethal astrocytosis

- Rare
- Autosomal recessive inheritance is suspected
- Signs are seen from 3–4 weeks
- Affected puppies have hair abnormalities, progressive weakness and recumbency by 5–6 weeks of age
(Yaeger *et al.*, 2000)

Ocular conditions

Progressive retinal atrophy (PRA)

- Rod–cone degeneration 4
- Affected dogs suffer night blindness progressing to total blindness
- Autosomal recessive inheritance
- Late onset
(Downs *et al.*, 2014a)

Retinopathy (rod–cone dystrophy)

- Affected dogs suffer day blindness
(Good *et al.*, 2016)

GRAND BLEU DE GASCOGNE HOUND

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaires relating to the 10 years prior to 2004
- Prevalence ratio for GDV mortality was 20.6 (95% CI 5.1–82.1; $p < 0.001$) compared to all other breeds combined
- Prevalence ratio for GDV morbidity was 31.1 (95% CI 11.32–85.45; $p < 0.001$) compared to all other breeds combined
(Evans & Adams, 2010b)

GREAT DANE

Cardiovascular conditions

Atrial fibrillation (AF)

- Breed at increased risk in a North American retrospective study of 109 cases
- Great Danes were over-represented both in the group with AF but no structural or

functional heart disease ($p < 0.05$) and in the group with AF, structural or functional heart disease and ascites or pulmonary oedema ($p < 0.001$)

- Males were over-represented: 80/109 (73.4%; $p < 0.01$)

(Menaut *et al.*, 2005)

Dilated cardiomyopathy (DCM)

- Autosomal dominant inheritance proposed
- In a UK study of a population of 107 Great Danes prevalence was 35.5%
- Males were predisposed
- Median age of onset was 5 years
- Median survival time following diagnosis was 5 weeks for Great Danes in one study
- Hypothyroidism has been seen associated with myocardial failure in some Great Danes. Heart function improved with management of the hypothyroidism (Meurs *et al.*, 2001; Phillips & Harkin, 2003; Martin *et al.*, 2010; Stephenson *et al.*, 2012)

Mitral valve dysplasia

- Rare
- In one study of 29 dogs with mitral valve dysplasia, 13 were Great Danes (Liu & Tilley, 1975; Matic, 1988)

Dermatological conditions

Cryptococcosis

See under *Infectious conditions*

Demodicosis

- Great Danes are at increased risk of juvenile-onset, generalized demodicosis: OR 2.6 (95% CI 1.2–5.5) based on a large retrospective case control study of a US practice database
- Juvenile onset was defined as < 18 months old
- Other risk factors associated with demodicosis included pyoderma, short hair and concurrent coccidiosis

(Plant *et al.*, 2011)

Epidermolysis bullosa acquisita

- Rare immune-mediated skin disease seen in humans and dogs
- In a USA-based retrospective study of 20 cases (1994–2014) 11 were Great Danes
- 10 of these were male (intact or neutered)
- Median age was 1.2 years

(Bizikova *et al.*, 2015)

Ichthyosis

- Inherited disorder; mutation in Great Danes has been identified
- Great Danes suffer from a severe form (Metzger *et al.*, 2015)

Idiopathic sterile granuloma and pyogranuloma

- Uncommon
- No age or sex predisposition
- Great Danes were over-represented compared to the general hospital population in a retrospective study in New York (Panich *et al.*, 1991)

Endocrine conditions

Hypoadrenocorticism (Addison's disease)

- Great Danes were at increased risk ($p = 0.001$) in a retrospective study of cases at University of Pennsylvania. 7/79 cases were Great Danes, compared to 1/200 dogs in the control population
- Median age was 3.2 years
- Females may be predisposed (Peterson *et al.*, 1996; Adler *et al.*, 2007)

Hypothyroidism (lymphocytic thyroiditis)

- Breed at increased risk of having thyroglobulin autoantibodies predisposing to lymphocytic thyroiditis
- May occur at a younger age in at-risk breeds (2–3 years)
- Neutered animals (both sexes) at increased risk
- Hypothyroidism has been seen associated with myocardial failure in some Great Danes. Heart function improved with management of the hypothyroidism (Nesbitt *et al.*, 1980; Haines *et al.*, 1984; Phillips & Harkin, 2003)

Gastrointestinal conditions

Chronic hepatitis

- Breed at increased risk in a UK-based study of 551 cases: OR 4.0 (95% CI 1.9–8.9; $p = 0.004$) compared to a control population
- Median age was 6 years 2 months
- Females were over-represented across the whole study

(Bexfield *et al.*, 2012a)

Gastric dilatation/volvulus (bloat, GDV)

- Highest incidence was found in Great Danes in a prospective cohort study in the USA involving 1914 dogs of 11 large and giant breeds
- Incidence in Great Danes was 53 cases per 1000 DYAR, compared to 26 cases per 1000 DYAR in all giant breeds (>45 kg body weight) and 23 cases per 1000 DYAR in all large breeds (23–45 kg body weight)
- Males slightly over-represented
- Risk increased with age
(Glickman *et al.*, 2000b)

Splenic torsion

- Breed at increased risk in a retrospective series of 102 cases (1992–2014)
- Great Danes accounted for 15 cases (14.7%)
(DeGroot *et al.*, 2016)

Infectious conditions**Cryptococcosis**

- Breed at increased risk in case series
- Usually seen in dogs < 4 years
- Living outdoors and exposure to bird droppings are risk factors
(Malik *et al.*, 1995; O'Brien *et al.*, 2004)

Infectious skin diseases

See under *Dermatological conditions*

Musculoskeletal conditions**Calcium phosphate deposition**

- Familial disease seen in Great Danes
- Paraplegia and incoordination are seen in very young pups, caused by mineral deposits in the vertebral bodies
- Periarticular and soft-tissue mineralization is also seen
(Woodard *et al.*, 1982)

Hip dysplasia

- Breed at increased risk in case series
- OR 1.6 (95% CI 1.2–2.1) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995
- Neutered male dogs predisposed
(LaFond *et al.*, 2002)

Inherited myopathy of Great Danes

- Previously known as central core myopathy
- Rare condition
- Affects dogs < 1 year of age

- Autosomal recessive inheritance likely
- Mutation identified
(Feliu-Pascual *et al.*, 2006; Böhm *et al.*, 2013)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males may be predisposed
- OR 189.8 (95% CI 82.7–435.6) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.* 2002, Safra *et al.*, 2016)

Osteochondrosis – elbow

- Breed at increased risk in case series
- Males predisposed
- OR 87.0 (95% CI 32.9–230.3) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002;

Kirberger & Stander, 2007)

Osteochondrosis – shoulder

- Breed at increased risk in case series
- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 32.8 (95% CI 23.8–45.3) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995
(Rudd *et al.*, 1990; LaFond *et al.*, 2002)

Osteochondrosis – stifle

- Breed at increased risk in case series
- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 309.4 (95% CI 123.4–775.5) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002)

Osteosarcoma

See under *Neoplastic conditions*

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- OR 2.3 (95% CI 1.8–3.0) compared to mixed breeds in a large retrospective series

based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Osteosarcoma

- Common bone tumour accounting for 85% of primary bone tumours and 5% of all canine neoplasms. Breed at increased risk in several case series
- OR 12.0 (95% CI 5.8–24.5; $p < 0.001$) compared to mixed-breed dogs in a retrospective study of 179 cases seen at the University of Florida, 1996–2005
- Mean age in this study was 7.8 years for the Great Dane
- Across all breeds there was an increased incidence with increasing age, plateauing at 10 years
- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes
- Breed at high risk of malignant bone tumours in a study of Swedish insurance data (1995–2002). Great Danes had an overall incidence of 45 cases (95% CI 25–65) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Median age was 7.2 years in Great Danes in this study
(Ru *et al.*, 1998; Egenvall *et al.*, 2007; Rosenberger *et al.*, 2007)

Neurological conditions

Cervical spondylomyelopathy (cervical vertebral malformation, wobbler syndrome)

- Common condition in large and giant breeds
- Breed at increased risk in case series. 15/27 dogs were Great Danes in one USA-based retrospective case series (2000–2012)
- Median age at presentation was 2 years
- Great Danes are predisposed to the osseous form, with C5–C6 and C6–C7 being the most commonly affected locations
(Delamaide Gasper *et al.*, 2014; Martin-Vaquero & da Costa, 2014)

Congenital deafness

- Great Danes homozygous for the merle gene are at increased risk of deafness
(Strain *et al.*, 2009)

Discospondylitis

- Breed at increased risk in a case–control study of 513 USA-based cases (1980–2001)
- Great Danes were at the highest risk: OR 7.3 (95% CI 4.3–12.6) compared to mixed breeds
- Males twice as likely to be affected as females
- Risk increased with age: dogs > 10 years had an OR of 4.1 (95% CI 2.9–5.8)
(Burkert *et al.*, 2005)

Primary orthostatic tremor

- Uncommon condition of unknown cause
- Muscle tremor while standing at rest
(Garosi *et al.*, 2005)

Ocular conditions

Eversion of the cartilage of the nictitating membrane

- Breed at increased risk
- In a study of 12 cases in 10 dogs, 3 were Great Danes
- Seen in young dogs (6–19 months)
(Allbaugh & Stuhr, 2013)

Glaucoma – primary

- Breed at increased risk of glaucoma associated with goniodysgenesis
- Age of onset 1–9 years
- Glaucoma is also seen associated with uveal cysts in Great Danes
(Speiss *et al.*, 1998; Wood *et al.*, 2001, 2004)

Multiple ocular defects

- Congenital
- Seen in homozygous merles with predominantly white coats
- Also referred to as merle ocular dysgenesis
- Defects may include microphthalmia, microcornea, cataract, equatorial staphylomas and retinal dysplasia; these dogs may also be deaf
(Bauer *et al.*, 2015)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk in a retrospective review of 155 cases in Europe (2001–2008)
- 75.4% of cases (all breeds) occurred before 1 year of age
- Great Danes are at increased risk of a bilateral occurrence with a short interval between prolapses
(Mazzucchelli *et al.*, 2012)

Uveal cysts

- Great Danes have been found to have uveal cysts associated with glaucoma (Speiss *et al.*, 1998)

Reproductive conditions**Dystocia**

- Breed at increased risk in a retrospective study of Swedish insurance data of 195 931 bitches (1995–2002)
- Incidence in Great Danes was 15.6 cases per 1000 DYAR (eighth-highest), compared to 5.7 cases per 1000 DYAR for the overall population (Bergström *et al.*, 2006)

GREAT PYRENEES

See *Pyrenean Mountain Dog*

GREATER SWISS MOUNTAIN DOG

See *Swiss Mountain Dog*

GREYHOUND**Cardiovascular conditions****Hypertension**

- Breed at increased risk
- In a USA-based cross-sectional study of 47 retired racing Greyhounds, 62% were hypertensive (systolic blood pressure > 160 mmHg)
- One study suggests Greyhounds have an increased risk of ischaemic stroke, which may result from hypertension
- One report suggests that the tendency to higher blood pressure in greyhounds results from the particular profile of eicosanoid metabolites of arachidonic acid which promote vascular dysfunction
- Hypertension may be associated with microalbuminaemia (Cox *et al.*, 1976; Surman *et al.*, 2012; Kent *et al.*, 2014; Martinez *et al.*, 2016)

Vascular ring anomaly – persistent right aortic arch (PRAA)

See under *Gastrointestinal conditions*

Dermatological conditions**Corns (on the foot pad)**

- In a study of 30 cases, 20 were greyhounds
- Males were predisposed (25/30 dogs)
- Corns affected the foot pads of the forelimbs more frequently than those of the hindlimbs
- One study suggested corns may be associated with papilloma virus infection (Guilliard *et al.*, 2010; Anis *et al.*, 2016)

Cutaneous and renal glomerular vasculopathy

- Uncommon condition with unknown aetiology, also known as Alabama rot
- Affected dogs have cutaneous ulcers on the extremities, thrombocytopenia and acute renal failure
- The condition is often fatal (Cowan *et al.*, 1997)

Cutaneous neoplasia

See under *Neoplastic conditions*

Pattern baldness (follicular dysplasia, bald thigh syndrome)

- Bilateral hair loss on the caudal thighs
- Cause unknown (Schoning & Cowan, 2000; Freeman, 2005)

Ventral comedone syndrome

- Common cosmetic syndrome
- Comedones on the ventral thorax (Burkett, 2000; Freeman, 2005)

Drug reactions**Thiobarbiturates**

- Reduced hepatic clearance results in an increased sensitivity to thiobarbiturates in Greyhounds
- Prolonged recovery times are seen
- Use of this drug is not recommended in Greyhounds (Robinson *et al.*, 1986; Freeman, 2005)

Gastrointestinal conditions**Vascular ring anomaly – persistent right aortic arch (PRAA)**

- Inheritance suspected in Greyhounds
- 4/52 cases were Greyhounds in one study
- Symptoms of regurgitation are seen at weaning (Gunby *et al.*, 2004; Krebs *et al.*, 2014)

Haematological/immunological conditions

Grey eosinophils

- Eosinophils with non-staining granules are common in Greyhounds (53% of 49 Greyhounds in one study)
- This may reflect different staining properties and is of no clinical significance (Iazbik & Couto, 2005)

Hyperhomocysteinaemia

- Frequent occurrence in Greyhounds
- May be associated with low serum folate and cobalamin (vitamin B₁₂) (Heilmann *et al.*, 2017)

Postoperative bleeding

- Noted frequently in retired Greyhounds. In one study 26% of retired racing Greyhounds were affected, compared to 0–2% in other breeds
- Primary or secondary haemostatic defects are not detected in these cases
- Bleeding may be related to altered fibrinolysis (Lara-Garcia *et al.*, 2008)

Infectious conditions

Babesiosis

- 46% of Greyhounds from 10 racing kennels in Florida tested positive in one study (Taboada *et al.*, 1992)

Musculoskeletal conditions

Avulsion of the tibial tuberosity

- Growth plate fracture
- Seen in dogs < 12 months old (Power, 1976; Skelly *et al.*, 1997)

Distal limb fractures and dislocations

- Common in this breed due to the counter-clockwise direction of racing, the repetitive nature of racing and the track conditions
- Include spontaneous tibial fracture, accessory carpal bone fracture, calcaneoquartal subluxation due to plantar tarsal ligament rupture, calcaneus fracture, central tarsal bone fracture and superficial digital flexor tendon luxation
- Many fractures include or are distal to the carpus and tarsus (Anderson *et al.*, 1995a, 1995b)

Medial displacement of the biceps brachii tendon

- Uncommon
- Occurs secondary to rupture of the transverse humeral ligament
- Biceps brachii or brachialis tendon avulsions have also been documented as acute injuries in racing Greyhounds (Boerno & Eaton-Wells, 1995; Schaaf *et al.*, 2009)

Osteosarcoma

See under *Neoplastic conditions*

Polyarthritis

- Erosive
- Affects animals aged 3–30 months
- Distal joints are affected
- Idiopathic (Huxtable & Davis, 1976; Ralphs & Beale, 2000)

Sacrocaudal fusion

- In a retrospective study of anatomical specimens and x-rays, 41% of 81 racing greyhounds were affected, compared to 15% of 91 non-Greyhound dogs
- Possibly due to selection of greyhounds for speed
- Clinical significance unknown (Oheida *et al.*, 2016)

Sesamoid disease

- Sesamoid fractures are common in racing greyhounds
- Believed to result from the action of the digital flexor tendon on the sesamoids at high impact (Harasen, 2009)

Neoplastic conditions

Cutaneous haemangiosarcoma

- Breed at increased risk of cutaneous haemangiosarcoma in a retrospective study of VMDB data, 1964–2002: OR 3.08 (95% CI 1.74–5.45) compared to all other dogs
- Mean age 9.7 years for cutaneous haemangiosarcoma in another study (Schultheiss, 2004; Villamil *et al.*, 2011)

Osteosarcoma

- Breed at increased risk in a USA-based retrospective case series of 179 dogs
- OR 17.3 (95% CI 9.3–32.1) compared to mixed-breed dogs

- Greyhounds were older at diagnosis than other predisposed breeds. Mean age 9.9 years
- Most common sites: proximal humerus, distal radius and proximal femur
- Also breed at high risk of malignant bone tumours in a study of Swedish insurance data, 1995–2002. Greyhounds had an overall incidence of 30 cases (95% CI 14–46) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Males over-represented
- Median age was 6.2 years in Greyhounds in this study; risk increases with increasing age (Egenvall *et al.*, 2007; Rosenberger *et al.*, 2007; Karlsson *et al.*, 2013)

Neurological conditions

Ischaemic stroke

- Breed at increased risk in a retrospective study
- OR 6.6 (95% CI 4.2–10.7) compared to all other breeds combined
- Hypertension is believed to be a contributing factor

(Kent *et al.*, 2014)

Meningoencephalitis

- Seen in Greyhounds in Ireland
- Age at presentation 5–18 months
- No infectious organism found
- Demonstration of an association with the DLA class 2 haplotype suggests an immunogenetic risk

(Shiel *et al.*, 2010a, 2014)

Polyneuropathy

- Seen in juvenile Greyhounds
- Uncommon
- Monogenic autosomal recessive inheritance; mutation identified

(Drögemüller *et al.*, 2008a)

Ocular conditions

Chronic superficial keratitis (pannus)

- Breed at increased risk in case series
- Age of onset 2 years
- Prevalence and severity increase at higher altitude

(Chavkin *et al.*, 1994; Cheng *et al.*, 2016)

Physiological conditions

Biochemical differences

- Several biochemical differences have been noted in normal Greyhounds compared to other breeds

- Reduced serum globulin concentration: IgM and IgA concentrations are significantly lower than other breeds, but IgG is unaffected
- Greyhounds have significantly higher creatinine levels than non-Greyhound dogs. This is believed to be due to their higher muscle mass
- In a retrospective study of 28 675 serum cobalamin tests, Greyhounds had an increased incidence of values below the lower limit of the reference interval (251 ng/l), OR 2.5 (95% CI 2.1–3.1)

(Drost *et al.*, 2006; Clemente *et al.*, 2010; Grützner *et al.*, 2012)

Blood group

- In a study of 206 Greyhounds, 13.1% were DEA-1.1 positive, which was significantly lower than the 66 non-Greyhound dogs, of which 60.6% were positive
- A majority of Greyhounds in the study met the criteria for universal donor

(Iazbik *et al.*, 2010)

Cardiac hypertrophy

- Greyhounds have a higher vertebral heart score (mean 10.5) and thicker left ventricular wall than other breeds
- Additionally, systemic arterial pressure is higher, stroke volume is higher and systemic vascular resistance is lower in racing Greyhounds than non-racing Greyhounds and other breeds

(Pape *et al.*, 1986; Marin *et al.*, 2007)

Haematological differences

- Several haematological differences have been noted in normal Greyhounds compared to other breeds
- Higher haematocrit (packed cell volume, PCV), haemoglobin and red cell counts
- Red blood cells are larger (higher mean corpuscular volume, MCV), and mean red cell haemoglobin concentration (MCHC) is higher
- Total and individual white blood cell counts (WCC) are lower
- Platelet counts are lower

(Shiel *et al.*, 2007b; Campora *et al.*, 2011)

Heart murmur

- Greyhounds commonly have a low-grade left-sided basilar systolic murmur of no clinical significance
- Slightly increased aortic velocities noted

(Fabrizio *et al.*, 2006)

NT-proBNP measurement

- NT-proBNP is a cardiac biomarker elevated in some dogs with congestive heart failure
- Healthy greyhounds have naturally higher levels than healthy non-Greyhound dogs; in one study 946 vs. 632 pmol/l ($p < 0.005$). 46% of greyhounds had levels > 1000 pmol/l in this study

(Couto *et al.*, 2015)**Polyodontia**

- Present in 36.4% of dogs in a study of 55 Greyhounds
- Usually the first premolar and usually in the upper arcade

(Dole & Spurgeon, 1998)

Thyroid hormones

- Total T_4 and free T_4 are significantly lower in healthy Greyhounds than in other breeds, leading to lower reference ranges being developed for these values in Greyhounds
- Mean circulating TSH is the same as for other breeds

(Shiel *et al.*, 2007a;

Chastain & Panciera, 2002)

Renal and urinary conditions**Renal arteriosclerosis**

- Seen with high frequency in young race-trained Greyhounds
- Higher blood pressure and sheer stresses may be contributing factors

(Bjotvedt *et al.*, 1988)**Reproductive conditions****Dystocia**

- Breed at increased risk in a study of Swedish insurance data (1995–2002) involving 3894 reimbursed claims for dystocia in an overall population of 195 931 bitches
- Incidence rate for greyhounds was 10.4 per 1000 DYAR, compared to 5.7 per 1000 DYAR for the overall population

(Bergström *et al.*, 2006)**GREYHOUND – SPANISH**See *Spanish Greyhound***GRIFFON BRUXELLOIS****Neurological conditions****Chiari malformation/syringomyelia**

- Breed at increased risk
- In a US prospective study of 84 American Kennel Club registered Griffon Bruxellois dogs, 65% had Chiari malformation, 52% had syringomyelia, 28% had neurologic deficits and 20% had neck pain

(Rusbridge *et al.*, 2009;Freeman *et al.*, 2014)**Ocular conditions****Cataract**

- Prevalence of primary cataract 5.41%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Prevalence decreased over the 40 years of the study
- Trend towards an increased number of males affected

(Gelatt & MacKay, 2005)

Vitreoretinopathy

- Vitreal liquefaction predisposes to retinal detachment in Griffon Bruxellois dogs

(Papaioannou & Dubielzig, 2013)

**HAMILTON HOUND
(HAMILTONSTÖVARE)****Endocrine conditions****Diabetes mellitus**

- Breed at increased risk in a retrospective review of Swedish insurance data, 1995–2004
- Incidence in the Hamilton Hound was 29 cases per 10 000 DYAR, compared to an overall incidence for all breeds of 13 cases per 10 000 DYAR
- Females at significantly increased risk

(Fall *et al.*, 2007)**Neurological conditions****Idiopathic epilepsy**

- Breed at increased risk of mortality due to epilepsy in a retrospective review of Swedish insurance data, 1995–2006
- Mortality rate in the Hamilton Hound was 24.9 cases per 10 000 DYAR, compared to an

overall incidence for all breeds of 11 cases per 10 000 DYAR

- Males at increased risk (1.4:1) across all breeds (Heske *et al.*, 2014)

HAVANESE

Cardiovascular conditions

Heart murmur

- One study suggests Havanese are predisposed to heart murmurs, though detail for the underlying cause is lacking (Starr *et al.*, 2007)

Dental conditions

Abnormal dentition

- May be inherited alongside cataracts, cardiac abnormalities and osteochondrodysplasia (Starr *et al.*, 2007)

Dermatological conditions

Sebaceous adenitis

- 35% of Havanese dogs were affected in a retrospective study of dogs presented at a dermatology referral service in the USA
- Seen in young adults (Frazer *et al.*, 2011; Bensignor & Guaguere, 2012)

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed at increased risk in a retrospective case study of 2400 cases from VMDB data (1980–2002)
- OR 64.9 (95% CI 8.9–234.3) compared to mixed-breed dogs
- Female:male ratio was 1.14:1 across all breeds
- Clinical signs are usually seen in young dogs < 1 year
- Extrahepatic shunts more common in the Havanese (Tobias & Rohrbach, 2003; Starr *et al.*, 2007)

Musculoskeletal conditions

Osteochondrodysplasia

- May be inherited alongside cataracts, cardiac abnormalities and abnormal dentition (Starr *et al.*, 2007)

Various skeletal defects

- Including bowed forelimbs, avascular necrosis of the femoral head (Legg–Calvé–Perthes

disease), patellar luxation: suggested as frequent findings in one study

(Starr *et al.*, 2007)

Ocular conditions

Cataract

- Prevalence of primary cataract 11.57%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- May be inherited alongside osteochondrodysplasia, cardiac abnormalities and abnormal dentition (Gelatt & MacKay, 2005; Starr *et al.*, 2007)

Vitreoretinopathy

- Vitreal liquefaction predisposes to retinal detachment in Havanese dogs (Papaioannou & Dubielzig, 2013)

Reproductive conditions

Dystocia

- Breed at increased risk in a retrospective study of Swedish insurance data (195 931 bitches, 1995–2002)
- Incidence in Havanese dogs was 12.2 cases per 1000 DYAR, compared to 5.7 cases per 1000 DYAR in the overall population (Bergström *et al.*, 2006)

HOKKAIDO

Ocular conditions

Collie eye anomaly

- Inherited; mutation identified
- In a study of 17 Hokkaido dogs, 5 were affected and 12 were heterozygous carriers (Mizukami *et al.*, 2012)

HOVAVART

Dermatological conditions

Sebaceous adenitis

- Breed suggested to be at increased risk in a number of studies (Linek *et al.*, 2005; Lortz *et al.*, 2010)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Heritability described as moderate to high in one study of Finnish Hovawarts

- High levels of autoantibodies to thyroglobulin found in 12.6% of this breed, predisposing to autoimmune lymphocytic thyroiditis (Ferm *et al.*, 2009; Bianchi *et al.*, 2015; Ahlgren & Uimari, 2016)

Gastrointestinal conditions

Intestinal adenocarcinoma

- In a Czech study of gastrointestinal biopsies, Hovawart dogs were found to be predisposed to intestinal adenocarcinoma ($p < 0.05$)
- Age at risk (across all affected breeds) was 7–8 years (Frgelecová *et al.*, 2013)

Haematological/immunological conditions

Selective IgA deficiency

- Breed at increased risk in case series
- 32% of 19 Hovawarts were found to have low serum IgA (< 0.07 g/l) in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease (Olsson *et al.*, 2014)

Musculoskeletal conditions

Bone tumour

See under *Neoplastic conditions*

Neoplastic conditions

Bone tumour

- Breed at high risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Hovawarts had an overall incidence of 28 cases (95% CI 12–43) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Males over-represented
- Median age was 6.9 years in Hovawarts, risk increases with increasing age (Egenvall *et al.*, 2007)

Intestinal adenocarcinoma

See under *Gastrointestinal conditions*

Neurological conditions

Degenerative myelopathy

- Breed at increased risk in a retrospective Swiss study of 50 cases
- OR 44.1 (95% CI 17.9–108.7; $p < 0.0001$) compared to the overall hospital population

- Mean age at diagnosis across all breeds was 9.1 years

(Kathmann *et al.*, 2006)

HUNGARIAN KUVASZ

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Breed with a higher prevalence of thyroid hormone autoantibodies (THAA)
- In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, Kuvasz had an OR of 2.18 ($p = 0.001$) of being affected compared to dogs of all other breeds
- Across the study, females were over-represented and the highest prevalence was in dogs 2–4 years old

(Nachreiner *et al.*, 2002)

Musculoskeletal conditions

Hip dysplasia

- Breed at increased risk
- OR 10.2 (95% CI 4.3–23.9) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Neutered male dogs predisposed (LaFond *et al.*, 2002)

Osteochondrosis – shoulder

- Breed at increased risk
- OR 29.1 (95% CI 12.8–66.3) compared to mixed breeds in a large retrospective case series based on cases recorded in the VMDB, 1986–1995
- Males predisposed
- Age of onset usually 4–7 months, but can be older (Rudd *et al.*, 1990; LaFond *et al.*, 2002)

Neurological conditions

Degenerative myelopathy

- Breed at increased risk in case series ($p < 0.05$)
- Mean age at diagnosis across breeds was 9.1 years

(Kathmann *et al.*, 2006)

Lumbosacral disease

- Breed at increased risk in case series
- Average age of onset across all breeds was 6.3 years

(Jaggy *et al.*, 1987)

HUNGARIAN PULI (HUNGARIAN WATER DOG)

Neoplastic conditions

Mammary neoplasia

- In a Hungarian study of 521 cases, Pulis were over-represented
- The number of mammary tumours was found to be highest in spring and autumn (Boldizsar *et al.*, 1992)

Ocular conditions

Cataract

- Prevalence of primary cataract 3.40%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005)

HUNGARIAN VIZSLA

Dermatological conditions

Atopic dermatitis (atopy)

- Breed at increased risk in Swiss and Hungarian case series compared to the general population
- Inheritance suspected (Tarpataki *et al.*, 2006; Picco *et al.*, 2008; Jaeger *et al.*, 2010)

Cutaneous neoplasia

See under *Neoplastic conditions*

Demodicosis

- Vizsla was described as a predisposed breed in a Swedish study (Holm, 2003)

Sebaceous adenitis

- Uncommon condition
- Affects young to middle-aged dogs
- Vizslas have been seen with a severe ulcerative form affecting the pinnae (Zur & Botero-Anug, 2011; Bensignor & Guaguere, 2012)

Urticaria

- Over-represented breed in a Swiss study of 24 cases
- Females over-represented across all breeds 2.4:1 (Rostaher *et al.*, 2017)

Haematological/immunological conditions

Haemophilia A

- Severe factor VIII deficiency is seen in Vizslas (Brooks, 1999)

Immune-mediated haemolytic anaemia (IMHA)

- Breed at increased risk in an Australian retrospective study of 110 cases: OR 10.0 (95% CI 1.3–74.7) (McAlees, 2010)

Musculoskeletal conditions

Hip dysplasia

- In a study of data from the Belgian Kennel club (2002–2006), prevalence in the Vizsla was 38%, compared to an overall prevalence of 20% (Coopman *et al.*, 2008)

Polymyopathy

- Uncommon condition. Immune-mediated aetiology suspected
- Dogs present with dysphagia, regurgitation and temporal muscle atrophy
- Mean age of onset 2.4 years
- Male dogs slightly over-represented (Massey *et al.*, 2013; Tauro *et al.*, 2015)

Neoplastic conditions

Lymphoma

- Breed at increased risk in case series: OR 2.7 (95% CI 2.0–3.5) compared with a US referral population (Villamil *et al.*, 2009)

Mast cell tumour (MCT)

- Breed at increased risk of cutaneous MCTs in a retrospective study of VMDB data, 1964–2002: OR 4.84 (95% CI 3.23–7.25) compared to all other dogs
- Risk higher in neutered dogs (Villamil *et al.*, 2011; Zink *et al.*, 2014)

Melanoma

- Breed at increased risk of cutaneous melanoma in a retrospective study of VMDB data, 1964–2002: OR 17.34 (95% CI 11.21–26.83) compared to all other dogs (Villamil *et al.*, 2011)

Neoplasia – overall

- Hungarian Wire-haired Vizsla had the third-highest proportional mortality from cancer

among pedigree breeds in the UK: 46.7% (95% CI 21.4–71.9)

(Adams *et al.*, 2010)

Neurological conditions

Idiopathic epilepsy

- Partial onset seizure disorder seen in the Vizsla
- Inherited as an autosomal recessive trait (Patterson *et al.*, 2003; Ekenstedt *et al.*, 2011)

Ocular conditions

Glaucoma – primary

- Breed at increased risk in a retrospective Swiss study of 129 cases (1995–2009)
- Mean age of onset in the Vizsla was 10.6 years
- M:F ratio 1:1.7

(Strom *et al.*, 2011a)

HUNGARIAN WATER DOG

See *Hungarian Puli*

HUSKY

See *Alaskan Husky*; *Siberian Husky*

IRISH RED AND WHITE SETTER

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaires relating to the 10 years prior to 2004
- Prevalence ratio for GDV mortality was 3.3 (95% CI 2.0–5.5; $p < 0.001$) compared to all other breeds combined

(Evans & Adams, 2010b)

Haematological/immunological conditions

Canine leucocyte adhesion deficiency (CLAD)

- Autosomal recessive inheritance
- 13% of Irish Red and White Setters tested in one study were found to be carriers of the mutation

(Debenham *et al.*, 2002; Foureman *et al.*, 2002)

IRISH SETTER (IRISH RED SETTER)

Cardiovascular conditions

Patent ductus arteriosus

- The Irish Setter is often suggested to be predisposed, but not all studies confirm this
- Females predisposed (Matic, 1988; Darke, 1989; Oliveira *et al.*, 2011)

Vascular ring anomaly – persistent right aortic arch (PRAA)

- The Irish Setter is often suggested to be predisposed, but not all studies confirm this

See also under *Gastrointestinal conditions*

(Matic, 1988; Krebs *et al.*, 2014)

Dermatological conditions

Anal furunculosis (perianal fistula)

- Breed at increased risk in a number of case series
- Middle-aged dogs are predisposed; mean age in one study was 5.2 years
- In the same study males were over-represented (2:1) and intact dogs of both sexes were at increased risk compared with their neutered counterparts

(Killingsworth *et al.*, 1988; Day & Weaver, 1992)

Atopic dermatitis (atopy)

- Breed at increased risk in a US series of 268 cases (1981–1984)
- Also reported as a breed at increased risk (OR 2.2 compared to the general hospital population) in a US study of 383 cases
- Some studies show no sex predilection, others show females predisposed (Scott, 1981; Schick & Fadok, 1986)

Cutaneous neoplasia

See under *Neoplastic conditions*

Endocrine conditions

Hypothyroidism

- Breed at increased risk in a North American study of 3206 cases (1964–1978)
- In another large US study the Irish Setter was in the top five breeds affected, with an incidence of 7.69% compared to an incidence of 1.54% in mixed breeds

- May occur at a younger age in breeds at risk (2–3 years)
(Nesbitt *et al.*, 1980; Milne & Hayes, 1981; Bellumori *et al.*, 2013)

Insulinoma

- Uncommon condition
- In a retrospective study of 73 cases, 7 were Irish Setters
- Middle-aged/older dogs affected. Average age across all breeds was 9.1 years in one study
(Kruth *et al.*, 1982; Caywood *et al.*, 1988)

Gastrointestinal conditions

Acquired megaesophagus

- Breed at increased risk in case series
- Megaesophagus can be idiopathic or secondary to peripheral neuropathies, myasthenia gravis, oesophagitis and chronic or recurrent gastric dilatation
(Gaynor *et al.*, 1997)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in many case series
- In one retrospective study of 1934 cases recorded in the VMDB (1980–1989), OR 3.5 (95% CI 2.4–5.0)
- Risk increases with higher thoracic depth: width ratios
- Risk increases by 33% with each year of age
- In a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaires relating to the 10 years prior to 2004, prevalence ratio for GDV morbidity was 12.6 (95% CI 9.31–17.06; $p < 0.0001$) compared to all other breeds combined, making the Irish Setter one of the four breeds with highest morbidity. Prevalence ratio for GDV mortality was also increased at 2.3 (1.5–3.4; $p < 0.001$) compared to all other breeds combined
(Glickman *et al.*, 1994; Schellenberg *et al.*, 1998; Evans & Adams, 2010b)

Gluten-sensitive enteropathy

- Autosomal recessive inheritance
- Clinical signs develop at 4–7 months
(Batt *et al.*, 1984; Garden *et al.*, 2000)

Vascular ring anomaly – persistent right aortic arch (PRAA)

- Clinical signs appear at the time of weaning
- Not all studies suggest a predisposition
- In one study of 44 cases, 5 were Irish Setters
(Matic, 1988; Buchanan, 2004)

Haematological/immunological conditions

Canine leucocyte adhesion deficiency (CLAD)

- Autosomal recessive inherited mutation
- Frequency of carriers of the mutation in Germany was estimated at 11%
(Kijas *et al.*, 2000; Pfeiffer & Brenig, 2005)

Haemophilia A

- Factor VIII deficiency
- X-linked recessive inheritance
(Brooks, 1999)

Musculoskeletal conditions

Irish Setter hypochondroplasia

- Rare condition
- Inherited as an autosomal recessive trait
(Hanssen *et al.*, 1998)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males may be predisposed
- OR 14.3 (95% CI 5.6–36.7) compared to mixed-breed dogs in a large retrospective case series based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002; Safra *et al.*, 2016)

Osteosarcoma

See under *Neoplastic conditions*

Neoplastic conditions

Insulinoma

See under *Endocrine conditions*

Melanoma

- Breed at increased risk of cutaneous melanoma in a retrospective study of VMDB data, 1964–2002: OR 2.23 (95% CI 1.47–3.38) compared to all other dogs
(Villamil *et al.*, 2011)

Osteosarcoma

- Breed at increased risk in a number of case series
- OR 3.5 (95% CI 2.5–4.8) compared to German Shepherd Dogs in a retrospective study of VMDB data, 1980–1994
- Across all breeds there was an increased risk with increasing age, plateauing at 10 years

- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes (Heyman *et al.*, 1992; Ru *et al.*, 1998)

Neurological conditions

Amblyopia and quadriplegia

- Rare condition – no recent reports in the literature
- Autosomal recessive inheritance
- Congenital paralysis and blindness with no visible ocular cause (amblyopia) (Palmer *et al.*, 1973; Sakai *et al.*, 1994)

Cerebellar hypoplasia

- Rare congenital condition. The Irish Setter is one of the few breeds in which it is reported
- Seen with lissencephaly in Irish Setters (Coates *et al.*, 2002; Heckler *et al.*, 2011)

Degenerative myelopathy

- Overall prevalence among all dogs in the VMDB (1990–1999) was 0.19%; prevalence for the Irish Setter over the same period was 0.68% (Coates *et al.*, 2007)

Globoid cell leucodystrophy (Krabbe disease)

- Rare condition. The Irish Setter is one of the few breeds in which it is reported
- Signs develop by 3 months of age (McGraw & Carmichael, 2006)

Idiopathic epilepsy

- Irish Setters have been described as being predisposed to both cluster seizures and refractory epilepsy (Raw & Gaskell, 1985; Forrester *et al.*, 1989)

Lissencephaly

- Rare congenital condition. The Irish Setter is one of the few breeds in which it is reported
- Seen with cerebellar hypoplasia in Irish Setters (Coates *et al.*, 2002; Heckler *et al.*, 2011)

Ocular conditions

Amblyopia (seen with quadriplegia)

See under *Neurological conditions*

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance
- Rod–cone dysplasia type I (rcd1)
- Early onset; most dogs are blind at 6 months old

- A second form of PRA, occurring at 4–5 years, may be seen in this breed (late-onset PRA or rcd4) (Aguirre & Rubin, 1975; Downs *et al.*, 2014a)

Renal and urinary conditions

Urethral sphincter mechanism incompetence

- Breed at increased risk in a case series (3/30 cases were Irish Setters in one study)
- Neutered females predisposed (Holt & Thrusfield, 1993; Martinoli *et al.*, 2014)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Idiopathic
- Males are more commonly affected, 2.4:1 in one study
- May be inherited by an autosomal dominant mode

(White, 1989; Burbidge, 1995)

IRISH TERRIER

Cardiovascular conditions

Aortic stenosis – subaortic stenosis (SAS)

- Irish terriers were in the top five breeds affected in a large USA-based retrospective case study of 90 004 dogs presented to a university teaching hospital. 3.13% of Irish Terriers were affected, compared to 0.15% of mixed breeds (Oliveira *et al.*, 2011; Bellumori *et al.*, 2013)

Dermatological conditions

Footpad hyperkeratosis

- Age of onset 6 months
- Autosomal recessive inheritance (Binder *et al.*, 2000; Drögemüller *et al.*, 2014)

Musculoskeletal conditions

Muscular dystrophy – X-linked

- Rare condition – occurs in Irish Terriers (Wentink *et al.*, 1972; Shelton, 1999)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Irish Terriers have been reported with tubular transport dysfunction (Picut & Lewis, 1987a)

Urolithiasis – cystine

- Sex-linked inheritance is suggested
- Young dogs are affected (2–5 years)

- Almost all cases are male
- In one study of 1731 calculi in West Germany, 10 Irish Terriers were included; all had cystine calculi and all were male

(Tsan *et al.*, 1972; Lewis & Morris, 1984; Hesse, 1990)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk in case series
- Common disease of older entire bitches (mean age 7.25 years)
- Most cases present within 12 weeks of oestrus (Chastain *et al.*, 1999; Smith, 2006)

IRISH WATER SPANIEL

Dermatological conditions

Pattern baldness (follicular dysplasia)

- Dominant mode of inheritance suggested
- Dietary factors may play a role
- Hair loss begins at 2–4 years of age, and occurs mainly on the flanks
- Hair loss is due to fracture of the hair in Irish Water Spaniels
- Eventually the whole of the trunk is involved (Cerundolo *et al.*, 2000)

IRISH WOLFHOUND

Cardiovascular conditions

Atrial fibrillation (AF)

- One study suggested 10.5% of the breed are affected
- Often associated with dilated cardiomyopathy (DCM), but in one study rhythm disturbances without evidence of DCM were found in 48/500 Irish Wolfhounds presented for examination at a cardiology practice
- Ventricular premature complexes may also be seen in this breed (Vollmar, 2000; Menaut *et al.*, 2005)

Dilated cardiomyopathy (DCM)

- Irish wolfhounds were in the top five breeds affected in a large USA-based retrospective case study of 90 004 dogs presented to a university teaching hospital. 6.08% of Irish Wolfhounds were affected, compared to 0.16% of mixed breeds

- Prevalence increases with increasing age
- Approximately twice as common in males as females
- Inheritance does not follow a simple pattern but involves multiple loci

(Vollmar, 2000; Philipp *et al.*, 2012; Bellumori *et al.*, 2013)

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed at increased risk in case series; digenic inheritance suspected
- Clinical signs usually seen in young dogs < 1 year
- Shunts are usually intrahepatic in the Irish Wolfhound
- Irish Wolfhound puppies may have a transient asymptomatic hyperammonaemia at 6–8 weeks in the absence of a portosystemic shunt. Ammonia levels normalize by 4 months (Hunt, 2004; Zandvliet & Rothuizen, 2007; van Steenbeek *et al.*, 2009)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaires relating to the 10 years prior to 2004
- Prevalence ratio for GDV mortality was 4.5 (95% CI 2.6–7.7; $p < 0.001$) compared to all other breeds combined

(Evans & Adams, 2010b)

Haematological/immunological conditions

Immunodeficiency syndrome

See under *Respiratory conditions* (Rhinitis/bronchopneumonia)

von Willebrand's disease (vWD)

- Irish Wolfhounds suffer from type I disease, but do not seem to exhibit a clinical bleeding tendency

(Clark & Parry, 1995; Brooks, 1999)

Musculoskeletal conditions

Bone tumour

See under *Neoplastic conditions*

Elbow dysplasia

- Breed at increased risk of fragmentation of the medial coronoid process: OR 93.4 (95% CI 39.6–220.3) compared to mixed-breed dogs in

a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002; Clements *et al.*, 2007)

Hip dysplasia

- Breed incidence of 10% in one study in Norway
- Incidence is affected by growth rate and exercise

(Krontveit & Moe, 2013)

Osteochondrosis – shoulder

- Males predisposed, 2.24:1 in one study
- Age of onset is usually 4–7 months, but can be older
- OR 47.1 (95% CI 26.4–84.0) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(Rudd *et al.*, 1990; LaFond *et al.*, 2002)

Osteochondrosis – stifle

- Age of onset usually 5–12 months, but may be earlier
- OR 523.5 (95% CI 165.6–1655.0) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.3 (95% CI 1.2–4.2) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Bone tumour

- Irish Wolfhounds had the highest risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Irish Wolfhounds had an overall incidence of 99 cases (95% CI 59–160) per 10 000 DYAR, compared to 5.5 cases per 10 000 DYAR for all breeds combined
- Males over-represented
- Mean age was 6.6 years in Irish Wolfhounds; the risk increases with increasing age

(Egenvall *et al.*, 2007; Anfinssen *et al.*, 2011;

Karlsson *et al.*, 2013)

Neurological conditions

Idiopathic epilepsy

- Suspected to be inherited as an autosomal recessive trait with incomplete penetrance
- One 2006 report suggested that the risk is increasing in Irish Wolfhounds
- In a retrospective study, idiopathic epilepsy was diagnosed in 18.3% of 796 Irish Wolfhounds
- Males were at increased risk (61.6% were male)
- First seizure occurred by 3 years of age in 73% of cases

(Casal *et al.*, 2006)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance
- Early onset. Blindness at 2–3 years of age

(Gould *et al.*, 1997)

Reproductive conditions

Dystocia

- Breed at increased risk in a retrospective study of Swedish insurance data (195 931 bitches 1995–2002)
- Incidence in Irish Wolfhounds was 16.5 cases per 1000 DYAR, compared to 5.7 cases per 1000 DYAR for the general population

(Bergström *et al.*, 2006)

Low fertility

- High incidence of low libido, small testicles and poor semen quality in Irish Wolfhounds

(Dahlbom *et al.*, 1997)

Respiratory conditions

Aspiration pneumonia

- High incidence noted in Irish Wolfhounds at one referral hospital in Australia
- The overall hospital incidence was 0.5%. Irish Wolfhounds had the highest breed incidence, with 9/25 dogs affected

(Greenwell & Brain, 2014)

Rhinitis/bronchopneumonia syndrome

- Uncommon
- Many but not all cases have been < 1 year old
- Seen in Europe and North America
- Underlying cause is unclear; cell-mediated immunodeficiency, low levels of IgA, and primary ciliary dyskinesia have all been investigated with no conclusion drawn

(Wilkinson, 1969; Leisewitz *et al.*, 1997;

Clercx *et al.*, 2003)

ITALIAN GREYHOUND

Dermatological conditions

Colour dilution alopecia

- Affects blue Italian Greyhounds
- Onset at 4–18 months, the condition starts at the dorsal midline

(Jae *et al.*, 2005)

Cutaneous neoplasia

See under *Neoplastic conditions*

Endocrine conditions

Multiple autoimmune diseases syndrome

- Resembles autoimmune polyendocrine syndrome type 2 (APS-2) in humans
- Seen in Italian Greyhounds in the USA
- Affects mainly glandular tissue, e.g. adrenals, thyroid, pancreas, gonads and skin
- Females are over-represented
- Typically occurs in middle age

(Pedersen *et al.*, 2012a, 2015)

Gastrointestinal conditions

Enamel hypoplasia

- Enamel is roughened and thinnish with brownish mottling
- Permanent and deciduous teeth uniformly affected
- Prevalence 14%, familial

(Gandolfi *et al.*, 2013)

Haematological/immunological conditions

Grey eosinophils

- Eosinophils with clear granules
- Detected in 34.5% of Italian Greyhounds in one study
- Some haematology analysers may not count or may underestimate grey eosinophils

(Giori *et al.*, 2011)

Neoplastic conditions

Cutaneous haemangiosarcoma

- Breed at increased risk in a case series of non-visceral haemangiosarcomas
- Mean age 9.7 years

(Schultheiss, 2004)

Ocular conditions

Cataract

- Prevalence of primary cataract 2.42%, compared to 1.61% in mixed-breed dogs, in a

retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Vitreous syneresis

- Breed at increased risk, familial
- May result in anterior displacement of the vitreous, and glaucoma
- May also predispose to retinal detachment

(Papaioannou & Dubielzig, 2013)

Physiological conditions

Alanine aminotransferase (ALT) levels

- Higher in Italian Greyhounds than other sight hounds

(Uhríková *et al.*, 2013)

ITALIAN MASTIFF (CANE CORSO)

Musculoskeletal conditions

Hip dysplasia

- In a French retrospective study over 14 years (1993–2006), of the 31 breeds represented by at least 100 affected dogs, the Italian Mastiff had the highest prevalence (59.7%)

(Genevois *et al.*, 2008)

Ocular conditions

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk in a retrospective review of 155 cases in Europe, 2001–2008
- 75.4% cases (across all breeds) occurred before 1 year of age
- Italian Mastiffs are at increased risk of bilateral occurrence, with a short interval between prolapses

(Mazzucchelli *et al.*, 2012)

ITALIAN SPINONE (SPINONE ITALIANO)

Neoplastic conditions

Neoplasia – overall

- Italian Spinone had the sixth-highest proportional mortality from cancer among pedigree breeds in the UK: 44.7% (95% CI 30.5–58.9)

(Adams *et al.*, 2010)

Neurological conditions

Idiopathic epilepsy

- Breed at increased risk in case series
- Prevalence of 5.3% (95% CI 4.03–6.57%) among Italian Spinone registered with the UK Kennel Club, compared to an estimated prevalence of 0.6% in the UK general dog population
- Can be severe in this breed. Early and aggressive treatment may improve survival (Kearsley-Fleet *et al.*, 2013; De Risio *et al.*, 2015)

Spinocerebellar ataxia (hereditary ataxia)

- Autosomal recessive inheritance
- A GAA repeat expansion in intron 35 ITPR1 identified in Italian Spinone in the UK
- Progressive ataxia, tremors and hypermetria seen from 4 months, usually leading to death or euthanasia by 1 year of age due to an inability to walk

(Forman *et al.*, 2015)

JACK RUSSELL TERRIER

Behavioural conditions

Aggression

- Breed scoring highly for aggression towards both people and other dogs in a study based on Canine Behavioural Assessment and Research Questionnaires (C-BARQ) completed by owners in the USA

(Duffy *et al.*, 2008)

Dermatological conditions

Black hair follicular dysplasia

- Thought to be inherited as an autosomal recessive trait
- Changes usually seen by 4 weeks of age (Schmutz *et al.*, 1998)

Demodicosis

- Jack Russell Terriers are at increased risk of juvenile-onset, generalized demodicosis compared to a control hospital population: OR 1.6 (95% CI 1.019–2.4) based on a large retrospective case-control study of a US practice database
- Juvenile onset was defined as < 18 months old
- Other risk factors associated with demodicosis included pyoderma, short hair and concurrent coccidiosis

(Plant *et al.*, 2011)

Dermatophytosis (ringworm)

- No sex predisposition
- Dogs < 1 year old predisposed (Sparkes *et al.*, 1993)

Ichthyosis

- Autosomal recessive inheritance
- Causative mutation identified (Credille *et al.*, 2009; Hartley *et al.*, 2012)

Vasculitis

- Alopecia and ulcers of the extremities and bony prominences resembling dermatomyositis of Collies and Shetland Sheepdogs (Parker & Foster, 1996)

Drug reactions

Vaccine-associated adverse effect

- In a large retrospective study in the USA (2002–2003) Jack Russell Terriers had a mean rate of 54.4 adverse reactions per 10 000 dogs vaccinated, compared to an overall rate of 38.2 reactions per 10 000 dogs vaccinated
- Risk seemed to decrease with increasing body weight

(Moore *et al.*, 2005)

Endocrine conditions

Hyperadrenocorticism (Cushing's syndrome)

- In a retrospective study of data from UK first-opinion practice (VetCompass) Jack Russell Terriers had an OR of 1.6 (95% CI 1.0–2.5) compared to mixed breeds
- Dogs > 12 years were at higher risk (OR 5.7) compared to dogs aged 6–8.9 years (O'Neill *et al.*, 2016d)

Gastrointestinal conditions

Chronic hepatitis

- In a UK-based study of 200 post-mortem samples of liver, Jack Russell Terriers were found to have an increased risk ratio of 3.6 (95% CI 1.3–10.1) (Watson *et al.*, 2010)

Congenital portosystemic shunt

- Breed at increased risk in case series in the UK and Australia
- Clinical signs usually seen in young dogs < 1 year
- Usually extrahepatic in Jack Russell Terriers (Hunt, 2004; Adam *et al.*, 2012)

Haematological/immunological conditions

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance (Brooks, 1999)

Severe combined immunodeficiency

- Inherited as an autosomal recessive trait (Meek *et al.*, 2001; Bell *et al.*, 2002)

Musculoskeletal conditions

Congenital myasthenia gravis

See under *Neurological conditions*

Myokymia and neuromyotonia

- Hereditary
- Age of clinical onset reported as 2 months to 3 years
- Myokymia (rippling muscle contractions), neuromyotonia (muscle stiffness) and hyperthermia are seen in affected dogs
- May be associated with hereditary ataxia
- Multiple forms may exist (Hartley & Palmer, 1973; Vanhaesebrouck *et al.*, 2010; Bhatti *et al.*, 2011; Gilliam *et al.*, 2014)

Neurological conditions

Congenital deafness

- Prevalence in Jack Russell Terriers 3.57% (unilateral) and 0.5% (bilateral) in one study
- Significant association with white coat colour (Comito *et al.*, 2012)

Congenital myasthenia gravis

- Autosomal recessive inheritance suspected
- Rare
- Age of clinical onset: 2–6 months (Wallace & Palmer, 1984; Wilkes *et al.*, 1987; Shelton, 1999)

Paroxysmal dyskinesia

- Breed at increased risk in one study
- Median age at onset of 23 Jack Russell Terriers was 4 years and 8 months; 15 of the 23 were male (Lowrie & Garosi, 2016)

Spinocerebellar ataxia (hereditary ataxia)

- Age of clinical onset reported as 2 months to 3 years
- Symptoms include ataxia, hypermetria and seizures

- Myokymia, neuromyotonia and hyperthermia may be associated with hereditary ataxia (see under *Musculoskeletal conditions*)

- Multiple forms may exist (Hartley & Palmer, 1973; Wessmann *et al.*, 2004; Vanhaesebrouck *et al.*, 2010; Bhatti *et al.*, 2011; Gilliam *et al.*, 2014)

Steroid-responsive meningitis–arteritis (SRMA)

- Breed at increased risk in case series
- OR 6.91 (95% CI 2.2–22.66; $p = 0.001$) in a retrospective UK-based case–control study
- Age of onset 3 month to 9 years (Rose *et al.*, 2014)

Ocular conditions

Cataract

- Prevalence of primary cataract 2.26%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Oberbauer *et al.*, 2008)

Glaucoma – primary

- Breed at increased risk compared to the overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
- Prevalence was 1.37% in Jack Russell Terriers for the years 1994–2002, compared to an overall prevalence in all dogs of 0.89% over the same period
- Prevalence was 1.31% in Jack Russell Terriers for the years 1984–1993, compared to an overall prevalence in all dogs of 0.76% over the same period (Gelatt & MacKay, 2004a)

Glaucoma – secondary

- Breed at increased risk of glaucoma secondary to lens luxation (Strom *et al.*, 2011b)

Lens luxation – primary

- Inherited; mutation identified
- Age of onset: 3–6 year (Oberbauer *et al.*, 2008; Gould *et al.*, 2011)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk. OR 2.33 (95% CI 2.05–2.65; $p < 0.0001$) compared to the

national insurance database in a retrospective study of 14 008 uroliths from dogs in the UK (1997–2006)

- Males predisposed

(Roe *et al.*, 2012)

Urolithiasis – cysteine

- Breed at increased risk. OR 2.32 (95% CI 1.59–3.6; $p < 0.0001$) compared to the national insurance database in a retrospective study of 14 008 uroliths from dogs in the UK (1997–2006)

(Roe *et al.*, 2012)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk. OR 1.84 (95% CI 1.64–2.07; $p < 0.0001$) compared to the national insurance database in a retrospective study of 14 008 uroliths from dogs in the UK (1997–2006)
- Females predisposed

(Roe *et al.*, 2012)

JÄMTHUND

See *Swedish Elkhound*

JAPANESE AKITA INU

Behavioural conditions

Aggression

- Breed scoring highly for aggression towards other dogs in a study based on Canine Behavioural Assessment and Research Questionnaires (C-BARQ) completed by owners in the USA

(Duffy *et al.*, 2008)

Dermatological conditions

Calcinosis cutis

- Breed at increased risk compared to the general population in a USA-based retrospective report of 46 cases
- Median age at diagnosis 7.6 years (across all breeds)
- Male dogs over-represented (across all breeds)
- 78% of all cases were associated with excess exogenous or endogenous steroids

(Doerr *et al.*, 2013)

Sebaceous adenitis

- Autosomal recessive mode of inheritance suspected
- Young to middle-aged dogs affected
- No apparent sex predisposition
- Generalized, greasy changes seen in Akitas
- Some animals may show systemic signs
- Mean age at diagnosis 4.8 years
- 10/104 cases were Akitas in a Swedish study (Reichler *et al.*, 2001; Tevell *et al.*, 2008; Szczepanik *et al.*, 2012)

Pemphigus foliaceus

- Uncommon
- Breed at increased risk in case series
- OR 37.8 (95% CI 13.1–98.8; $p = 0.0001$) in a histopathological study of 50 cases in the USA, 1986–1991
- No sex predisposition noted
- Mean age of onset 4.2 years (Kuhl *et al.*, 1994)

Uveodermatologic syndrome

(Vogt–Koyanagi–Harada-like syndrome)

- Akitas account for 80% of diagnosed cases

See also under *Ocular conditions*

(Angles *et al.*, 2005)

Gastrointestinal conditions

Exocrine pancreatic insufficiency (EPI)

- Breed at increased risk in a retrospective study of 635 dogs in the USA (Williams & Minnich, 1990)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased in case series (Glickman *et al.*, 2000a, 2000b)

Haematological/immunological conditions

Haemophilia A

- Factor VIII deficiency
- X-linked recessive inheritance (Brooks, 1999)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Breed at increased risk in a case series
- Neutered dogs predisposed
- Heavier dogs may be predisposed (Duval *et al.*, 1999)

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- OR 1.8 (95% CI 1.2–2.7) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Patellar luxation

- OR 6.7 (95% CI 4–11.3) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Males predisposed (1.8:1)
- Mostly affects dogs < 2 years of age (LaFond *et al.*, 2002; Gibbons *et al.*, 2006)

Polyarthritis

- Seen in young Akitas < 1 year old
- Affected dogs have fever, joint pain, lymphadenopathy and signs of meningeal pain
- The condition seems resistant to immunosuppressive therapy (Dougherty & Center, 1990)

Neoplastic conditions

Canine acanthomatous ameloblastoma

- Oral tumour most common in the rostral maxilla
- Breed at increased risk in a case series
- Neutered dogs of both sexes at increased risk
- Mean age at presentation 8.8 years (Fiani *et al.*, 2011)

Neurological conditions

Acquired myasthenia gravis

- Breed at increased risk compared to mixed-breed dogs
- Intact male dogs and dogs < 1 year old had some protection from risk
- Two age peaks, at 2–4 years and 9–12 years, have been observed (Shelton *et al.*, 1997)

Ocular conditions

Glaucoma – primary

- Breed at increased risk compared to the overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
- Prevalence was 1.39% in Akitas for the years 1994–2002, compared to an overall prevalence in all dogs of 0.89% over the same period

- Prevalence was 1.11% in Akitas for the years 1984–1993, compared to an overall prevalence in all dogs of 0.76% over the same period (Gelatt & MacKay, 2004a)

Glaucoma – secondary

- Akitas have an increased risk of glaucoma secondary to uveitis which occurs with uveodermatologic syndrome (Gelatt & MacKay, 2004b)

Multiple ocular defects

- Congenital defects; autosomal recessive inheritance suspected
- May include microphthalmia, cataract, posterior lenticonus and retinal dysplasia (Laratta *et al.*, 1985)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance
- Night blindness is present at 1–3 years, complete blindness at 3–5 years (O'Toole & Roberts, 1984)

Uveodermatologic syndrome

(Vogt–Koyanagi–Harada-like syndrome)

- Breed at increased risk
- Ocular signs include uveitis, which can progress to glaucoma, retinal detachment and blindness
- Young adults (1.5–4 years) affected (Angles *et al.*, 2005; Carter *et al.*, 2005)

Physiological conditions

Pseudohyperkalaemia

- Red blood cells in some Akitas contain more potassium than in other breeds; haemolysis may therefore cause false findings of hyperkalaemia (Degen, 1987; Conrado *et al.*, 2014)

Red cell microcytosis

- Red blood cells may be small in Akitas without clinical disease
- Mean corpuscular volume (MCV) may be as low as 55–64 fl (normal 85–95) (Degen, 1987; Fujise *et al.*, 1997; Gookin *et al.*, 1998)

Serum cobalamin levels

- In a retrospective study of 28 675 serum cobalamin tests, Akitas had an increased incidence of values below the lower limit of the reference interval (251 ng/l) OR 2.8 (95% CI 1.9–4.1) (Grützner *et al.*, 2012)

JAPANESE CHIN

Musculoskeletal conditions

Patellar luxation

- OR 4.8 (95% CI 1.1–20.5) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Neurological conditions

Lysosomal storage disease – GM₂ gangliosidosis

- Rare
- Mutation identified in Japanese Chins (Sanders *et al.*, 2013)

Ocular conditions

Cataract

- Breed at increased risk in case series
- In a retrospective study of 561 small-breed dogs in Korea, OR 13.1 (95% CI 6.0–28.7) compared to the general hospital population. Female dogs at significantly lower risk than male dogs in the Japanese Chin (OR 0.1)
- Prevalence of primary cataract 4.89%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003). Male predisposition was replicated in this study (Gelatt & MacKay, 2005; Park *et al.*, 2009)

JAPANESE SHIBA INU

Behavioural conditions

Aggression

- Shiba Inu was reported with high aggression to dogs and snapping at children in Japan
- Polymorphism c.471 T>C associated with some types of aggressive behaviour in the Shiba Inu in the USA (Takeuchi & Mori, 2006; Takeuchi *et al.*, 2009)

Tail-chasing

- Repetitive behaviour and pathological behaviour
- Dogs originating from pet stores predisposed
- Shiba Inu was significantly predisposed ($p < 0.001$) compared with other breeds in Japan, with 28% of Shiba Inu affected (Goto *et al.*, 2012)

Musculoskeletal conditions

Patellar luxation

- 6.2% of US Shiba Inu affected
- Mainly medial luxation observed, often bilateral (Orthopedic Foundation for Animals, 2015)

Neoplastic conditions

Oral epulis

- Shiba Inu accounted for 12.1% of a fibromatous epulis caseload in Japan (Yoshida *et al.*, 1999)

Neurological conditions

Intervertebral disc disease (IVDD)

- Shiba Inu accounted for 2.3% of a cervical IVDD caseload and 2.4% of a thoracolumbar IVDD caseload at a Japanese referral hospital (Itoh *et al.*, 2008)

Lysosomal storage disease – GM₁ gangliosidosis

- Inherited as autosomal recessive traits
- Described in Shiba Inu in Japan
- Age at onset of 5–6 months and survival period of 14–15 months (Yamato *et al.*, 2003)

Ocular conditions

Glaucoma

- 29.0% of Shiba Inu referred for eye or neurological conditions in Japan were affected (Kato *et al.*, 2006)

Physiological conditions

Litter size

- Smaller litters associated with older bitches and smaller breeds
- Shiba Inu had the eighth smallest mean litter size (3.3 puppies) among registered breeds in Norway (Borge *et al.*, 2011)

JAPANESE TOSA

Infectious conditions

Babesiosis

- High incidence reported in this breed in Japan (unknown if this is due to a true breed susceptibility or increased risk of exposure via ticks or possibly bites)

- Often subclinical
- Babesiosis is tick-borne but possibly also transmitted by fighting
(Miyama *et al.*, 2005)

JINDO

See *Korean Jindo*

KEESHOND (WOLFSPITZ)

Cardiovascular conditions

Patent ductus arteriosus

- Breed at increased risk
- Females predisposed
- Mode of inheritance is polygenic
(Matic, 1988; Buchanan, 2001; Oliveira *et al.*, 2011)

Tetralogy of Fallot

- Congenital
- One of a group of defects known as conotruncal defects
(Matic, 1988; Patterson, 1989; Werner *et al.*, 2005)

Ventricular septal defect

- Congenital
(Matic, 1988; Patterson, 1989; Werner *et al.*, 2005)

Dermatological conditions

Alopecia X

- Males may be predisposed
- Clinical signs seen at any age but often 1–2 years
(Lothrup, 1988; Chastain & Panciera, 2004; Mausberg *et al.*, 2007)

Endocrine conditions

Diabetes mellitus

- Autosomal recessive inheritance
- Insulin-dependent diabetes mellitus due to hypoplasia of the cells of the Islet of Langerhans
- Onset commonly before 6 months of age
- OR 2.45 (95% CI 1.47–4.12) compared to mixed-breed dogs in a retrospective report of 6860 dogs with diabetes mellitus from the VMDB, 1970–1999. Females predisposed: OR 1.37
(Kramer *et al.*, 1980, 1988; Guptill *et al.*, 2003)

Hypothyroidism

- Keeshonden were in the top five breeds affected in a large USA-based retrospective case study of 90 004 dogs presented to a university teaching hospital. 6.63% of Keeshonden were affected, compared to 1.54% of mixed breeds
(Bellumori *et al.*, 2013)

Primary hyperparathyroidism

- Keeshonden represent at least 25% of dogs diagnosed with primary hyperparathyroidism
- Autosomal dominant inheritance
- Older dogs affected. Mean age 9.8 years
(Berger & Feldman, 1987; Chastain & Panciera, 2007; Goldstein *et al.*, 2007)

Musculoskeletal conditions

Hip dysplasia

- OR 1.8 (95% CI 1.1–2.8) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002)

Patellar luxation

- OR 4.4 (95% CI 2.5–8.0) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002)

Neoplastic conditions

Parathyroid tumours (resulting in primary hyperparathyroidism)

See under *Endocrine conditions*

Neurological conditions

Idiopathic epilepsy

- Inherited
- Bias towards males reported
- Age of onset 6 months to 6 years
(Cunningham & Farnbach, 1988; Hall & Wallace, 1996; Ekenstedt & Oberbauer, 2013)

Ocular conditions

Spontaneous chronic corneal epithelial defects (refractory corneal ulceration, indolent ulcers)

- Persistent corneal erosions with no obvious cause
- Breed at increased risk in case series; 11.1% of 45 dogs were Keeshonden in one US study
(Murphy *et al.*, 2001)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- OR 4.6 (95% CI 3.5–5.8) compared to mixed-breed dogs in a USA-based retrospective study of 10 673 calcium oxalate cases
- Males at increased risk ($p < 0.001$), representing 69% of submissions across all breeds
- In a Canadian case series of urolith submissions, 1998–2001, the OR was 3.00 (95% CI 1.26–7.17) compared to mixed-breed dogs (Ling *et al.*, 2003; Low *et al.*, 2010)

KELPIE

See *Australian Kelpie*

KERRY BLUE TERRIER**Dermatological conditions**

Cutaneous neoplasia

See under *Neoplastic conditions*

Footpad hyperkeratosis

- Familial
- Develops at 4–6 months (Duclos, 2013)

Spiculosis

- Rare
- Affects young, entire male Kerry Blues (McKeever *et al.*, 1992)

Gastrointestinal conditions

Hypodontia

- Condition identified in Kerry Blues in Russia
- Reduced numbers of premolars
- Litter sizes were smaller in those with affected pups
- Inheritance not established (Aksenovich *et al.*, 2004)

Haematological/immunological conditions

Factor XI deficiency

- Inherited as an autosomal trait with incomplete penetrance
- Affected dogs have mild bleeding tendencies. Some are asymptomatic (Knowler *et al.*, 1994; Brooks, 1999; Tcherneva & Giger, 2007)

Neoplastic conditions

Pilomatrixoma

- Uncommon (1% of all cutaneous neoplasms in the dog)
- Breed at increased risk in case series
- Mean age 6.6 years (Scott & Anderson, 1991; Toma & Noli, 2005)

Neurological conditions

Cerebellar degeneration

- Rare
- Signs seen at 9–16 weeks of age and progress until unable to stand at 10–12 months (de Lahunta & Averill, 1976; Montgomery, 1982)

Canine multiple system degeneration

- Rare
- Presents as a movement disorder
- Autosomal recessive inheritance (O'Brien *et al.*, 2005)

Ocular conditions

Cataract

- Prevalence of primary cataract 2.58%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005)

KING CHARLES SPANIEL (ENGLISH TOY SPANIEL)**Cardiovascular conditions**

Mitral valve disease

- Breed at increased risk: OR 5.36 (95% CI 3.36–8.55) in a UK study (Thrusfield *et al.*, 1985)

Neurological conditions

Primary brain tumour – oligodendroglioma

- Breed at significantly increased risk compared to the general population in a study of post-mortem results of 435 dogs with intracranial neoplasia (Song *et al.*, 2013)

Chiari malformation/syringomyelia

- Breed at increased risk in a UK-based retrospective study of VetCompass data, 2009–2014 (Sanchis-Mora *et al.*, 2016)

Ocular conditions

Corneal ulceration (ulcerative keratitis)

- In a UK-based study of first-opinion practice the prevalence in the King Charles was 2.22% (95% CI 1.11–3.93), compared to an overall prevalence of 0.8% (95% CI 0.75–0.86)

(O'Neill *et al.*, 2017b)

KLEE KAI

See *Alaskan Klee Kai*

KOOIKER DOG (KOOIKERHONDJE) (SMALL DUTCH WATERFOWL DOG)

Haematological/immunological conditions

von Willebrand's disease (vWD)

- Inherited as an autosomal recessive trait
- Type III disease is seen in this breed
- Identification of the mutations involved and development of a genetic test have enabled elimination of the mutation from breeding stock approved by the Dutch breeding club

(Slappendel *et al.*, 1998;

van Oost *et al.*, 2004)

Musculoskeletal conditions

Patellar luxation

- Patellar luxation was present in 24% of a cohort of 842 screened Dutch Kooikers
- Males and females similarly affected

(Wangdee *et al.*, 2014)

Neurological conditions

Hereditary necrotizing myelopathy

- Clinical signs begin at 3–12 months age
- No sex predisposition

(Mandigers *et al.*, 1993)

Renal and urinary conditions

Renal dysplasia

- Rare condition
- Reported in 3 young adult Dutch Kookier dogs

(Schulze *et al.*, 1998)

KOREAN JINDO

Physiological conditions

Pseudohyperkalaemia

- Korean Jindos may have red blood cells containing high levels of potassium, leading to an increased risk of pseudohyperkalaemia
- In a study of 35 Korean Jindos in Korea, 25.7% were found to be affected

(Yamato *et al.*, 1999)

KROMFOHRLÄNDER

Dermatological conditions

Footpad hyperkeratosis

- Autosomal recessive inheritance
- Age of onset 6 months

(Drögemüller *et al.*, 2014)

KUVASZ

See *Hungarian Kuvasz*

LABRADOR RETRIEVER

Behavioural conditions

Attraction to water

- Labrador Retrievers were found to be more attracted to water than to social stimuli (humans or other dogs) in one study

(Tavares *et al.*, 2015)

Cardiovascular conditions

Accessory pathway arrhythmia

- Presence of abnormal pathways of conduction connecting the atria to the ventricles
- Result most commonly in atrial fibrillation or orthodromic reciprocating tachycardia
- Males predisposed

(Wright *et al.*, 1996; Finster *et al.*, 2008; Santilli, 2010)

Atrioventricular (AV) block

- Breed at increased risk in case series
- Labrador Retrievers have a high risk of both second- and third-degree AV block

(Schrope & Kelch, 2006; Wess *et al.*, 2006)

Pericardial effusion

- Male predisposition suggested
- Studies have demonstrated a predisposition to both idiopathic pericardial effusion and primary cardiac haemangiosarcoma (MacDonald *et al.*, 2009; Kumar *et al.*, 2011; Yamamoto *et al.*, 2013)

Tricuspid valve dysplasia

- Breed at increased risk in several case series
- OR 13.81 (95% CI 3.89–49.05) compared to the general hospital population, in an Australian retrospective study of 10 dogs with tricuspid valve dysplasia seen between 2001 and 2012
- In an Italian retrospective case series of 976 dogs with congenital heart defects, Labrador Retrievers had an OR of 11.13 ($p < 0.0001$) compared to the general hospital population (Famula *et al.*, 2002; Oliveira *et al.*, 2011; Aherne & Beijerink, 2013)

Dermatological conditions**Atopic dermatitis (atopy)**

- In an Australian retrospective case series, Labradors Retrievers had an OR of 2.2 (95% CI 1.65–3.0) compared to the overall hospital population
- Breed at increased risk in a US series of 268 cases (1981–1984)
- Breed at increased risk in a study of Munich cases: OR 3.7 (95% CI 1.6–8.5) compared to the general hospital population
- Birth in autumn or summer is a predisposing factor
- Dogs between 1 and 2 years of age have the highest probability of an insurance claim for atopy
- Some studies show no sex predilection, others show females predisposed (Shaw *et al.*, 2004; Nødtvedt *et al.*, 2006; Jaeger *et al.*, 2010; Mazrier *et al.*, 2016)

Blastomycosis

See under *Infectious conditions*

Calcinosis cutis

- Breed at increased risk in case series
- Males over-represented
- Occurs most commonly secondary to excess endogenous or exogenous corticosteroid (Doerr *et al.*, 2013)

Cutaneous neoplasia

See under *Neoplastic conditions*

Eosinophilic dermatitis and oedema

- Rare
- Hypersensitivity reaction suspected
- 4/9 dogs in a case series were Labrador Retrievers (Holm *et al.*, 1999; Mauldin *et al.*, 2006)

Hereditary nasal parakeratosis in Labrador Retrievers

- Autosomal recessive inheritance
- Affects Labradors and their crosses
- Age of onset 6–12 months (Pagé *et al.*, 2003; Jagannathan *et al.*, 2013)

Mural folliculitis and parakeratosis

- Rare condition identified in Labrador Retrievers
- Lesions can be diffuse, starting as comedones and progressing to large dense crusted plaques
- An immune-mediated condition is suspected (Hargis *et al.*, 2013)

Pemphigus foliaceus

- Predisposed breed in a number of studies
- Males over-represented (Gomez *et al.*, 2004; Mueller *et al.*, 2006)

Endocrine conditions**Diabetes mellitus**

- Breed at increased risk in case series
- Neutered dogs of both sexes at increased risk
- More frequently diagnosed in the winter months (Chastain *et al.*, 2001; Davison *et al.*, 2005)

Hypothyroidism

- Breed at increased risk
- Predisposition to hyperlipidaemia, atherosclerosis and neurological signs in affected dogs (Vitale & Olby, 2007)

Gastrointestinal conditions**Acquired megaesophagus**

- Breed at increased risk
- May be secondary to myopathy (see under *Musculoskeletal conditions*) (Bedu *et al.*, 2012)

Chronic hepatitis

- Breed at increased risk in a UK-based study of 551 cases: OR 2.0 (95% CI 1.6–2.5; $p < 0.001$) compared to a control population. Median age was 8 years 3 months, and females were over-represented across the whole study
- Additionally at increased risk of copper-associated hepatitis due to an inherited defect in copper storage. Middle-aged females predisposed. Higher levels of dietary copper can increase the risk

(Hoffmann *et al.*, 2006; Shih *et al.*, 2007; Bexfield *et al.*, 2012a; Fietan *et al.*, 2014, 2015)

Congenital portosystemic shunt

- Breed at increased risk of intrahepatic shunts in several case series. 28/100 cases were Labrador Retrievers in one retrospective case series
- Clinical signs usually seen in young dogs < 1 year (Tobias & Rohrbach, 2003; Adam *et al.*, 2012; Weisse *et al.*, 2014)

Spirocercosis

- Oesophageal infection with *Spirocerca lupi*
- Breed at increased risk in a case series in Israel
- Median age was 5 years
- May be associated with oesophageal sarcomas (Mazaki-Tovi *et al.*, 2002; Ranen *et al.*, 2008)

Haematological/immunological conditions**Haemophilia A**

- Factor VIII deficiency
- X-linked recessive inheritance (Brooks, 1999; Aslanian *et al.*, 2014)

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance (Mischke *et al.*, 2011)

Selective IgA deficiency

- Breed at increased risk in case series
- 12% of 141 Labrador Retrievers were found to have low serum IgA (< 0.07 g/l) in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease (Olsson *et al.*, 2014)

Infectious conditions**Blastomycosis**

- Breed at increased risk in case series. 29.6% of 125 cases were Labrador Retrievers in one retrospective study in Minnesota
- Young (2–4 years) entire males predisposed
- Geographic distribution: mainly North America (Mississippi, Missouri, Ohio River valleys, mid-Atlantic States, Quebec, Manitoba and Ontario)
- Proximity to a body of water is a risk factor in endemic areas (Rudmann *et al.*, 1992; Kerl, 2003; Crews *et al.*, 2008)

Borreliosis (Lyme disease)

- Breed at increased risk of Lyme nephritis
- See also under *Renal and urinary conditions* (Littman, 2013)

Infectious skin diseases

See under *Dermatological conditions*

Spirocercosis

See under *Gastrointestinal conditions*

Metabolic conditions**Overweight/obesity**

- In a study of 21 754 dogs carried out in 1995 in the USA, Labrador Retrievers had an OR of 1.6 compared to all other dogs
- Across all breeds, dogs aged 6–10 years and neutered dogs (OR 1.5) were at greatest risk
- A genetic basis is believed to exist in the Labrador Retriever (Lund *et al.*, 2006; Raffan *et al.*, 2016; Mankowska *et al.*, 2017)

Musculoskeletal conditions**Cranial cruciate ligament (CCL) disease**

- Breed at increased risk in case series. 16% of 426 dogs with cranial cruciate disease were Labrador Retrievers in one retrospective study
- OR 2.56 (95% CI 2.47–2.64) compared to all dogs in a separate retrospective study of data from the VMDB, 1964–2003. In this study neutered males, neutered females and dogs > 4 years old of all breeds combined were at increased risk (Comerford *et al.*, 2006; Witsberger *et al.*, 2008; Guthrie *et al.*, 2012)

Discospondylitis

- Labrador Retrievers were at higher risk in a study of 513 cases in North America: OR 1.5 (95% CI 1.0–2.2) compared to mixed-breed dogs
- Higher risk with increasing age
- Males twice as likely to be affected as females: OR 2.0 (95% CI 1.7–2.4)

(Burkert *et al.*, 2005)

Elbow dysplasia

- Breed at increased risk
- OR 20.5 (95% CI 13.9–30.3) for fragmented coronoid process, OR 109.4 (95% CI 54.4–219.9) for osteochondrosis and OR 8.5 (95% CI 6.5–11.1) for ununited anconeal process, all compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Male dogs were 1.7 times more likely to have osteoarthritis in affected joints in one study
- Polygenic inheritance suspected (LaFond *et al.*, 2002; Woolliams *et al.*, 2011; Lavrijsen *et al.*, 2012)

Hip dysplasia

- Breed at increased risk in case series
- OR 2.34 (95% CI 2.28–2.41; $p < 0.001$) compared to all dogs in a retrospective study of data from the VMDB, 1964–2003. In this study, neutered males and dogs < 4 years old of all breeds combined were at increased risk
- One study of 1018 Labrador Retrievers estimated the prevalence in the breed as 12.6% (Witsberger *et al.*, 2008; Sánchez-Molano *et al.*, 2014; Lavrijsen *et al.*, 2014)

Limber tail

- Breed at increased risk in case series
- Swimming (OR 4.7) and being a working dog (OR 5.1) were risk factors
- Affected related dogs more than might be expected purely by chance

(Pugh *et al.*, 2016)

Lumbosacral transitional vertebrae

- Inheritance suspected
- Females more frequently affected (Morgan *et al.*, 1999; Moeser & Wade, 2017)

Metaphyseal osteopathy (hypertrophic osteodystrophy)

- Affects dogs aged 2–6 months
- Males may be predisposed

- OR 5.9 (95% CI 2.8–12.4) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Muscular stiffness

- A muscular stiffness which persists at rest has been reported in male Labrador Retrievers
- X-linked inheritance suggested
- Onset of clinical signs 2–16 months
- Signs stabilize in adulthood

(Vanhaesebrouck *et al.*, 2011)

Myopathy

- Several different types are inherited in Labrador Retrievers
- Inherited as either an autosomal recessive or an X-linked condition
- Onset at 1–7 months of age
- X-linked myotubular myopathy (XLMTM) and X-linked dystrophin-deficient muscular dystrophy (XLMD) present as severe generalized muscle weakness which is rapidly progressive
- Autosomal recessive centronuclear myopathy is a less severe and less rapidly progressive form
- A mild form of XLMD has been seen in older male dogs which have raised muscle enzymes and histological changes in their muscles but do not have clinical signs of weakness and muscle atrophy
- Sarcolemmal-specific collagen VI deficiency results in progressive gait abnormalities and joint deformity

(Bley *et al.*, 2002; Maurer *et al.*, 2012;

Snead *et al.*, 2015; Cerda-Gonzalez *et al.*, 2016)

Oculoskeletal dysplasia

See under *Ocular conditions*

Osteochondrosis – elbow

See *Elbow dysplasia*

Osteochondrosis – shoulder

- Breed at increased risk in case series
- OR 13.1 (95% CI 10.4–16.4) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Males predisposed
- Age of onset usually 4–7 months, but can be older (Rudd *et al.*, 1990; LaFond *et al.*, 2002; Biezyński *et al.*, 2012)

Osteochondrosis – stifle

- Breed at increased risk in case series
- OR 27.6 (95% CI 13.2–58.0) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Males predisposed
- Age of onset usually 5–12 months (LaFond *et al.*, 2002)

Osteochondrosis – tarsus

- Breed at increased risk in case series
- OR 45.9 (95% CI 26.8–78.6) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Age at presentation 6 months to 1 year, but some present later (Montgomery *et al.*, 1994; LaFond *et al.*, 2002)

Osteosarcoma

See under *Neoplastic conditions*

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- OR 1.6 (95% CI 1.4–1.9) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002)

Patellar luxation

- Breed at increased risk in case series
- In one study of 70 large-breed dogs with patellar luxation, 25 cases were in Labrador Retrievers (RR 3.3)
- Males predisposed
- Mainly medial luxation observed in Labrador Retrievers (Arthurs & Langley-Hobbs, 2006; Gibbons *et al.*, 2006; Bound *et al.*, 2009)

Polyarthritis

- Breed at increased risk in case series
- Usually affects dogs at 1–3 years, but any age can be affected (Clements *et al.*, 2004; Stull *et al.*, 2008; Foster *et al.*, 2014)

Neoplastic conditions**Cutaneous haemangioma**

- Breed at increased risk in case series (Schultheiss, 2004)

Haemangiosarcoma

- Breed at increased risk in case series
- At risk of both visceral (including splenic and cardiac) and non-visceral (mostly skin) haemangiosarcomas
- Mean age reported as 10 years (Schultheiss, 2004; Kim *et al.*, 2007; Yamamoto *et al.*, 2013)

Histiocytic sarcoma complex – histiocytic sarcoma and disseminated histiocytoma

- Breed at increased risk in case series
 - Age of onset: > 5 years
- See also under *Ocular conditions* (Schultz *et al.*, 2007; Takahashi *et al.*, 2014; Dervisis *et al.*, 2016)

Lingual neoplasia

- Breed at increased risk in case series
- OR 2.41 (95% CI 1.46–3.98) for lingual squamous cell carcinoma in a retrospective study of 1196 dogs with lingual lesions
- Females at increased risk (OR 2.26) (Dennis *et al.*, 2006; Culp *et al.*, 2013)

Lipoma

- Breed at increased risk of infiltrative lipoma in a case series
- 8/15 dogs were Labrador Retrievers in this study
- Ratio of female:male (all breeds) was 4:1 (Bergman *et al.*, 1994)

Mast cell tumour (MCT)

- Breed at increased risk in case series
- OR 2.9 (95% CI 2.17–3.89) compared to the general hospital population in a retrospective UK study of 222 cases
- May be seen at any age (from 4 months onwards), but usually seen in older animals
- Predilection sites include hindlimb, perineum and scrotum
- An association with low vitamin D levels noted in this breed

See also under *Ocular conditions*

(Wakshlag *et al.*, 2011; Warland & Dobson, 2013)

Melanoma

- Breed at increased risk of limbal, canine uveal anterior, oral and skin/nail bed melanoma in case series
- Labradors may have a mutation predisposing them to ocular melanomas

- Limbal melanomas are usually benign, oral melanomas highly malignant (Donaldson *et al.*, 2006; Schultheiss, 2006; Featherstone *et al.*, 2009; Boston *et al.*, 2014; Tuohy *et al.*, 2014; Kawabe *et al.*, 2015)

Osteosarcoma

- Breed at increased risk in case series
- OR 1.3 (95% CI 1.0–1.7) compared to German Shepherd Dogs in a retrospective study of VMDB data, 1980–1994
- Across all breeds there was an increased risk with increasing age, plateauing at 10 years
- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes (Ru *et al.*, 1998)

Sarcoma – soft tissue

- 14/87 cases were Labrador Retrievers in a USA-based retrospective case series
- Breed at increased risk of cutaneous soft-tissue sarcoma in a retrospective study of VMDB data, 1964–2002: OR 1.48 (95% CI 1.25–1.74) compared to all other dogs (Heller *et al.*, 2005; Stefanello *et al.*, 2011; Villamil *et al.*, 2011)

Squamous cell carcinoma – digit

- Breed at increased risk in case series
- Middle-aged to older dogs predisposed
- Labrador Retrievers seem to be at risk of digital tumours generally (O'Brien *et al.*, 1992; Henry *et al.*, 2005; Capak *et al.*, 2007)

Thymoma

- Uncommon tumour
- Breed at increased risk (30/116 cases were Labrador Retrievers in a USA-based retrospective study)
- Mean age 9.5 years (Day, 1997a; Robat *et al.*, 2013; Carette *et al.*, 2014)

Neurological conditions

Exercise-induced collapse

- Onset at 7 months to 2 years of age
- Autosomal recessive inheritance, mutation identified
- In one retrospective genetic survey of different populations of Labrador Retrievers in the USA and Canada, 17.9–38.0% of tested dogs

were carriers, and 1.8–13.6% were homozygous. Of the latter, 83.6% had collapsed by 4 years of age

(Minor *et al.*, 2011; Furrow *et al.*, 2013)

Idiopathic epilepsy

- Polygenic recessive inheritance suggested
- Age of onset 6 months to 6 years
- 3.1% prevalence in a study of Danish Labrador Retrievers, compared to 1% in the general canine population (Jaggy *et al.*, 1998; Berendt *et al.*, 2002)

Leucodystrophy

- Rare conditions involving a disorder of myelin synthesis and maintenance
- Labrador Retriever central axonopathy affects spinal cord myelin; pups from 4–6 weeks are unable to walk; autosomal recessive inheritance
- Fibrinoid leucodystrophy affects myelin of the cerebrum and spinal cord; ataxia and weakness seen at 6–9 months of age; similar to Alexander disease in humans
- Spongiform leucodystrophy affects myelin diffusely (cerebellum, cerebrum and spinal cord); ataxia and tremor seen as early as 2 weeks of age; similar to Canavan's disease in humans

(Zachary & O'Brien, 1985;

de Lahunta *et al.*, 1994; Sisó *et al.*, 2005)

Narcolepsy–cataplexy

- Rare condition
- Autosomal recessive inheritance, mutation identified
- Age of clinical onset < 1 year (Hungs *et al.*, 2001; Tonokura *et al.*, 2007)

Paroxysmal dyskinesia

- Breed at increased risk in case series
- Median age at onset in 36 Labrador Retrievers was 2.25 years; 29 of the 36 were male. (Lowrie & Garosi, 2016)

Ocular conditions

Cataract

- Dominant inheritance with incomplete penetrance suggested
- Age of onset 6–18 months
- Localization: posterior polar subcapsular; slowly progressive; rarely proceeds to blindness

- Other types have been reported less commonly
- In a large-scale retrospective review of ocular examinations of Labrador Retrievers in the Netherlands (1980–2000), the prevalence of cataract was 8%

(Curtis & Barnett, 1989;
Kraijer-Huwer *et al.*, 2008)

Entropion

- Polygenic inheritance suspected
 - Lateral lower lids affected
- (Read & Broun, 2007)

Histiocytic sarcoma – ocular

- Breed at increased risk in case series
- Mean age 8.61 years

See also under *Neoplastic conditions*
(Naranjo *et al.*, 2007)

Mast cell tumour (MCT) – conjunctival

- Breed at increased risk of conjunctival mast cell tumours in case series (12/32 dogs were Labrador Retrievers)

See also under *Neoplastic conditions*
(Fife *et al.*, 2011)

Melanoma

See under *Neoplastic conditions*

Oculoskeletal dysplasia

- Dogs may be affected with retinal dysplasias (complete, geographic or multifocal) and varying degrees of developmental skeletal abnormalities (short-limbed dwarfism)
- Autosomal recessive inheritance, mutation identified
- Retinal dysplasia may also be seen without skeletal abnormality in the Labrador Retriever (Carrig *et al.*, 1988; Pellegrini *et al.*, 2002; Goldstein *et al.*, 2010)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance
 - Progressive rod–cone degeneration
 - Late onset. Night blindness at 4–6 years, severe visual impairment at 6–8 years
- (Gentilini *et al.*, 2009)

Retinal pigment epithelial dystrophy (RPED, central progressive retinal atrophy)

- Dominant inheritance with incomplete penetrance has been suggested

- More prevalent in the UK than in the USA. Becoming less prevalent following the introduction of control schemes
 - Ophthalmoscopic signs seen at 2–3 years of age; visual problems noticed at 5–7 years
- (Barnett, 1988)

Uveal cysts

- Breed at increased risk in a retrospective study of 28 cases (1989–1991)
 - Mean age was 9.1 years in Labrador Retrievers
- (Corcoran & Koch, 1993)

Physiological conditions

Serum cobalamin levels

- In a retrospective study of 28 675 serum cobalamin tests, Labradors had an increased incidence of values below the lower limit of the reference interval (251 ng/l): OR 1.4 (95% CI 1.3–1.5)
- (Grützner *et al.*, 2012)

Vertebral heart score (VHS)

- The healthy Labrador had a higher mean score than other dogs (except the Boxer and Cavalier King Charles Spaniel) in one study
- (Lamb *et al.*, 2001)

Renal and urinary conditions

Ectopic ureter

- Breed at increased risk in case series
 - Usually presents < 1 year of age
 - More commonly diagnosed in females. May present later, and is possibly underdiagnosed in males
- (Holt & Moore, 1995; Reichler *et al.*, 2012)

Lyme nephritis

- Breed at increased risk of post-borrelial-infection immune-mediated glomerulonephritis
 - Seen in 1–2% of Lyme-seropositive dogs
- (Littman, 2013)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Breed at increased risk in many case series
 - In a retrospective study in the USA of 232 cases seen at a university hospital, 1987–2012, 107 were Labrador Retrievers. Across all breeds 65% were male
 - Middle-aged to older dogs affected
- (Burbidge, 1995; Wilson & Monet, 2016)

LAKELAND TERRIER

Ocular conditions

Cataract

- Prevalence of primary cataract 7.46%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

LANCASHIRE HEELER

Gastrointestinal conditions

Parvovirus enteritis

See under *Infectious conditions*

Infectious conditions

Parvovirus enteritis

- Breed at increased risk of infection in a study of 15 442 confirmed cases seen in Banfield practices in the USA, 2010/2011
- Females at lower risk than males
- Puppies <6 months old were 11.40 times more likely to be infected than older dogs

(Lefebvre, 2013)

Ocular conditions

Collie eye anomaly

- Congenital condition
- Incidence of 13.7% demonstrated in Lancashire Heelers in the UK in 1996

(Bedford, 1998; Parker *et al.*, 2007)

Lens luxation – primary

- Mutation identified
- Age of onset: 2–6 years
- Usually progresses to glaucoma in affected dogs

(Sargan *et al.*, 2007)

LAPLAND REINDEER DOG (LAPINPOROKOIRA)

Neurological conditions

Lysosomal enzyme disease – glycogen storage disease type II (Pompe disease)

- Rare condition
- Age of onset from 6 months, death around 2 years

(Walvoort, 1985)

LAPPHUND

See *Finnish Lapphund*; *Swedish Lapphund*

LEONBERGER

Musculoskeletal conditions

Bone tumour

See under *Neoplastic conditions*

Diffuse idiopathic skeletal hyperostosis (DISH)

- Breed at increased risk in retrospective radiographic study of 2041 purebred dogs in the Netherlands, 2003–2008 (OR 9.88; $p < 0.001$)
- Prevalence was 12% (of 25 Leonbergers) compared to an overall prevalence in all dogs in the study of 3.8%

(Kranenburg *et al.*, 2011)

Hip dysplasia

- Incidence of 25% in a prospective study of Norwegian Leonbergers born 1998–2001

(Krontveit *et al.*, 2010)

Neoplastic conditions

Bone tumour

- Breed at high risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Leonbergers had an overall incidence of 53 cases (95% CI 35–70) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Males over-represented
- Median age was 7.2 years in Leonbergers; risk increases with increasing age

(Egenvall *et al.*, 2007)

Mammary neoplasia

- 46% of insured Swedish Leonbergers had developed mammary tumours at 10 years of age (ranked first), compared to 13% of the overall population

(Jitpean *et al.*, 2012)

Neurological conditions

Inherited polyneuropathy

- X-linked inheritance suspected
- Age of onset between 1 and 9 years of age

(Shelton *et al.*, 2003; Ekenstedt *et al.*, 2014)

Ocular conditions

Cataract

- In a survey of 365 Leonbergers, 1990–1998, 90 dogs had cataracts
- Localization: posterior polar subcapsular, or nuclear

(Heinrich *et al.*, 2006)

Pectinate ligament dysplasia (PLD)

- 22% of 232 dogs in a retrospective study of UK Leonbergers were affected
- Some cases developed glaucoma
- Females were more commonly affected

(Fricker *et al.*, 2016)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 61% of insured Swedish Leonbergers had developed pyometra at 10 years of age (ranked third), compared to 19% for the general population

(Chastain *et al.*, 1999; Jitpean *et al.*, 2012)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- May be seen in Leonbergers with inherited polyneuropathy
- May be the only symptom in some cases

See also under *Neurological conditions* (*Inherited polyneuropathy*)

(Granger, 2011)

LHASA APSO

Dermatological conditions

Atopic dermatitis (atopy)

- Breed at increased risk in a US series of 268 cases (1981–1984)
- Also reported as a breed at increased risk in another US study of 383 cases: OR 3.4 compared to the general hospital population
- Some studies show no sex predilection, others show females predisposed

(Scott, 1981; Schick & Fadok, 1986)

Cutaneous neoplasia

See under *Neoplastic conditions*

Haematological/immunological conditions

Haemophilia A

- Severe factor VIII deficiency is seen in the Lhasa Apso

- X-linked recessive inheritance

(Brooks, 1999)

Haemophilia B

- Factor IX deficiency
- X-linked recessive inheritance

(Mausser *et al.*, 1996)

Musculoskeletal conditions

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- Breed at increased risk in case series
- OR 6.7 (95% CI 2.6–17.1) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Patellar luxation

- Breed at increased risk in case series
- OR 3.4 (95% CI 2.5–4.8) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Sebaceous gland tumours

- Breed at increased risk of sebaceous epithelioma in a case series
- Mean age 10.7 years
- Common site: head and eyelids

(Scott & Anderson, 1990)

Neurological conditions

Hydrocephalus

- Congenital
- Breed at increased risk in a retrospective series of 564 cases
- Onset of clinical signs < 3 months

(Selby *et al.*, 1979)

Intervertebral disc disease (IVDD)

- Breed at significantly increased risk in a study of 8117 cases in North America in 1976
- Peak incidence at 4–6 years of age

(Priester, 1976)

Lissencephaly

- Rare developmental disease
 - Age of onset < 1 year
- (Saito *et al.*, 2002)

Ocular conditions**Cataract**

- Breed at increased risk in two large studies
 - OR 5.9 (95% CI 1.2–29.1) compared to the hospital population in a retrospective study of 561 dogs of small breeds in Korea 2002–2007
 - Prevalence of primary cataract 4.61%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- (Gelatt & MacKay, 2005; Park *et al.*, 2009)

Corneal ulceration (ulcerative keratitis)

- In a UK-based study of first-opinion practice, the prevalence in the Lhasa Apso was 2.13% (95% CI 1.29–3.31), compared to an overall prevalence of 0.8% (95% CI 0.75–0.86)
- (O'Neill *et al.*, 2017b)

Glaucoma – primary

- Breed at increased risk compared to the overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
 - Prevalence was 1.33% in Lhasas for the years 1994–2002, compared to an overall prevalence in all dogs of 0.89% over the same period
 - Prevalence was 1.00% in Lhasas for the years 1984–1993, compared to an overall prevalence in all dogs of 0.76% over the same period
- (Gelatt & MacKay, 2004a)

Keratoconjunctivitis sicca

- Breed at increased risk in case series
- (Kaswan & Salisbury, 1990; Berdoulay *et al.*, 2005)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk in case series. Second most commonly affected breed, representing 11/89 cases
 - Usually presents in the first 1–2 years of life
- (Morgan *et al.*, 1993)

Proptosis

- Breed at significantly increased risk of ocular proptosis in a study of 66 cases
 - Most cases are associated with trauma
 - Males are at increased risk
- (Gilger *et al.*, 1995)

Renal and urinary conditions**Familial renal disease (familial nephropathy)**

- Cases present with chronic renal failure from a few months of age
- (O'Brien *et al.*, 1982; Picut & Lewis, 1987b; Nash, 1989)

Urolithiasis – apatite

- Breed at increased risk in a case series: OR 4.1 (95% CI 3.7–4.6) compared to mixed-breed dogs in a retrospective study of 25 499 laboratory stone submissions in the USA, 1985–2006
 - Females were at significantly increased risk ($p < 0.001$), representing 77.6% of submissions across all breeds
- (Low *et al.*, 2010)

Urolithiasis – calcium oxalate

- Breed at increased risk in case series. OR 10.1 (95% CI 9.2–11.2) compared to mixed-breed dogs in one retrospective study of 25 499 laboratory stone submissions in the USA, 1985–2006. Males were at increased risk ($p < 0.001$), representing 69% of submissions across all breeds
 - In a Canadian study of urolith submissions, 1998–2001, OR was 5.19 (95% CI 3.69–7.31) compared to mixed-breed dogs
 - In a retrospective study of 14 008 uroliths from dogs in the UK, 1997–2006, OR was 7.2 (95% CI 6.03–8.61; $p < 0.0001$) compared to the national insurance database. Males predisposed
- (Ling *et al.*, 2003; Houston *et al.*, 2004; Low *et al.*, 2010; Roe *et al.*, 2012)

Urolithiasis – silica

- Breed at increased risk in case series. OR 6.7 (95% CI 5.3–8.4) compared to mixed-breed dogs in a retrospective study of 25 499 laboratory stone submissions in the USA, 1985–2006
 - Males were at significantly increased risk ($p < 0.001$), representing 88.8% of submissions across all breeds
- (Low *et al.*, 2010)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in a retrospective study of 16 000 laboratory stone submissions, 1998–2003. Average age at diagnosis: males 6.0 years, females 5.7 years. Females predisposed (16:1 in Lhasa Apsos and other predisposed breeds)
- In a retrospective study of 14 008 uroliths from dogs in the UK, 1997–2006, OR was 4.2 (95% CI 3.51–5.02; $p < 0.0001$) compared to the national insurance database. Females predisposed

(Houston *et al.*, 2004; Roe *et al.*, 2012)

LOWCHEN (LITTLE LION DOG)

Ocular conditions

Vitreoretinopathy

- Vitreal liquefaction predisposes to retinal detachment in Lowchens

(Papaioannou & Dubielzig, 2013)

MAH THAI

See *Thai Ridgeback*

MALAMUTE

See *Alaskan Malamute*

MALTESE

Cardiovascular conditions

Mitral valve disease

- Breed at increased risk in case series
- Increased prevalence with increasing age

(Heejin Oui *et al.*, 2015)

Patent ductus arteriosus

- OR 4.14 ($p < 0.0001$) compared to hospital population in a study of 976 cases of congenital heart disease in Italy
- Females predisposed (OR 2.7)

(Oliveira *et al.*, 2011)

Ventricular septal defect

- OR 9.70 (95% CI 2.98–31.57) compared to the general hospital population, in an Australian

retrospective study of 13 dogs with ventricular septal defect seen between 2001 and 2012

- Median age at diagnosis 6 months (range 2–60)

(Aherne & Beijerink, 2013)

Drug reactions

Vaccine-associated adverse effect

- In a large retrospective study in the USA (2002–2003), Maltese dogs had a mean rate of 66.8 adverse reactions per 10 000 dogs vaccinated, compared to an overall rate of 38.2 reactions per 10 000 dogs vaccinated
- Risk seemed to decrease with increasing body weight

(Moore *et al.*, 2005)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Breed with an increased prevalence of thyroid hormone autoantibodies (THAA)
- In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, Maltese had an odds ratio of being affected of 2.25 ($p < 0.001$) compared to dogs of all other breeds
- Across the study, females were over-represented and the highest prevalence was in dogs 2–4 years old

(Nachreiner *et al.*, 2002)

Gastrointestinal conditions

Antral pyloric hypertrophy (pyloric stenosis)

- Breed at increased risk in case series
- Mean age at diagnosis 8.2 years
- Males may be predisposed

(Bellenger *et al.*, 1990)

Atresia ani

- Breed at increased risk of this rare disease: OR 13.385 compared to mixed-breed dogs
- Females predisposed: OR 1.796
- Symptoms seen at weaning

(Vianna & Tobias, 2005)

Congenital portosystemic shunt

- Maltese were in the top five breeds affected in a large USA-based retrospective study of 90 004 dogs presented to a university teaching hospital. 5.87% were affected, compared to 0.35% of mixed breeds
- Clinical signs usually seen in young dogs < 1 year

- Usually extrahepatic in the Maltese (Tobias & Rohrbach, 2003; Hunt, 2004; Bellumori *et al.*, 2013; O'Leary *et al.*, 2014)

Haemorrhagic gastroenteritis

- Breed at increased risk in case series
- In a study of 108 dogs in Germany, 2006–2009, Maltese represented 4.6% of the affected dogs, but only 1.3% of the general hospital population
- Increased incidence in the winter months (Mortier *et al.*, 2015)

Haematological/immunological conditions

Haemophilia B

- Severe factor IX deficiency in the Maltese
- X-linked recessive inheritance (Brooks, 1999)

Immune-mediated haemolytic anaemia (IMHA)

- Breed at increased risk in case series
- OR 2.8 (95% CI 1.5–4.9) in a study of 110 cases in Australia
- Median age was 6 years (McAlees, 2010)

Musculoskeletal conditions

Patellar luxation

- Mainly medial luxation observed in the Maltese
- Breed at increased risk in case series
- OR 6.5 (95% CI 4.1–10.2) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995 (LaFond *et al.*, 2002; Alam *et al.*, 2007)

Neoplastic conditions

Mammary neoplasia

- Breed at increased risk in case series in Japan
- 28/101 dogs in the study were Maltese (higher than the breed prevalence in the area) (Itoh *et al.*, 2005)

Testicular neoplasia

See under *Reproductive conditions*

Neurological conditions

Granulomatous meningoencephalitis

- Breed at increased risk
- Dogs over 8 years less likely to be diagnosed than younger dogs (Granger *et al.*, 2010; Wouda, 2015)

Hydrocephalus

- Congenital
- Breed at increased risk in a retrospective series of 564 cases
- Onset of clinical signs < 3 months (Selby *et al.*, 1979)

Necrotizing meningoencephalitis

- Breed at increased risk
- Genetic risk loci identified (Granger *et al.*, 2010; Schrauwen *et al.*, 2014)

Shaker dog disease

- Uncommon
- Age of clinical onset < 2 years (Bagley *et al.*, 1993)

Ocular conditions

Cataract

- Prevalence of primary cataract 3.21%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- However, at lower risk (OR 0.7) compared to the hospital population in a retrospective study of 561 dogs of small breeds in Korea, 2002–2007 (Gelatt & MacKay, 2005; Park *et al.*, 2009)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series. OR 5.2 (95% CI 4.4–6.0) compared to mixed-breed dogs, in one retrospective case series of 25 499 laboratory stone submissions in the USA, 1985–2006
- Males at increased risk ($p < 0.001$), representing 69% of submissions across all breeds
- In a Canadian case series of urolith submissions, 1998–2001, OR was 6.77 (95% CI 3.63–12.62) compared to mixed-breed dogs (Ling *et al.*, 2003; Low *et al.*, 2010)

Reproductive conditions

Cryptorchidism

- Breed at increased risk in case series
- Congenital defect believed to be inherited as a sex-limited, autosomal recessive trait
- Retained testicles are at a higher risk of neoplasia (Hayes *et al.*, 1985; Romagnoli, 1991)

Dystocia

- Breed at increased risk in a retrospective study of Swedish insurance data (including 195 931 bitches, 1995–2002)
- Incidence in Maltese was 11.5 cases per 1000 DYAR, compared to 5.7 cases per 1000 DYAR for the general population (Bergström *et al.*, 2006)

Testicular neoplasia

- Breed at increased risk in case series
- May be associated with high risk of cryptorchidism (Liao *et al.*, 2009)

MANCHESTER TERRIER**Cardiovascular conditions****Dilated cardiomyopathy (DCM)**

- Rapidly progressive form identified in juvenile Toy Manchester terriers
- Affected dogs are 10–58 weeks old
- May die suddenly without prior clinical signs (Legge *et al.*, 2013)

Haematological/immunological conditions**Haemophilia A**

- Factor VIII deficiency
- X-linked recessive inheritance (Brooks, 1999)

von Willebrand's disease (vWD)

- Possibly inherited as an autosomal recessive trait
- Mainly type I disease seen in the Manchester Terrier (Brooks, 1999)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- Test matings have shown that this condition is inherited with high heritability in the Manchester Terrier (Vasseur *et al.*, 1989)

Ocular conditions**Cataract**

- Prevalence of primary cataract 3.81%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at

North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Glaucoma – primary

- Breed at increased risk compared to overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
- Prevalence of 2.63% in Manchester Terriers for the years 1984–1993 compared to an overall prevalence in all dogs of 0.76% in the same period (Gelatt & MacKay, 2004a)

Glaucoma – secondary

- Breed at increased risk in case series of 217 cases in Switzerland 1995–2009 (Strom *et al.*, 2011b)

MASTIFF

See *Dogue de Bordeaux*; *Italian Mastiff*; *Neapolitan Mastiff*; *Old English Mastiff*; *Savoy Sheepdog*

MASTIFF (UNSPECIFIED)**Cardiovascular conditions****Atrial fibrillation (AF)**

- Breed at increased risk in two case series
- In a retrospective report of cases from the VMDB, 1969–2007, OR was 8.34 compared to all breeds
- Males were significantly over-represented across all breeds in the series (2549 vs. 977 females) (Menaut *et al.*, 2005; Westling *et al.*, 2008)

Drug reactions**Vaccine-associated adverse effect**

- In a large retrospective study in the USA (June 2012 to March 2013), Mastiffs had a rate of 83 reactions per 10 000 dogs vaccinated, compared to an overall rate of 26.3 reactions per 10 000 dogs vaccinated (Peng *et al.*, 2015)

Gastrointestinal conditions**Gastric dilatation/volvulus (bloat, GDV)**

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred

dogs conducted by questionnaires relating to the 10 years prior to 2004

- Prevalence ratio for GDV mortality was 6.8 (95% CI 4.1–11.3; $p < 0.001$) compared to all other breeds combined

(Evans & Adams, 2010b)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Breed at increased risk in a study of 201 dogs < 2 years of age

(Duval *et al.*, 1999)

Elbow dysplasia

- Breed at increased risk in case series
- Fragmented coronoid process: OR 48.4 (95% CI 26.3–89.1) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Ununited anconeal process: OR 20.2 (95% CI 8.9–45.9) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Hip dysplasia

- Neutered male dogs predisposed
- OR 3.5 (95% CI 2.2–5.6) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Osteochondrosis – shoulder

- Males predisposed
- Age of onset usually 4–7 months, but can be older
- OR 11.9 (95% CI 6.1–23.2) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Osteochondrosis – stifle

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 1006.9 (95% CI 339.1–2989.5) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Osteosarcoma

See under *Neoplastic conditions*

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- OR 3.5 (95% CI 2.2–5.6) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Osteosarcoma

- OR 5.1 (95% CI 1.8–14.5) compared to a control population in a case–control study of 3062 cases from the VMDB, 1980–1994
- Across all breeds there was an increased incidence with age, plateauing at 10 years
- Across all breeds there was a slight predisposition for males, but an almost twofold increase for neutered dogs of both sexes

(Ru *et al.*, 1998)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal dominant inheritance

(Kijas *et al.*, 2003)

Renal and urinary conditions

Urolithiasis – cystine

- Breed at increased risk in case series
- Young dogs affected (2–5 years)
- Almost all cases are male

(Case *et al.*, 1992)

Reproductive conditions

Delivery by caesarean section

- In the top 10 breeds in a cross-sectional study conducted by the Kennel Club in the UK. The study found a caesarean rate of 64.9% in 79 litters of Mastiffs born over a 10-year period

(Evans & Adams, 2010a)

MCNAB SHEPHERD

Drug reactions

Ivermectin and milbemycin

- High doses can cause tremors, ataxia, coma and death
- Associated with the mutant *MDR1* allele

(Dowling, 2006)

MEXICAN HAIRLESS

Dermatological conditions

Acne

- Comedones commonly occur spontaneously in this breed

(Kimura & Doi, 1996)

Canine ectodermal dysplasia (hairlessness)

- Monogenic, autosomal semi-dominant inheritance

(Drögemüller *et al.*, 2008b)

Reproductive conditions

High perinatal mortality

- Hairless pups have a markedly lower survival rate than haired pups
- Raising the temperature of cages to a minimum of 25°C improved survival rates

(Kimura *et al.*, 1993)

MINIATURE BULL TERRIER

Ocular conditions

Lens luxation – primary

- Estimated prevalence 7.3–15.2%
- Mutation identified
- Age of onset > 3 years

(Curtis *et al.*, 1983; Gould *et al.*, 2011; Gharahkhani *et al.*, 2012)

Reproductive conditions

Delivery by caesarean section

- In the top 10 breeds in a cross-sectional study conducted by the Kennel Club in the UK. The study found a caesarean rate of 52.4% in 42 litters of Miniature Bull Terriers born over a 10-year period

(Evans & Adams, 2010a)

MINIATURE PINSCHER

Cardiovascular conditions

Mitral valve disease

- Breed at increased risk: OR 8.75 (95% CI 3.29–23.28) in a UK study

(Thrusfield *et al.*, 1985)

Dermatological conditions

Colour dilution alopecia

- Reported in blue or fawn Dobermanns and Miniature Pinschers

- Frequency of the disease may be as high as 93% in blues and 75% in fawns
- Coat-colour genes play a role in the inheritance of this condition

(Miller, 1990a)

Demodicosis

- Miniature Pinschers are at increased risk of juvenile-onset, generalized demodicosis compared to a control hospital population, OR 2.0 (95% CI 1.3–3.1) based on a large retrospective case-control study of a US practice database
- Juvenile onset was defined as < 18 months old
- Other risk factors associated with demodicosis included pyoderma, short hair and concurrent coccidiosis

(Plant *et al.*, 2011)

Drug reactions

Vaccine-associated adverse effect

- In a large retrospective study in the USA (2002–2003), Miniature Pinschers had a mean rate of 74.6 adverse reactions per 10 000 dogs vaccinated, compared to an overall rate of 38.2 reactions per 10 000 dogs vaccinated. The risk seemed to decrease with increasing body weight
- In a second large retrospective study in the USA (June 2012 to March 2013), Miniature Pinschers had a rate of 50.8 reactions per 10 000 dogs vaccinated, compared to an overall rate of 26.3 reactions per 10 000 dogs vaccinated

(Moore *et al.*, 2005; Peng *et al.*, 2015)

Gastrointestinal conditions

Congenital portosystemic shunt

- Breed at increased risk in a retrospective case study of 2400 cases from VMDB data, 1980–2002
- OR 7.0 (95% CI 1.2–22.7) compared to mixed-breed dogs
- Female:male ratio was 1.14:1 across all breeds
- Clinical signs usually seen in young dogs < 1 year

(Tobias & Rohrbach, 2003)

Haemorrhagic gastroenteritis

- Breed at increased risk in a case-control study of 108 cases
- More cases seen in the winter months

(Mortier *et al.*, 2015)

Haematological/immunological conditions

Immune-mediated haemolytic anaemia (IMHA)

- Breed at increased risk in a study of 33 cases: OR 7.4 (95% CI 1.2–47.1) compared to control population
- Usually affects young adult and middle-aged dogs
- Females at increased risk: OR 2.1 in this study (Miller *et al.*, 2004)

Musculoskeletal conditions

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- Breed at increased risk in case series
- OR 71.5 (95% CI 31.1–164.4) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Patellar luxation

- Breed at increased risk in case series
- OR 14.4 (95% CI 5.8–35.9) compared to mixed breeds in a large retrospective series based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Ocular conditions

Cataract

- Prevalence of primary cataract 4.58%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in a Canadian study of urolith submissions, 1998–2001: OR 5.03 (95% CI 2.12–11.93) compared to mixed-breed dogs

(Ling *et al.*, 2003)

Urolithiasis – cystine

- Autosomal dominant inheritance, mutation identified
- Breed at increased risk in case series. OR 9.3 (95% CI 4.0–22.0) compared to mixed-breed dogs in one retrospective case series of 25 499 laboratory stone submissions in the USA,

1985–2006. Males at increased risk ($p < 0.001$), representing 97.8% of submissions across all breeds

(Case *et al.*, 1992; Low *et al.*, 2010; Brons *et al.*, 2013)

Reproductive conditions

Eclampsia (puerperal tetany)

- Breed at increased risk in case series
- Usually seen in the first 6 weeks postpartum (Chastain *et al.*, 2001)

MOUNTAIN DOG

See *Entlebucher Mountain Dog*; *Estrela Mountain Dog*; *Swiss Mountain Dog*

MÜNSTERLÄNDER – POINTER

Dermatological conditions

Black hair follicular dysplasia

- Seen in large Münsterländers
- Autosomal recessive inheritance
- Grey and white fur is noted at birth
- Grey hair falls out over first few months of life (Philipp *et al.*, 2005; Von Bomhard *et al.*, 2006)

Musculoskeletal conditions

Osteochondrosis – shoulder

- OR 83.1 (95% CI 28.2–245.0) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Males predisposed
- Age of onset usually 4–7 months, but can be older

(LaFond *et al.*, 2002)

Renal and urinary conditions

Urolithiasis – cystine

- Mutation identified

(Karmi *et al.*, 2010a)

NEAPOLITAN MASTIFF

Cardiovascular conditions

Dilated cardiomyopathy (DCM)

- Neapolitan Mastiffs were in the top five breeds affected in a large USA-based retrospective

case study of 90 004 dogs presented to a university teaching hospital. 6.52% were affected, compared to 0.16% of mixed breeds

(Bellumori *et al.*, 2013)

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaire, relating to the 10 years prior to 2004
- Prevalence ratio for GDV mortality was 11.7 (95% CI 3.6–38; p 0.0012) compared to all other breeds combined

(Evans & Adams, 2010b)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Breed at increased risk in a study of 201 dogs < 2 years of age
- Neutered individuals are predisposed

(Duval *et al.*, 1999)

Hip dysplasia

- OR 4.7 (95% CI 1.7–13.2) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Neutered male dogs predisposed

(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- OR 2.0 (95% CI 1.2–3.5) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Ocular conditions

Ectropion

- High prevalence seen in an Italian study
- Associated with macroblepharon

(Guandalini *et al.*, 2016)

Entropion

- High prevalence seen in an Italian study
- Associated with macroblepharon

(Guandalini *et al.*, 2016)

Macroblepharon (diamond eye)

- High prevalence seen in an Italian study

(Guandalini *et al.*, 2016)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Breed at increased risk in case series

(Prémont *et al.*, 2012; Guandalini *et al.*, 2016)

NEW ZEALAND HUNTAWAY

Cardiovascular conditions

Dilated cardiomyopathy (DCM)

- In one study, a statistically significant predisposition to this disease was observed; however, only small numbers were involved
- All the Huntaway cases in this study were male
- Average age at diagnosis was 4 years

(Munday *et al.*, 2006)

Neurological conditions

Mucopolysaccharidosis IIIA

- Autosomal recessive inheritance
- Causes severe progressive CNS disease

(Yogalingam *et al.*, 2002)

NEWFOUNDLAND

Cardiovascular conditions

Aortic stenosis – subaortic stenosis (SAS)

- Breed at increased risk (OR 7.0; p < 0.0001) compared to the general hospital population in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
- This study found a male predisposition (OR 1.7). Previous studies have found no gender difference
- In a separate retrospective review of 195 cases (1967–1991), the OR was 88.1 (95% CI 59.7–130) for the Newfoundland compared to the hospital population
- Autosomal dominant inheritance, mutation identified

(Kienle *et al.*, 1994; Oliveira *et al.*, 2011; Stern *et al.*, 2014)

Atrial fibrillation (AF)

- Breed at increased risk in a series of cases of atrial fibrillation without cardiac disease or failure
- In a separate retrospective study of VMDB data, 1969–2007, Newfoundlands had an OR

of 11.60 compared to all dogs. Male dogs were over-represented

(Menaut *et al.*, 2005; Westling *et al.*, 2008)

Dilated cardiomyopathy (DCM)

- Autosomal dominant inheritance with late onset
- Males predisposed
- Median age of onset 5 years
(Tidholm & Jonsson, 1996, 1997; Wiersma *et al.*, 2008)

Patent ductus arteriosus

- Breed at increased risk in case series
- OR 4.65 ($p < 0.001$) compared to the general hospital population in a retrospective study of the medical records of 976 dogs diagnosed with congenital heart disease at a cardiology clinic in Italy
(Oliveira *et al.*, 2011)

Dermatological conditions

Atopic dermatitis (atopy)

- In a study of insured Swedish dogs (1995–2002), Newfoundlands had an incidence of 3.8 cases per 1000 DYAR, compared to 1.7 cases per 1000 DYAR for all dogs in the study
- Birth in autumn or summer is a predisposing factor
- Dogs between 1 and 2 years of age have the highest probability of an insurance claim for atopy
- Some studies show no sex predilection, others show females predisposed
(Nødtvedt *et al.*, 2006)

Pemphigus foliaceus

- Breed at increased risk in case series
- Mean age of onset 4.2 years
(Ihrke *et al.*, 1985)

Gastrointestinal conditions

Acquired megaesophagus

- May be seen secondary to myasthenia gravis and inflammatory myopathy in Newfoundlands
(Evans *et al.*, 2004)

Exocrine pancreatic insufficiency (EPI)

- Breed at increased risk in case series
(Williams & Minnich, 1990)

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaire, relating to the 10 years prior to 2004
- Prevalence ratio for GDV mortality was 2.3 (95% CI 1.4–3.8; $p = 0.0016$) compared to all other breeds combined
(Evans & Adams, 2010b)

Musculoskeletal conditions

Acquired myasthenia gravis

See under *Neurological conditions*

Bone tumour

See under *Neoplastic conditions*

Cranial cruciate ligament (CCL) disease

- Breed at increased risk in a study of 201 dogs < 2 years of age
- Breed at highest risk in a separate retrospective study of VMDB cases, 1964–2003: OR 3.77 (95% CI 3.4–4.18; $p < 0.001$) compared to all other dogs. Neutered dogs of both sexes and dogs > 4 years old of all breeds combined were found to be at increased risk
- Newfoundlands were also found to be at highest risk of having both cranial cruciate disease and hip dysplasia: OR 10.01 (95% CI 8.33–12.02)
- An autosomal recessive mode of inheritance with 51% penetrance has been suggested
(Duval *et al.*, 1999; Wilke *et al.*, 2006; Witsberger *et al.*, 2008)

Elbow dysplasia

- Increased risk of osteochondrosis of the elbow: OR 261 (95% CI 107.1–635.8) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Increased risk of fragmented coronoid process: OR 10.9 (95% CI 5.0–24.0) compared to mixed breeds in the same study
- Increased risk of ununited anconeal process: OR 13.8 (95% CI 7.1–26.8) compared to mixed breeds in the same study
(LaFond *et al.*, 2002)

Hip dysplasia

- Breed at highest risk in case series: OR 5.77 (95% CI 5.35–6.21; $p < 0.001$) compared to all

other dogs in a retrospective study of VMDB cases 1964–2003. Neutered male dogs and dogs < 4 years of age of all breeds combined were found to be at increased risk

- Newfoundlands were also found to be at the highest risk of having both cranial cruciate disease and hip dysplasia: OR 10.01 (95% CI 8.33–12.02)

(Witsberger *et al.*, 2008;
Krontveit *et al.*, 2010)

Myopathy – inflammatory

- Significantly over-represented in a study of 200 cases in the USA. Newfoundlands represented 9.5% of the cases but only 0.26% of AKC registrations
- Mean age at diagnosis 2.1 years

(Evans *et al.*, 2004)

Osteochondrosis – shoulder

- Breed at increased risk in case series: OR 18.7 (95% CI 12–29.2) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Young males predisposed

(LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Seen in rapidly growing young dogs
- Breed at increased risk in case series: OR 1.9 (95% CI 1.3–2.8) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Neoplastic conditions

Bone tumour

- Breed at high risk of malignant bone tumours in one study of Swedish insurance data (1995–2002). Newfoundlands had an overall incidence of 22 cases (95% CI 11–32) per 10 000 DYAR, compared to the risk in all dogs combined of 5.5 cases per 10 000 DYAR
- Males over-represented
- Median age was 8.2 years in Newfoundlands; risk increases with increasing age

(Egenvall *et al.*, 2007)

Neurological conditions

Acquired myasthenia gravis

- Familial predisposition

(Lipsitz *et al.*, 1999)

Ocular conditions

Eversion of the cartilage of the nictitating membrane

- Inheritance suspected
- Usually occurs in young dogs
(Allbaugh & Stuhr, 2013)

Glaucoma – primary

- Breed at increased risk in a retrospective study of 123 cases in Switzerland, 1995–2009
- Age at presentation 6.8 ± 2.5 years
(Strom *et al.*, 2011a)

Renal and urinary conditions

Ectopic ureter

- Breed at increased risk in a case series: OR 12.6 (95% CI 5.0–29.6) compared to a control population in a retrospective study of cases in North American teaching hospitals, 1964–1981
- Usually presents < 1 year of age
- More commonly diagnosed in females
(Hayes, 1984)

Urolithiasis – cystine

- Breed at increased risk in case series: OR 12.6 (95% CI 6.9–22.6) compared to mixed-breed dogs in a retrospective case series of 25 499 uroliths in the USA, 1985–2006
- Presents earlier in Newfoundlands than in other breeds (from 6 months to 1 year)
- Males and females are both affected in this breed, but only males are seen with urinary obstruction
(Case *et al.*, 1992; Casal *et al.*, 1995; Low *et al.*, 2010)

NORFOLK TERRIER

Cardiovascular conditions

Mitral valve disease

- In a study of 48 apparently healthy Norfolk Terriers 6 years and older, 48% had a heart murmur and 85% had echocardiographic evidence of degenerative mitral valve disease
(Trafny *et al.*, 2012)

Dermatological conditions

Epidermal hyperkeratosis

- Autosomal recessive mode of inheritance
- Clinical signs noted at birth
(Barnhart *et al.*, 2004)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk. OR 9.69 (95% CI 7.02–13.37; $p < 0.0001$) compared to the national insurance database in a retrospective study of 14 008 uroliths from dogs in the UK, 1997–2006
- Males predisposed

(Roe *et al.*, 2012)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk. OR 2.39 (95% CI 1.55–3.69; $p < 0.0001$) compared to the national insurance database in a retrospective study of 14 008 uroliths from dogs in the UK, 1997–2006
- Females predisposed

(Roe *et al.*, 2012)**Reproductive conditions**

Dystocia

- Breed at increased risk in a retrospective study of Swedish insurance data including 195 931 bitches, 1995–2002
- Incidence in Norfolk Terriers was 13.6 cases per 1000 DYAR, compared to 5.7 cases per 1000 DYAR for the general population

(Bergström *et al.*, 2006)**NORWEGIAN BUKUND****Behavioural conditions**

Noise sensitivity

- Highest frequency of noise sensitivity of 17 breeds in a web-based survey of owners in Norway
- Fearfulness increased with age, and was greater in females (OR 1.3) and in neutered dogs (OR 1.73)

(Storengen & Lingaas, 2015)

Ocular conditions

Cataract

- Autosomal dominant inheritance
- In studies, approximately 50% of the breed have been found to be affected
- Age of onset 6 weeks. Rate of progression varies
- Localization: starts as small dots and progresses to involve the whole fetal nucleus at 4–5 years. Pulverulent ('candyfloss') appearance

- Norwegian Sheepdogs also suffer with posterior polar and cortical cataracts (Bjerkås & Haaland, 1995; Kristiansen *et al.*, 2017)

NORWEGIAN ELKHOUND**Endocrine conditions**

Diabetes mellitus

- Breed at increased risk of diabetes in dioestrus and pregnancy
- Almost all cases in the Norwegian Elkhound are female

(Fall *et al.*, 2007)**Gastrointestinal conditions**

Gastric carcinoma

See under *Neoplastic conditions***Haematological/immunological conditions**

Selective IgA deficiency

- Breed at increased risk in case series
- 21% of 14 Norwegian Elkhounds were found to have low serum IgA (< 0.07 g/l) in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease

(Olsson *et al.*, 2014)**Musculoskeletal conditions**

Hip dysplasia

- Breed at increased risk in case series: OR 2.3 (95% CI 1.5–3.6) compared to mixed-breed dogs in a large retrospective study based on cases recorded in the VMDB, 1986–1995

(LaFond *et al.*, 2002)

Norwegian Elkhound chondrodysplasia

- Inherited as an autosomal recessive trait
- Shortened body and disproportionately short limbs

(Bingel & Sande, 1982; Kyöstiä *et al.*, 2013)**Neoplastic conditions**

Gastric carcinoma

- Breed at increased risk in a retrospective study of data from the Norwegian Canine Cancer Register, 1998–2009
- Males at increased risk: OR 2.3 (all breeds combined)
- Median age 10 years (all breeds combined)

(Seim-Wikse *et al.*, 2013)

Intracutaneous cornifying epithelioma (keratoacanthoma)

- Predisposed to the generalized form (up to 40 growths)
- Mean age 7.3 years (Stannard & Pulley, 1975)

Ocular conditions

Glaucoma – primary

- Inheritance suspected, mutation identified
- Most primary cases appear to be open-angle
- Lens luxation may occur secondary to glaucoma
- Often only diagnosed in middle-aged and older dogs
- Breed at increased risk compared to overall population of dogs in a retrospective study of 9778 dogs with primary glaucoma from the VMDB, 1964–2002
- Prevalence was 1.98% in Norwegian Elkhounds for the years 1994–2002, compared to an overall prevalence in all dogs of 0.89% over the same period
- Prevalence was 2.34% in Norwegian Elkhounds for the years 1984–1993, compared to an overall prevalence in all dogs of 0.76% over the same period

(Gelatt & MacKay, 2004a, Oshima *et al.*, 2004; Ahonen *et al.*, 2014)

Lens luxation – secondary

- Occurs secondary to glaucoma (Ahonen *et al.*, 2014)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance
- Early retinal degeneration
- Night blindness at 6 weeks followed by total vision loss at 12–18 months (Acland & Aguirre, 1987)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Mode of inheritance unknown
- Norwegian Elkhounds suffer with periglomerular and interstitial fibrosis. Dogs may present in renal failure from a few months to 5 years of age. Primary renal glycosuria is present in some cases

(Finco *et al.*, 1977; Wiersma *et al.*, 2005)

NORWEGIAN LUNDEHUND

Dermatological conditions

Alopecia

- A non-infectious spontaneous multifocal alopecia has been seen in approximately 11.7% of Norwegian Lundehund in Sweden
- Median age of onset 1.5 years (Bergvall & Shokrai, 2014)

Gastrointestinal conditions

Chronic atrophic gastritis

- Associated with gastric neoplasia and lymphangiectasia (see below) (Qvigstad *et al.*, 2008)

Gastric neoplasia

- Breed at increased risk of gastric neuroendocrine carcinoma
- Seen associated with chronic atrophic gastritis and lymphangiectasia (see below) (Qvigstad *et al.*, 2008)

Inflammatory bowel disease

- Associated with lymphangiectasia (see below) (Landsverk & Gamlem, 1984; Kolbjørnsen *et al.*, 1994)

Lymphangiectasia (resulting in protein-losing enteropathy)

- Breed at increased risk
- Specific syndrome in this breed associated with chronic atrophic gastritis, gastric neoplasia and lymphocytic-plasmacytic enteritis (Landsverk & Gamlem, 1984; Kolbjørnsen *et al.*, 1994)

Musculoskeletal conditions

Joint flexibility

- Extreme flexibility seen in all joints (including the spine) (Pfahler & Distl, 2015)

Polydactyly

- Affects all four limbs. Often six toes per foot (Pfahler & Distl, 2015)

NORWEGIAN SHEEPDOG

See *Norwegian Buhund*.

NORWICH TERRIER

Gastrointestinal conditions

Congenital portosystemic shunt

- Norwich Terriers were among the top five breeds affected in a large USA-based retrospective study of 90 004 dogs presented to a university teaching hospital. 7.4% of Norwich Terriers were affected, compared to 0.35% of mixed breeds

(Bellumori *et al.*, 2013)

Musculoskeletal conditions

Paroxysmal dyskinesia

- In a study of 195 Norwich Terriers, 13% were affected
- Mean age at first episode was 3 years

(De Risio *et al.*, 2016a)

Ocular conditions

Cataract

- Prevalence of primary cataract 4.02%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005)

Respiratory conditions

Brachycephalic obstructive airway syndrome (BOAS)

- Breed at increased risk
- In a prospective study of 16 dogs, 12 had upper respiratory symptoms, and 11 of these were found to have abnormalities of the laryngeal opening including everted laryngeal sacculles, laryngeal collapse and narrow laryngeal opening

(Johnson *et al.*, 2013)

NOVA SCOTIA DUCK TOLLING RETRIEVER

Endocrine conditions

Hypoadrenocorticism (Addison's disease)

- Autosomal recessive inheritance
- Autoimmune basis suspected
- Early onset (median age 3 years) in this breed
- Females may be predisposed

(Hughes *et al.*, 2007, 2010)

Haematological/immunological conditions

Selective IgA deficiency

- Breed at increased risk in case series
- 20% of 11 Nova Scotia Duck Tolling Retrievers were found to have a low serum IgA (<0.07 g/l) in one 2014 study
- Immunodeficiency is associated with recurrent infections of mucosal sites and increased susceptibility to immune-mediated disease

(Olsson *et al.*, 2014)

Musculoskeletal conditions

Immune-mediated rheumatic disease (IMRD)

- Breed at increased risk. In a study of Swedish insurance data, 1995–2006 this breed was 18 times more likely to suffer with IMRD than all other breeds combined
- Incidence 6.8 cases per 10 000 DYAR
- In one study, 70% of cases were antinuclear antibody (ANA)-positive, and some showed immune-mediated skin changes
- Dogs were mostly middle-aged

(Hansson-Hamlin & Lilliehöök, 2009; Bremer *et al.*, 2015)

Neoplastic conditions

Lymphoma

- Breed at increased risk. In a study of Swedish insurance data, 1995–2006, this breed was 2.8 times more likely to suffer with lymphoma than all other breeds combined
- Incidence 15 cases per 10 000 DYAR

(Bremer *et al.*, 2015)

Neurological conditions

Degenerative encephalopathy

- Autosomal recessive inheritance suspected
- Symptoms are present at 2 months to 5 years, are progressive, and include movement during sleep, increased anxiety, noise phobia and gait abnormalities

(Barker *et al.*, 2016)

Steroid-responsive meningitis–arteritis (SRMA)

- Breed at increased risk. In a study of Swedish insurance data, 1995–2006, this breed was 12 times more likely to suffer with steroid-responsive meningitis–arteritis than all other breeds combined

- Incidence 20 cases per 10 000 DYAR
- Dogs were mostly young
(Hansson-Hamlin & Lilliehöök, 2013; Bremer *et al.*, 2015)

OGAR POLSKI

See *Polish Hound*

OLD ENGLISH MASTIFF

Neurological conditions

Primary brain tumour – oligodendroglioma

- Breed at significantly increased risk compared to the general population in a study of post-mortem results of 435 dogs with intracranial neoplasia
(Song *et al.*, 2013)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal dominant inheritance
(Vilboux *et al.*, 2008)

OLD ENGLISH SHEEPDOG (OESD)

Cardiovascular conditions

Atrial fibrillation (AF)

- Breed at increased risk: OR 3.0 in a review of VMDB data, 1969–2007, compared to all dogs
- Males were significantly over-represented across all breeds
(O'Grady *et al.*, 2008; Westling *et al.*, 2008)

Dermatological conditions

Demodicosis

- Breed at increased risk: RR 28.9 in a Canadian study, 1987–1988, compared to hospital population
- Lesions may be confined to the paws in OESDs
(Scott & Paradis, 1990)

Drug reactions

Ivermectin and milbemycin

- In a UK study, 11% of OESDs were found to have the *MDR1* mutation which has been associated with multiple drug sensitivity
(Tappin *et al.*, 2012)

Endocrine conditions

Hypothyroidism (lymphocytic thyroiditis)

- Breed with a higher prevalence of thyroid hormone autoantibodies (THAA)
- In a cohort study of 287 948 serum samples from dogs in the USA with clinical signs of hypothyroidism, OESD had an OR of 2.65 ($p < 0.001$) of being affected compared to dogs of all other breeds
(Nachreiner *et al.*, 2002)

Haematological/immunological conditions

Immune-mediated haemolytic anaemia (IMHA)

- Breed at increased risk. In a study of 24 cases, 33% were OESDs (however other studies have failed to confirm a predisposition in this breed)
- Most cases were ANA-positive, suggesting a multi-system autoimmune syndrome
- Usually affects young adult and middle-aged animals
- May be more common in bitches
(Mills *et al.*, 1985)

Musculoskeletal conditions

Hip dysplasia

- Breed at increased risk in case series
- OR 3.46 (95% CI 3.21–3.74) compared to all other dogs in a large retrospective study of VMDB cases, 1964–2003. Male neutered dogs and dogs < 4 years of age of all breeds combined were found to be at increased risk
- OR 7.1 (95% CI 4.1–12.4) compared to mixed breeds in a separate retrospective study based on cases recorded in the VMDB, 1986–1995
(LaFond *et al.*, 2002; Witsberger *et al.*, 2008)

Osteochondrosis – shoulder

- Breed at increased risk in case series. OR 7.1 (95% CI 4.1–12.4) compared to mixed breeds in a large retrospective study based on cases recorded in the VMDB, 1986–1995
- Young males predisposed
(LaFond *et al.*, 2002)

Neoplastic conditions

Oral neoplasia

- Breed at increased risk (compared to the hospital population) in a retrospective case series in the USA, 1980–1987
- Median age across all breeds was 10 years

- 62% of all affected dogs were > 20 kg, and 54% of all affected dogs were female (Schwarz *et al.*, 1991)

Thyroid neoplasia

- Breed affected more frequently than expected in a series of 30 cases in the UK (Sullivan *et al.*, 1987a)

Primary brain tumour

See under *Neurological conditions*

Neurological conditions

Cerebellar degeneration

- Autosomal recessive inheritance
- Late onset
- Slowly progressive (Steinberg *et al.*, 2000)

Primary brain tumour

- Breed at increased risk in case series. In a USA retrospective study of 86 cases, 1983–1988, OESDs accounted for 4% of cases but < 1% of the hospital population. In more recent studies of 173 cases and 435 cases, an increased risk was not observed
- Older dogs affected; median age 9 years (Heidner *et al.*, 1991; Snyder *et al.*, 2006; Song *et al.*, 2013)

Ocular conditions

Cataract

- Breed at increased risk in case series
- Prevalence of primary cataract 2.61%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Koch, 1972; Gelatt & MacKay, 2005)

Renal and urinary conditions

Urethral sphincter mechanism incompetence

- Breed at increased risk in case series; OESDs were the most commonly affected breed (8/60) in a retrospective review in the UK 1973–1983
- Spayed females predisposed (Holt, 1985; Holt & Thrusfield, 1993)

Urolithiasis – silica

- Breed at increased risk in case series
- Males significantly over-represented (Aldrich *et al.*, 1997)

Reproductive conditions

Cryptorchidism

- Congenital defect believed to be inherited as a sex-limited, autosomal recessive trait
- Breed at increased risk in case series (Hayes *et al.*, 1985; Romagnoli, 1991)

Respiratory conditions

Primary ciliary dyskinesia

- Recessively inherited, mutation identified
- One study of 578 OESDs including 28 affected dogs suggested a carrier frequency of 19% in European dogs and 7% in non-European dogs (Merveille *et al.*, 2014)

OTTERHOUND

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- Breed at increased risk in a cross-sectional study of data from a health survey of purebred dogs conducted by questionnaire, relating to the 10 years prior to 2004
- Prevalence ratio for GDV morbidity was 13.1 (95% CI 5.601–30.40; $p < 0.0001$) compared to all other breeds combined, making it one of the four breeds with highest morbidity (Evans & Adams, 2010b)

Haematological/immunological conditions

Hereditary thrombopathy

- A genetic platelet defect inherited as an autosomal recessive trait. Mutation identified
- Similar to Glanzmann's thrombasthenia in humans (Boudreaux & Lipscomb, 2001)

PAPILLON

Musculoskeletal conditions

Patellar luxation

- Mainly medial luxation observed, often bilateral
- 3.1% of US referred Papillons affected
- Papillon has an OR of 8.4 (95% CI 3.4–20.7) in the USA and is over-represented in Korea (LaFond *et al.*, 2002; Alam *et al.*, 2007; Orthopedic Foundation for Animals, 2015)

Neoplastic conditions

Mammary neoplasia

- Papillon had an incidence of 158 cases 10 000 DYAR compared with 116 per 10 000 DYAR in crossbreeds among insured entire bitches in Sweden

(Egenvall *et al.*, 2005)

Neurological conditions

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Papillon had an incidence of 23.4 per 10 000 DYAR (95% CI 17.8–29.0) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
- Papillon had 3.40% prevalence, compared with 0.91% for crossbreeds in a US referral study

(Bellumori *et al.*, 2013;

Heske *et al.*, 2014)

Intervertebral disc disease (IVDD)

- Males and older dogs predisposed
- Papillon had an incidence rate of 39.5 per 10 000 DYAR (95% CI 32.2–46.9), compared with 27.8 for an overall population of insured dogs in Sweden

(Bergknut *et al.*, 2012)

Necrotizing meningoencephalitis

- Identified in a US Papillon

(Cooper *et al.*, 2014b)

Neuroaxonal dystrophy

- Group of neurodegenerative disorders characterized by severe degeneration of neuronal cells and their processes; can be familial
- Onset from 8 weeks, with euthanasia or death before 12 months
- Several reports in Papillons

(Franklin *et al.*, 1995; Diaz *et al.*, 2007;

Nibe *et al.*, 2007, 2009)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal recessive disease in Papillons
- Late onset reported in Papillons in Sweden
- Frameshift mutation identified in the Papillon and its Phalene variant
- 17.8% of Papillons affected in a Swedish study

(Narfström & Ekesten, 1998;

Narfström & Wrigstad, 1999;

Ahonen *et al.*, 2013; Winkler *et al.*, 2013)

Physiological conditions

Litter size

- Smaller litters associated with older bitches and smaller breeds
- Papillon had the eleventh smallest mean litter size (3.3 puppies) among registered breeds in Norway

(Borge *et al.*, 2011)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- Papillon had OR 9.85 compared to crossbreeds in the USA

(Lekcharoensuk *et al.*, 2000a)

Kidney disease

- Papillon had an incidence of 28 (95% CI 22–34) cases per 10 000 DYAR, compared with 15.8 overall among insured dogs in Sweden

(Pelander *et al.*, 2015)

PARSON RUSSELL TERRIER

Neoplastic conditions

Mast cell tumour (MCT)

- Parson Russell Terrier had the highest breed RR (15.29) using laboratory records in the USA
- Parson Russell Terrier had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland

(Grüntzig *et al.*, 2016; Mochizuki *et al.*, 2016)

Neurological conditions

Spinocerebellar ataxia (hereditary ataxia)

- Incoordination noticed from 2–9 months of age
- Progressive and often leads to euthanasia
- Missense mutation in the *CAPN1* gene, and mutation in *KCNJ10*, identified in the Parson Russell Terrier

(Forman *et al.*, 2013; Rohdin *et al.*, 2015)

Ocular conditions

Glaucoma – secondary

- High intraocular pressure due to primary intraocular diseases that impede aqueous humor flow

- Parson Russell Terrier has the highest standardized morbidity ratio (7.1) compared with an overall referral population in the USA

(Johnsen *et al.*, 2006)

Lens luxation – primary

- *ADAMTS17* mutation has been identified: 24.53% of UK and 22.40% of US Parson Russell Terriers were carriers

(Gould *et al.*, 2011)

Renal and urinary conditions

Hyperuricosuria

- Parson Russell Terrier has 7.75% prevalence of carriers for the mutation in the USA

(Karmi *et al.*, 2010a)

PEKINGESE

Behavioural conditions

Aggression

- Pekingese had 1.56 times the risk of attacking children in Austria compared with the overall population

(Schalamon *et al.*, 2006)

Dental conditions

Abnormal dentition

- The most common specific problems among affected dogs overall were polyodontia (33.2%), retained deciduous teeth (19.7%) and prognathism (10.0%)
- Pekingese comprised the third-highest proportion (15.4%) of an affected caseload aged 7–18 months in Bulgaria

(Borissov *et al.*, 2004)

Gastrointestinal conditions

Congenital portosystemic shunt

- Referred Pekingese (0.36% prevalence) had an OR of 7.1 (95% CI 3.7–13.8) compared with an overall US referral population

(Tobias & Rohrbach, 2003)

Metabolic conditions

Overweight/obesity

- 51.9% of Pekingese attending veterinary clinics in China were obese

(Courcier *et al.*, 2010; Mao *et al.*, 2013;

Raffan *et al.*, 2016)

Musculoskeletal conditions

Chondrodysplasia (short-limbed or disproportional dwarfism)

- Defining Pekingese breed characteristic
- An *fgf4* retrogene identified as a cause of chondrodysplasia

(Israel *et al.*, 2009; Parker *et al.*, 2009)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- OR 3.4 (95% CI 2.7–5.9) in the USA compared with crossbreeds

(LaFond *et al.*, 2002;

Orthopedic Foundation for Animals, 2015)

Neurological conditions

Atlantoaxial subluxation/instability

- Congenital or developmental condition
- Young small-breed dogs affected
- Pekingese comprised 10.5% and 8.7% of two US case-study populations

(Beaver *et al.*, 2000; Platt *et al.*, 2004)

Hemivertebrae (wedge-shaped vertebrae)

- Assumed to be highly heritable
- 100% of Pekingese affected

(Schlensker & Distl, 2013)

Intervertebral disc disease (IVDD)

- Pekingese had 20.59% prevalence, compared with 4.43% for crossbreeds in a US referral caseload
- Pekingese comprised 3.8% of a cervical IVDD caseload and 3.6% of a thoracolumbar IVDD caseload at a Japanese referral hospital
- Pekingese comprised 3.8% and 2.9% of two US referral IVDD caseloads, and 31.9% of a referral population in Turkey
- Pekingese had RR 3.5 ($p < 0.01$) compared with an overall referral population in the USA
- Pekingese was significantly ($p < 0.01$) over-represented, comprising 4.67% of referral cases, in the Czech Republic (Priester, 1976; Nečas, 1999; Bartels *et al.*, 2003; Besalti *et al.*, 2006; Itoh *et al.*, 2008; Israel *et al.*, 2009; Bellumori *et al.*, 2013)

Ocular conditions

Bacterial keratitis

- Pekingese had OR 122.4 (95% CI 24.1–620.5) compared with a US referral population, and comprised 13.2% of referral cases in Serbia (Tolar *et al.*, 2006; Hadži-Milić *et al.*, 2013)

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.14%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Corneal ulceration (ulcerative keratitis)

- Pekingese showed 19.3% prevalence of corneal ulceration among ophthalmic referrals in South Africa
- Pekingese was the second commonest breed, comprising 25% of a referral caseload, in South Korea (Petrick, 1996; Kim *et al.*, 2009)

Epiphora

- Associated with anatomical features such as medial canthal entropion, trichiasis, tight medial palpebral ligament and close apposition of the eyelids to the globe
- Pekingese comprised 17.4% of a surgical referral caseload in Korea (Yi *et al.*, 2006)

Glaucoma – primary

- Pekingese had 1.22% prevalence in a US referral population, 1994–2002 (Gelatt & MacKay, 2004a)

Renal and urinary conditions**Renal calculi**

- May be associated with urinary tract infections
- OR 8.27 ($p < 0.001$) in females and OR 3.97 ($p = 0.054$) in males compared with crossbred dogs in a US referral study (Ling *et al.*, 1998c)

Urolithiasis – apatite

- OR 5.5 (95% CI 4.6–6.5) compared with crossbred dogs in the USA
- Females are predisposed (Low *et al.*, 2010)

Urolithiasis – silica

- OR 4.4 (95% CI 2.8–7.0) compared with crossbred dogs in the USA, and OR 3.3 ($p < 0.001$) compared with all dogs in another US study (Aldrich *et al.*, 1997; Low *et al.*, 2010)

Urolithiasis – struvite (magnesium ammonium phosphate)

- OR 5.3 (95% CI 4.5–6.1) compared with crossbred dogs in the USA, and OR 12.3 (95% CI 8.8–17.2) compared with all dogs in the UK
- Small breeds and males are predisposed (Low *et al.*, 2010; Roe *et al.*, 2012)

Urolithiasis – urate

- OR 7.2 (95% CI 5.8–9.0) compared with crossbred dogs in the USA
- Younger and female animals predisposed (Low *et al.*, 2010)

Reproductive conditions**Cryptorchidism**

- Pekingese had RR 1.9 (95% CI 1.5–2.5) compared with an overall US referral population (Hayes *et al.*, 1985)

Dystocia

- 43.8% of Pekingese litters born by caesarean section in the UK (Evans & Adams, 2010a)

Respiratory conditions**Brachycephalic obstructive airway syndrome (BOAS)**

- Pekingese had a prevalence of 20.0% among referral dogs in Belgium (Njikam Nsangou *et al.*, 2009; Roedler *et al.*, 2013; Marchant *et al.*, 2017)

Tracheal collapse

- Common cause of cough in mature, small-breed dogs
- Pekingese comprised 15.5% of a referral caseload in Korea (Eom *et al.*, 2008)

Soft-tissue conditions**Perineal herniation**

- Pekingese comprised 10.0% of referral cases in Croatia (Vnuk *et al.*, 2008)

PETIT BASSET GRIFFON VENDEEN**Endocrine conditions****Hypothyroidism**

- Females and younger dogs are predisposed to having serum thyroid hormone autoantibodies (THAA) that are associated with hypothyroidism

- Petit Basset Griffon Vendeen had OR 2.16 ($p=0.036$) for THAA compared with all other breeds in the USA

(Nachreiner *et al.*, 2002)

Neoplastic conditions

Mast cell tumour (MCT)

- Mean age at presentation 7.5–9 years, but can occur at any age
- OR 5.1 (95% CI 1.4–18.1) compared with crossbred dogs in the USA

(Goldschmidt & Mcmanus, 2000)

Ocular conditions

Glaucoma – primary

- Prevalence of 10.4% among Petit Basset Griffon Vendeen formally screened in the UK
- Presents from 3 years of age onwards
- Considered to be genetically determined in this breed

(Bedford, 2017)

Neurological conditions

Idiopathic epilepsy

- Prevalence of 8.9% (95% CI 6.3–11.5) in Petit Basset Griffon Vendeen registered with Danish Kennel Club
- Median age at onset of 24 months in this breed
- Hereditary component supported in this breed

(Gulløv *et al.*, 2011)

Physiological conditions

High prevalence of electrocardiographic J waves

- The J wave is a positive deflection at the J point in the ECG and can be classified as notched, slurred or undetermined
- J waves may be considered a normal variant on the canine ECG and should not be interpreted as cardiac disease
- The overall prevalence of J waves in Petit Basset Griffon Vendeen was 91% was significantly higher ($p<0.05$) than the 43% in other breeds in Denmark

(Rudling *et al.*, 2016)

PHU QUOC RIDGEBACK

Physiological conditions

Webbed feet

- Breed comes from Phu Quoc Island in Vietnam's southern Kien Giang Province

- Has webbing between all main digits
- May be an adaptation for an aquatic environment

(Dang *et al.*, 2017)

PINSCHER – MINIATURE

See *Miniature Pinscher*

POINTER (INCLUDING GERMAN SHORT-HAIRED, GERMAN WIRE-HAIRED, ENGLISH)

Dermatological conditions

Blastomycosis

See under *Infectious conditions*

Cutaneous neoplasia

See under *Neoplastic conditions*

Exfoliative cutaneous lupus erythematosus (ECLE)

- Age of onset from 6 months to 2.75 years
- The prognosis is poor because of failure to respond or complications associated with treatment
- Characterized in German Short-haired Pointers from primary and referral veterinary practices in the USA, UK and Australia

(Bryden *et al.*, 2005)

Familial cutaneous lupus erythematosus (CLE)

- Typically presents before 10 months of age
- Therapy is rarely rewarding and cases often lead to euthanasia
- Simple autosomal recessive mode of inheritance identified
- 8.5% of a US targeted sample of German Short-haired Pointers were affected

(Wang *et al.*, 2011)

Hereditary lupoid dermatosis (exfoliative cutaneous lupus erythematosus)

- Autosomal recessive or polygenic recessive inheritance suggested
- Characterized in German Short-haired Pointer across several countries

(Bryden *et al.*, 2005)

Junctional epidermolysis bullosa (JEB)

- Recessive inherited blistering disorder of the skin and mucous membranes

- Genetic basis identified in the German Short-haired Pointer

(Capt *et al.*, 2005)

Malassezia dermatitis

- German Short-haired Pointer had the seventh-highest prevalence (4.1%) in a dermatology referral caseload in Romania

(Mircean *et al.*, 2010)

Drug reactions

Anaesthetic-related complications

- German Short-haired Pointer had 5% incidence among general small animal practices in Canada

(Dyson *et al.*, 1998)

Endocrine conditions

Hypothyroidism

- Females and younger dogs are predisposed to having serum thyroid hormone autoantibodies (THAA) that are associated with hypothyroidism
- Pointer (unspecified) had OR 3.61 ($p=0.001$), English Pointer OR 3.31 ($p=0.01$), and German Wire-haired Pointer OR 2.72 ($p=0.01$) for THAA compared with all other breeds in the USA

(Nachreiner *et al.*, 2002)

Gastrointestinal conditions

Primary hepatitis

- Multiple causes including microorganisms, toxins and drugs, immune-mediated reactions and breed-associated metabolic errors
- Females predisposed ($p<0.01$)
- German Pointer was over-represented in a referral population in the Netherlands

(Poldervaart *et al.*, 2009)

Haematological/immunological conditions

von Willebrand's disease (vWD)

- German Short-haired and German Wire-haired Pointers affected by type II disease
- Polymerase chain reaction (PCR) test developed

(Kramer *et al.*, 2004)

Infectious conditions

Blastomycosis

- Young, large-breed dogs predisposed
- Pointer (unspecified) was at increased risk compared with referral crossbred dogs in the USA

(Rudmann *et al.*, 1992; Arceneaux *et al.*, 1998)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Estimated heritability of 0.48 and a highly polygenic complex trait
- German Short-haired Pointer had an OR of 1.20 (95% CI 1.06–1.35) compared with an overall referral population in the USA
- German Short-haired Pointer had 5.63% prevalence, compared with 1.58% overall in referred dogs in the Czech Republic

(Nečas *et al.*, 2000; Witsberger *et al.*, 2008; Baker *et al.*, 2017)

Enchondrodystrophy

- Inherited dwarfism in English Pointers
- 3.3% of English Pointers affected in an older UK study

(Whitbread *et al.*, 1983)

Hip dysplasia

- German Wire-haired Pointer had 9.2% prevalence in the USA
- German Wire-haired Pointer had OR 7.1 (95% CI 3.3–15.1) and Pointer (unspecified) OR 2.4 (95% CI 1.4–4.2) compared with crossbred dogs in the USA

(LaFond *et al.*, 2002; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Osteochondrosis – shoulder

- The coefficient of heritability varies from 0.25 to 0.45
- Age of onset usually 4–7 months, the period of rapid growth
- Males predisposed
- German Short-haired Pointer had OR 5.5 (95% CI 3.0–10.3) and German Wire-haired Pointer OR 38.8 (95% CI 16.7–90.3) compared to crossbred dogs in the USA

(Rudd *et al.*, 1990; Nečas *et al.*, 1999; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- German Short-haired Pointer had an OR of 1.6 (95% CI 1.1–2.4) compared with crossbred dogs in the USA

(LaFond *et al.*, 2002)

Neoplastic conditions

Cutaneous haemangiosarcoma

- English Pointer had an OR of 2.3 (95% CI 1.3–3.9) compared to a US referral population

(Villamil *et al.*, 2011)

Cutaneous neoplasia

- Median age of 9 years
- English Pointer comprised 7.5% of purebred cases in a referral hospital in Greece (Kaldrymidou *et al.*, 2002)

Haemangiosarcoma

- English Pointer had OR 9.03 (95% CI 1.38–59.10) of haemangiosarcoma compared with crossbred dogs in the USA
- Two studies showed English Pointer had OR 2.25 (95% CI 1.30–3.90) and OR 3.65 (95% CI 1.50–8.86) of cutaneous haemangiosarcoma compared with crossbred dogs in the USA (Hargis *et al.*, 1992; Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011)

Mammary neoplasia

- Pointer (unspecified) had an incidence of 18.4 benign mammary cases and 12.9 malignant mammary cases per 1000 dogs in the USA, significantly higher than the general population ($p < 0.05$) (MacVean *et al.*, 1978)

Mast cell tumour (MCT)

- Mean age at presentation 7.5–9 years, but can occur at any age
- Two US studies showed Pointer had OR 1.7 (95% CI 1.1–2.6) and English Pointer OR 1.84 (95% CI 1.31–2.60) compared with crossbred dogs (Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011)

Prostate neoplasia

- Neutering associated with 3.56 times increased odds
- German Short-haired Pointer had OR 1.89 ($p = 0.01$) compared with an overall referral population in the USA
- German Pointer had OR 2.19 (95% CI 1.16–4.06) compared with an overall referral population in the Netherlands (Teske *et al.*, 2002; Bryan *et al.*, 2007)

Neurological conditions**Acquired myasthenia gravis**

- German Short-haired Pointers had significantly higher risk than crossbred dogs in the USA (Shelton *et al.*, 1997)

Discospondylitis

- Males and older dogs predisposed
- Pointer (unspecified) had OR 6.2 (95% CI 2.4–15.9) compared with referral crossbred dogs in the USA (Burkert *et al.*, 2005)

Hemivertebrae (wedge-shaped vertebrae)

- Assumed to be highly heritable
- 34.5% of German Short-haired Pointers affected (Kramer *et al.*, 1982)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- German Smooth and Wire-haired Pointer had an incidence of 17.7 cases per 10 000 DYAR (95% CI 12.8–22.7) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds (Heske *et al.*, 2014)

Lipoidystrophy (amaurotic idiocy)

- Affects young males from 10–18 months
- Generally leads to euthanasia
- Recessive sex-linked mode of transmission suspected in male German Short-haired Pointers in the USA (Karbe & Schiefer, 1967)

Sensory neuropathy (acral mutilation syndrome)

- Affects English Pointer and German Short-haired Pointer
- Presents at about 4 months, when affected pups suddenly begin to self-mutilate their paws (Cummings *et al.*, 1983, 1984)

Ocular conditions**Cone degeneration (hemeralopia or day blindness)**

- Autosomal recessive inheritance
- A missense mutation in exon 6 (D262N, nucleotide 784) identified in German Short-haired Pointers (Sidjanin *et al.*, 2002)

Eversion of the cartilage of the nictitating membrane

- Described in a line of German Short-haired Pointers
- Recessive inheritance suggested (Martin & Leach, 1970)

Physiological conditions

Litter size

- Larger litters associated with younger bitches and larger breeds
- German Short-haired Pointer had the fourth largest mean litter size (8.3 puppies) and the German Wire-haired Pointer had the thirteenth largest mean litter size (7.3 puppies) among registered breeds in Norway (Borge *et al.*, 2011)

Reproductive conditions

Dystocia

- 26% of Pointer (unspecified) litters born by caesarean section in the UK (Evans & Adams, 2010a)

XX Sex reversal

- Highly likely to be inherited disorder in German Short-haired Pointers (Meyers-Wallen *et al.*, 1995)

Respiratory conditions

Primary ciliary dyskinesia

- Clinical signs are apparent early in life
- Described in related English Pointers in the USA (Morrison *et al.*, 1987)

POLISH HOUND (POLISH OGAR DOG, OGAR POLSKI)

Musculoskeletal conditions

Elbow dysplasia

- Polish hound had 3.6% prevalence among radiographed dogs in Poland, compared with 0.7% in the overall population (Narojek *et al.*, 2008; Michelsen, 2013)

Ocular conditions

Distichiasis

- Prevalence of 40.5% in Polish Hounds in Poland (Ksiazek *et al.*, 2014)

Entropion

- Prevalence of 23.0% in Polish Hounds in Poland (Ksiazek *et al.*, 2014)

Physiological conditions

Low thrombocyte count (physiological thrombocytopenia)

- Mean platelet count in normal Polish Hounds ($167 \pm 11.6 \times 10^9/l$) was lower than in other breeds combined ($344.4 \pm 6.85 G/l$)

- Associated with genetic bottleneck after the Second World War

(Micun *et al.*, 2009)

POLISH LOWLAND SHEEPDOG (POLSKI OWCZAREK NIZINNY)

Cardiovascular conditions

Atrioventricular (AV) block

- Polish Lowland Sheepdog had the highest prevalence (35.7%) of first-degree atrioventricular blocks among referred cardiology cases in Poland (Noszczyk-Nowak *et al.*, 2017)

Patent ductus arteriosus

- Polish Lowland Sheepdog comprised 30.8% of a referral caseload in Germany
- Median age at presentation was 5.7 months (Schneider *et al.*, 2001)

Musculoskeletal conditions

Short tail (bobtail)

- 64.3% of Polish Lowland Sheepdogs tested in Finland affected (Hytönen *et al.*, 2009)

Neurological conditions

Neuronal ceroid lipofuscinosis

- Described in related Polish Lowland Sheepdogs in Sweden
- Appeared normal at birth, with progressive neurological signs developing from 6 months (Narfström *et al.*, 2007)

Ocular conditions

Progressive retinal atrophy (PRA)

- The rcd4 mutation in the *C2ORF71* gene was associated with the majority of PRA in the Polish Lowland Sheepdog in Sweden (Svensson *et al.*, 2016)

POMERANIAN

Cardiovascular conditions

Patent ductus arteriosus

- Small breeds (median body weight 5 kg) and females predisposed
- Median age was 4 months

- Pomeranian comprised 7.3% and 7.7% of two US referral caseloads
(Bureau *et al.*, 2005; Goodrich *et al.*, 2007)

Dermatological conditions

Alopecia X

- Pomeranian comprised 79.3% of a referral caseload in Japan
- Pomeranian was the most commonly affected breed, comprising 26.5% of an endocrine alopecia caseload of neutered dogs in the USA
(Takada *et al.*, 2002; Frank *et al.*, 2003, 2004)

Demodicosis

- An inheritance pathway has been described
- Short-haired breeds and dogs aged <1 year predisposed
- Pomeranian had a 26.19% prevalence among dermatitis cases that were skin scraped in India
(Solanki *et al.*, 2007; It *et al.*, 2010)

Drug reactions

Anaesthetic-related complications

- Pomeranian had 4% incidence among general small animal practices in Canada
(Dyson *et al.*, 1998)

Endocrine conditions

Hypothyroidism

- Pomeranian was at increased risk in a US referral study
(Milne & Hayes, 1981)

Gastrointestinal conditions

Congenital portosystemic shunt

- Referred Pomeranians (0.29% prevalence) had an OR of 5.6 (95% CI 2.7–11.6) compared with an overall US referral population
(Tobias & Rohrbach, 2003)

Gallbladder mucocoele

- Associated with hyperlipidaemia
- Older dogs predisposed
- Pomeranian had an OR of 7.74 (95% CI 5.43–21.03) compared with an overall referral population in Japan
(Kutsunai *et al.*, 2014)

Infectious conditions

Salmonellosis

- 7.7% of dogs may harbour *Salmonella* when apparently healthy, subclinically affected, as latent infection or as carriers

- Pomeranians had 45.56% seroprevalence among apparently healthy dogs in India
(Verma *et al.*, 2011)

Metabolic conditions

Overweight/obesity

- 54.6% of Pomeranians attending veterinary clinics in China were obese
(Courcier *et al.*, 2010; Mao *et al.*, 2013; Raffan *et al.*, 2016)

Musculoskeletal conditions

Elbow dysplasia

- OR 3.7 (95% CI 1.7–8.1) for ununited anconeal process compared to crossbred dogs in the USA
(LaFond *et al.*, 2002; Michelsen, 2013)

Patellar luxation

- Genetic mechanism suggested
- 6.5% of Pomeranians affected in the UK
- Mainly medial luxation observed, often bilateral
- OR 6.5 (95% CI 4.0–10.7) in the UK and OR 18.6 (95% CI 13.1–26.4) in the USA, compared to crossbreeds
- Pomeranian was over-represented in Korea and comprised 28.9% of a caseload in Thailand and 9.3% of a US caseload
- Pomeranian had a 75% prevalence in Thailand
(LaFond *et al.*, 2002; Alam *et al.*, 2007; Campbell *et al.*, 2010; Nganvongpanit & Yano, 2011; Soontornvipart *et al.*, 2013; Orthopedic Foundation for Animals, 2015; O'Neill *et al.*, 2016c)

Neurological conditions

Atlantoaxial subluxation/instability

- Congenital or developmental condition
- Young small-breed dogs affected
- In case studies, Pomeranian comprised 10.6% in Japan, 46.7% in the UK, and 10.5% and 6.5% of two US populations
(Denny *et al.*, 1988; Beaver *et al.*, 2000; Platt *et al.*, 2004; Aikawa *et al.*, 2013)

Intervertebral disc disease (IVDD)

- Pomeranian comprised 6.1% of a cervical IVDD caseload and 1.2% of a thoracolumbar IVDD caseload at a Japanese referral hospital
(Itoh *et al.*, 2008)

Physiological conditions

Litter size

- Smaller litters associated with older bitches and smaller breeds
- Pomeranians had the second smallest mean litter size (2.4 puppies) among registered breeds in Norway

(Borge *et al.*, 2011)

Vertebral heart score (VHS)

- Pomeranians from US referral practices had significantly greater ($p=0.0014$) radiographic vertebral heart score (size) than a previously established average of 9.7 ± 0.5

(Buchanan & Bücheler, 1995;
Jepsen-Grant *et al.*, 2013)

Renal and urinary conditions

Hyperuricosuria

- Pomeranian had 1.13% prevalence of carriers for the mutation in the USA

(Karmi *et al.*, 2010a)

Urolithiasis – calcium oxalate

- OR 7.2 (95% CI 6.3–8.2) compared with crossbred dogs in the USA
- OR 9.85 compared with crossbred dogs in the USA

(Lekcharoensuk *et al.*, 2000a; Low *et al.*, 2010;
Okafor *et al.*, 2014)

Reproductive conditions

Cryptorchidism

- Pomeranian had RR 3.1 (95% CI 2.3–4.0) and OR 2.9 (1.94–4.36) compared with overall referral populations in two US studies

(Pendergrass & Hayes, 1975;
Hayes *et al.*, 1985)

Dystocia

- Pomeranian had the fourth-highest incidence (29.5 per 1000 DYAR), compared with 5.7 per 1000 DYAR in the overall population of insured bitches in Sweden

(Bergström *et al.*, 2006)

Eclampsia (puerperal tetany)

- Small-sized bitches predisposed
- Pomeranian was over-represented and comprised 6% of a US referral caseload

(Drobatz & Casey, 2000)

Respiratory conditions

Tracheal collapse

- Common cause of cough in mature small-breed dogs
- Pomeranian comprised 6%, 13.5% and 16.7% of three US referral caseloads, and 11.9% of a referral caseload in Korea
- Pomeranian had a 9.3% prevalence in a referral population in Australia

(Amis, 1974; Johnson & Fales, 2001;
Macready *et al.*, 2007; Marolf *et al.*, 2007;
Eom *et al.*, 2008)

POODLE (UNSPECIFIED VARIANT)

Behavioural conditions

Aggression

- Poodle (unspecified) was significantly over-represented ($p<0.01$), comprising 5.7% of cases compared with 2.9% of an underlying hospital population in Australia

(Blackshaw, 1991)

Anxiety

- Poodle (unspecified) had a significantly higher risk than other breeds in Denmark

(Lund *et al.*, 1996)

Victim of dog-to-dog aggression

- Poodle (unspecified) was over-represented as the victim in dog–dog conflict in Germany, representing 6.8% of victims and 2.8% of the general population

(Roll & Unshelm, 1997)

Cardiovascular conditions

Left systolic apical murmur

- Poodle (unspecified) had 22.7% (± 6.2) prevalence compared with 14.4% (± 2.2) in an overall population of small-sized breeds in France

(Serfass *et al.*, 2006)

Mitral valve disease

- Poodle (unspecified) had OR 3.9 (95% CI 1.9–8.1) compared with crossbred dogs in UK primary-care veterinary population
- Poodle (unspecified) comprised 18% of an affected caseload, compared with 8% of healthy controls, from a general population in France

(Serres *et al.*, 2008;
Parker & Kilroy-Glynn, 2012;
Mattin *et al.*, 2015a, 2015b)

Ruptured chordae tendineae

- Recorded among dogs diagnosed with mitral valve disease
- Poodle (unspecified) comprised 34.2% of a referral caseload in France

(Serres *et al.*, 2007)**Dental conditions****Abnormal dentition**

- The most common specific problems among affected dogs overall were polyodontia (33.2%), retained deciduous teeth (19.7%) and prognathism (10.0%)
- Poodle (unspecified) comprised the second-highest proportion (21.2%) of an affected caseload aged 7–18 months in Bulgaria

(Borissov *et al.*, 2004)**Periodontal disease**

- Poodle (unspecified) had the most severe periodontal destruction of breeds necropsied from general practice in Sweden

(Hamp *et al.*, 1997)**Dermatological conditions****Atopic dermatitis (atopy)**

- Poodles (unspecified) were over-represented in Hungary

(Tarpataki *et al.*, 2006)**Endocrine alopecia**

- Poodle (unspecified) was the most second commonly affected breed, comprising 6.3% of an endocrine alopecia caseload of neutered dogs in the USA

(Frank *et al.*, 2003)**Endocrine conditions****Diabetes mellitus**

- Poodle (unspecified) had an OR of 2.80 ($p < 0.001$) compared with a referral population in Italy

(Fracassi *et al.*, 2004)**Gastrointestinal conditions****Colitis**

- Poodle (unspecified) was the third most commonly affected breed (7.5% of cases) in a referral caseload in the Netherlands

(van der Gaag, 1988)

Sialocoele

- Poodle (unspecified) was over-represented ($p < 0.01$), comprising 18.3% of cases compared with 3.1% of a referral population in Australia (Bellenger & Simpson, 1992)

Haematological/immunological conditions**Immune-mediated thrombocytopenia (IMTP)**

- Poodles (unspecified) comprised 21% of the caseload and had RR 4.0 ($p < 0.001$) compared to a US referral hospital population

(Grindem *et al.*, 1991)**Musculoskeletal conditions****Lumbosacral transitional vertebrae**

- Poodle (unspecified) had 19.0% prevalence, compared with an overall 10.0% prevalence, among dogs screened for pelvic problems in the Czech Republic

(Fialová *et al.*, 2014)**Patellar luxation**

- Mainly medial luxation observed, often bilateral
- Poodle (unspecified) was over-represented in Korea
- Poodle (unspecified) comprised 34.3% of a caseload in Thailand
- Poodle (unspecified) comprised 44.4% of a medial patellar luxation caseload in Brazil

(Alam *et al.*, 2007; Mortari *et al.*, 2009; Nganvongpanit & Yano, 2011)**Neoplastic conditions****Adenoma/adenocarcinoma**

- Poodle (unspecified) had an OR significantly above 1.0 ($p < 0.05$) compared with cross-breeds from a cancer registry in Switzerland

(Grüntzig *et al.*, 2016)**Haemangiosarcoma**

- Poodle (unspecified) was the second most commonly affected breed in the USA

(Brown *et al.*, 1985)**Mammary neoplasia**

- Poodle (unspecified) had OR 2.5 for mammary gland tumour and OR 2.7 for malignant mammary gland tumour compared with an overall referral population in the Czech Republic ($p < 0.01$)

(Zatloukal *et al.*, 2005)

Mast cell tumour (MCT)

- Median age 9 years
- Poodle (unspecified) comprised 7% of a US referral caseload of cutaneous MCT

(Patnaik *et al.*, 1984a)**Melanoma**

- Poodle (unspecified) had an OR significantly above 1.0 ($p < 0.05$) compared with cross-breeds from a cancer registry in Switzerland

(Grüntzig *et al.*, 2016)**Squamous cell carcinoma – lingual**

- Females predisposed (OR 2.26; 95% CI 1.45–3.54)
- Poodle (unspecified) had 4.61 (95% CI 1.91–11.13) OR compared with other breeds with lingual neoplasia at a US diagnostic laboratory

(Dennis *et al.*, 2006)**Neurological conditions****Atlantoaxial subluxation/instability**

- Congenital or developmental condition
- Young small-breed dogs affected
- Poodle (unspecified) comprised 26.1% of a US caseload

(Beaver *et al.*, 2000)**Discospondylitis**

- Males and older dogs predisposed
- Poodle (unspecified) had OR 6.7 (95% CI 2.1–21.4) compared with referral crossbreeds in the USA

(Burkert *et al.*, 2005)**Intervertebral disc disease (IVDD)**

- Poodle (unspecified) was significantly over-represented ($p < 0.01$), comprising 1.67% of referral cases in the Czech Republic

(Nečas, 1999)

Ocular conditions**Cataract**

- Poodles (unspecified) showed 41.3% prevalence of cataract among ophthalmic referrals in South Africa

(Petrick, 1996)

Glaucoma – secondary

- High intraocular pressure due to primary intraocular diseases that impede aqueous humor flow

- Poodle (unspecified) had the second-highest standardized morbidity ratio (4.7) compared with an overall referral population in the USA

(Johnsen *et al.*, 2006)**Myopia (short-sightedness)**

- Poodle (unspecified) had a low mean refractive error (−0.38) in the USA

(Murphy *et al.*, 1992)**Renal and urinary conditions****Urolithiasis**

- Poodle (unspecified) was the third most common breed submitted for urolith analysis overall in Germany (9.1%) and the fourth most common breed from Spain and Portugal (5.1%)

(Hesse, 1990; Vrabelova *et al.*, 2011)**POODLE – MINIATURE****Cardiovascular conditions****Mitral valve disease**

- Miniature Poodle comprised 9.0% of a referral caseload in Italy

(Borgarelli *et al.*, 2008)**Patent ductus arteriosus**

- Median age at presentation was 4 months
- Small breeds and females appear predisposed
- Miniature Poodle comprised 8.6% of a US referral caseload
- Miniature and Toy Poodle comprised 5.4% of a US referral caseload

(Goodrich *et al.*, 2007; Selmic *et al.*, 2013)**Endocrine conditions****Diabetes mellitus**

- Familial and inherited
- Older entire females are predisposed
- Miniature Poodle showed OR 1.79 (95% CI 1.55–2.06) compared with crossbred dogs in the USA
- Miniature Poodle had OR 4.0 (95% CI 2.1–7.1) compared with crossbred dogs in a US referral study
- Miniature and Toy Poodle had 11th-highest breed incidence in Sweden: 24 cases per 10 000 DYAR (95% CI 16–32)

(Hess *et al.*, 2000; Guptill *et al.*, 2003; Fall *et al.*, 2007; Catchpole *et al.*, 2008)

Hyperadrenocorticism (Cushing's syndrome)

- Miniature Poodle comprised 15.9% of a referral caseload of pituitary-dependent hyperadrenocorticism in Spain (Alenza *et al.*, 2006)

Gastrointestinal conditions**Cleft lip and/or palate**

- Birth defect, small breeds predisposed
- Miniature and Toy Poodle had RR 1.3 compared with all referral dogs in the USA (Mulvihill *et al.*, 1980)

Portal vein hypoplasia

- May occur in conjunction with portosystemic shunt (PSS)
- Cases present from 6 months to 6 years
- Miniature and Toy Poodle comprised 10.7% of a US referral caseload (Christiansen *et al.*, 2000)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- Inherited as an autosomal recessive trait
- Miniature Poodle had an OR of 12.1 (95% CI 6.5–22.4) compared to crossbreeds in the USA (Brenig *et al.*, 1999; LaFond *et al.*, 2002)

Cranial cruciate ligament (CCL) disease

- Estimated heritability of 0.48 and is a highly polygenic complex trait
- Miniature Poodle had an OR of 1.17 (95% CI 1.11–1.24) compared with an overall referral population in the USA (Witsberger *et al.*, 2008; Baker *et al.*, 2017)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- Miniature Poodle had OR 4.1 (95% CI 3.3–5.5) in the USA
- Miniature and Toy Poodles had OR 5.62 (95% CI 1.93–16.41) compared with all other small breeds in Austria (LaFond *et al.*, 2002; Vidoni *et al.*, 2006)

Neoplastic conditions**Cutaneous neoplasia**

- Median age 9 years
- Miniature Poodle comprised 19.4% of pure-bred cases in a referral hospital in Greece (Kaldrymidou *et al.*, 2002)

Mammary neoplasia

- Miniature Poodle had OR 2.5 (95% CI 2.3–2.9) for mammary gland carcinoma compared with an overall US referral population (Langenbach *et al.*, 1998)

Neurological conditions**Atlantoaxial subluxation/instability**

- Congenital or developmental condition
- Young small-breed dogs affected
- Miniature Poodle comprised 25% of a US referral caseload (Sanders *et al.*, 2004)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Medium and Miniature Poodle had an incidence of 23.3 per 10 000 DYAR (95% CI 23.3–31.7) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds (Heske *et al.*, 2014)

Intervertebral disc disease (IVDD)

- Miniature Poodle comprised 26.1% of a referral population in Turkey (Besalti *et al.*, 2006)

Mucopolysaccharidosis IV

- Autosomal recessive inheritance suspected
- Congenital disorder, and many malformed pups may be euthanized without investigation
- Described in a Miniature Poodle in the USA (Jolly *et al.*, 2012)

Ocular conditions**Cataract**

- Prevalence 10.79%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Prevalence of primary cataract 13.8% in Brazil
- Miniature/Toy Poodles had OR 2.6 (95% CI 2.1–3.2) for cataract compared to a hospital population in Korea
- Miniature Poodle had 21.49% prevalence, compared with 4.04% for crossbreeds, in a US referral study
- Miniature Poodles had OR 4.3 compared with an overall referral population in the USA (Adkins & Hendrix, 2005; Gelatt & MacKay, 2005; Baumworcel *et al.*, 2009; Park *et al.*, 2009; Bellumori *et al.*, 2013)

Distichiasis

- Heritability of 0.043
- Miniature Poodle comprised 20% of a distichiasis caseload in France
(Ketteritzsch *et al.*, 2004; Raymond-Letron *et al.*, 2012)

Glaucoma – primary

- Miniature Poodle had 1.68% prevalence in a US referral population, 1994–2002
(Gelatt & MacKay, 2004a)

Keratoconjunctivitis sicca

- Miniature Poodle was the third most common breed in a European multicentre study, accounting for 11% of cases
(Ofri *et al.*, 2009)

Lens luxation – primary

- Miniature Poodle had RR 4.51 (95% CI 3.67–5.55) compared with overall referral dogs in the USA
(Sargan *et al.*, 2007)

Optic nerve hypoplasia

- Described in related Miniature Poodles
- Congenital, inheritance suspected
(Kern & Riis, 1981)

Progressive retinal atrophy (PRA)

- Genetic mechanism identified in Miniature Poodle
- PRA mutation identified in Miniature Poodle in the UK
(Goldstein *et al.*, 2006; Downs *et al.*, 2014a)

Progressive rod–cone degeneration (PRCD)

- PRCD gene mapped to the centromeric region of canine chromosome 9 (CFA9)
- Miniature Poodle has 0.45 allele frequency of the disease-causing mutation
(Dostal *et al.*, 2011)

Physiological conditions**Litter size**

- Smaller litters associated with older bitches and smaller breeds
- Miniature Poodle had the fourth smallest mean litter size (3.0 puppies) among registered breeds in Norway
(Borge *et al.*, 2011)

Renal and urinary conditions**Ectopic ureter**

- Miniature/Toy Poodle bitches had RR 3.6 (95% CI 2.5–4.8) and RR 3.26 (95% CI 2.26–7.93) compared with overall referral populations in two US studies
(Hayes, 1974b, 1984)

Urinary tract infections (UTI)

- Miniature and Toy Poodle (7.8%) was the second most commonly affected breed among referral dogs in the USA for recurrent or persistent UTI
(Norris *et al.*, 2000)

Urolithiasis – calcium oxalate

- Small breeds and males predisposed
- Miniature Poodle had OR 15.2 (95% CI 10.7–21.6) compared with all dogs in the UK
- Miniature and Toy Poodle had OR 3.32 in the USA
(Lekcharoensuk *et al.*, 2000a; Houston *et al.*, 2004; Roe *et al.*, 2012)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Smaller breeds and females are predisposed
- Miniature Poodle had OR 12.7 (95% CI 9.2–17.8) compared with all dogs in the UK
(Houston & Moore, 2009; Roe *et al.*, 2012)

Reproductive conditions**Cryptorchidism**

- Miniature Poodle had RR 1.8 (95% CI 1.5–2.1) compared with an overall US referral population
- Miniature and Toy Poodles had OR 2.8 (95% CI 2.43–3.23) compared with an overall US referral population
- Miniature Poodle had a 9.0% prevalence in the UK, which was significantly higher than for crossbred dogs ($p=0.024$)
(Pendergrass & Hayes, 1975; Hayes *et al.*, 1985; Yates *et al.*, 2003)

Testicular neoplasia

- Miniature and Toy Poodle had a proportional morbidity ratio of 1.9 (95% CI 1.1–3.2) compared with all other dogs in a Norwegian cancer registry
(Nødtevd *et al.*, 2011)

Eclampsia (puerperal tetany)

- Small-sized bitches predisposed
- Miniature Poodle was over-represented and comprised 6% of a US referral caseload (Drobatz & Casey, 2000)

Respiratory conditions**Bronchiectasis**

- Miniature Poodle had OR 2.88 (95% CI 1.71–4.84) compared with an overall US referral population (Hawkins *et al.*, 2003)

Tracheal collapse

- Miniature Poodles comprised 8.1% of a US referral caseload and 14.3% of a referral caseload in Korea (Johnson & Fales, 2001; Eom *et al.*, 2008)

POODLE – STANDARD**Cardiovascular conditions****Dilated cardiomyopathy (DCM)**

- Males predisposed
- Standard Poodle was over-represented (Tidholm & Jonsson, 1997)

Dermatological conditions**Sebaceous adenitis**

- Autosomal recessive inheritance with variable expression
- Median age at diagnosis 5 years
- Standard Poodle had the third-highest breed-relative risk in Sweden and comprised 20.2% of a general veterinary caseload (Tevell *et al.*, 2008; Pedersen *et al.*, 2012b)

Endocrine conditions**Hypoadrenocorticism (Addison's disease)**

- Complex inheritance pattern likely: estimated heritability 0.75 in Standard Poodle
- Females predisposed
- Standard Poodle had OR 8.90 (95% CI 4.43–17.32) compared with an overall US referral population
- Standard Poodle comprised 9.8% of a US referral caseload
- Standard Poodle had second-highest breed incidence rate in insured dogs in Sweden: 33.4 (95% CI 26.6–41.9) cases per 10 000

DYAR, compared with 2.26 (95% CI 2.07–2.46) across all dogs

(Melián & Peterson, 1996; Peterson *et al.*, 1996; Famula *et al.*, 2003; Oberbauer *et al.*, 2006; Hanson *et al.*, 2016)

Gastrointestinal conditions**Congenital portosystemic shunt**

- Referred Standard Poodles (0.17% prevalence) had OR 3.4 (95% CI 1.6–7.5) compared with an overall US referral population (Tobias & Rohrbach, 2003)

Gastric dilatation/volvulus (bloat, GDV)

- Pedigree Standard Poodles had 3.1% prevalence in the UK
- Standard Poodle had an incidence of 24 cases per 1000 DYAR (95% CI 6–42) in the USA, with 2.4% of dogs affected per year
- Standard Poodles represented 7.8% of US referral caseloads in two studies and were over-represented in two other US studies (Glickman *et al.*, 2000a; Beck *et al.*, 2006; Evans & Adams, 2010b; Mackenzie *et al.*, 2010; Green *et al.*, 2012; Sartor *et al.*, 2013)

Musculoskeletal conditions**Osteochondrosis – shoulder**

- The coefficient of heritability varies from 0.25 to 0.45
- Age of onset usually 4–7 months, the period of rapid growth
- Males predisposed
- Standard Poodle had an OR of 2.6 (95% CI 1.3–5.3) compared to crossbreeds in the USA (Rudd *et al.*, 1990; Nečas *et al.*, 1999; LaFond *et al.*, 2002)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- Standard Poodle had OR 3.2 (95% CI 2.2–4.7) compared to crossbreeds in the USA (LaFond *et al.*, 2002; Alam *et al.*, 2007)

Neoplastic conditions**Gastric carcinoma**

- Median age at diagnosis of 10 years
- Standard Poodle had a proportional morbidity ratio of 7.6 (95% CI 2.3–25.0) compared with all breeds in a Norwegian cancer registry (Seim-Wikse *et al.*, 2013)

Squamous cell carcinoma – digit

- Middle-aged to older dogs and dogs with black coats predisposed
- Standard Poodle comprised 9.1% and 14.3% of two US affected caseloads
(O'Brien *et al.*, 1992; Henry *et al.*, 2005)

Vertebral tumours

- Mean age of 7 years at diagnosis
- Standard Poodles were heavily represented
(Morgan *et al.*, 1980)

Neurological conditions**Idiopathic epilepsy**

- Hereditary basis established in several breeds
- Standard Poodle had an incidence of 20.9 per 10 000 DYAR (95% CI 15.2–26.6) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
- Standard Poodle had 3.19% prevalence compared with 0.91% for crossbreeds in a US referral study
(Bellumori *et al.*, 2013; Heske *et al.*, 2014)

Neonatal encephalopathy with seizures (NEWS)

- Autosomal recessive inheritance
- NEWS locus mapped to a 2.87 Mb segment of CFA36 containing the canine ortholog of ATF2 in the Standard Poodle in the USA
(Chen *et al.*, 2008)

Ocular conditions**Cataract**

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 7.00%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Progressive retinal atrophy (PRA)

- PRA mutation identified in Standard Poodle in the UK
(Downs *et al.*, 2014a)

Renal and urinary conditions**Kidney disease**

- Standard Poodle had an incidence of 28 (95% CI 21–35) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall, among insured dogs in Sweden
(Pelander *et al.*, 2015)

Reproductive conditions**Cryptorchidism**

- Standard Poodle had RR 1.6 (95% CI 1.1–2.3) compared with an overall US referral population
(Pendergrass & Hayes, 1975)

POODLE – TOY**Cardiovascular conditions****Patent ductus arteriosus**

- Small breeds and females appear predisposed
- Median age at presentation was 4 months
- Toy Poodle was the second most common breed, comprising 9.6% of a US referral caseload
- Toy and Miniature Poodle comprised 5.4% of a US referral caseload
(Bureau *et al.*, 2005; Goodrich *et al.*, 2007)

Endocrine conditions**Diabetes mellitus**

- Familial and inherited
- Peak incidence age 7–9 years
- Older entire females are predisposed
- Toy Poodle had OR 3.3 (95% CI 1.4–6.7) compared with crossbreeds in a US referral study
- Miniature and Toy Poodle had 11th-highest breed incidence in Sweden: 24 cases per 10 000 DYAR (95% CI 16–32)
(Guptill *et al.*, 2003; Hess *et al.*, 2003; Fall *et al.*, 2007; Catchpole *et al.*, 2008)

Primary hypoparathyroidism

- Toy Poodles comprised 17.9% of a US referral case series
(Bruyette & Feldman, 1988)

Gastrointestinal conditions**Cleft lip and/or palate**

- Birth defect, small breeds predisposed
- Toy and Miniature Poodle had RR 1.3 compared with all referral dogs in the USA
(Mulvihill *et al.*, 1980)

Congenital portosystemic shunt

- Referred Toy Poodles (0.16% prevalence) had OR 3.1 (95% CI 1.6–6.0) compared with an overall US referral population
(Tobias & Rohrbach, 2003)

Portal vein hypoplasia

- May occur in conjunction with portosystemic shunt (PSS)
- Cases present from 6 months to 6 years
- Toy and Miniature Poodle comprised 10.7% of a US referral caseload

(Christiansen *et al.*, 2000)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- Inherited as an autosomal recessive trait
- Toy Poodle had OR 22.4 (95% CI 11.7–42.8) compared to crossbreeds in the USA in the USA (Pidduck & Webbon, 1978; Brenig *et al.*, 1999; LaFond *et al.*, 2002)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- Toy and Miniature Poodles had OR 5.62 (95% CI 1.93–16.41) compared with all other small breeds in Austria
- Toy Poodle had OR 9.7 (95% CI 7.6–12.3) in the USA and was over-represented in Korea (LaFond *et al.*, 2002; Vidoni *et al.*, 2006; Alam *et al.*, 2007)

Neurological conditions**Atlantoaxial subluxation/instability**

- Congenital or developmental condition
- Young small-breed dogs affected
- Toy Poodle comprised 12.8% of a Japanese caseload (Aikawa *et al.*, 2013)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Toy Poodles comprised 6.4% of epilepsy cases and 7.6% of idiopathic epilepsy cases among referred dogs in Japan (Hamamoto *et al.*, 2016)

Intervertebral disc disease (IVDD)

- Toy Poodles comprised 1.5% of a cervical IVDD caseload and 1.2% of a thoracolumbar IVDD caseload at a Japanese referral hospital (Itoh *et al.*, 2008)

Lysosomal storage disease – GM₂ gangliosidosis

- Inherited as an autosomal recessive trait
- Described in related Toy Poodles in Japan (Tamura *et al.*, 2010)

Ocular conditions**Cataract**

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 10.21%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Toy/Miniature Poodle had OR 2.6 (95% CI 2.1–3.2) for cataract compared to a hospital population in Korea
- Toy Poodle had OR 6.1 compared with an overall referral population in the USA (Adkins & Hendrix, 2005; Gelatt & MacKay, 2005; Baumworcel *et al.*, 2009; Park *et al.*, 2009; Donzel *et al.*, 2016)

Glaucoma

- Toy Poodle had 1.20% prevalence in a US referral population, 1994–2002
- 12.9% of Toy Poodles referred for eye or neurological conditions in Japan were affected (Gelatt & MacKay, 2004a; Kato *et al.*, 2006)

Lens luxation – primary

- Toy Poodle had RR 3.10 (95% CI 2.37–4.06) compared with overall referral dogs in the USA (Sargan *et al.*, 2007)

Myopia (short-sightedness)

- The Toy Poodle had the highest mean myopic refractive error of all breeds tested ($p < 0.001$), with a 76.6% prevalence among privately owned dogs in the USA (Kubai *et al.*, 2008)

Progressive retinal atrophy (PRA)

- Genetic mechanism identified in Toy Poodle (Goldstein *et al.*, 2006)

Progressive rod–cone degeneration (PRCD)

- PRCD gene mapped to the centromeric region of canine chromosome 9 (CFA9)
- Toy Poodle has 0.45 allele frequency of the disease-causing mutation (Dostal *et al.*, 2011)

Physiological conditions**Litter size**

- Smaller litters are associated with older bitches and smaller breeds

- Toy Poodle had the smallest mean litter size (2.4 puppies) among registered breeds in Norway

(Borge *et al.*, 2011)

Renal and urinary conditions

Ectopic ureter

- Toy/Miniature Poodle bitches had RR 3.6 (95% CI 2.5–4.8) and RR 3.26 (95% CI 2.26–7.93) compared with overall referral populations in two US studies

(Hayes, 1974b, 1984)

Urinary tract infections (UTI)

- Toy and Miniature Poodle (7.8%) was the second most commonly affected breed among referral dogs in the USA for recurrent or persistent UTI

(Norris *et al.*, 2000)

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- Toy and Miniature Poodle had OR 3.32 in the USA

(Lekcharoensuk *et al.*, 2000a)

Reproductive conditions

Cryptorchidism

- Toy Poodle had RR 6.0 (95% CI 5.1–7.0) compared with an overall US referral population
- Toy and Miniature Poodles had OR 2.8 (95% CI 2.43–3.23) compared with an overall US referral population

(Pendergrass & Hayes, 1975;

Hayes *et al.*, 1985)

Testicular neoplasia

- Toy and Miniature Poodle had a proportional morbidity ratio of 1.9 (95% CI 1.1–3.2) compared with all other dogs in a Norwegian cancer registry

(Nødtvedt *et al.*, 2011)

PORTUGUESE PODENGO PEQUENO (SMALL PORTUGUESE PODENGO)

Musculoskeletal conditions

Patellar luxation

- 7.2% of US Portuguese Podengo Pequeno affected
- Mainly medial luxation observed, often bilateral (Orthopedic Foundation for Animals, 2015)

Neurological conditions

Cerebellar degeneration

- Born normal but then develop cerebellar deficits from 2 weeks of age
- Autosomal recessive inheritance suspected
- Described in Portuguese Podengo Pequeno in the Netherlands

(van Tongeren *et al.*, 2000)

PORTUGUESE WATER DOG (PORTIE CÃO DE AGUA)

Cardiovascular conditions

Dilated cardiomyopathy (DCM)

- Early onset and no sex predisposition in Portuguese Water Dog
- Often causes sudden death or is rapidly fatal
- Autosomal recessive inheritance suspected
- May be associated with taurine deficiency

(Dambach *et al.*, 1999;

Alroy *et al.*, 2000, 2005)

Dermatological conditions

Canine follicular dysplasia (seasonal flank alopecia)

- Familial tendency with early age of onset suggests an inherited basis
- Portuguese Water Dog comprised 28.6% of a US referral population

(Miller & Scott, 1995;

Rothstein *et al.*, 1998)

Improper coat

- Does not meet the American Kennel Club breed standard for having 'a profuse thickly planted coat of strong healthy hair, which covers the body evenly'
- Causal allelic variation at R-spondin 2 (*RSPO2*) gene identified
- 4.7% of Portuguese Water Dogs in the USA affected

(Parker *et al.*, 2010)

Endocrine conditions

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely: estimated heritability 0.49 (± 0.16)
- Prevalence of 1.5% in Portuguese Water Dog
- OR 46.66 (95% CI 3.73–245.9) compared with an overall US referral population
- Portuguese Water Dog had highest breed incidence rate in insured dogs in Sweden: 64.5

(95% CI 34.4–122.0) cases per 10 000 DYAR, compared with 2.26 (95% CI 2.07–2.46) across all dogs

(Peterson *et al.*, 1996; Chase, 2006; Oberbauer *et al.*, 2006; Mitchell & Pearce, 2012; Hanson *et al.*, 2016)

Musculoskeletal conditions

Hip dysplasia

- Portuguese Water Dog prevalence was 13.2% in the USA
- OR 3.1 (95% CI 1.3–7.4) compared to cross-breeds in the USA
(LaFond *et al.*, 2002; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Neurological conditions

Lysosomal storage disease – GM₁ gangliosidosis

- Autosomal recessive inheritance
- Signs seen at 5–6 months and usually fatal
- Reported in Portuguese Water Dogs in the USA
(Saunders *et al.*, 1988; Shell *et al.*, 1989)

Ocular conditions

Progressive retinal atrophy (PRA)

- Genetic mechanism identified in Portuguese Water Dog
(Goldstein *et al.*, 2006)

Progressive rod–cone degeneration (PRCD)

- PRCD gene mapped to the centromeric region of canine chromosome 9 (CFA9)
- Portuguese Water Dog has 0.33 allele frequency of the disease-causing mutation
(Dostal *et al.*, 2011)

PRESA CANARIO (PERRO DE PRESA CANARIO, DOGO CANARIO, CANARY DOG)

Behavioural conditions

Aggression

- Presa Canario had higher levels of territorial aggression in Spain
(Perez-Guisado & Munoz-Serrano, 2009)

Cardiovascular conditions

Dilated cardiomyopathy (DCM)

- Males of large and giant breeds predisposed
- Generally occurs from 5–8 years

- Median survival of 19 weeks
- High prevalence in the Presa Canario reported in Gran Canaria

(Morales *et al.*, 2001)

Infectious conditions

Dirofilariasis

- Large breeds and dogs living outdoors predisposed
- Among dogs attending first-opinion veterinary clinics in the Canaries, Presa Canario had 76% prevalence compared with 58.9% prevalence across all breeds on Gran Canaria, and 34% prevalence compared with 21% prevalence across all breeds on Tenerife
(Montoya *et al.*, 1998, 2006)

PUG

Dental conditions

Dentigerous cyst

- Brachycephalic breeds over-represented
- Pug comprised 17.2% of a US referral caseload and was significantly ($p < 0.001$) over-represented compared with the overall hospital population
(Verstraete *et al.*, 2011)

Dermatological conditions

Canine pigmented epidermal naevus

- Characterized in Pugs in Japan
(Narama *et al.*, 2005)

Demodicosis

- An inheritance pathway has been described
- Short-haired breeds predisposed
- OR 1.5 (95% CI 1.0–2.1) compared with an overall US first-opinion population
- Pugs were the most commonly affected breed (16.7%) in a pan-European study
- Pugs were significantly over-represented in Russia ($p < 0.001$)
(Mueller *et al.*, 2009; It *et al.*, 2010; Plant *et al.*, 2011; Kuznetsova *et al.*, 2012)

Drug reactions

Vaccine-associated adverse effect

- Smaller breeds predisposed
- Pug had the second-highest incidence (93.0 cases per 10 000 dogs within 3 days of vaccine administration; 95% CI 80.2–107.2), compared with 38.2 per 10 000 (95% CI 37.1–39.3) overall in a US primary-care population
(Moore *et al.*, 2005)

Endocrine conditions

Diabetes mellitus

- Familial and inherited
- Older entire females are predisposed
- OR 3.9 (95% CI 1.2–9.7) compared with cross-breeds in a US referral study (Hess *et al.*, 2000)

Gastrointestinal conditions

Congenital portosystemic shunt

- Smaller breeds predisposed
- Pug had 5.88% prevalence for portosystemic shunt, compared with 0.35% for crossbreeds, in a US referral study, and comprised 9.4% of another US caseload
- Referred Pugs (1.3% prevalence) had OR 26.2 (95% CI 15.7–42.5) compared with an overall US referral population (Tobias & Rohrbach, 2003; Winkler *et al.*, 2003; Bellumori *et al.*, 2013)

Metabolic conditions

Overweight/obesity

- 70.7% of Pugs attending veterinary clinics in China were obese
- Overweight/obesity was the most common disorder in UK Pugs (prevalence 13.2%, 95% CI 11.1–15.4)
- The Pug had significantly higher body condition score (6.00) than other show dogs in Holland (Courcier *et al.*, 2010; Corbee, 2013; Mao *et al.*, 2013; O'Neill *et al.*, 2016a; Raffan *et al.*, 2016)

Musculoskeletal conditions

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

- Inherited as an autosomal recessive trait
- OR 65.6 (95% CI 28.1–152.9) compared to crossbreeds in the USA in the USA (Brenig *et al.*, 1999; LaFond *et al.*, 2002)

Hip dysplasia

- Pug had second-highest prevalence in a US study: 61% (Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Patellar luxation

- 3.5% of UK Pugs affected, and 5.4% of US referred Pugs affected
- Mainly medial luxation observed, often bilateral
- OR 3.7 (95% CI 2.3–5.8) compared to cross-breeds under UK primary veterinary care

- OR 3.3 (95% CI 1.9–5.6) compared to cross-breeds in the USA (LaFond *et al.*, 2002; Alam *et al.*, 2007; Orthopedic Foundation for Animals, 2015; O'Neill *et al.*, 2016c)

Spina bifida

- 38.2% of clinically normal Pugs affected in a UK referral study, and Pugs were significantly more likely to be affected ($p < 0.0001$) than French Bulldogs or British Bulldogs (Ryan *et al.*, 2017)

Thoracic vertebral malformations

- 73.5% of clinically normal Pugs affected with at least one thoracic vertebral malformation in a UK referral study: 25% had no lumbar vertebrae and 17.6% had hemivertebrae (Ryan *et al.*, 2017)

Neoplastic conditions

Mast cell tumour (MCT)

- Multiple cutaneous tumours were recorded in 56% of Pug cases
- Mean age at presentation 7.5–9 years, but can occur at any age
- OR 10.0 (95% CI 1.5–64.5) compared with crossbreeds in the UK
- OR 5.2 (95% CI 4.3–6.3) and OR 3.17 (95% CI 1.47–6.82) compared with crossbreeds in two US studies
- OR 2.28 (95% CI 1.81–2.86) compared with an overall referral population in the USA
- RR 4.38 compared with all breeds in laboratory records in the USA, but these tended to be low-grade tumours (Goldschmidt & Mcmanus, 2000; McNie *et al.*, 2006; White *et al.*, 2011; Shoop *et al.*, 2015; Mochizuki *et al.*, 2016)

Squamous cell carcinoma – cornea

- Rare tumour type
- Brachycephalic breeds and animals with keratoconjunctivitis predisposed
- Pug was over-represented and comprised 36.6% of US referral cases (Dreyfus *et al.*, 2011)

Neurological conditions

Hemivertebrae (wedge-shaped vertebrae)

- Assumed to be highly heritable
- 100% of Pugs affected (Schlensker & Distl, 2013)

Intervertebral disc disease (IVDD)

- Pugs comprised 0.8% of a cervical IVDD caseload and 2.4% of a thoracolumbar IVDD caseload at a Japanese referral hospital (Itoh *et al.*, 2008)

Necrotizing meningoencephalitis

- Strong heritability in Pugs
- Genetic mechanism via dog leucocyte antigen (DLA) system reported in Pugs in the USA
- Pugs comprised 90.9% of a US caseload, and were predisposed in Switzerland
- Median age of onset 18 months (Tipold, 1995; Suzuki *et al.*, 2003; Schatzberg *et al.*, 2005; Levine *et al.*, 2008; Greer *et al.*, 2009, 2010; Safra *et al.*, 2011)

Ocular conditions**Bacterial keratitis**

- OR 45.9 (95% CI 7.1–302.3) compared with a US referral population (Tolar *et al.*, 2006)

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.28%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Conjunctivitis

- 43.1% of a Pug referral ophthalmology caseload in Austria affected (Krecny *et al.*, 2015)

Corneal pigmentation

- A local biological response to various irritating stimuli
- Detected in at least one eye of 82.4% (95% CI 77.6–86.3%) of Pugs from the general population in the USA
- 68.5% of a Pug referral ophthalmology caseload in Austria affected (Labelle *et al.*, 2013; Krecny *et al.*, 2015)

Corneal ulceration (ulcerative keratitis)

- Pug showed 23.8% prevalence of corneal ulceration among ophthalmic referrals in South Africa

- Pug had the highest primary-care veterinary predisposition, with OR 19.05 (95% CI 13.45–26.97) compared with crossbreeds in the UK (Petrick, 1996; O'Neill *et al.*, 2017b)

Distichiasis

- Heritability of 0.043
- 26.9% of a Pug referral ophthalmology caseload in Austria affected (Ketteritzsch *et al.*, 2004; Krecny *et al.*, 2015)

Ectopic cilia

- 4.6% of a Pug referral ophthalmology caseload in Austria affected (Krecny *et al.*, 2015)

Entropion

- 100% of a Pug referral ophthalmology caseload in Austria affected (Krecny *et al.*, 2015)

Iris hypoplasia

- Recorded in 71.0% (95% CI 65.3–76.0%) of left eyes and 72.1% (95% CI 66.5–77.1%) of right eyes of Pugs from the general population in the USA (Labelle *et al.*, 2013)

Keratoconjunctivitis sicca

- 30% of a Pug referral ophthalmology caseload in Austria affected
- 1.9% of UK general Pug population affected (Krecny *et al.*, 2015; O'Neill *et al.*, 2016a)

Macroblepharon (diamond eye)

- 100% of a Pug referral ophthalmology caseload in Austria affected (Krecny *et al.*, 2015)

Persistent pupillary membranes

- Recorded in 83.8% (95% CI 78.9–87.7%) of left eyes and 85.3% (95% CI 80.6–89.0%) of right eyes of Pugs from the general population in the USA (Labelle *et al.*, 2013)

Sudden acquired retinal degeneration (SARD, amaurosis)

- Small breeds predisposed
- Median age 8.1 years
- Pug comprised 5% of one US caseload, and 8.9% of a separate US referred caseload (Montgomery *et al.*, 2008; Heller *et al.*, 2017)

Physiological conditions

Vertebral heart score (VHS)

- Pugs from referral US practice had significantly greater ($p < 0.00001$) radiographic vertebral heart score (size) than a previously established average of 9.7 ± 0.5

(Buchanan & Bücheler, 1995;
Jepsen-Grant *et al.*, 2013)

Renal and urinary conditions

Renal calculi

- May be associated with urinary tract infections
- OR 9.01 ($p < 0.001$) in female Pugs compared with crossbreeds in a US referral study

(Ling *et al.*, 1998c)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Small breeds and males are predisposed
- OR 4.7 (95% CI 3.3–6.8) compared with all dogs in the UK

(Roe *et al.*, 2012)

Reproductive conditions

Dystocia

- 27.4% of Pug litters born by caesarean section in the UK
- Pug had the fifth-highest incidence (24.1 per 1000 DYAR) among insured bitches in Sweden, compared with 5.7 per 1000 DYAR overall
- OR 11.3 (95% CI 7.1–17.9) compared with first-opinion emergency-care crossbred bitches in the UK

(Bergström *et al.*, 2006;
Evans & Adams, 2010a; O'Neill *et al.*, 2017c)

High perinatal mortality

- Older dams and increasing litter size associated with increased perinatal mortality
- Overall perinatal mortality risk in Norway was 8.0%
- Pug had the fourth-highest perinatal mortality (16.9%) among registered purebred litters in Norway

(Tønnessen *et al.*, 2012)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 48% of insured Pug bitches develop the condition by 10 years of age, compared with 19% for bitches overall

(Jitpean *et al.*, 2012)

Respiratory conditions

Aspiration pneumonia

- Pugs comprised 4.4% of cases but only 1.6% of the underlying referral population in the USA

(Kogan *et al.*, 2008)

Brachycephalic obstructive airway syndrome (BOAS)

- Pugs comprised 17.9% and 26% of two UK BOAS caseloads, and 21% and 50% of two US BOAS caseloads
- Pugs had the highest breed prevalence (60.0%) among referral dogs in Belgium

(Torrez & Hunt, 2006;

Riecks *et al.*, 2007;

De Lorenzi *et al.*, 2009;

Njikam Nsangou *et al.*, 2009;

Fonfara *et al.*, 2011; Roedler *et al.*, 2013;

Marchant *et al.*, 2017)

Lung lobe torsion

- Pugs comprised 30.4% of all referral cases in Canada and had a breed prevalence of 2.6%

(Murphy & Brisson, 2006)

Nasopharyngeal turbinates

- Associated with brachycephalic obstructive airway syndrome (BOAS)
- 53% of Pugs affected in a US study

(Ginn *et al.*, 2008)

Tracheal collapse

- Small dog breeds predisposed
- Pug comprised 7.1% of a referral caseload in Korea

(Eom *et al.*, 2008)

PULI

See *Hungarian Puli*

PYRENEAN MOUNTAIN DOG (GREAT PYRENEES)

Cardiovascular conditions

Myocardial infarction (MI)

- Pyrenean Mountain Dogs (9%) were over-represented among MI cases compared with an overall US referral caseload (0.1%)

(Driehuys *et al.*, 1998)

Haematological/immunological conditions

Hereditary thrombopathy

- Inherited intrinsic platelet function defect causing excessive bleeding
- Described in Pyrenean Mountain Dogs in the USA

(Boudreaux *et al.*, 1996;
Boudreaux & Lipscomb, 2001)

Musculoskeletal conditions

Chondrodysplasia (short-limbed or disproportional dwarfism)

- Described in related Pyrenean Mountain Dogs in the USA
- Simple autosomal recessive inheritance
- Radiographic abnormalities restricted to the metaphyses of long bones and vertebrae in Pyrenean Mountain Dog

(Bingel & Sande, 1994)

Hip dysplasia

- Pyrenean Mountain Dog prevalence was 9.6% in the USA and 23% in Belgium
- OR 3.3 (95% CI 1.9–5.9) compared to crossbreeds in a US study
(LaFond *et al.*, 2002; Coopman *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Osteochondrosis – shoulder

- The coefficient of heritability varies from 0.25 to 0.45
- Age of onset usually 4–7 months, the period of rapid growth
- Males predisposed
- OR 42.7 (95% CI 24.4–74.8) compared to crossbreeds in the USA
(Rudd *et al.*, 1990; Nečas *et al.*, 1999; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 5.3 (95% CI 3.5–8.0) compared to crossbreeds in the USA
(LaFond *et al.*, 2002)

Patellar luxation

- OR 64.0 (95% CI 22.0–185.9) compared to crossbreeds in the USA
(LaFond *et al.*, 2002)

Ocular conditions

Multifocal retinopathy

- Two spontaneous mutations in the canine *VMD2* gene identified in Pyrenean Mountain Dogs
- Autosomal recessive inheritance
- Pyrenean Mountain Dogs had 43.2% prevalence in Canada
(Grahn *et al.*, 1998; Grahn & Cullen, 2001; Guziewicz *et al.*, 2007)

Physiological conditions

Hypercholesterolaemia

- Healthy Pyrenean Mountain Dogs had high plasma cholesterol compared with other breeds tested in Italy
(Pasquini *et al.*, 2008)

Reproductive conditions

Dystocia

- 28.9% of Pyrenean Mountain Dog litters born by caesarean section in the UK
(Evans & Adams, 2010a)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 43% of insured Pyrenean Mountain Dog bitches develop the condition by 10 years of age, compared with 19% for bitches overall
(Jitpean *et al.*, 2012)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Autosomal recessive mode of inheritance suspected
- Described in a family of Pyrenean Mountain Dogs
(Gabriel *et al.*, 2006)

PYRENEAN SHEPHERD (BERGER DES PYRÉNÉES, PASTOR DE LOS PIRINEOS, LABRIT, PETIT BERGER)

Gastrointestinal conditions

Cleft lip and/or palate

- Monogenic autosomal recessive inheritance in Pyrenean Shepherds
- Pyrenean Shepherd had 28.8% prevalence in Germany
(Kemp *et al.*, 2009)

Musculoskeletal conditions

Short tail (bobtail)

- 10.9% of Pyrenean Shepherds tested in Finland affected

(Hytönen *et al.*, 2009)

RAT TERRIER**Endocrine conditions**

Congenital hypothyroidism

- Simple autosomal recessive inheritance in Rat Terriers
- Associated with central nervous system hypomyelination
- A homozygous nonsense mutation in the thyroid peroxidase gene identified in Rat Terriers in the USA

(Pettigrew *et al.*, 2007)

Neurological conditions

Primary brain tumour

- Rat Terrier had a significantly increased risk ($p=0.047$) compared with an overall referral post-mortem population in the USA

(Song *et al.*, 2013)

Ocular conditions

Lens luxation – primary

- *ADAMTS17* mutation has been identified: 37.65% of US Rat Terriers were carriers
- RR 4.83 (95% CI 2.69–8.68) and RR 17.80 (95% CI 2.53–125.10) compared with overall referral dogs in the USA

(Sargan *et al.*, 2007; Gould *et al.*, 2011)

RED AND WHITE SETTER

See *Irish Red and White Setter*

RED KELPIE

See *Australian Kelpie*

RED SETTER

See *Irish Setter*

RETRIEVER

See *Flat-coated Retriever*; *Golden Retriever*; *Labrador Retriever*

RHODESIAN RIDGEBACK**Dermatological conditions**

Atopic dermatitis (atopy)

- Rhodesian Ridgeback had the eighth-highest incidence in insured dogs in Sweden (5.0 cases per 1000 DYAR; 95% CI 3.8–6.2) and was over-represented in Switzerland

(Nødtvedt *et al.*, 2006; Picco *et al.*, 2008)

Endocrine conditions

Hypothyroidism

- Females and younger dogs are predisposed to having serum thyroid hormone autoantibodies (THAA) that are associated with hypothyroidism
- Novel genes and pathways reported for Rhodesian Ridgebacks from the USA
- OR 1.72 ($p=0.001$) for THAA compared with all other breeds in the USA
- Rhodesian Ridgeback was strongly over-represented in the UK

(Nachreiner *et al.*, 2002; Kennedy *et al.*, 2006b; Bianchi *et al.*, 2015)

Musculoskeletal conditions

Elbow dysplasia

- Rhodesian Ridgeback had 14.3% prevalence in the UK, 6.3% in the USA, 21.1% in South Africa and 8% in Belgium

(Kirberger & Stander, 2007; Coopman *et al.*, 2008; Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Lumbosacral transitional vertebrae

- Large breeds predisposed
- Rhodesian Ridgeback had 15.7% prevalence, compared with an overall 10.0% prevalence among dogs screened for pelvic problems, in the Czech Republic

(Fialová *et al.*, 2014)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.0 (95% CI 1.1–3.4) compared to cross-breeds in the USA

(LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Neoplastic conditions

Cutaneous soft-tissue sarcoma

- Rhodesian Ridgeback had OR 4.8 (95% CI 2.7–8.7) compared with a US referral population

and was significantly over-represented ($p=0.016$) in another US referral study

(Baker-Gabb *et al.*, 2003; Villamil *et al.*, 2011)

Mammary neoplasia

- 26% of insured Rhodesian Ridgeback bitches in Sweden developed the condition by 10 years of age, compared with 13% for bitches overall (Jitpean *et al.*, 2012)

Mast cell tumour (MCT)

- Mean age at presentation between 7.5 and 9 years, but can occur at any age
- Three US studies showed OR 34.44 (95% CI 5.17–229.23), OR 3.6 (95% CI 2.8–4.7) and OR 5.07 (95% CI 3.22–7.98) compared with crossbreeds

(Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011; White *et al.*, 2011)

Neurological conditions

Cerebellar Purkinje cell degeneration and coat colour dilution

- Clinical signs evident by 2 weeks of age and generally fatal
- Autosomal recessive inheritance suspected
- Reported in related Rhodesian Ridgebacks in the USA

(Chieffo *et al.*, 1994)

Degenerative myelopathy

- Insidious, progressive ataxia and paresis of the hindlimbs that ultimately leads to paraplegia and euthanasia
- Rhodesian Ridgeback had 0.74% prevalence, compared with 0.15% in referral crossbreeds in the USA
- Homozygosity for the A allele of the canine *SOD1* gene associated with degenerative myelopathy in Rhodesian Ridgeback

(Coates *et al.*, 2007; Awano *et al.*, 2009)

Dermoid sinus

- Prevalence of 8–10% in Rhodesian Ridgebacks in Sweden
- Complex dihybrid mode of inheritance suggested
- Causative mutation is a duplication of *FGF3*, *FGF4*, *FGF19* and *ORAOV1* genes that cause hair ridge and predisposition to dermoid sinus in Rhodesian Ridgeback dogs

(Hillbertz, 2005; Salmon Hillbertz *et al.*, 2007)

Ocular conditions

Glaucoma – secondary

- High intraocular pressure due to primary intraocular diseases that impede aqueous humor flow
- Rhodesian Ridgeback had the third-highest standardized morbidity ratio (4.1) compared with an overall referral population in the USA

(Johnsen *et al.*, 2006)

Physiological conditions

Dorsal ridge

- Breed-defining ridge of Rhodesian Ridgebacks comes from a duplication of a specific region on chromosome 18 called the Ridge allele
- Dominant inheritance
- Homozygosity for the Ridge allele predisposes to dermoid sinus (see under *Neurological conditions*)
- Polymerase chain reaction (PCR) test now developed to accurately determining the copy number genotype for the Ridge allele of Rhodesian Ridgebacks

(Waldo & Diaz, 2015)

Litter size

- Larger litters associated with younger bitches and larger breeds
- Rhodesian Ridgeback had the largest mean litter size (8.9 puppies) among registered breeds in Norway

(Borge *et al.*, 2011)

Reproductive conditions

Dystocia

- Rhodesian Ridgeback had an incidence of 14.1 per 1000 DYAR among insured bitches in Sweden, compared with 5.7 per 1000 DYAR overall

(Bergström *et al.*, 2006)

High perinatal mortality

- Older dams and increasing litter size associated with increased perinatal mortality
- Overall perinatal mortality risk in Norway was 8.0%
- Rhodesian Ridgeback had the third-highest perinatal mortality (17.9%) among registered purebred litters in Norway

(Tønnessen *et al.*, 2012)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Larger breeds and overweight individuals predisposed
- Rhodesian Ridgeback comprised 6% of an Australian caseload

(Snelling & Edwards, 2003)

ROTTWEILER

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- Rottweiler reported with higher risk of aggression in Denmark, Italy, UK and Spain
- Rottweilers were the commonly reported breed involved in fatal human attacks in the USA
- Rottweiler was over-represented as the aggressor in dog–dog conflict in Germany, representing 3.4% of aggressors and 2.8% of the general population
- Rottweiler had a bite-risk index of 3.9 ($p < 0.001$) compared with an overall general population of dogs in the Netherlands

(Bradshaw *et al.*, 1996;

Lund *et al.*, 1996;

Roll & Unshelm, 1997; Bradshaw &

Goodwin, 1999; Sacks *et al.*, 2000;

Notari & Goodwin, 2007; Perez-Guisado &

Munoz-Serrano, 2009; Cornelissen &

Hopster, 2010; Zapata *et al.*, 2016)

Cardiovascular conditions

Aortic or cardiac mineralization

- Condition had a prevalence of 0.61% in an overall referral population in the UK
- Older dogs predisposed
- Rottweilers were significantly ($p = 0.003$) among a UK referral caseload

(Schwarz *et al.*, 2002)

Aortic stenosis – subaortic stenosis (SAS)

- Rottweilers was significantly over-represented, comprising 17.4% of a US referral caseload

(Kienle *et al.*, 1994)

Atrial fibrillation (AF)

- Large and giant breeds predisposed
- Rottweiler comprised 4% of the caseload and was significantly over-represented ($p < 0.01$)

in the US and Canada among AF cases without structural and functional disease

(Menaut *et al.*, 2005)

Dental conditions

Abnormal dentition

- The most common specific problems among affected dogs overall were polyodontia (33.2%), retained deciduous teeth (19.7%) and prognathism (10.0%)
- Rottweiler was the most commonly affected large breed, comprising 5.3% of an overall affected caseload aged 7–18 months in Bulgaria

(Borissov *et al.*, 2004)

Dermatological conditions

Pyotraumatic folliculitis (acute moist dermatitis, hot spot, wet eczema)

- Breeds (and males) with a heavy pelage and dense undercoat predisposed
- Associated with hot weather
- Rottweiler had the highest breed relative risk (3.38) compared with a hospital population in Sweden

(Holm *et al.*, 2004)

Endocrine conditions

Diabetic ketoacidosis

- Rottweiler comprised 12.9% of a referral caseload in Belgium

(De Causmaecker *et al.*, 2009)

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely
- Females predisposed
- OR 2.60 (95% CI 1.22–5.05) compared with an overall US referral population

(Peterson *et al.*, 1996)

Gastrointestinal conditions

Gastroduodenal perforation

- Large breeds are predisposed
- Rottweilers were over-represented and younger in a US referral study, comprising 26.7% of the caseload

(Hinton *et al.*, 2002)

Inflammatory bowel disease

- Mode of inheritance is incompletely understood and may differ between breeds
- OR 2.97 (95% CI 1.76–5.02) compared with referred crossbreeds in the UK

(Kathrani *et al.*, 2011)

Haematological/immunological conditions

Eosinophilia

- Rottweiler had the highest incidence and the highest median eosinophil count among laboratory submissions in Sweden but is only the seventh most common breed in Sweden (Lilliehöök *et al.*, 2000)

Immunodeficiency syndrome

- Described in related Rottweiler puppies in the UK
- Complex inheritance suggested (Day, 1999)

Infectious conditions

Giardiasis

- Dogs aged <1 year significantly predisposed ($p < 0.0001$)
- OR 2.12 (95% CI 1.03–4.34) compared with an overall rescue population in the UK (Upjohn *et al.*, 2010)

Leishmaniasis

- Rottweiler was significantly over-represented ($p < 0.001$) among referral animals in Spain, representing 13.1% of affected dogs and 3.8% of the referral population (Miranda *et al.*, 2008)

Parvovirus enteritis

- Rottweiler had an OR of 6.0 (95% CI 1.4–25.3) compared with an overall referral US population in one study, and was at significantly increased risk in another US study
- Suggested due to defective immune system in Rottweilers (Glickman *et al.*, 1985; Seton & Seton, 1990; Houston *et al.*, 1996)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Estimated heritability of 0.48 and is a highly polygenic complex trait
- Rottweiler had 5.63% prevalence, compared with 1.58% overall in referred dogs, in the Czech Republic, and was the most commonly affected breed in a UK referral caseload, comprising 12.6% of the total
- OR 3.58 (95% CI 3.40–3.78) compared with an overall US referral population, and OR 5.12 (95% CI 2.28–11.49) compared with an overall UK referral population

- Referral Rottweilers in the UK had significantly increased incidence of bilateral CCL disease ($p = 0.05$) and were significantly younger (median 977 days; $p < 0.0001$) compared with other breeds studied

(Bennett *et al.*, 1988; Nečas *et al.*, 2000; Witsberger *et al.*, 2008; Adams *et al.*, 2011; Guthrie *et al.*, 2012; Baker *et al.*, 2017)

Distal tibial valgus deformity

- Median age at presentation 9 months
- RR 12.2 (95% CI 1.5–30.4) compared with an overall referral population in the USA (Jaeger *et al.*, 2007)

Elbow dysplasia

- Rottweiler had 0.34 ± 0.02 heritability in Sweden
- Rottweiler had 54.9% prevalence in the UK, 39.1% in the USA, 54.7% in South Africa, 54.2% in Germany and 33% in Belgium
- Rottweiler had 6.31% prevalence, versus 0.16% prevalence in crossbreeds, in a US referral caseload
- OR 36.1 (95% CI 23.6–55.2) for fragmented coronoid process, OR 174.0 (95% CI 78.2–387.1) for osteochondrosis of the elbow, and OR 27.4 (95% CI 20.5–36.6) for ununited anconeal process compared to crossbreeds in the USA
- Rottweiler had 2.0% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland
- Rottweiler comprised 17.5% of a referral caseload in Germany and 12.25% of a UK referral caseload

(Beuing *et al.*, 2000; LaFond *et al.*, 2002; Meyer-Lindenberg *et al.*, 2002; Kirberger & Stander, 2007; Coopman *et al.*, 2008; Malm *et al.*, 2008; Narojek *et al.*, 2008; Fitzpatrick *et al.*, 2009; Bellumori *et al.*, 2013; Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Hip dysplasia

- Rottweiler had 0.38 ± 0.02 heritability in Sweden
- Rottweiler prevalence was 21.3% and 35.4% among officially submitted radiographs in the USA, and ranged from 41% to 69% among client-owned dogs in the USA

- Two studies reported Rottweiler with OR 6.5 (95% CI 5.5–7.5) and OR 3.33 (95% CI 3.18–3.49) compared to crossbreeds in the USA

(LaFond *et al.*, 2002;

Rettenmaier *et al.*, 2002;

Paster, 2005; Malm *et al.*, 2008;

Witsberger *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Juvenile-onset distal myopathy

- Considered as a type of muscular dystrophy
- Described in related Rottweiler puppies in the USA

(Hanson *et al.*, 1998)

Osteochondrosis – hock

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 206.2 (95% CI 108.2–393.2) compared to crossbreeds in the USA

(Nečas *et al.*, 1999; LaFond *et al.*, 2002)

Osteochondrosis – shoulder

- The coefficient of heritability varies from 0.25 to 0.45
- Age of onset usually 4–7 months, the period of rapid growth
- Males predisposed
- OR 22.8 (95% CI 17.4–29.9) compared to crossbreeds in the USA

(Rudd *et al.*, 1990; Nečas *et al.*, 1999;

LaFond *et al.*, 2002)

Osteochondrosis – stifle

- Males predisposed
- Age of onset usually 5–12 months, but may be earlier
- OR 66.3 (95% CI 29.4–149.6) compared to crossbreeds in the USA

(Nečas *et al.*, 1999;

LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.4 (95% CI 1.2–1.6) compared to crossbreeds in the USA
- Rottweiler comprised 9.74% of a referral caseload in Romania

(LaFond *et al.*, 2002;

Trostel *et al.*, 2003; Igna *et al.*, 2016)

Sacroiliac joint biomechanics

- Rottweiler had the highest sacroiliac joint (SIJ) loading forces among breeds tested in Austria

(Breit & Künzel, 2001)

Sesamoid disease

- Rottweilers showed a radiographic prevalence of 73% but an attributable lameness prevalence of 21.8%

(Read *et al.*, 1992)

Neoplastic conditions

Bone tumour

- The majority of bone tumours are thought to be osteosarcoma
- Rottweiler had the fifth-highest incidence rate (36 per 10 000 DYAR; 95% CI 29–44) among insured dogs in Sweden

(Egenvall *et al.*, 2007)

Canine cutaneous histiocytoma

- OR 1.79 (95% CI 1.60–2.01) compared with crossbreeds in the USA

(Goldschmidt & Mcmanus, 2000; Fulmer & Mauldin, 2007)

Fibroma/fibrosarcoma

- Rottweiler had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland

(Grüntzig *et al.*, 2016)

Histiocytic sarcoma complex – histiocytic sarcoma and disseminated histiocytoma

- Median age at diagnosis 10 years
- Rottweilers were highly represented, comprising 28.2% of a US referral caseload
- Rottweiler was the most commonly affected breed with disseminated histiocytoma, comprising 33.3% of a US referral caseload

(Affolter & Moore, 2002; Schultz *et al.*, 2007)

Lymphoma

- Odds ratios 2.7 (95% CI 2.4–3.0) in the USA, 6.0 (95% CI 2.2–9.9) in Poland and 4.52 (2.09–9.73) in Australia compared with overall referral populations
- RR 2.14 (95% CI 1.65–4.31) compared with an overall referral population in the Netherlands and RR 2.81 in France
- Rottweiler comprised 7.4% of a referral caseload in Brazil and was over-represented in a UK referral caseload

- Rottweiler had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland
(Teske *et al.*, 1994; Jagielski *et al.*, 2002; Sueiro *et al.*, 2004; Pastor *et al.*, 2009; Villamil *et al.*, 2009; Di Bella *et al.*, 2013; Grüntzig *et al.*, 2016; Yau *et al.*, 2017)

Melanoma

- Rottweiler had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland
(Grüntzig *et al.*, 2016)

Mast cell tumour (MCT)

- Rottweiler had OR 2.46 for high-grade MCT tumours compared with all breeds, and showed approximately twofold increase in proportion of high-grade MCT tumours compared with all breeds (19.6 vs. 8.2%)
(Mochizuki *et al.*, 2016)

Neoplasia – overall

- Rottweiler had the fifth-highest proportional mortality from cancer among pedigree breeds in the UK: 45.3% (95% CI 36.9–53.6)
(Adams *et al.*, 2010)

Osteosarcoma

- Highly heritable
- Rottweilers neutered before 1 year old had RR 3.8 (95% CI 1.5–9.2) in males and 3.1 (95% CI 1.1–8.3) in females in the USA
- Rottweiler showed 5.3% prevalence in the USA
- Two US studies reported OR 3.9 (95% CI 2.6–6.0) and 14.6 (95% CI 8.9–24.0) compared to crossbreeds
- OR 4.6 (95% CI 3.6–5.8) for skeletal osteosarcoma compared with an overall US referral population
- Rottweiler had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland
- Increased incidence reported in Rottweilers in Romania, comprising 19.2% of a general caseload
(Langenbach *et al.*, 1998; Ru *et al.*, 1998; Cooley *et al.*, 2002; Rosenberger *et al.*, 2007; Muste *et al.*, 2010; Karlsson *et al.*, 2013; Grüntzig *et al.*, 2016)

Osteosarcoma – soft-tissue

- A malignant, osteoid-producing, mesenchymal neoplasm without primary periosteal or bone involvement
- Median survival time across all breeds was 26 days
- Rottweiler had OR 2.6 (95% CI 1.1–11.7) compared with an overall US referral population
(Langenbach *et al.*, 1998)

Squamous cell carcinoma – digit

- Middle-aged to older dogs and dogs with black coats predisposed
- Rottweiler comprised 9.1% of a US referral caseload
(Henry *et al.*, 2005)

Neurological conditions

Discospondylitis

- Males and older dogs predisposed
- OR 3.1 (95% CI 2.1–4.7) compared with referral crossbreeds in the USA
(Burkert *et al.*, 2005)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Rottweiler had an incidence of 24.3 cases per 10 000 DYAR (95% CI 19.5–29.1) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
- Median age at onset of seizures was 36 months (range 8–84 months) in Rottweilers in Sweden and Norway
(Heske *et al.*, 2014, 2015)

Intervertebral disc disease (IVDD)

- Males and older dogs predisposed
- Rottweiler had an incidence rate of 38.3 (95% CI 32.3–44.4), compared with 27.8 for an overall population of insured dogs in Sweden
(Bergknut *et al.*, 2012)

Leucoencephalomyelopathy

- Described in adult Rottweilers in Australia
- Autosomal recessive inheritance suspected
(Slocombe *et al.*, 1989; Davies & Irwin, 2003)

Neuroaxonal dystrophy

- Group of neurodegenerative disorders characterized by severe degeneration of neuronal cells and their processes; can be familial
- Onset from 8 weeks, with euthanasia or death before 12 months

- Several reports in Rottweilers in the Netherlands, Brazil and the USA
(Chrisman, 1992; Kortz *et al.*, 1997; van den Ingh *et al.*, 1998; Jardim *et al.*, 1999; Davies & Irwin, 2003)

Progressive degenerative polyneuropathy

- Described in Rottweilers in France
(Braund *et al.*, 1994b)

Spinal arachnoid pseudocysts

- Described in Rottweilers in Germany
- Mean age at diagnosis was 3.5 years
- Males predisposed
(Jurina & Grevel, 2004)

Ocular conditions

Entropion

- Clear breed predispositions and polygenic inheritance suspected
- Rottweiler comprised 19.4% of referral case-load in Australia
(Read & Broun, 2007)

Myopia (short-sightedness)

- Rottweiler had 43.5% prevalence among privately owned dogs in the USA
- Rottweiler had the lowest mean refractive error (−1.77) among breeds tested in the USA
(Murphy *et al.*, 1992; Kubai *et al.*, 2008)

Uveitis

- Rottweiler showed 12.8% prevalence of uveitis among ophthalmic referrals in South Africa
(Petrick, 1996)

Physiological conditions

Pronounced eosinophilic response

- Rottweiler were over-represented, with the highest incidence of eosinophilia and the highest median eosinophil count, among dogs with disease in Sweden
(Lilliehöök *et al.*, 2000)

Hypercholesterolaemia

- Healthy Rottweilers had the highest plasma cholesterol among the breeds tested in Italy
(Pasquini *et al.*, 2008)

Litter size

- Larger litters associated with younger bitches and larger breeds

- Rottweiler had the tenth largest mean litter size (7.4 puppies) among registered breeds in Norway

(Borge *et al.*, 2011)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Described in related Rottweilers in the USA
(Wakamatsu *et al.*, 2007)

Kidney disease

- Rottweiler had an incidence of 26 (95% CI 21–31) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall, among insured dogs in Sweden
(Pelander *et al.*, 2015)

Urethral sphincter mechanism incompetence

- Larger breeds had 7.2 OR compared with smaller breeds in the USA
- Rottweiler over-represented in the UK
(Holt & Thrusfield, 1993; Forsee *et al.*, 2013)

Urolithiasis – cystine

- OR 7.0 (95% CI 4.3–11.4) compared with all dogs in the UK
(Roe *et al.*, 2012)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Rottweiler showed RR 4.4 (95% CI 3.3–5.4) compared with all bitches insured by Agria for veterinary care in Sweden
- 58% of insured Rottweiler bitches develop the condition by 10 years of age, compared with 19% for bitches overall
- OR 2.29 (95% CI 1.68–3.12) compared with all entire bitches in Finland
(Niskanen & Thrusfield, 1998; Egenvall *et al.*, 2001; Smith, 2006; Jitpean *et al.*, 2012)

Respiratory conditions

Laryngeal paralysis–polyneuropathy syndrome

- Larger breeds and overweight individuals predisposed
- Autosomal recessive mode of inheritance suspected
- Described in unrelated Rottweiler puppies in the USA
- RR 2.07 (95% CI 1.02–4.19) compared with an overall referral population in New Zealand
(Mahony *et al.*, 1998; Broome *et al.*, 2000)

Soft-tissue conditions**Calcinosis circumscripta**

- Rottweiler was over-represented, comprising 13% of a US referral caseload (Tafti *et al.*, 2005)

Perineal herniation

- Rottweiler accounted for 12.5% of referral cases in Croatia (Vnuk *et al.*, 2008)

ROUGH COLLIE

See *Collies – Rough and Smooth*

RUSSIAN BLACK TERRIER (TCHIORNY, RUSSIAN BEAR SCHNAUZER, CHORNYI)**Musculoskeletal conditions****Elbow dysplasia**

- Russian Black Terrier had 1.9% prevalence, compared with 0.7% in the overall population among radiographed dogs in Poland (Narojek *et al.*, 2008; Michelsen, 2013)

Neurological conditions**Polyneuropathy with ocular abnormalities and neuronal vacuolation (POANV)**

- Autosomal recessive inheritance
- RAB3GAP1:c.743delC mutation identified in Russian Black Terrier (Mhlanga-Mutangadura *et al.*, 2016)

Renal and urinary conditions**Hyperuricosuria**

- Russian Black Terriers in the USA had 0.51 allele frequency for the *SLC2A9* mutation associated with hyperuricosuria, and 27% of these dogs were estimated to be hyperuricosuric (Karmi *et al.*, 2010b)

ST BERNARD**Cardiovascular conditions****Dilated cardiomyopathy (DCM)**

- Males of large and giant breeds predisposed
- Generally occurs from 5–8 years
- Median survival 19 weeks
- St Bernard comprised 10% of a referral caseload in Italy

- St Bernard was commonly recorded in a UK study and was over-represented in a US study (Tidholm & Jonsson, 1997; Borgarelli *et al.*, 2006; Martin *et al.*, 2009)

Heart disease

- St Bernard had the fourth-highest breed incidence of death from heart disease in insured dogs in Sweden: 151 (95% CI 103–199) deaths per 10 000 DYAR (Egenvall *et al.*, 2006)

Dermatological conditions**Dermal arteritis of the nasal philtrum**

- Described in related St Bernards in the USA (Torres *et al.*, 2002)

Endocrine conditions**Hypoadrenocorticism (Addison's disease)**

- Complex inheritance pattern likely
- St Bernard was over-represented, comprising 8.6% of dogs with Addison's disease but only 4.6% of a total referral population in the USA (Thompson *et al.*, 2007)

Primary hypoparathyroidism

- St Bernard was the most common breed (17.6%) affected in Australia
- Successfully treatable with synthetic vitamin D (Jones & Alley, 1985; Russell *et al.*, 2006)

Gastrointestinal conditions**Palate agenesis, anotia and polydactyly**

- Described in related St Bernards in Mexico
- Fully penetrant autosomal recessive inheritance (Villagómez & Alonso, 1998)

Gastric dilatation/volvulus (bloat, GDV)

- 15.1% of UK pedigree St Bernard dogs died of the condition
- Pedigree St Bernard had seventh-highest breed prevalence (4.6%) in the UK
- St Bernard had an incidence of 6 cases per 1000 DYAR (95% CI 0–14) in the USA, with 0.6% of dogs affected per year
- St Bernard had 3.76% prevalence, compared with 0.20% for crossbreeds, in a US referral study, and were over-represented in two US referral studies (Glickman *et al.*, 2000a; Evans & Adams, 2010b; Beer *et al.*, 2012; Green *et al.*, 2012; Bellumori *et al.*, 2013)

Musculoskeletal conditions

Cranial cruciate ligament (CCL) disease

- Estimated heritability of 0.48 and is a highly polygenic complex trait
- OR 1.42 (95% CI 1.28–1.58) compared with an overall referral population in the USA
- St Bernard had 15.0% prevalence, compared with 1.58% overall in referred dogs in the Czech Republic
(Nečas *et al.*, 2000; Witsberger *et al.*, 2008; Baker *et al.*, 2017)

Polydactyly

- Large and giant breeds predisposed
- Ruled as an undesirable characteristic in pedigree St Bernards
(Alberch, 1985)

Elbow dysplasia

- St Bernard had 19.4% prevalence in the UK and 15.3% in the USA
- OR 53.4 (95% CI 28.1–101.6) for fragmented coronoid process and OR 14.2 (95% CI 6.7–30.3) for ununited anconeal process compared to crossbreeds in the USA
- St Bernard had 1.8% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland
(LaFond *et al.*, 2002; Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Hip dysplasia

- Two studies report St Bernard with OR 6.1 (95% CI 4.2–8.8) and OR 4.87 (95% CI 4.62–5.13) compared to crossbreeds in the USA
- St Bernard had sixth-highest prevalence in the USA: 49.2%
(LaFond *et al.*, 2002; Witsberger *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Osteochondrosis – shoulder

- The coefficient of heritability varies from 0.25 to 0.45
- Age of onset usually 4–7 months, the period of rapid growth
- Males predisposed
- OR 12.2 (95% CI 6.4–23.2) compared to crossbreeds in the USA
(Rudd *et al.*, 1990; Nečas, *et al.*, 1999; LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.5 (95% CI 1.6–3.9) compared to crossbreeds in the USA
(LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Neoplastic conditions

Bone tumour

- The majority of bone tumours are thought to be osteosarcoma
- St Bernards had the second-highest incidence rate (78 per 10 000 DYAR; 95% CI 43–113) among insured dogs in Sweden
(Egenvall *et al.*, 2007)

Haemangiosarcoma

- OR 2.82 (95% CI 1.16–6.83) for cutaneous haemangiosarcoma compared with crossbreeds in the USA
(Hargis *et al.*, 1992; Goldschmidt & Mcmanus, 2000)

Lymphoma

- OR 2.3 (95% CI 2.0–2.7) compared with a US referral population
- RR 2.63 (95% CI 1.09–6.32) compared with an overall referral population in the Netherlands
(Teske *et al.*, 1994; Villamil *et al.*, 2009)

Osteosarcoma

- Highly heritable
- St Bernard had the second-highest incidence in Sweden: 78 (95% CI 42–113) per 10 000 DYAR
- OR 11.9 (95% CI 6.9–20.6) for malignant bone tumour compared to crossbreeds in the USA
(Ru *et al.*, 1998; Egenvall *et al.*, 2007; Karlsson *et al.*, 2013)

Neurological conditions

Dysplasia of the cerebellar cortex

- Associated with hydrocephalus
- Described in St Bernards in the UK
- Neurological signs evident at 4 weeks of age
(Franklin *et al.*, 1997)

Fibrocartilaginous embolic myelopathy (FCEM)

- St Bernard comprised 9.1% of a US referral caseload
(Cauzinille & Kornegay, 1996)

Reproductive conditions**Dystocia**

- 41.2% of St Bernard litters born by caesarean section in the UK (Evans & Adams, 2010a)

High perinatal mortality

- Older dams and increasing litter size associated with increased perinatal mortality
- Overall perinatal mortality risk in Norway was 8.0%
- St Bernard had the fifth-highest perinatal mortality (16.9%) among registered purebred litters in Norway (Tønnessen *et al.*, 2012)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- OR 3.29 (95% CI 1.78–6.07) compared with all entire bitches in Finland (Niskanen & Thrusfield, 1998; Egenvall *et al.*, 2001; Smith, 2006)

SALUKI (PERSIAN GREYHOUND, PERSIAN SIGHTHOUND, TANJI, GAZELLE HOUND, ARABIAN HOUND)**Cardiovascular conditions****Dilated cardiomyopathy (DCM)**

- Prevalence increases with age, male and body size
- Saluki had 5.88% prevalence, compared with 0.16% in crossbreeds, in a US referral caseload
- Autosomal dominant transmission suspected (Dukes-McEwan *et al.*, 2003; Bellumori *et al.*, 2013)

Dermatological conditions**Black hair follicular dysplasia**

- Changes usually seen by 4 weeks of age
- Described in Saluki puppies (Hargis *et al.*, 1991; Schmutz *et al.*, 1998)

Neoplastic conditions**Cardiac tumour**

- Uncommon, and can occur both as primary and as metastatic lesions
- Saluki had the highest incidence (1.5%) and RR 7.75 (95% CI 3.92–15.38) compared with an overall referral population in the USA (Ware & Hopper, 1999)

Haemangiosarcoma

- OR 9.19 (95% CI 1.40–60.17) compared with crossbreeds in the USA (Hargis *et al.*, 1992)

Mammary neoplasia

- 26% of insured Saluki bitches in Sweden develop the condition by 10 years of age, compared with 13% for bitches overall (Jitpean *et al.*, 2012)

Physiological conditions**Thyroid hormones**

- Healthy Salukis had lower thyroid hormone values than non-sighthounds in Ireland (Shiel *et al.*, 2010b)

SAMOYED**Dermatological conditions****Endocrine alopecia**

- Samoyed comprised 3.4% of an endocrine alopecia caseload of neutered dogs in the USA (Frank *et al.*, 2003)

Drug reactions**Sulfonamide-associated hypersensitivity**

- Neutered female dogs predisposed in the USA ($p = 0.004$)
- Samoyed comprised 7.5% of a US referral caseload and were significantly over-represented ($p < 0.001$) (Trepanier *et al.*, 2003)

Endocrine conditions**Diabetes mellitus**

- Familial and inherited
- Older entire females are predisposed
- OR 21.7 (95% CI 14.7–31.9) compared with crossbreeds in the UK
- OR 11.8 (95% CI 5.5–23.3) compared with crossbreeds in a US referral study
- Samoyed had second-highest breed incidence in Sweden: 104 cases per 10 000 DYAR (95% CI 72–136)
- Samoyed comprised 4.6% of cases, compared with <0.1% of the overall population of insured dogs, in the UK (Hess *et al.*, 2000; Kimmel *et al.*, 2002; Davison *et al.*, 2005; Fall *et al.*, 2007; Catchpole *et al.*, 2008)

Gastrointestinal conditions

Chronic hepatitis

- Multiple possible causes, so breed predisposition may change with time and geographic location due to genetic and environmental factors
- Median age at diagnosis in the UK was 8 years
- Samoyed had OR 12.6 (95% CI 5.3–29.9) compared with an overall laboratory population in the UK

(Bexfield *et al.*, 2012a)

Congenital portosystemic shunt

- Referred Samoyeds (0.15% prevalence) had OR 2.9 (95% CI 1.1–7.6) compared with an overall US referral population

(Tobias & Rohrbach, 2003)

Musculoskeletal conditions

Hip dysplasia

- Two studies report odds ratios of 4.5 (95% CI 3.4–5.9) and 2.70 (95% CI 2.50–2.92) compared to crossbreeds in the USA
- Samoyed had 11.4% prevalence in the USA (LaFond *et al.*, 2002; Witsberger *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Neoplastic conditions

Squamous cell carcinoma – lingual

- Females predisposed: OR 2.26 (95% CI 1.45–3.54)
- OR 24.63 (95% CI 2.85–212.92) compared with other breeds with lingual neoplasia at a US diagnostic laboratory

(Dennis *et al.*, 2006)

Neurological conditions

Hypomyelination syndrome

- Reported in related Samoyed dogs
- Sex-linked inheritance suspected
- Signs seen at 2–8 weeks

(Cummings *et al.*, 1986)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.42%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Glaucoma – primary

- Evidence for a hereditary component in the Samoyed
- Samoyed had 1.59% prevalence in a US referral population, 1994–2002

(Ekestén & Torräng, 1995; Gelatt & MacKay, 2004a)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Familial glomerular diseases which may progress to renal failure
- Presents from 3 months onwards, with males predisposed
- 25% of one experimental group affected in the USA
- Described in Samoyeds in Canada

(Bernard & Valli, 1977; Jansen *et al.*, 1987)

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- OR 4.69 compared with crossbreeds in the USA

(Lekcharoensuk *et al.*, 2000a)

Urolithiasis – silica

- OR 4.2 (95% CI 2.6–6.6) compared with crossbred dogs in the USA

(Low *et al.*, 2010)

SAVOY SHEEPDOG (FRENCH ALPINE MASTIFF)

Musculoskeletal conditions

Short tail (bobtail)

- 11.8% of Savoy Sheepdogs tested in Finland affected

(Hytönen *et al.*, 2009)

SCHAPENDOES (DUTCH SHEEPDOG)

Ocular conditions

Progressive retinal atrophy (PRA)

- Genetics of generalized PRA (GPRA) reported in Schapendoes in Germany: gPRA locus on canine chromosome 20 identified
- Autosomal recessive inheritance described in Schapendoes

(Lippmann *et al.*, 2007; Dekomien *et al.*, 2010)

SCHIPPERKE

Dermatological conditions

Pemphigus foliaceus

- Schipperke had significantly higher risk than an overall referral population in the USA (Ihrke *et al.*, 1985)

Endocrine conditions

Diabetes mellitus

- Familial and inherited
- Older entire females are predisposed
- Schipperke had RR 4.9 (95% CI 2.4–10.8) compared with an overall US referral population (Marmor *et al.*, 1982)

Musculoskeletal conditions

Short tail (bobtail)

- 66.7% of Schipperke tested in Finland affected (Hytönen *et al.*, 2009)

Neurological conditions

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Genetic pathway implicating the *ADAM23* gene shown in Schipperke
- Schipperke had 3.42% prevalence, compared with 0.91% for crossbreeds, in a US referral study (Bellumori *et al.*, 2013; Koskinen *et al.*, 2015)

Mucopolysaccharidosis IIIB

- Autosomal recessive inheritance suspected
- Progressive and fatal
- Schipperke can be an animal model for the human variant of this disease (Ellinwood *et al.*, 2003)

Ocular conditions

Cataract

- Schipperke showed 9.1% prevalence of cataract among ophthalmic referrals in South Africa (Petrick, 1996)

Reproductive conditions

Dystocia

- 28.9% of Schipperke litters born by caesarean section in the UK (Evans & Adams, 2010a)

SCHNAUZER (UNSPECIFIED VARIANT)

Behavioural conditions

Aggression

- Schnauzer (unspecified) had 1.33 times the risk of attacking children in Austria (Schalamon *et al.*, 2006)

Gastrointestinal conditions

Congenital portosystemic shunt

- Referred Schnauzers (2.0% incidence) had an OR of 3.6 (95% CI 1.4–9.2) compared with an overall Australian referral population (Hunt, 2004)

Metabolic conditions

Overweight/obesity

- 35.6% of Schnauzers (unspecified) attending veterinary clinics in China were obese (Courcier *et al.*, 2010; Mao *et al.*, 2013; Raffan *et al.*, 2016)

Neoplastic conditions

Cutaneous melanoma

- Schnauzers (unspecified) had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland (Grüntzig *et al.*, 2016)

Mast cell tumour (MCT)

- Median age of 9 years
- Schnauzer (unspecified) comprised 10% of a US referral caseload of cutaneous MCT (Patnaik *et al.*, 1984a)

Squamous cell carcinoma

- Schnauzers (unspecified) had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland (Grüntzig *et al.*, 2016)

SCHNAUZER – GIANT

Drug reactions

Low thiopurine methyltransferase (TPMT) activity in red blood cells

- Azathioprine is a cytotoxic thiopurine anti-metabolite used to treat neoplasia, control immune-mediated diseases and prevent organ transplant rejection

- Giant Schnauzers in the USA had much lower TPMT activity ($p < 0.001$) than other breeds
- Low TPMT levels may predispose to myelotoxicity with thiopurine drugs (Kidd *et al.*, 2004)

Endocrine conditions

Hypothyroidism

- A genetic pathway for the Giant Schnauzer has been identified
- Congenital hypothyroid dwarfism with suspected autosomal inheritance reported in Giant Schnauzers in the USA
- Giant Schnauzer had a prevalence of 10% in Sweden, and 11.45% prevalence, compared with 1.54% for crossbreeds, in a US referral caseload
- Giant Schnauzer had OR 1.72 ($p = 0.001$) for serum thyroid hormone autoantibodies (THAA) compared with all other breeds in the USA (Greco *et al.*, 1991; Nachreiner *et al.*, 2002; Ferm *et al.*, 2009; Wilbe *et al.*, 2010a; Bellumori *et al.*, 2013)

Metabolic conditions

Overweight/obesity

- Giant Schnauzer had significantly higher body condition score (5.60) than other show dogs in Holland (Courcier *et al.*, 2010; Corbee, 2013; Raffan *et al.*, 2016)

Musculoskeletal conditions

Elbow dysplasia

- Giant Schnauzer had 5.3% prevalence in the UK, 8.2% in the USA, and 25.0% in South Africa
- Giant Schnauzer had 1.9% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland (Kirberger & Stander, 2007; Narojek *et al.*, 2008; Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Hip dysplasia

- Giant Schnauzer prevalence was 18.2% in the USA and 16% in Belgium
- OR 3.0 (95% CI 1.5–6.0) compared to crossbreeds in the USA (LaFond *et al.*, 2002; Coopman *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 3.4 (95% CI 1.7–6.7) compared to crossbreeds in the USA (LaFond *et al.*, 2002)

Neoplastic conditions

Cutaneous haemangioma

- OR 2.7 (95% CI 1.1–6.6) compared to crossbreeds in the USA (Goldschmidt & Mcmanus, 2000)

Neoplasia – overall

- Giant Schnauzer had the tenth-highest proportional mortality from cancer among pedigree breeds in the UK: 41.0% (95% CI 25.6–56.6%) (Adams *et al.*, 2010)

Osteosarcoma

- Highly heritable
- Increased incidence reported in large Schnauzers in Romania, comprising 19.2% of a general caseload (Muste *et al.*, 2010)

Squamous cell carcinoma – digit

- Associated with black coloration in middle-aged to older large-breed dogs
- Reported in related Giant Schnauzers (Paradis *et al.*, 1989)

Renal and urinary conditions

Hyperuricosuria

- Giant Schnauzer had 11.2% prevalence of carriers for the mutation in the USA (Karmi *et al.*, 2010a)

Kidney disease

- Giant Schnauzer had an incidence of 29 (95% CI 19–39) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall, among insured dogs in Sweden (Pelander *et al.*, 2015)

Urethral sphincter mechanism incompetence

- Larger breeds had OR 7.2 compared with smaller breed in the USA
- Giant Schnauzer had increased risk in Germany (Arnold, 1997; Forsee *et al.*, 2013)

SCHNAUZER – MINIATURE

Cardiovascular conditions

Atherosclerosis

- Across all breeds, mean age at diagnosis was 8.5 years, with males over-represented
- Miniature Schnauzer had a higher prevalence among necropsy referral cases in the USA (Liu *et al.*, 1986)

Mitral valve disease

- OR 2.5 (95% CI 1.0–6.4) compared with crossbreeds in a UK primary-care veterinary population (Parker & Kilroy-Glynn, 2012; Mattin *et al.*, 2015a, 2015b)

Patent ductus arteriosus

- Miniature Schnauzer comprised 30.8% of a referral caseload in Germany
- Median age at presentation was 5.7 months (Schneider *et al.*, 2001)

Sick sinus syndrome

- Females predisposed
- Miniature Schnauzer was the most commonly affected breed among referral dogs undergoing pacemaker implantation in the USA
- Characterized in Miniature Schnauzer in the USA (Jochman-Edwards *et al.*, 2002; Wess *et al.*, 2006)

Dental conditions

Canine odontogenic parakeratinized cyst

- Miniature Schnauzer comprised 22.2% of a US referral caseload and was significantly ($p < 0.001$) over-represented compared with the overall hospital population (Verstraete *et al.*, 2011)

Drug reactions

Vaccine-associated adverse effect

- Smaller breeds predisposed
- Miniature Schnauzer had the seventh-highest incidence (64.7 per 10 000 dogs within 3 days of vaccine administration; 95% CI 52.6–78.7), which was significantly higher than the overall incidence in a US primary-care population (38.2 per 10 000 dogs; 95% CI 37.1–39.3) (Moore *et al.*, 2005)

Sulfonamide-associated hypersensitivity

- Neutered female dogs predisposed in the USA ($p = 0.004$)
- Miniature Schnauzers comprised 12.5% of a US referral caseload and were significantly over-represented ($p < 0.001$) (Trepanier *et al.*, 2003)

Endocrine conditions

Diabetes mellitus

- Familial and inherited
- Older entire females are predisposed
- Miniature Schnauzer had odds ratios of 3.13 (95% CI 2.61–3.76) in the USA and 4.1 (95% CI 2.5–6.8) in the UK, compared with crossbreeds
- OR 9.9 (95% CI 6.0–16.0) compared with crossbreeds in a US referral study
- Miniature Schnauzer had an incidence of 20 cases per 10 000 DYAR (95% CI 10–30) in Sweden
- OR 2.6 (95% CI 1.9–3.5) compared with an overall US referral population (Marmor *et al.*, 1982; Hess *et al.*, 2000; Guptill *et al.*, 2003; Fall *et al.*, 2007; Catchpole *et al.*, 2008)

Hypothyroidism

- Miniature Schnauzer was at increased risk in a US referral study (Milne & Hayes, 1981)

Primary hypoparathyroidism

- Miniature Schnauzer comprised 14.3% of a US referral case series (Bruyette & Feldman, 1988)

Gastrointestinal conditions

Cleft lip and/or palate

- Birth defect; small breeds predisposed
- Miniature Schnauzers had RR 11.0 compared with all referral dogs in the USA (Mulvihill *et al.*, 1980)

Congenital megaesophagus

- Affects puppies, but most recover by 6 months of age
- Autosomal dominant or a 60% penetrance autosomal recessive mode of inheritance suspected, but polygenic inheritance in outcrosses
- Described in Miniature Schnauzer puppies from a colony (Cox *et al.*, 1980)

Congenital portosystemic shunt

- Small breeds predisposed
- Usually presents in younger dogs
- Referred Miniature Schnauzers (1.0% prevalence) had OR 19.8 (95% CI 14.0–28.0) compared with an overall US referral population
- Miniature Schnauzers comprised 8.0% and 14.0% of two US referral caseloads
(Tobias & Rohrbach, 2003; Winkler *et al.*, 2003; Toulza *et al.*, 2006)

Gallbladder mucocoele

- Associated with hyperlipidaemia
- Older dogs predisposed
- Miniature Schnauzers had OR 3.80 (95% CI 2.21–12.37) compared with overall referral population in Japan
(Kutsunai *et al.*, 2014)

Pancreatitis

- Middle-aged to older, overweight and dogs with diabetes mellitus predisposed
- May be associated with variants of the *SPINK1* gene in Miniature Schnauzers
- OR 4.1 (95% CI 1.9–9.2) compared with an overall referral population in the USA
(Jaeger *et al.*, 2003; Lem *et al.*, 2008; Bishop *et al.*, 2010)

Haematological/immunological conditions**Immune-mediated haemolytic anaemia (IMHA)**

- Females and neutered animals predisposed
- Miniature Schnauzer (4% of cases) was significantly ($p < 0.001$) over-represented compared with a US control referral population (1% were Miniature Schnauzers)
(Weinkle *et al.*, 2005)

Musculoskeletal conditions**Myotonia**

- Puppies show myotonia and skeletal muscle hypertrophy from a few weeks of age
- Associated with severe prognathism in Miniature Schnauzers
- Missense mutation in the *CLC-1* allele identified that may be transmitted as an autosomal dominant (Thomsen's disease) or recessive (recessive generalized myotonia) trait
(Rhodes *et al.*, 1999; Bhalerao *et al.*, 2002)

Spondylocostal dysostosis

- Reported in three puppies of a litter of eight Miniature Schnauzers in Australia
- Consistent with a highly penetrant autosomal recessive monogenic trait
- Frameshift mutation with a single-base deletion in the coding region of *HES7* identified
(Willet *et al.*, 2015)

Neoplastic conditions**Cutaneous melanoma**

- Miniature Schnauzer had OR 7.5 (95% CI 6.0–9.5) compared with a US referral population
(Villamil *et al.*, 2011)

Histiocytic sarcoma

- Miniature Schnauzers had OR 4.8 (95% CI 2.4–88) compared with an overall referral population in the USA
(Lenz *et al.*, 2017)

Meningioma

- Meningiomas represent 51.5% of primary intracranial neoplasms in the USA
- Older dogs predisposed
- Meningioma was more common in Miniature Schnauzers necropsied (4.85% prevalence) than in the overall referral population in the USA ($p = 0.0089$)
(Song *et al.*, 2013)

Neurological conditions**Catechol-O-methyltransferase gene polymorphism**

- Miniature Schnauzer had allele frequencies of 18.8% and 25.0% for two abnormal polymorphisms
(Masuda *et al.*, 2004)

Fibrocartilaginous embolic myelopathy (FCEM)

- Miniature Schnauzer comprised 11.5% of a referral caseload in Japan
- FCEM is the most frequent cause of myelopathy in Miniature Schnauzers
- Mortality is lower in affected Miniature Schnauzers (22%) than in giant breeds (64%)
(Hawthorne *et al.*, 2001; Nakamoto *et al.*, 2009)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Miniature Schnauzer had an incidence of 22.5 per 10 000 DYAR (95% CI 16.9–28.0) among

insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds (Heske *et al.*, 2014)

Intervertebral disc disease (IVDD)

- Males and older dogs predisposed
- Miniature Schnauzer had an incidence rate of 35.83 (95% CI 28.8–42.8), compared with 27.8 for an overall population of insured dogs in Sweden
- Miniature Schnauzer was significantly ($p < 0.01$) over-represented, comprising 1.67% of referral cases, in the Czech Republic (Nečas, 1999; Bergknut *et al.*, 2012)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 4.98%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Miniature Schnauzer had an OR of 3.7 compared with an overall referral population in the USA (Adkins & Hendrix, 2005; Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Progressive retinal atrophy (PRA)

- 23% of a US ophthalmological Miniature Schnauzers caseload affected with photoreceptor dysplasia
- PRA mutation identified in Miniature Schnauzer in the UK (Parshall *et al.*, 1991; Zhang *et al.*, 1998, 1999; Downs *et al.*, 2014a)

Myopia (short-sightedness)

- Miniature Schnauzer had 40.5% prevalence among privately-owned dogs in the USA
- Miniature Schnauzer had a low mean refractive error (−0.67) in the USA (Murphy *et al.*, 1992; Kubai *et al.*, 2008)

Persistent hyperplastic primary vitreous (PHPV)

- Prevalence of 14.2% in Miniature Schnauzers in Canada
- Autosomal recessive mode of inheritance (Grahn *et al.*, 2004)

Photoreceptor dysplasia

- Autosomal recessive disease of Miniature Schnauzers causing retinal degeneration
- Missense mutation disorder (Arg82Gly) identified but other causal mutations are also likely (Zhang *et al.*, 1998)

Retinal dysplasia

- Prevalence of 22.6% in Miniature Schnauzers in Canada
- Autosomal recessive mode of inheritance (Grahn *et al.*, 2004)

Solid intraocular xanthogranuloma

- Associated with concurrent diabetes mellitus, hyperlipidaemia, cataract formation and lens-induced uveitis in Miniature Schnauzers
- Seen in older dogs (9–13 years)
- Reported in Miniature Schnauzers in the USA (Zarfoss & Dubielzig, 2007)

Sudden acquired retinal degeneration (SARD, amaurosis)

- Small breeds predisposed
- Median age 8.1 years
- Miniature Schnauzers comprised 10% of one US caseload, and 7.9% of another US referred caseload (Montgomery *et al.*, 2008; Heller *et al.*, 2017)

Physiological conditions

Increased C-reactive protein (CRP)

- Serum CRP concentrations are moderately to highly inheritable (35–45%) in humans and are associated with atherosclerosis
- Median serum CRP concentrations in healthy Miniature Schnauzer dogs were slightly, but significantly, higher than in healthy non-Miniature Schnauzer dogs in Canada (Wong *et al.*, 2011)

Primary hyperlipidaemia (primary hypertriglyceridaemia, idiopathic hypertriglyceridaemia)

- Associated with insulin resistance and pancreatitis
- Miniature Schnauzers had significantly higher plasma lipids than crossbred dogs in Japan and the USA (Ford, 1993; Xenoulis *et al.*, 2007, 2010; Mori *et al.*, 2010)

Urine composition

- Miniature Schnauzers (0.6–1.8 urinations per 24 hours) urinated significantly less often ($p=0.002$) than Labrador Retrievers (1.5–4.5 urinations per 24 hours) in the UK
- Miniature Schnauzers produced significantly ($p=0.04$) lower volume of urine and significantly ($p=0.007$) higher urine pH than Labrador Retrievers in the UK
- Low urine frequency and volume may predispose to urolithiasis
(Stevenson & Markwell, 2001)

Renal and urinary conditions**Juvenile renal disease**

- Median age 8 months, invariably fatal
- Reported in a series of 8 cases among related Miniature Schnauzers in the USA and Canada
(Morton *et al.*, 1990)

Kidney disease

- Miniature Schnauzer had an incidence of 39 (95% CI 31–46) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden
(Pelander *et al.*, 2015)

Renal calculi

- May be associated with urinary tract infections
- Miniature Schnauzer had OR 14.55 ($p<0.001$) in females and OR 11.62 ($p<0.001$) in males compared with crossbreeds in a US referral study
(Ling *et al.*, 1998c)

Urinary tract infections (UTI)

- Miniature Schnauzer (3.9%) was the sixth most commonly affected breed among referral dogs in the USA for recurrent or persistent UTI
(Norris *et al.*, 2000)

Urolithiasis

- Miniature Schnauzer was the second most common breed for uroliths overall submitted for laboratory testing from Spain and Portugal (11.5%) and the eleventh most common in the UK (2.5%)
(Rogers *et al.*, 2011; Vrabelova *et al.*, 2011)

Urolithiasis – apatite

- Miniature Schnauzer had OR 9.5 (95% CI 8.8–10.4) compared with crossbred dogs in the USA
- Females are predisposed
(Low *et al.*, 2010)

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- Miniature Schnauzer was the most common breed with submissions for calcium oxalate uroliths in Canada
- Miniature Schnauzer had odds ratios of 21.6 (95% CI 19.9–23.4) and 3.9 (95% CI 3.1–4.9) compared with crossbred dogs in two US studies, and 8.3 (95% CI 6.8–10.1) compared with all dogs in the UK
- OR 14.1 (95% CI 10.2–19.5) compared with crossbreeds in the USA
(Lekcharoensuk *et al.*, 2000a; Ling *et al.*, 2003; Houston & Moore, 2009; Low *et al.*, 2010; Roe *et al.*, 2012; Okafor *et al.*, 2014)

Urolithiasis – silica

- Miniature Schnauzer had OR 10.1 (95% CI 8.4–12.2) compared with crossbred dogs and OR 8.3 ($p<0.001$) compared with all dogs in two US studies
(Aldrich *et al.*, 1997; Low *et al.*, 2010)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Small breeds and females are predisposed
- Miniature Schnauzer was the third most common breed with submissions for struvite uroliths in Canada
- Miniature Schnauzer had OR 8.0 (95% CI 7.5–8.7) compared with crossbred dogs in the USA and OR 6.3 (95% CI 5.3–7.6) compared with all dogs in the UK
(Houston & Moore, 2009; Low *et al.*, 2010; Roe *et al.*, 2012; Okafor *et al.*, 2014)

Urolithiasis – urate

- Miniature Schnauzer had OR 12.5 (95% CI 11.3–14.0) compared with crossbred dogs in the US and OR 3.1 (95% CI 2.0–4.9) compared with all dogs in the UK
- Younger and female animals predisposed
(Low *et al.*, 2010; Roe *et al.*, 2012)

Reproductive conditions**Cryptorchidism**

- Miniature Schnauzer had RR 1.8 (95% CI 1.5–2.3) and OR 2.2 (95% CI 1.52–3.03) compared with overall referral populations in two US studies
- Evidence of inheritance in US Miniature Schnauzers
(Pendergrass & Hayes, 1975; Cox *et al.*, 1978; Hayes *et al.*, 1985)

Dystocia

- Miniature Schnauzer had an incidence of 10.1 cases per 1000 DYAR among insured bitches in Sweden, compared with 5.7 per 1000 DYAR overall

(Bergström *et al.*, 2006)

Male pseudohermaphroditism (persistent Müllerian duct syndrome)

- Affected Miniature Schnauzers are 78,XY males with both male and female reproductive organs
- Inherited as an autosomal recessive trait with expression limited to homozygous males
- Caused by C-to-T transition in exon 3 of the Müllerian inhibiting substance type II receptor (MISRII)
- The mutation is identical by descent in affected Miniature Schnauzers in the USA

(Pujar & Meyers-Wallen, 2009;

Vegter *et al.*, 2010)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- OR 2.74 (95% CI 1.74–4.29) compared with all entire bitches in Finland

(Niskanen & Thrusfield, 1998;

Egenvall *et al.*, 2001; Smith, 2006)

SCHNAUZER – STANDARD**Cardiovascular conditions****Congenital heart disease**

- Includes a range of congenital heart disorders
- Standard Schnauzer had OR 7.1 compared with all referral dogs in Italy

(Oliveira *et al.*, 2011)

Endocrine conditions**Diabetes mellitus**

- Familial and inherited
- Peak incidence age 7–9 years
- Older entire females are predisposed
- OR 4.78 (95% CI 2.91–7.82) compared with crossbreeds in the USA

(Guptill *et al.*, 2003; Catchpole *et al.*, 2008)

Gastrointestinal conditions**Congenital portosystemic shunt**

- Referred Standard Schnauzers (0.82% prevalence) had an OR of 16.1 (95% CI 8.0–30.0) compared with an overall US referral population

(Tobias & Rohrbach, 2003)

Neoplastic conditions**Haemangiosarcoma**

- Standard Schnauzer had OR 10.08 (95% CI 0.72–141.95) compared with crossbreeds in the USA

(Hargis *et al.*, 1992)

Osteosarcoma

- Highly heritable
- Standard Schnauzer had incidence rate of 9 (95% CI 2–15) per 10 000 DYAR in Sweden

(Egenvall *et al.*, 2007; Karlsson *et al.*, 2013)

Ocular conditions**Cataract**

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 4.73%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005;

Donzel *et al.*, 2016)

Renal and urinary conditions**Urolithiasis – calcium oxalate**

- Small breeds and males are predisposed
- Standard Schnauzer had OR 18.06 compared with crossbreeds in the USA

(Lekcharoensuk *et al.*, 2000a)

SCOTTISH DEERHOUND (DEERHOUND)**Musculoskeletal conditions****Arthrosis of cervical articular facet joints**

- Described in 64.3% of Scottish Deerhounds tested in Germany

(Kinzel *et al.*, 2003)

Osteochondrodysplasia

- Described in related Scottish Deerhound pups
- Single autosomal recessive mode of inheritance suspected
- Pups appear normal at birth but show exercise intolerance and retarded growth by 4–5 weeks of age

(Breur *et al.*, 1989)

Neoplastic conditions**Osteosarcoma**

- Highly heritable: heritability estimated at 0.69

- May be at least two different genetic risk factors for osteosarcoma in Scottish Deerhounds, with differing inheritance patterns (Phillips, 2007; Phillips *et al.*, 2010; Karlsson *et al.*, 2013; Dillberger & McAtee, 2017)

Physiological conditions

Unusual haematological and biochemical normal values

- Scottish Deerhounds in the USA and Canada shared clinicopathological similarities with other sighthound breeds that are not found in other breeds: low platelet counts, high mean corpuscular volume (MCV), low total calcium concentration, high alanine aminotransferase (ALT) activity
- High serum cholesterol appeared unique to Scottish Deerhounds: high serum cholesterol (Sheerer *et al.*, 2013)

SCOTTISH TERRIER

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- A survey of UK veterinarians classified Scottish Terrier as having high aggression (Bradshaw & Goodwin, 1999)

Dermatological conditions

Demodicosis

- Scottish Terrier was described as a predilected breed in a Swedish referral study (Holm, 2003)

Familial vasculopathy of the nasal planum, nostrils and nasal mucosa

- Described in Scottish Terrier puppies
- Probably autosomal dominant inheritance (Pedersen & Scott, 1991)

Superficial necrolytic dermatitis

- Males and older dogs may be predisposed
- Scottish Terrier suggested as predisposed, and comprised 8.3% of a US referral caseload (Outerbridge *et al.*, 2002)

Endocrine conditions

Primary hypoparathyroidism

- Scottish Terrier comprised 7.1% of a US referral case series (Bruyette & Feldman, 1988)

Gastrointestinal conditions

Chronic hepatitis

- Scottish Terrier was 10 times over-represented ($p < 0.001$) in laboratory samples in Sweden (Andersson & Sevelius, 1991)

Congenital portosystemic shunt

- Referred Scottish Terriers (0.23% prevalence) had an OR of 4.4 (95% CI 1.4–10.8) compared with an overall US referral population (Tobias & Rohrbach, 2003)

Haematological/immunological conditions

Low immunological response to vaccination

- Scottish Terrier was in the lowest quartile for titre response to rabies vaccination in the UK (Kennedy *et al.*, 2007)

Raised serum alkaline phosphatase (ALP)

- Scottish Terriers had significantly ($p < 0.001$) higher mean serum ALP activity (mean \pm SD, 1520 ± 2010 U/l) than control dogs (306 ± 697 U/l)
- Scottish Terriers also had OR 2.4 (95% CI 1.3–4.3) to have a disease associated with high serum ALP activity compared with a control population (Nestor *et al.*, 2006)

von Willebrand's disease (vWD)

- Scottish Terriers are affected by type III vWD, and the genetic mechanism in this breed in the USA has been elucidated
- Scottish Terriers in Australia had a prevalence of 27% (Stokol *et al.*, 1995; Venta *et al.*, 2000)

Musculoskeletal conditions

Patellar luxation

- 6.1% of US Scottish Terriers affected
- Mainly medial luxation observed, often bilateral (Orthopedic Foundation for Animals, 2015)

Neoplastic conditions

Cardiac tumour

- Uncommon, and can occur both as primary and as metastatic lesions
- Scottish Terrier had the highest incidence (0.49% incidence) and RR 2.50 (95% CI 1.55–4.03) compared with an overall referral population in the USA (Ware & Hopper, 1999)

Cutaneous melanoma

- OR 3.1 (95% CI 1.7–5.6) compared with a US referral population (Villamil *et al.*, 2011)

Lower urinary tract neoplasia

- Scottish Terrier was over-represented for urinary bladder neoplasia in the UK, comprising 5.7% of a referral caseload, and also comprised 22.6% of a US referral caseload
- OR 11.6 for lower urinary tract tumours compared with an overall referral population in Canada (Burnie & Weaver, 1983; Norris *et al.*, 1992; Schrempp *et al.*, 2013)

Lymphoma

- Scottish Terrier had odds ratios of 3.4 (95% CI 3.0–3.9) and 5.6 (95% CI 3.4–9.0) compared with referral populations in two US studies, and was significantly ($p < 0.05$) over-represented in another US referral study
- RR 9.16 (95% CI 3.05–27.21) compared with an overall referral population in the Netherlands (Keller *et al.*, 1993; Teske *et al.*, 1994; Villamil *et al.*, 2009, 2011)

Primary brain tumour

See under *Neurological conditions*

Prostate neoplasia

- Neutering associated with 3.56 times increased odds
- OR 3.81 ($p < 0.0001$) compared with an overall referral population in the USA
- OR 7.43 (95% CI 2.79–18.6) compared with an overall referral population in the Netherlands (Teske *et al.*, 2002; Bryan *et al.*, 2007)

Neurological conditions**Acquired myasthenia gravis**

- Scottish Terrier had significantly higher risk than crossbred dogs in the USA (Shelton *et al.*, 1997)

Central axonopathy

- Described in related Scottish Terriers in Switzerland
- Heritability suspected
- Tremors occur at 10–12 weeks (van Ham *et al.*, 1994)

Cerebellar degeneration

- Cases identified from several countries around the world
- Litter analysis consistent with fully penetrant simple autosomal recessive inheritance
- Minimum prevalence of 0.07% among Scottish Terriers registered with the American Kennel Club
- Characterized in in Scottish Terriers in the USA (Urkasemsin *et al.*, 2010)

Muscle cramping ('Scottie cramp')

- Autosomal recessive inheritance
- Age of onset 6 weeks to 18 months
- Females over-represented
- Characterized in in Scottish Terriers in the USA (Urkasemsin & Olby, 2015)

Primary brain tumour

- Scottish Terriers were over-represented in the USA, comprising 5% of cases but < 1% of the overall referral population (Heidner *et al.*, 1991)

Renal and urinary conditions**Urolithiasis – struvite (magnesium ammonium phosphate)**

- OR 4.8 (95% CI 3.7–6.2) compared with all dogs in the UK
- Small breeds and males predisposed (Roe *et al.*, 2012)

Urolithiasis – urate

- OR 4.2 (95% CI 3.2–5.5) compared with crossbred dogs in the USA
- Younger and female animals predisposed (Low *et al.*, 2010)

Reproductive conditions**Dystocia**

- 59.8% of Scottish Terrier litters born by caesarean section in the UK
- Scottish Terrier had the highest incidence (38.3 cases per 1000 DYAR) among insured bitches in Sweden, compared with 5.7 per 1000 DYAR overall (Bergström *et al.*, 2006; Evans & Adams, 2010a)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 41% of insured Scottish Terrier bitches develop the condition by 10 years of age, compared with 19% for bitches overall

(Jitpean *et al.*, 2012)

SEALYHAM TERRIER

Ocular conditions

Keratoconjunctivitis sicca

- Sealyham Terriers were the second most commonly affected breed in a European multi-centre study, comprising 17% of the cases

(Ofri *et al.*, 2009)

Lens luxation – primary

- *ADAMTS17* mutation has been identified: 49.0% of UK and 36.0% of US Sealyham Terriers were carriers

(Gould *et al.*, 2011)

SENNENHUND

See *Swiss Mountain Dog*

SHAR PEI (CHINESE SHAR PEI)

Dermatological conditions

Acute febrile neutrophilic vasculitis

- Acute widespread dermatopathy associated with fever and malaise
- Described in 3 Shar Pei in Australia

(Malik *et al.*, 2002)

Atopic dermatitis (atopy)

- 41.7% prevalence among Shar Peis in the Czech Republic
- Shar Pei comprised 2.6% (95% CI 1.1–5.3) of the atopic caseload, compared with 0.7% of overall hospital population, in a US referral study
- OR 25.53 ($p=0.0005$) compared with a referral population in Greece

(Saridomichelakis *et al.*, 1999;

Zur *et al.*, 2002; Počta & Svoboda, 2007)

Cutaneous mucinosis

- Condition is associated with the distinctive wrinkled, thickened skin selected for in Shar Pei

- Associated with mast cell activity, hyaluronic acid and a consequence of over-transcription or increased activity of the HAS2 allele

(Welle *et al.*, 1999; Zanna *et al.*, 2008, 2009;

Docampo *et al.*, 2011)

Demodicosis

- An inheritance pathway has been described
- Short-haired breeds predisposed
- Shar Pei had the third-highest breed OR of 7.2 (95% CI 2.9–17.6) compared with an overall US first-opinion population

(It *et al.*, 2010; Plant *et al.*, 2011)

Folliculitis

- Shar Pei had 36.4% prevalence in the USA

(Miller *et al.*, 1992)

Malassezia dermatitis

- Shar Pei had the third-highest prevalence (6.8%) among a dermatology referral caseload in Romania

(Mircean *et al.*, 2010)

Otitis externa

- Shar Pei was significantly over-represented ($p<0.001$) in Israel

(Zur *et al.*, 2011)

Pemphigus foliaceus

- OR 7.9 (95% CI 2.2–30.1) compared with a referral population in the USA

(Kuhl *et al.*, 1994)

Gastrointestinal conditions

Canine autoinflammatory disease (AID)

- Results from the dysregulation of mediators of the innate immune system
- Associated with selection for the heavily thickened and wrinkled skin traits in Shar Pei
- Genetic signature flanking the chromosome 13 AID locus identified

(Olsson *et al.*, 2013)

Hiatal hernia

- Can be congenital or acquired
- Congenital form described in related Shar Pei dogs in France, and in unrelated Shar Pei in the USA
- Ages 2–11 months at diagnosis
- Acquired form often associated with severe upper respiratory disease

(Callan *et al.*, 1993; Guiot *et al.*, 2008)

Haematological/immunological conditions

Immunodeficiency syndrome

- Associated with recurrent infections, autoimmune disease and neoplasia
- Described in ten Shar Pei dogs in the USA (Rivas *et al.*, 1995)

Metabolic conditions

Hypocobalaminaemia

- Mutation on chromosome 13 suggested as causative in Shar Pei
- Shar Peis with or without gastrointestinal disease are predisposed
- Shar Pei had the highest breed OR (21.1, 95% CI 9.1–54.5) compared with a laboratory population in the UK
- 64.0% of Shar Pei serum samples in the USA had subnormal cobalamin concentrations, and Shar Pei had significantly lower values than all other breeds ($p < 0.05$) (Grützner *et al.*, 2010; Bishop *et al.*, 2012; Dandrieux *et al.*, 2013)

Familial Shar Pei fever

- Part of the overall autoinflammatory disease syndrome of Shar Pei
- A 16.1 kb duplication located approximately 350 kb upstream of HAS identified and proposed as a causative mutation for both hyaluronanosis and Shar Pei fever (Olsson *et al.*, 2011)

Musculoskeletal conditions

Elbow dysplasia

- Shar Pei prevalence was 31% in Belgium
- OR 4.6 (95% CI 2.2–9.4) for ununited anconeal process compared to crossbreeds in the USA
- Shar Pei had 3.8% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland (LaFond *et al.*, 2002; Coopman *et al.*, 2008; Narojek *et al.*, 2008; Michelsen, 2013)

Hip dysplasia

- Shar Pei prevalence was 16.1% in US and 21% in Belgium (Coopman *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 3.5 (95% CI 2.8–4.5) compared to crossbreeds in the USA (LaFond *et al.*, 2002)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- OR 11.4 (95% CI 7.4–17.6) compared to crossbreeds in the USA (LaFond *et al.*, 2002)

Neoplastic conditions

Canine cutaneous histiocytoma

- OR 2.10 (95% CI 1.73–2.55) compared with crossbreeds in the USA (Goldschmidt & Mcmanus, 2000; Fulmer & Mauldin, 2007)

Histiocytic sarcoma

- OR 16.0 (95% CI 4.1–44.0) compared with an overall referral population in the USA (Lenz *et al.*, 2017)

Lymphoma – gastrointestinal

- Shar Pei comprised 13.6% of cases in a US study (Coyle & Steinberg, 2004)

Mast cell tumour (MCT)

- Mean age at presentation between 7.5 and 9 years, but can occur at any age
- Three US studies showed Shar Pei had odds ratios of 7.81 (95% CI 2.71–22.55), 2.4 (95% CI 2.0–2.9) and 3.84 (95% CI 2.89–5.11) compared with crossbreeds
- Shar Pei comprised 2.2% of MCTs but 1% of the submitting population to a US diagnostic laboratory (Miller, 1995; Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011; White *et al.*, 2011)

Melanoma- oral

- Large-breed dogs predisposed compared with small-breed dogs: OR 6.88 (95% CI 2.95–15.05)
- Shar Pei had OR 24.43 (95% CI 2.98–200.18) compared with other breeds with lingual neoplasia at a US diagnostic laboratory (Dennis *et al.*, 2006)

Ocular conditions

Entropion

- Clear breed predisposition, and polygenic inheritance suspected
- Shar Pei comprised 20.0% of referral caseload in Australia

(Read & Broun, 2007)

Glaucoma – primary

- Shar Pei had 4.4% prevalence in a US referral population, 1994–2002

(Gelatt & MacKay, 2004a)

Lens luxation – primary

- Autosomal recessive inheritance suspected
- Shar Pei comprised 20% and 23.5% of two US ophthalmology caseloads
- RR 7.18 (95% CI 5.66–9.12) and RR 44.66 (95% CI 16.94–117.73) compared with overall referral dogs in the USA

(Lazarus *et al.*, 1998; Binder *et al.*, 2007; Sargan *et al.*, 2007)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Usually presents in the first 1–2 years of life
- Inheritance is complex and potentially multigenic
- Shar Pei significantly over-represented in France

(Mazzucchelli *et al.*, 2012; Edelmann *et al.*, 2013)

Renal and urinary conditions

Renal amyloidosis

- Inherited disease with complicated inheritance pattern
- Median age at presentation 4.5 years and median survival time 5 days in Israel
- Reported in 16 related Shar Pei in the USA
- Amyloidosis was over-represented among Shar Pei with glomerulonephritis (88.9% of cases) in the USA

(DiBartola *et al.*, 1990; Rivas *et al.*, 1993; Segev *et al.*, 2012; Schneider *et al.*, 2013)

Respiratory conditions

Brachycephalic obstructive airway syndrome (BOAS)

- Shar Pei had a prevalence of 6.67% among referral dogs in Belgium

(Njikam Nsangou *et al.*, 2009; Roedler *et al.*, 2013; Marchant *et al.*, 2017)

Soft-tissue conditions

Hiatal hernia

See *Gastrointestinal conditions*

Perineal herniation

- Shar Pei suggested as predisposed by case reports from the USA, UK and Canada (Williams, 1990; Callan *et al.*, 1993; Auger & Riley, 1997)

SHETLAND SHEEPDOG (SHELTIE)

Behavioural conditions

Vocalization

- Shetland Sheepdog had 8.9% prevalence, compared with an overall prevalence of 1.5%, in a US hospital population (Bamberger & Houpt, 2006)

Cardiovascular conditions

Left atrial rupture

- Associated with chronic mitral valve insufficiency
- Shetland Sheepdog was the most commonly affected breed in a US referral study, comprising 21.4% of the caseload

(Reineke *et al.*, 2008)

Patent ductus arteriosus

- Small breeds (median body weight 5 kg) and females predisposed
- Median age was 4 months
- Shetland Sheepdog was the most common breed affected, comprising 13.5% of a US referral caseload

(Bureau *et al.*, 2005)

Dermatological conditions

Familial canine dermatomyositis

- Shetland Sheepdogs and collie breeds predisposed
- Described in Shetland Sheepdog in the UK
- Inherited as an autosomal dominant trait with incomplete penetrance
- Linkage disequilibrium identified a microsatellite marker FH3570 on chromosome 35 in Shetland Sheepdogs in the USA

(Ferguson *et al.*, 2000; Clark *et al.*, 2005; Wahl *et al.*, 2008)

Superficial necrolytic dermatitis

- Males and older dogs may be predisposed
- Shetland Sheepdogs suggested as predisposed; comprised 13.9% of a US referral caseload (Outerbridge *et al.*, 2002)

Vesicular cutaneous lupus erythematosus

- Associated with summertime and sun exposure
- Autoimmune pathogenesis suggested
- Described in Shetland Sheepdogs in the USA (Jackson, 2004; Jackson *et al.*, 2004)

Drug reactions**Multiple drug sensitivity**

- High doses of many drugs (e.g. ivermectin or milbemycin) cause neurological signs
- Associated with the nt230(del4) *MDR1* mutation
- Shetland Sheepdog had a 30.3% allelic frequency with 9% recessive mutant allele frequency in Europe, and a 36% allelic frequency with 12% recessive mutant allele frequency in the UK
- Shetland Sheepdog had a 7.9% allelic frequency in Brazil (Geyer *et al.*, 2005; Dowling, 2006; Gramer *et al.*, 2011; Tappin *et al.*, 2012; Monobe *et al.*, 2015; Firdova *et al.*, 2016)

Endocrine conditions**Hypothyroidism**

- Shetland Sheepdog was at increased risk in a US referral study
- OR 1.69 ($p=0.001$) for serum thyroid hormone autoantibodies (THAA) compared with all other breeds in the USA (Milne & Hayes, 1981; Nachreiner *et al.*, 2002)

Gastrointestinal conditions**Congenital portosystemic shunt**

- Referred Shetland Sheepdogs (0.26% prevalence) had OR 5.2 (95% CI 3.1–8.8) compared with an overall US referral population (Tobias & Rohrbach, 2003)

Gallbladder mucocoele

- Mutation identified for gallbladder mucocoele in Shetland Sheepdog
- OR 7.2 ($p<0.001$) compared with the overall referral population in the USA: 10.2% of Shetland Sheepdogs affected, compared with 1.6% of the overall population

- OR 4.52 (95% CI 2.43–15.86) compared with overall referral population in Japan
- Shetland Sheepdog comprised 8.8% of a US referral caseload

(Aguirre *et al.*, 2007; Crews *et al.*, 2009; Mealey *et al.*, 2010; Kutsunai *et al.*, 2014)

Haematological/immunological conditions**Hypercholesterolaemia**

- Plasma cholesterol was above normal in 43.8% of Shetland Sheepdogs, compared with 12.2% of control dogs, in Japan (Sato *et al.*, 2000)

Low immunological response to vaccination

- Shetland Sheepdog was in the lowest quartile for titre response to rabies vaccination in the UK (Kennedy *et al.*, 2007)

von Willebrand's disease (vWD)

- Shetland Sheepdog had a 23% prevalence among tested dogs in the USA
- This breed affected by severe type III disease (Raymond *et al.*, 1990; Pathak, 2004)

Metabolic conditions**Overweight/obesity**

- Shetland Sheepdogs had an OR of 1.9 (95% CI 1.2–2.8) for obesity in the USA, and were significantly associated with obesity in the UK (Edney & Smith, 1986; Lund *et al.*, 2006; Courcier *et al.*, 2010; Raffan *et al.*, 2016)

Musculoskeletal conditions**Distal tibial valgus deformity**

- Median age at presentation was 9 months
- RR 12.3 (95% CI 1.5–30.5) compared with an overall referral population in the USA (Jaeger *et al.*, 2007)

Elbow dysplasia

- Shetland Sheepdog had 3.0% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland (Narojek *et al.*, 2008; Michelsen, 2013)

Lateral luxation of the superficial digital flexor tendon

- Simple autosomal recessive inheritance
- Genetics characterized in Shetland Sheepdogs in Finland (Solanti *et al.*, 2002)

Neoplastic conditions

Lower urinary tract neoplasia

- Shetland Sheepdog comprised 16.1% of a US referral caseload

(Schrempp *et al.*, 2013)

Oral epulis

- Shetland Sheepdog comprised 19.6% of a fibromatous epulis caseload in Japan

(Yoshida *et al.*, 1999)

Prostate neoplasia

- Neutering associated with 3.56 times increased odds
- Shetland Sheepdog had OR 1.82 ($p=0.0005$) compared with an overall referral population in the USA

(Bryan *et al.*, 2007)

Thyroid neoplasia

- Large breeds and older dogs predisposed
- Shetland Sheepdog had OR 1.7 (95% CI 1.05–2.77) compared with an overall referral population in the USA

(Wucherer & Wilke, 2010)

Neurological conditions

Canine spongiform leucoencephalomyelopathy

- Described in related Shetland Sheepdogs in the USA
- Clinical signs develop from 7 days to 3 weeks
- Missense mutation in cytochrome b identified, and maternal inheritance suggested

(Li *et al.*, 2006)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Shetland Sheepdog had an incidence of 19.6 per 10 000 DYAR (95% CI 15.2–24.0) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
- Cliniconeuropathology described in Shetland Sheepdogs in Japan

(Morita *et al.*, 2002; Heske *et al.*, 2014)

Intervertebral disc disease (IVDD)

- Shetland Sheepdog comprised 1.5% of a cervical IVDD caseload at a Japanese referral hospital

(Itoh *et al.*, 2008)

Leucodystrophy

- Described in related Shetland Sheepdogs in the USA

- Mode of inheritance unknown
- Clinical signs began at 7 days to 3 weeks of age (Wood & Patterson, 2001)

Ocular conditions

Chronic superficial keratitis (pannus)

- Shetland Sheepdog was significantly over-represented in the USA, comprising 2.38% of referral cases

(Slatter *et al.*, 1977)

Collie eye anomaly

- Congenital; autosomal recessive inheritance with nearly 100% penetrance
- Genetic mechanism in Shetland Sheepdog identified

(Lowe *et al.*, 2003; Parker *et al.*, 2007)

Physiological conditions

Low Schirmer tear test (STT)

- Shetland Sheepdog had lower normal STT results than other breeds tested
- Mean value for STT was 15.8 ± 1.8 mm/min in Shetland Sheepdogs in the USA

(Hamor, 2000)

Primary hyperlipidaemia (primary hypertriglyceridaemia, idiopathic hypertriglyceridaemia)

- Shetland Sheepdog had significantly higher plasma lipids than crossbred dogs in Japan

(Mori *et al.*, 2010)

Renal and urinary conditions

Kidney disease

- Shetland Sheepdog had an incidence of 31 (95% CI 25–37) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden

(Pelander *et al.*, 2015)

Reproductive conditions

Cryptorchidism

- Shetland Sheepdog had RR 1.8 (95% CI 1.4–2.4) and OR 2.0 (95% CI 1.24–3.10) compared with overall referral populations in two US studies

- Shetland Sheepdog had 3.2% prevalence, compared with 2.1% in an overall population in the Netherlands

(Pendergrass & Hayes, 1975;

Hayes *et al.*, 1985; Gubbels *et al.*, 2009)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- RR 2.4 (95% CI 1.5–3.3) compared with all bitches insured by Agria for veterinary care in Sweden (Egenvall *et al.*, 2001; Smith, 2006)

Testicular neoplasia

- Shetland Sheepdog had a proportional morbidity ratio of 5.7 (95% CI 3.8–8.6) compared with all other dogs in a Norwegian cancer registry (Nødtvedt *et al.*, 2011)

SHIBA INU

See *Japanese Shiba Inu*

SHIH TZU

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- A Canadian veterinary survey showed that 22.0% of Shih Tzu aged ≥ 1 year had bitten (Guy *et al.*, 2001; Zapata *et al.*, 2016)

Cardiovascular conditions

Mitral valve disease

- OR 2.9 (95% CI 1.5–5.5) compared with crossbreeds in a UK primary-care veterinary population (Parker & Kilroy-Glynn, 2012; Mattin *et al.*, 2015a, 2015b)

Dental conditions

Dentigerous cyst

- Brachycephalic breeds over-represented
- Shih Tzu comprised 6.9% of a US referral caseload and was significantly ($p < 0.001$) over-represented compared with the overall hospital population (Verstraete *et al.*, 2011)

Periodontal disease

- Shih Tzu comprised 15.4% of a dental referral caseload of small breeds in South Korea (Kim *et al.*, 2013)

Dermatological conditions

Demodicosis

- Shih Tzu was over-represented in the UK: 6.45% of laboratory diagnoses versus 0.45% of normal clinic population (Day, 1997a)

Malassezia dermatitis

- RR 6.46 compared to a general laboratory population in the USA (Mauldin *et al.*, 1997)

Drug reactions

Anaesthetic-related complications

- Shih Tzu had 5% incidence among general small animal practices in Canada (Dyson *et al.*, 1998)

Gastrointestinal conditions

Chronic hypertrophic pyloric gastropathy

- Shih Tzu was the most commonly affected breed in a referral case series in Australia
- Mean age was 8.2 years and mean body weight was 6.5 kg
- Males may be predisposed (Bellenger *et al.*, 1990)

Congenital portosystemic shunt

- Usually presents in young dogs
- Referred Shih Tzu (0.78% prevalence) had OR 15.4 (95% CI 10.1–23.4) compared with an overall US referral population
- Referred Shih Tzu (3.4% incidence) had OR 3.9 (95% CI 1.8–8.2) compared with an overall Australian referral population
- Shih Tzu comprised 5.3% and 10.9% of two US referral caseloads (Tobias & Rohrbach, 2003; Winkler *et al.*, 2003; Hunt, 2004; Toulza *et al.*, 2006)

Portal vein hypoplasia

- May occur in conjunction with portosystemic shunt (PSS)
- Cases present from 6 months to 6 years
- Shih Tzu comprised 10.7% of a US referral caseload (Christiansen *et al.*, 2000)

Metabolic conditions

Overweight/obesity

- 32.2% of Shih Tzu attending veterinary clinics in China were obese (Courcier *et al.*, 2010; Mao *et al.*, 2013; Raffan *et al.*, 2016)

Musculoskeletal conditions

Chondrodysplasia (short-limbed or disproportional dwarfism)

- Part of the Shih Tzu breed description
- An *fgf4* retrogene identified as a cause of chondrodysplasia (Israel *et al.*, 2009; Parker *et al.*, 2009)

Patellar luxation

- Mainly medial luxation observed, often bilateral
- Shih Tzu had 2.3 OR (95% CI 1.6–3.4) compared to crossbreeds in the USA (LaFond *et al.*, 2002)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.4 (95% CI 1.1–1.8) compared to crossbreeds in the USA (LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Neoplastic conditions

Mast cell tumour (MCT)

- Shih Tzu had an OR of 3.58 for high-grade MCT tumours compared with all breeds and showed an approximately twofold increase in the proportion of high-grade MCT tumours compared with all breeds (16.9 vs. 8.2%) (Mochizuki *et al.*, 2016)

Oral epulis

- Shih Tzu comprised 9.3% of a fibromatous epulis caseload in Japan (Yoshida *et al.*, 1999)

Neurological conditions

Atlantoaxial subluxation/instability

- Congenital or developmental condition
- Young small breeds affected
- Shih Tzu comprised 6.5% and 16.7% of two US referral caseloads (Beaver *et al.*, 2000; Sanders *et al.*, 2004)

Intervertebral disc disease (IVDD)

- Males and older dogs predisposed
- Shih Tzu comprised 11.4% of a cervical IVDD caseload and 4.8% of a thoracolumbar IVDD caseload at a Japanese referral hospital
- Shih Tzu comprised 5.1% of a US thoracolumbar caseload, and 2.2% of a US referral population
- RR 3.9 ($p < 0.01$) compared with an overall referral population in the USA

- Shih Tzu had an incidence rate of 39.6 (95% CI 30.6–48.6), compared with 27.8 for an overall population of insured dogs in Sweden (Priester, 1976; Bartels *et al.*, 2003; Itoh *et al.*, 2008; Israel *et al.*, 2009; Bergknut *et al.*, 2012)

Necrotizing meningoencephalitis

- Identified in a US Shih Tzu (Cooper *et al.*, 2014b)

Ocular conditions

Bacterial keratitis

- OR 44.7 (95% CI 8.5–234.6) compared with a US referral population (Tolar *et al.*, 2006)

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 4.14%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Corneal ulceration (ulcerative keratitis)

- Shih Tzu was the commonest breed, comprising 50% of a referral caseload in Korea
- Shih Tzu had the third-highest primary-care veterinary predisposition in the UK, with an OR of 10.04 (95% CI 7.30–13.80) compared with crossbreeds (Kim *et al.*, 2009; O'Neill *et al.*, 2017b)

Epiphora

- Associated with anatomical features such as medial canthal entropion, trichiasis, tight medial palpebral ligament and close apposition of the eyelids to the globe
- Shih Tzu comprised 65.2% of a surgical referral caseload in Korea (Yi *et al.*, 2006)

Glaucoma

- Shih Tzu had 1.58% prevalence in a US referral population, 1994–2002
- 9.6% of Shih Tzu referred for eye or neurological conditions in Japan were affected (Gelatt & MacKay, 2004a; Kato *et al.*, 2006)

Keratoconjunctivitis sicca

- Shih Tzu was the fourth most common breed in a UK referral study (Sanchez *et al.*, 2007)

Renal and urinary conditions**Kidney disease**

- Shih Tzu had an incidence of 30 (95% CI 23–38) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden (Pelander *et al.*, 2015)

Renal calculi

- May be associated with urinary tract infections
- Shih Tzu had odds ratios of 11.74 ($p < 0.001$) in females and 15.75 ($p < 0.001$) in males compared with crossbreeds in a US referral study (Ling *et al.*, 1998c)

Familial renal disease (familial nephropathy)

- No sex predisposition in Shih Tzu
- Shih Tzu comprised 15.6% of a US referral caseload (Picut & Lewis, 1987b; Hoppe *et al.*, 1990)

Urolithiasis

- Shih Tzu was the fifth most common breed for uroliths overall submitted for laboratory testing from Spain and Portugal (2.6%) and was the third most common in the UK (6.0%) (Rogers *et al.*, 2011; Vrabelova *et al.*, 2011)

Urolithiasis – apatite

- OR 8.6 (95% CI 7.8–9.5) compared with crossbred dogs in the USA
- Females are predisposed (Low *et al.*, 2010)

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- Shih Tzu was the second most common breed with submissions for calcium oxalate uroliths in Canada
- Odds ratios of 10.2 (95% CI 9.2–11.3) and 1.4 (95% CI 1.1–1.9) compared with crossbred dogs in two US studies, and 5.6 (95% CI 4.7–6.7) compared with all dogs in the UK
- OR 4.49 compared with crossbreeds in the USA (Lekcharoensuk *et al.*, 2000a; Ling *et al.*, 2003; Houston & Moore, 2009; Low *et al.*, 2010; Roe *et al.*, 2012; Okafor *et al.*, 2014)

Urolithiasis – silica

- OR 6.7 ($p < 0.001$) compared with all dogs in a US study (Aldrich *et al.*, 1997)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Shih Tzu was the most common breed with submissions for struvite uroliths in Canada
- Odds ratios of 7.4 (95% CI 6.8–8.1) compared with crossbred dogs in the USA and 9.8 (95% CI 8.6–11.2) compared with all dogs in the UK
- Small breeds and females are predisposed (Houston & Moore, 2009; Low *et al.*, 2010; Okafor *et al.*, 2014)

Urolithiasis – urate

- OR 6.4 (95% CI 4.9–8.4) compared with all dogs in the UK
- Younger and female animals predisposed (Low *et al.*, 2010; Roe *et al.*, 2012)

Respiratory conditions**Brachycephalic obstructive airway syndrome (BOAS)**

- Shih Tzu had a prevalence of 13.33% among referral dogs in Belgium (Njikam Nsangou *et al.*, 2009; Roedler *et al.*, 2013; Marchant *et al.*, 2017)

Tracheal collapse

- Shih Tzu comprised 10.8% of a US referral caseload (Johnson & Fales, 2001)

Reproductive conditions**Cryptorchidism**

- Shih Tzu had a 9.5% prevalence in the UK, which was significantly higher than for crossbreeds ($p = 0.012$) (Yates *et al.*, 2003)

Dystocia

- OR 2.1 (95% CI 1.1–3.8) compared with first-opinion emergency-care crossbred bitches in the UK (O'Neill *et al.*, 2017c)

Eclampsia (puerperal tetany)

- Small-sized bitches predisposed
- Shih Tzu was over-represented and comprised 6% of a US referral caseload (Drobatz & Casey, 2000)

SIBERIAN HUSKY

Dermatological conditions

Endocrine alopecia

- Siberian Husky comprised 3.0% of an endocrine alopecia caseload of neutered dogs in the USA (Frank *et al.*, 2003)

Zinc-responsive dermatosis

- Northern-breed dogs (e.g. Husky, Malamute, Samoyed) predisposed
- Siberian Husky represented 75.6% of affected cases from the USA, France and Israel (White *et al.*, 2001)

Gastrointestinal conditions

Congenital portosystemic shunt

- Referred Siberian Huskies (0.13% prevalence) had an OR of 2.5 (95% CI 1.1–5.9) compared with an overall US referral population (Tobias & Rohrbach, 2003)

Infectious conditions

Canine distemper

- Younger dogs and dolicocephalic breeds predisposed
- Siberian Husky was over-represented, comprising 4.0% of a general caseload in Brazil (Headley & Graça, 2000)

Neoplastic conditions

Cutaneous soft-tissue sarcoma

- OR 2.7 (95% CI 2.0–3.6) compared with a US referral population (Villamil *et al.*, 2011)

Palpebral neoplasia

- Mean age 9.6 years
- Siberian Husky at higher risk than general population (Roberts *et al.*, 1986)

Thyroid neoplasia

- Large breeds and older dogs predisposed
- Siberian Husky had OR 2.47 (95% CI 1.50–4.07) compared with an overall referral population in the USA (Wucherer & Wilke, 2010)

Neurological conditions

Degenerative myelopathy

- Reported in three related Siberian Huskies
- Suspected to be inherited (Bichsel *et al.*, 1983)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 4.68%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Chronic superficial keratitis (pannus)

- Siberian Husky significantly over-represented, comprising 2.38% of referral cases in the USA
- Siberian Husky was disproportionately affected among referral dogs in another US study (Slatter *et al.*, 1977; Chavkin *et al.*, 1994)

Congenital achromatopsia (rod monochromacy, day blindness)

- Autosomal recessive disease
- Five casual mutations identified: *PDE6C*, *PDE6H*, *GNAT2*, *CNGA3* and *CNGB3*
- Siberian Huskies in the USA had 0.877% frequency of the *CNGB3* deletion mutation (Yeh *et al.*, 2013)

Corneal dystrophy

- Prevalence of 14% in screened Siberian Huskies
- Prevalence increased with age (MacMillan *et al.*, 1979)

Glaucoma – primary

- Siberian Husky had 1.88% prevalence in a US referral population, 1994–2002
- OR 5.06 ($p < 0.01$) compared with an overall referral population in Switzerland (Gelatt & MacKay, 2004a; Strom *et al.*, 2011a)

Progressive retinal atrophy (PRA)

- X-linked recessive inheritance in Siberian Husky (Acland *et al.*, 1994)

Physiological conditions

Benign familial hyperphosphatasaemia

- Described in related Siberian Husky pups
- Markedly high serum alkaline phosphatase (ALP) levels seen in affected Siberian Husky
- Familial and autosomal inheritance (Lawler *et al.*, 1996)

Renal and urinary conditions**Ectopic ureter**

- Siberian Husky bitches had risk ratios of 21.7 (95% CI 15.2–30.9) and 49.81 (95% CI 28.74–127.86) compared with overall referral populations in two US studies
- Siberian Husky comprised 9.1% of a US referral caseload
(Hayes, 1974b, 1984; Ho *et al.*, 2011)

Reproductive conditions**Cryptorchidism**

- Siberian Husky reported as over-represented in a US study
- OR 2.4 (95% CI 1.32–4.27) compared with an overall US referral population
(Pendergrass & Hayes, 1975; Zhao *et al.*, 2010)

Testicular neoplasia

- Siberian Husky had a proportional morbidity ratio of 2.0 (95% CI 1.0–4.3) compared with all other dogs in a Norwegian cancer registry
(Nødtvedt *et al.*, 2011)

Respiratory conditions**Bronchiectasis**

- Siberian Husky had 2.86 OR, (95% CI 1.47–5.55) compared with an overall US referral population
(Hawkins *et al.*, 2003)

Spontaneous pneumothorax

- Siberian Husky comprised 18.8% of a US referral caseload and had an OR of 28.8 (95% CI 6.3–277.4) compared with a control population
(Puerto *et al.*, 2002)

SILKEN WINDHOUND**Drug reactions****Multiple drug sensitivity**

- High doses of many drugs (e.g. ivermectin or milbemycin) cause neurological signs
- Associated with the nt230(del4) *MDR1* mutation
- Silken Windhound had a 28.1% allelic frequency with 3.8% recessive mutant allele frequency in Europe
(Firdova *et al.*, 2016)

SILKY TERRIER

See *Australian Silky Terrier*

SKYE TERRIER**Endocrine conditions****Hypothyroidism**

- Females and younger dogs are predisposed to having serum thyroid hormone autoantibodies (THAA) that are associated with hypothyroidism
- Skye Terrier had an OR of 3.04 ($p=0.001$) for THAA compared with all other breeds
(Nachreiner *et al.*, 2002)

Gastrointestinal conditions**Chronic hepatitis**

- Familial
- Associated with copper accumulation in Skye Terrier
(Haywood *et al.*, 1988; McGrotty *et al.*, 2003)

Musculoskeletal conditions**Elbow dysplasia**

- Skye Terrier had 7.2% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland
- Elbow joint incongruity was scored as grade 1 in 49%, grade 2 in 31% and grade 3 in 18% of Skye terriers in Finland. Lameness was reported at <1 year old in 33% of dogs and was so common that it was called 'Skye limp'
(Narojek *et al.*, 2008; Michelsen, 2013; Lappalainen *et al.*, 2016)

Renal and urinary conditions**Ectopic ureter**

- Skye Terrier bitches had an OR of 86.6 (95% CI 31.6–258.6) compared with a general UK referral population
(Holt & Moore, 1995; Holt *et al.*, 2000)

Reproductive conditions**Pyometra (cystic endometrial hyperplasia–pyometra complex)**

- Nulliparous Skye Terrier bitches had an OR of 3.9 (95% CI 1.84–8.29) compared with all dogs attending primary-care practice in Finland
(Niskanen & Thrusfield, 1998)

SLOUGHI (ARABIAN GREYHOUND, SLOUGHI MOGHREBI, LEVRIER MAROCAIN)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance with candidate gene identified in Sloughi (Dekomien *et al.*, 2000)

Physiological conditions

Reduced thyroxine levels

- Sighthounds predisposed to physiologically lower thyroxine levels
- Privately owned Sloughis from Germany, the Czech Republic and France had lower thyroid concentrations than healthy controls ($p < 0.0001$)

(Panakova *et al.*, 2008)

SMOOTH COLLIE

See *Collies – Rough and Smooth*

SOFT-COATED WHEATEN TERRIER

Behavioural conditions

Aggression

- Soft-coated Wheaten Terrier had a prevalence of 16.2%, compared with 10.7% in the study dogs overall in the USA

(Duffy *et al.*, 2008)

Dermatological conditions

Atopic dermatitis (atopy)

- Soft-coated Wheaten Terrier had the 13th-highest incidence in insured dogs in Sweden: 3.3 cases per 1000 DYAR (95% CI 2.6–4.0)

(Nødtvedt *et al.*, 2006)

Endocrine conditions

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely
- Females predisposed
- OR 6.68 (95% CI 1.11–29.49) compared with an overall US referral population

(Peterson *et al.*, 1996)

Gastrointestinal conditions

Lymphangiectasia (resulting in protein-losing enteropathy)

- Associated with protein-losing nephropathy
- Characterized in Soft-coated Wheaten Terriers in the USA
- Enhanced wheat gluten sensitivity may be one factor involved in the pathogenesis
- Familial
- Mean age at diagnosis 4.7 years

(Littman *et al.*, 2000;
Vaden *et al.*, 2000)

Neoplastic conditions

Mammary neoplasia

- Soft-coated Wheaten Terrier had an incidence of 199 cases per 10 000 DYAR, compared with 116 per 10 000 DYAR in crossbreeds among insured entire bitches in Sweden

(Egenvall *et al.*, 2005)

Physiological conditions

Perinuclear antineutrophilic cytoplasmic autoantibodies (pANCA)

- 20.7% of healthy Soft-coated Wheaten Terriers in the UK had positive results for pANCA

(Wieland *et al.*, 2012)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Associated with protein-losing enteropathy
- Described in Soft-coated Wheaten Terriers in Norway
- Complicated inheritance reported for protein-losing nephropathy, with multiple genes, variable expression and possibly environmental triggers
- Mutations causing a podocytopathy identified, associated with protein-losing nephropathy in Soft-coated Wheaten Terriers in the USA and Canada

(Eriksen & Grøndalen, 1984;
Littman *et al.*, 2013;
Vaden *et al.*, 2013)

Kidney disease

- Soft-coated Wheaten Terrier had an incidence of 30 (95% CI 23–37) cases per 10 000 DYAR compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden

(Pelander *et al.*, 2015)

SPANISH GREYHOUND (GALGOS ESPAÑOL)

Physiological conditions

Haematological differences

- Spanish Greyhounds have higher haematocrits, haemoglobin concentrations, red cell counts and blood pH than other dogs
- Spanish Greyhounds have lower platelet counts than other dogs

(Mesa-Sánchez *et al.*, 2012)

SPANISH MASTIFF

Behavioural conditions

Aggression

- Spanish Mastiff had higher levels of territorial aggression in Spain

(Perez-Guisado & Munoz-Serrano, 2009)

SPANISH WATER DOG

Musculoskeletal conditions

Hip dysplasia

- Spanish Water Dog had 22nd-highest prevalence in a US study: 27.0%

(Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Short tail (bobtail)

- 57.1% of Spanish Water Dog tested in Finland affected

(Hytönen *et al.*, 2009)

Renal and urinary conditions

Portal vein hypoplasia

- Can lead to urine supersaturation with uric acid (hyperuricosuria) and urate urolithiasis
- Associated with mutated *SLC2A9* gene
- Identified in a Spanish Water Dog in the UK

(Cosgrove *et al.*, 2015)

SPINONE ITALIANO

See *Italian Spinone*

SPRINGER SPANIEL (ENGLISH, WELSH, UNSPECIFIED VARIANTS)

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- 55.6% of Springer Spaniels aged < 1 year and 26.8% of those aged ≥ 1 year had bitten, in a Canadian veterinary survey
- Intense aggression was reported in 48.4% of English Springer Spaniels, and 26.3% had bitten a human in the USA
- English Springer Spaniel had a prevalence of 17.5% for dog-to-dog aggression, compared with 10.7% in study dogs overall in the USA

(Guy *et al.*, 2001; Reisner *et al.*, 2005; Duffy *et al.*, 2008; Zapata *et al.*, 2016)

Dental conditions

Canine odontogenic parakeratinized cyst

- English Springer Spaniel comprised 22.2% of a US referral caseload and was significantly ($p < 0.001$) over-represented compared with the overall hospital population

(Verstraete *et al.*, 2011)

Dermatological conditions

Atopic dermatitis (atopy)

- Welsh Springer Spaniel had the 14th-highest incidence in insured dogs in Sweden: 2.9 cases per 1000 DYAR (95% CI 2.0–3.8)
- English Springer Spaniel comprised 3.0% (95% CI 1.3–5.8) of an atopic caseload compared with 1.2% of the overall hospital population in a US referral study

(Zur *et al.*, 2002; Nødtvedt *et al.*, 2006)

Grass awn migration

See under *Soft-tissue conditions*

Pemphigus foliaceus

- English Springer Spaniel had OR 20.7 (95% CI 13.1–98.8) compared with a referral population in the USA

(Kuhl *et al.*, 1994)

Primary seborrhoea

- Described in English Springer Spaniel in the USA
- Half develop clinical signs by 2 years of age
- Familial occurrence reported in English Springer Spaniels

(Scott & Miller, 1996)

Sebaceous adenitis

- Autosomal recessive inheritance with variable expression
- Median age at diagnosis 5 years
- Springer Spaniel had the fifth-highest breed relative risk in Sweden and comprised 24.0% of a general veterinary caseload

(Tevell *et al.*, 2008)

Endocrine conditions**Diabetes mellitus**

- Familial and inherited
- Older entire females are predisposed
- English Springer Spaniel had RR 4.7 (95% CI 1.7–13.0) compared with an overall US referral population

(Marmor *et al.*, 1982)

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely: estimated heritability 0.49 (± 0.16)
- Association with alleles in genes *COL4A4*, *OSBPL9*, *CTLA4*, *PTPN22* and *STXBP5* identified in Springer Spaniel
- Welsh Springer Spaniel had incidence rate of 6.31 (95% CI 2.96–13.7) cases per 10 000 DYAR, compared with 2.26 (95% CI 2.07–2.46) overall in insured dogs in Sweden

(Short *et al.*, 2013; Hanson *et al.*, 2016)

Gastrointestinal conditions**Chronic hepatitis**

- Multiple possible causes, so breed predisposition may change with time and geographic location due to genetic and environmental factors
- Median age at diagnosis was 8 years and median survival time was 189 days in the UK
- Protective and risk-conferring alleles and haplotype identified in English Springer Spaniel in the UK
- English Springer Spaniel had OR 4.6 (95% CI 1.9–11.2) compared with overall unselected dogs from first-opinion practices in the UK and OR 5.3 (95% CI 4.2–6.7) compared with an overall laboratory population in the UK
- English Springer Spaniel had RR 3.7 (95% CI 1.1–11.9) among referral post-mortem cases in the UK

(Watson *et al.*, 2007, 2010; Bexfield *et al.*, 2011, 2012a, 2012b)

Gastrointestinal foreign bodies

- Mean age of UK affected dogs was 2.5 years
- Springer Spaniel had RR 8.3 (95% CI 4.1–16.8) compared with an overall general population in the UK

(Hayes, 2009)

Haematological/immunological conditions**Immune-mediated haemolytic anaemia (IMHA)**

- Females predisposed
- Springer Spaniel had OR 10.0 (95% CI 1.3–74.7) compared with an overall referral population in Australia
- English Springer Spaniel had OR 32.4 (95% CI 2.8–379.0) compared an overall referral population in the USA

(Reimer *et al.*, 1999; McAlees, 2010)

Phosphofructokinase deficiency

- Molecular basis of a nonsense mutation in the penultimate exon of the *M-PFK* gene identified in English Springer Spaniel
- Inherited as an autosomal recessive trait

(Giger *et al.*, 1986; Smith *et al.*, 1996a; Skibild *et al.*, 2001)

Infectious conditions**Parvovirus enteritis**

- English Springer Spaniel had OR 8.1 (95% CI 2.3–29.6) compared with an overall referral US population

(Glickman *et al.*, 1985)

Musculoskeletal conditions**Incomplete ossification of the humeral condyle (IOHC)**

- Associated with humeral condylar fractures following minimal trauma during normal activity
- Males predisposed
- Springer Spaniels comprised 85.0% of an affected caseload in the UK

(Carrera *et al.*, 2008)

Hip dysplasia

- English Springer Spaniel had 13.1% prevalence in the USA
- Two US reports show English Springer Spaniel had odds ratios of 1.4 (95% CI 1.1–1.8) and 1.26 (1.16–1.37) compared with crossbreeds

(LaFond *et al.*, 2002; Witsberger *et al.*, 2008; Ginja *et al.*, 2010; Orthopedic Foundation for Animals, 2016)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 2.8 (95% CI 2.2–3.5) compared to cross-breeds in the USA
(LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Neoplastic conditions

Anal sac adenocarcinoma

- Springer Spaniel had OR 3.7 (95% CI 2.5–5.3) in the UK
(Polton *et al.*, 2006)

Cutaneous haemangioma

- English Springer Spaniel had OR 1.6 (95% CI 1.2–2.2) compared with crossbreeds in the USA
(Goldschmidt & Mcmanus, 2000)

Mammary neoplasia

- English Springer Spaniel had an incidence of 319 cases per 10 000 DYAR, compared with 116 per 10 000 DYAR in crossbreeds among insured entire bitches in Sweden
- English Springer Spaniel had the third-highest relative risk ratio in Norway
(Moe, 2001; Egenvall *et al.*, 2005)

Neurological conditions

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Median age at onset of seizures in English Springer Spaniels was 3 years
- Partially penetrant autosomal recessive inheritance or polygenic inheritance in English Springer Spaniels
- English Springer Spaniel had an incidence of 17.3 per 10 000 DYAR (95% CI 14.0–20.7) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
(Patterson *et al.*, 2005; Heske *et al.*, 2014)

Lysosomal storage disease – fucosidosis

- Leads to undesirable behaviour in adolescent and adult dogs that may be misdiagnosed as a primary behavioural problem or acquired neurologic disease
- Reported in English Springer Spaniel in the USA
- Autosomal recessive inheritance
- Molecular defect identified in English Springer Spaniel
(Skelly *et al.*, 1996; Smith *et al.*, 1996b)

Steroid-responsive meningitis–arteritis (SRMA)

- Young, medium- to large-breed dogs predisposed
- Suspected to be immune-mediated
- English Springer Spaniels were significantly over-represented ($p < 0.001$) in a UK referral population, comprising 20% of the caseload
(Lowrie *et al.*, 2009)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.22% in English Springer Spaniel and 1.99% in Welsh Springer Spaniel, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Hereditary cataract described in Welsh Springer Spaniels in the UK
(Barnett, 1980; Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Corneal and anterior segment foreign body trauma

- Working dogs and dogs < 5 years predisposed
- English Springer Spaniel had OR 6.2 (95% CI 4.5–8.7) compared to other referral dogs in the UK
(Tetas Pont *et al.*, 2015)

Corneal ulceration (ulcerative keratitis)

- English Springer Spaniel had OR 1.93 (95% CI 1.17–3.19) compared with crossbreeds in the UK
(O'Neill *et al.*, 2017b)

Glaucoma – primary

- Female Welsh Springer Spaniels predisposed in the UK
- Welsh Springer Spaniel had OR 5.00 ($p = 0.04$) compared with an overall referral population in Switzerland
(Cottrell & Barnett, 1988; Strom *et al.*, 2011a)

Pectinate ligament dysplasia (PLD)

- Associated with glaucoma in many breeds
- English Springer Spaniel had 25.5% prevalence in a Norwegian screening programme
- Welsh Springer Spaniel had 61.2% prevalence among a UK general population
(Bjerkås *et al.*, 2002; Oliver *et al.*, 2016b)

Retinal dysplasia

- Clinically described in English Springer Spaniels in the USA (O'Toole *et al.*, 1983)

Sudden acquired retinal degeneration (SARD, amaurosis)

- Median age 8.1 years
- English Springer Spaniel comprised 5% of a US caseload (Montgomery *et al.*, 2008)

Physiological conditions**Litter size**

- Larger litters associated with younger bitches and larger breeds
- English Springer Spaniel had the twelfth largest mean litter size (7.3 puppies) among registered breeds in Norway (Borge *et al.*, 2011)

Renal and urinary conditions**Kidney disease**

- English Springer Spaniel had an incidence of 23 (95% CI 19–27) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden (Pelander *et al.*, 2015)

Urethral sphincter mechanism incompetence

- Larger breeds had OR 7.2 compared with smaller breed in the USA
- Springer Spaniel was over-represented in a UK referral caseload (Holt, 1985; Forsee *et al.*, 2013)

Urinary tract infections (UTI)

- Springer Spaniel (3.4%) was the seventh most commonly affected breed among referral dogs in the USA for recurrent or persistent UTI (Norris *et al.*, 2000)

Urolithiasis

- Springer Spaniel was the twelfth most common breed for uroliths overall submitted for laboratory testing in the UK (1.8%) (Rogers *et al.*, 2011)

Reproductive conditions**Pyometra (cystic endometrial hyperplasia-pyometra complex)**

- English Springer Spaniel showed RR 1.9 (95% CI 1.3–2.5) compared with all bitches insured by Agria for veterinary care in Sweden (Egenvall *et al.*, 2001; Smith, 2006)

Respiratory conditions**Aspiration pneumonia**

- English Springer Spaniel comprised 4.4% of cases but only 0.6% of the underlying referral population in the USA (Kogan *et al.*, 2008)

Bronchiectasis

- English Springer Spaniel had an OR of 2.39 (95% CI 1.23–4.64) compared with an overall US referral population (Hawkins *et al.*, 2003)

Pyothorax

- May be associated with initial inhalation of foreign bodies during the scenting habits and outdoor nature of some breeds
- Springer Spaniel was the most common breed, comprising 43.8% of a UK referral caseload (Johnson & Martin, 2007)

Soft-tissue conditions**Grass awn migration**

- Grass awns comprised 61% of all foreign-body-related cases
- Ear canal (51%) was the most commonly affected site
- Springer Spaniel was significantly over-represented in the USA (Brennan & Ihrke, 1983)

Oropharyngeal penetrating injury

- Medium- to large-breed dogs and males predisposed
- May be associated with stick-chasing activity in larger breeds and head posture during retrieving
- Springer Spaniels had 3.09 (95% CI 1.07–8.15) OR compared with a UK referral population (Griffiths *et al.*, 2000)

STAFFORDSHIRE BULL TERRIER (SBT)**Behavioural conditions****Aggression**

- A genetic basis for aggression has been described
- A survey of UK veterinarians classified SBT as having high aggression
- SBT was over-represented as the aggressor in dog-dog conflict in Germany, representing

5.8% of aggressors and 1.2% of the general population

(Roll & Unshelm, 1997;
Bradshaw & Goodwin, 1999)

Dermatological conditions

Atopic dermatitis (atopy)

- SBT had the fifth-highest incidence in insured dogs in Sweden: 8.0 cases per 1000 DYAR (95% CI 5.3–10.7)
(Nødtved *et al.*, 2006)

Demodicosis

- Short-haired breeds predisposed
- An inheritance pathway has been described
- SBT was described as a predilected breed in a Swedish referral study
- SBT had the second-highest breed OR of 17.12 (95% CI 2.2–133.4) compared with an overall US first-opinion population
(Holm, 2003; It *et al.*, 2010; Plant *et al.*, 2011)

Gastrointestinal conditions

Gastrointestinal foreign bodies

- Mean age of UK affected dogs was 2.5 years
- RR 3.3 (95% CI 2.2–5.1) compared with an overall general population in the UK
(Hayes, 2009)

Infectious conditions

Angiostrongylosis

- The median age of affected dogs was 10 months, and young dogs were predisposed ($p < 0.0001$)
- SBT were significantly over-represented ($p < 0.0001$), comprising 21.7% of a UK referral caseload
(Chapman *et al.*, 2004)

Metabolic conditions

Hypocobalaminaemia

- SBT had the second-highest breed OR of 2.6 (95% CI 1.7–3.9) compared with a laboratory population in the UK
(Dandrieux *et al.*, 2013)

Musculoskeletal conditions

Elbow dysplasia

- SBT had 33.3% prevalence in the UK, 16.2% in the USA, and 31.3% in South Africa
(Kirberger & Stander, 2007; Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Neoplastic conditions

Gastric carcinoma

- SBT significantly over-represented
(Sullivan *et al.*, 1987b)

Mammary neoplasia

- 25% of insured SBT bitches in Sweden develop the condition by 10 years of age, compared with 13% for bitches overall
(Jitpean *et al.*, 2012)

Mast cell tumour (MCT)

- Compared to crossbreeds, SBT had odds ratios of 4.2 (95% CI 2.2–8.2) in the UK and 4.1 (95% CI 2.4–7.1) in the USA
- OR 3.49 (95% CI 2.09–5.81) compared with an overall referral population in the UK
- SBT had the second-highest breed RR (9.97) compared with all breeds in laboratory records in the USA, and was significantly over-represented in a US referral study ($p = 0.044$)
(Goldschmidt & Mcmanus, 2000; Baker-Gabb *et al.*, 2003; Warland & Dobson, 2013; Shoop *et al.*, 2015; Mochizuki *et al.*, 2016)

Neoplasia – overall

- SBT had the eighth-highest proportional mortality from cancer among pedigree breeds in the UK: 44.4% (95% CI 35.4–53.4)
(Adams *et al.*, 2010)

Neurological conditions

L-2-Hydroxyglutaric aciduria (organic aciduria)

- Progressive neurological signs between 4 months and 7 years of age
- Described in SBT in the UK and a mutation in canine *L2HGDH* gene identified
(Abramson *et al.*, 2003; Penderis *et al.*, 2007; Short *et al.*, 2010)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- SBT showed 10.9% prevalence of cataract among ophthalmic referrals in South Africa
- Autosomal recessive mutation identified in SBT, and DNA test available
(Petrick, 1996; Mellersh *et al.*, 2006; Donzel *et al.*, 2016; Animal Health Trust, 2017)

Corneal ulceration (ulcerative keratitis)

- OR 2.50 (95% CI 1.81–3.45) compared with crossbreeds in the UK

(O'Neill *et al.*, 2017b)

Renal and urinary conditions**Urolithiasis – cystine**

- OR 8.7 (95% CI 6.8–11.2) compared with all dogs in the UK

(Roe *et al.*, 2012)

Reproductive conditions**Cryptorchidism**

- SBT had a 9.3% prevalence in the UK, which was significantly higher than for crossbreeds ($p=0.0006$)

(Yates *et al.*, 2003)

Dystocia

- SBT had the sixth-highest incidence (23.6 cases per 1000 DYAR) among insured bitches in Sweden, compared with 5.7 per 1000 DYAR overall
- OR 4.1 (95% CI 2.7–6.2) compared with first-opinion emergency-care crossbred bitches in the UK

(Bergström *et al.*, 2006;
O'Neill *et al.*, 2017c)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 54% of insured SBT bitches develop the condition by 10 years of age, compared with 19% for bitches overall

(Jitpean *et al.*, 2012)

Respiratory conditions**Brachycephalic obstructive airway syndrome (BOAS)**

- SBT comprised 5.5% of a US BOAS caseload

(Torrez & Hunt, 2006;
Marchant *et al.*, 2017)

Soft-tissue conditions**Non-accidental injury (NAI, battered pet)**

- Males ($p<0.001$) and dogs aged <2 years ($p<0.001$) significantly predisposed
- SBT had OR 7.2 (95% CI 3.5–13.4) and SBT-cross dogs had OR 48 (95% CI 11–18.1) compared with the general population of dogs under primary veterinary care in the UK

(Munro & Thrusfield, 2001)

STAFFORDSHIRE BULL TERRIER – AMERICAN

See *American Staffordshire Terrier*

SUSSEX SPANIEL**Metabolic conditions****Overweight/obesity**

- Sussex Spaniel had significantly higher body condition score (5.57) than other show dogs in Holland

(Courcier *et al.*, 2010; Corbee, 2013;
Raffan *et al.*, 2016)

Pyruvate dehydrogenase phosphatase 1 deficiency

- Causes severe exercise intolerance
- Null mutation in *PDP1* identified
- Classic Mendelian autosomal recessive inheritance
- 63.6% of Sussex Spaniels tested in the USA were carriers

(Abramson *et al.*, 2004; Cameron *et al.*, 2007)

Musculoskeletal conditions**Elbow dysplasia**

- Sussex Spaniel had 20.5% prevalence in the USA

(Michelsen, 2013; Orthopedic
Foundation for Animals, 2016)

Hip dysplasia

- Sussex Spaniel had tenth-highest prevalence in a US study: 41.2%

(Ginja *et al.*, 2010; Orthopedic
Foundation for Animals, 2016)

SWEDISH CATTLE DOG

See *Swedish Vallhund*

SWEDISH ELKHOUND (JÄMTHUND)**Endocrine conditions****Diabetes mellitus**

- Familial and inherited
- Peak incidence age 7–9 years

- Older entire females during dioestrus and pregnancy are predisposed
- Swedish Elkhound had fourth-highest breed incidence in Sweden: 45 cases per 10 000 DYAR (95% CI 32–58)

(Fall *et al.*, 2007, 2010; Catchpole *et al.*, 2008)

Neoplastic conditions

Mammary neoplasia

- 26% of insured Swedish Elkhound bitches in Sweden develop the condition by 10 years of age, compared with 13% for bitches overall (Jitpean *et al.*, 2012)

Neurological conditions

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Swedish Elkhound had an incidence of 16.5 per 10 000 DYAR (95% CI 12.8–20.1) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds (Heske *et al.*, 2014)

Ocular conditions

Progressive retinal atrophy (PRA)

- Described in closely related Swedish Elkhound in Sweden
- Compatible with a recessive inheritance (Hertil *et al.*, 2010)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- 40% of insured Swedish Elkhound bitches develop the condition by 10 years of age, compared with 19% for bitches overall (Jitpean *et al.*, 2012)

SWEDISH FARMDOG

See Danish/Swedish Farmdog

SWEDISH LAPPHUND

Endocrine conditions

Diabetes mellitus

- Familial and inherited
- Breed at increased risk in a retrospective review of Swedish insurance data, 1995–2004
- Peak incidence age 7–9 years

- Older entire females are predisposed
- Swedish Lapphund had third-highest breed incidence in Sweden: 72 cases per 10 000 DYAR (95% CI 33–111), compared to an overall incidence for all breeds of 13 cases per 10 000 DYAR

(Fall *et al.*, 2007; Catchpole *et al.*, 2008)

SWEDISH VALLHUND (SWEDISH CATTLE DOG)

Musculoskeletal conditions

Short tail (bobtail)

- 72.7% of Swedish Vallhund tested in Finland affected

(Hytönen *et al.*, 2009)

Ocular conditions

Progressive retinal atrophy (PRA)

- Autosomal recessive mode of inheritance with environmental disease modifiers suggested in Swedish Vallhund
- Genetics and clinical presentation described in Swedish Vallhund in several countries (Cooper *et al.*, 2014a)

SWISS MOUNTAIN DOG (SENNEHUND)

Haematological/immunological conditions

Platelet disorder

- Familial and hereditary in Greater Swiss Mountain Dogs
- Spontaneous haemorrhage is absent to mild in affected dogs, but excessive bleeding can occur following routine surgical procedures or trauma
- Platelet numbers, coagulation and von Willebrand assays are normal
- Mutation for the gene encoding the ADP receptor P2Y₁₂ identified in Greater Swiss Mountain Dogs in Canada (Boudreaux & Martin, 2011)

Metabolic conditions

Overweight/obesity

- Swiss Mountain Dogs had significantly higher body condition score than other show dogs in Holland (Courcier *et al.*, 2010; Corbee, 2013; Raffan *et al.*, 2016)

Musculoskeletal conditions

Lumbosacral transitional vertebrae

- Breed at increased risk in a Swiss retrospective study of x-rays of 4000 medium- and large-breed dogs, 1996–1998. Prevalence in the 64 Swiss Mountain dogs was 9.4%, compared to an overall prevalence of 3.5%
- There was no effect of gender
(Damur-Djuric *et al.*, 2006)

Osteochondrosis – shoulder

- Breed at increased risk in retrospective study of 626 cases from VMDB entries, 1982–1986. Incidence 7.69%
- Breed at increased risk of humeral head osteochondrosis. A Swiss study demonstrated a prevalence of 14%, 1993–2013. This was suggested to be in the mid to upper range compared to reported prevalences in other breeds
(Rudd *et al.*, 1990; Ohlerth *et al.*, 2016)

Neoplastic conditions

Fibroma/fibrosarcoma

- Swiss Mountain Dogs had an OR significantly above 1.0 ($p < 0.05$) compared with cross-breeds from a cancer registry in Switzerland
(Grüntzig *et al.*, 2016)

Lymphoma

- Swiss Mountain Dogs had an OR significantly above 1.0 ($p < 0.05$) compared with cross-breeds from a cancer registry in Switzerland
(Grüntzig *et al.*, 2016)

Mast cell tumour (MCT)

- Swiss Mountain Dogs had an OR significantly above 1.0 ($p < 0.05$) compared with cross-breeds from a cancer registry in Switzerland
(Grüntzig *et al.*, 2016)

TENTERFIELD TERRIER

Endocrine conditions

Congenital hypothyroidism

- Associated with other developmental abnormalities including dwarfism, delayed epiphyseal ossification, abnormal hair texture, unresponsiveness and lethargy
- Causal missense mutation in exon 9 (R593W) identified in Tenterfield Terrier
- Autosomal recessive inheritance
(Dodgson *et al.*, 2012)

Neurological conditions

Spinocerebellar ataxia (hereditary ataxia)

- Incoordination noticed from 2–9 months of age
- Progressive and often leads to euthanasia
- Mutation in the *KCNJ10* gene identified in the Tenterfield Terrier
(Rohdin *et al.*, 2015)

Ocular conditions

Lens luxation – primary

- Tenterfield Terriers had 12% prevalence in Australia
- Frequency of the A-422 haplotype was 0.41 in the Tenterfield Terrier in Australia
(Gharakhani *et al.*, 2012)

THAI RIDGEBACK (MAH THAI)

Neurological conditions

Dermoid sinus

- Complex dihybrid mode of inheritance suggested
- Causative mutation is a duplication of *FGF3*, *FGF4*, *FGF19* and *ORAOV1* genes that cause hair ridge and predisposition to dermoid sinus in Thai Ridgeback dogs
(Salmon Hillbertz *et al.*, 2007)

Physiological conditions

Dorsal ridge

- Breed-defining ridge of Thai Ridgebacks comes from a duplication of a specific region on chromosome 18 called the Ridge allele
- Dominant inheritance
- Homozygosity for the Ridge allele predisposes to dermoid sinus
(Waldo & Diaz, 2015)

TIBETAN MASTIFF

Musculoskeletal conditions

Elbow dysplasia

- Tibetan Mastiff had 22.8% prevalence in the UK and 14.3% in the USA
(Michelsen, 2013; Kennel Club, 2016; Orthopedic Foundation for Animals, 2016)

Neurological conditions

Hypertrophic neuropathy

- Autosomal recessive inheritance

- Described in related Tibetan Mastiffs
- Age of clinical onset 7–18 weeks
(Cummings *et al.*, 1981; Sponenberg & de Lahunta, 1981)

Physiological conditions

Adaptation to hypoxia

- Tibetan Mastiff has adapted to the extreme high-altitude environment of the Tibetan Plateau
- 12 candidate genes identified for adaptations to high altitude and roles in responses to hypoxia
- Tibetan Mastiff has lower blood haemoglobin levels and greater smooth muscle content of pulmonary blood vessel and pulmonary pleura than other breeds

(Jian-guo *et al.*, 2009; Li *et al.*, 2014; Wang *et al.*, 2014)

TIBETAN SPANIEL

Musculoskeletal conditions

Patellar luxation

- 12.6% of US referred Tibetan Spaniels affected
- Mainly medial luxation observed, often bilateral (Orthopedic Foundation for Animals, 2015)

Neoplastic conditions

Mast cell tumour (MCT)

- Mean age at presentation between 7.5 and 9 years, but can occur at any age
- OR 7.73 (95% CI 1.03–10.69) compared with crossbreeds in Austria
(Goldschmidt & Mcmanus, 2000; Leidinger *et al.*, 2014)

Neurological conditions

Intervertebral disc disease (IVDD)

- Males and older dogs predisposed
- Tibetan Spaniel had an incidence rate of 48.7 (95% CI 37.5–59.9), compared with 27.8 for an overall population of insured dogs in Sweden
(Bergknut *et al.*, 2012)

Ocular conditions

Progressive retinal atrophy (PRA)

- PRA mutation identified in Tibetan Spaniel in the UK
- Autosomal recessive mode inheritance and rapid progression over 1 year from initial

clinical signs to complete loss of vision in Tibetan Spaniel in Norway and Sweden
(Bjerkås & Narfström, 1994; Downs *et al.*, 2014a)

Renal and urinary conditions

Primary hyperoxaluria

- Reported in related Tibetan Spaniels in Norway
- Cases presented at 6 weeks and had an unusual coat colour
- May be part of an overall juvenile nephropathy syndrome in the Tibetan Spaniel
(Jansen & Arnesen, 1990)

TIBETAN TERRIER

Dermatological conditions

Atopic dermatitis (atopy)

- Tibetan Terrier comprised 1.9% (95% CI 0.6–4.3) of an atopic caseload, compared with 0.25% of the overall hospital population, in a US referral study
- Tibetan Terrier had 5.86% prevalence, compared with 1.08% for crossbreeds in a US referral caseload
(Zur *et al.*, 2002; Bellumori *et al.*, 2013)

Endocrine conditions

Diabetes mellitus

- Familial and inherited
- Older entire females are predisposed
- Tibetan Terrier comprised 5.3% of cases, compared with 0.3% of the overall population of insured dogs in the UK
(Davison *et al.*, 2005)

Haematological/immunological conditions

Haemophagocytic syndrome

- Benign proliferative disorder of activated macrophages associated with multiple blood cytopenias
- Tibetan Terriers were over-represented, comprising 12.5% of a US referral caseload
(Weiss, 2007)

Musculoskeletal conditions

Patellar luxation

- 5.2% of US referred Tibetan Terriers affected
- Mainly medial luxation observed, often bilateral (Orthopedic Foundation for Animals, 2015)

Neurological conditions

Neuronal ceroid lipofuscinosis

- Only nyctalopia (night blindness) noted until 5/6 years of age and then personality and learning changes develop afterwards
- Autosomal recessive inheritance of a truncating 376 mutation in canine *ATP13A2* identified in the Tibetan Terrier in the USA
- Application of DNA testing was associated with a drop in mutant allele frequencies from 0.20–0.28 before the introduction to 0.09–0.14 two years after the introduction

(Riis *et al.*, 1992; Farias *et al.*, 2011; Wöhlke *et al.*, 2011; Kluth *et al.*, 2014)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Tibetan Terrier had 4.71% prevalence and 13% heritability for non-congenital cataracts in Germany
- Prevalence of primary cataract 5.92%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
- Tibetan Terrier had 18.92% prevalence, compared with 4.04% for crossbreeds, in a US referral study (Ketteritzsch *et al.*, 2004; Gelatt & MacKay, 2005; Bellumori *et al.*, 2013; Donzel *et al.*, 2016)

Distichiasis

- Tibetan Terrier had 4.3% heritability and 11.4% prevalence in Germany (Ketteritzsch *et al.*, 2004)

Lens luxation – primary

- *ADAMTS17* mutation has been identified, with a simple autosomal recessive hypothesis
- Tibetan Terrier had 99.3% heritability and 1.29% prevalence in Germany
- Tibetan Terrier had relative risks (RR) of 4.94 (95% CI 1.87–13.04) and 3.69 (95% CI 1.75–7.80) compared with overall referral dogs in two US studies (Willis *et al.*, 1979; Ketteritzsch *et al.*, 2004; Sargan *et al.*, 2007; Gould *et al.*, 2011)

Persistent pupillary membranes

- Tibetan Terrier had 17.1% heritability and 12.8% prevalence in Germany (Ketteritzsch *et al.*, 2004)

Progressive retinal atrophy (PRA)

- Simple autosomal recessive mode of inheritance suspected in Tibetan Terrier
- Tibetan Terrier had 49.1% heritability and 1.4% prevalence in Germany
- PRA mutation identified in Tibetan Terrier in the UK

(Millichamp *et al.*, 1988; Ketteritzsch *et al.*, 2004; Downs *et al.*, 2014a)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- OR 2.3 (95% CI 1.6–3.4) compared with all dogs in the UK

(Roe *et al.*, 2012)

Urolithiasis – struvite (magnesium ammonium phosphate)

- OR 2.0 (95% CI 1.4–2.8) compared with all dogs in the UK

(Roe *et al.*, 2012)

TOSA

See *Japanese Tosa*

TREEING WALKER COONHOUND

See *Coonhound*

TURKISH SHEPHERD DOG (AKBASH, KANGAL AND KARS)

Ocular conditions

Retinal dysplasia

- 17.1% of Akbash and 5.4% of Kangal dogs from breeding or referral dogs in Turkey affected

(Saroğlu *et al.*, 2005)

Physiological conditions

Blood group

- DEA 1.1 is highly immunogenic and causes acute haemolytic transfusion reactions in sensitized dogs
- The frequency of positivity for DEA 1.1 was 71.2% in Kars, 67.9% in Kangal and 60.0% in Akbash

- 14.4–35.6% probability of sensitizing a DEA 1.1-negative recipient at the first transfusion with blood from a DEA 1.1-positive donor of any of the three Turkish Shepherd breeds (Arikan *et al.*, 2009; Ergul Ekiz *et al.*, 2011)

VALLHUND

See *Swedish Vallhund*

VIZSLA

See *Hungarian Vizsla*

VORSTEHHUND

See *Weimaraner*

WACHTELHUND (GERMAN SPANIEL, DEUTSCHER WACHTELHUND)

Dermatological conditions

Atopic dermatitis (atopy)

- Wachtelhund had the tenth-highest incidence in insured dogs in Sweden: 4.2 cases per 1000 DYAR (95% CI 3.3–5.1)

(Nødtvedt *et al.*, 2006)

Endocrine conditions

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely: estimated heritability 0.49 (± 0.16)
- Wachtelhund had an incidence of 4.93 (95% CI 2.54–9.72) cases per 10 000 DYAR, compared with 2.26 (95% CI 2.07–2.46) overall among insured dogs in Sweden

(Hanson *et al.*, 2016)

Haematological/immunological conditions

Phosphofructokinase deficiency

- Inherited as an autosomal recessive trait
- Deficiency causes haemolytic crises and exertional myopathy
- Missense point mutation (c.550C>T) for phosphofructokinase identified in Wachtelhunds in Sweden

(Inal Gultekin *et al.*, 2012b)

Neoplastic conditions

Mammary neoplasia

- Wachtelhund had an incidence of 149 cases per 10 000 DYAR, compared with 116 per 10 000 DYAR in crossbreeds among insured entire bitches in Sweden

(Egenvall *et al.*, 2005)

WEIMARANER (VORSTEHHUND)

Cardiovascular conditions

Congenital heart disease

- Includes a range of congenital heart disorders
- Weimaraner had an OR of 9.4 compared with all referral dogs in Italy

(Oliveira *et al.*, 2011)

Pulmonary artery dissection

- Weimaraner comprised 37.5% of a UK/US referral caseload and was suggested as predisposed

(Scansen *et al.*, 2015)

Dermatological conditions

Blastomycosis

See under *Infectious conditions*

Canine follicular dysplasia (seasonal flank alopecia)

- Familial tendency with early age at onset suggests an inherited basis
- Group of similar hair disorders
- Described in Weimaraners in France

(Laffort-Dassot *et al.*, 2002)

Drug reactions

Anaesthetic-related complications

- Weimaraner had the equal-highest incidence (25%) among general small animal practices in Canada

(Dyson *et al.*, 1998)

Metaphyseal osteopathy

See under *Musculoskeletal conditions*

Gastrointestinal conditions

Gastric dilatation/volvulus (bloat, GDV)

- 11.6% of UK pedigree Weimaraners died of the condition
- Weimaraner had sixth-highest pedigree prevalence (5.0%) in the UK and was highly represented in a referral study in Israel

- Weimaraner had an incidence of 21 cases per 1000 DYAR (95% CI 0–42) in the USA (Glickman *et al.*, 2000a; Evans & Adams, 2010b; Israeli *et al.*, 2012)

Inflammatory bowel disease

- Mode of inheritance is incompletely understood and may differ between breeds
- Weimaraner had an OR of 3.68 (95% CI 2.02–6.71) compared with referred crossbreeds in the UK

(Kathrani *et al.*, 2011)

Haematological/immunological conditions

Immunodeficiency syndrome

- Considered inherited in Weimaraner, but mode of inheritance unclear
- Median age at presentation of 4 months
- Clinical signs and low blood antibody levels often develop within 7 days of vaccination
- Described in Weimaraner in the UK (Couto *et al.*, 1989; Foale *et al.*, 2003; Cuthbert *et al.*, 2016)

Infectious conditions

Blastomycosis

- Weimaraner was at increased risk compared with referral crossbred dogs in the USA (Rudmann *et al.*, 1992; Arceneaux *et al.*, 1998)

Musculoskeletal conditions

Metaphyseal osteopathy

- Primarily young (2–6 months), rapidly growing large- and giant-breed dogs
- OR 21.4 (95% CI 8.2–55.8) compared to crossbreeds in the USA
- Some reports of a link with recent vaccination (LaFond *et al.*, 2002; Crumlish *et al.*, 2006; Safra *et al.*, 2013)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.9 (95% CI 1.3–2.7) compared to crossbreeds in the USA (LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Neoplastic conditions

Mast cell tumour (MCT)

- Mean age at presentation between 7.5 and 9 years, but can occur at any age

- Three US studies showed Weimaraner had odds ratios of 16.26 (95% CI 3.22–82.22), 3.2 (95% CI 2.6–3.9) and 3.96 (95% CI 3.02–5.21) compared with crossbreeds
- OR 5.28 (95% CI 2.09–6.21) compared with crossbreeds in Austria (Goldschmidt & Mcmanus, 2000; Villamil *et al.*, 2011; White *et al.*, 2011; Leidinger *et al.*, 2014)

Neurological conditions

Disco-spondylitis

- Males and older dogs predisposed
- OR 5.6 (95% CI 1.8–18.1) compared with referral crossbreeds in the USA (Burkert *et al.*, 2005)

Hypomyelination syndrome

- Described in related Weimaraners in Spain
- Mutation in the folliculin-interacting protein 2 (FNIP2) gene reported in Weimaraner in the USA (Kornegay *et al.*, 1987; Millán *et al.*, 2010; Pemberton *et al.*, 2014)

Spinal dysraphism

- Described in Weimaraners in the UK
- Inherited myelodysplasia possibly transmitted by a codominant lethal gene with variable expressivity (van den Broek *et al.*, 1991)

Ocular conditions

Progressive retinal atrophy (PRA)

- PRA is highly heterogeneous, with autosomal dominant, recessive or X-linked modes of inheritance reported in dogs
- Large deletion of first four exons of the X-linked retinitis pigmentosa GTPase regulator (*RPGR*) gene identified in Weimaraner in Germany
- Represents a de novo mutation concerning only recent generations of the Weimaraner breed in Germany (Kropatsch *et al.*, 2016)

Renal and urinary conditions

Hyperuricosuria

- The predisposing *SLC2A9* mutation in Dalmatians has also been described in Weimaraner
- Weimaraner has 25.41% prevalence of carriers for the mutation in the USA (Karmi *et al.*, 2010a; Westropp *et al.*, 2014)

Urethral sphincter mechanism incompetence

- Larger breeds had an OR of 7.2 compared with smaller breeds in the USA
- Weimaraner over-represented in the UK
(Holt & Thrusfield, 1993; Bacon *et al.*, 2002; Forsee *et al.*, 2013)

Soft-tissue conditions**Peritoneopericardial diaphragmatic hernia (PPDH)**

- Solely the result of congenital anomalies in the dog
- Weimaraner comprised 30.8% of affected dogs but only 1.1% of the hospital population in one study
- Weimaraner comprised 21.4% and 62.5% of cases in two US referral caseloads
(Evans & Biery, 1980; Banz & Gottfried, 2010; Burns *et al.*, 2013)

WELSH CORGI (PEMBROKE, CARDIGAN, UNSPECIFIED VARIANTS)**Behavioural conditions****Aggression**

- A genetic basis for aggression has been described
- A survey of UK veterinarians classified Welsh Corgi as having high aggression
(Bradshaw & Goodwin, 1999)

Cardiovascular conditions**Patent ductus arteriosus**

- Diagnosed at 12–24 weeks in Pembroke Welsh Corgi
- Described in related Pembroke Welsh Corgis
- Strong genetic predisposition reported
(Oswald & Orton, 1993)

Metabolic conditions**Exercise-induced collapse**

- Many potential underlying causes of weakness and collapse associated with exercise exist in dogs
- The mutation causing dynamin 1 (*DNM1*)-associated exercise-induced collapse (d-EIC) has been identified in Pembroke Welsh Corgi
(Minor *et al.*, 2011)

Musculoskeletal conditions**Elbow dysplasia**

- Pembroke Welsh Corgi had 6.7% prevalence, compared with 0.7% in the overall population, among radiographed dogs in Poland
(Narojek *et al.*, 2008; Michelsen, 2013)

Immune-mediated inflammatory myopathy with severe tongue muscular atrophy

- Characterized in Pembroke Welsh Corgis in Japan
- Mean age at diagnosis 3.4 years
- Many muscles affected, but tongue lesions most severe
(Toyoda *et al.*, 2010)

Short tail (bobtail)

- Characterized in Pembroke Welsh Corgi in Norway
(Haworth *et al.*, 2001; Indrebø *et al.*, 2008; Hytönen *et al.*, 2009)

Neoplastic conditions**Histiocytic sarcoma**

- Pembroke Welsh Corgi had an OR of 9.7 (95% CI 5.6–17.0) compared with a referral population in Japan
- Median survival time was 43 days
(Takahashi *et al.*, 2014)

Lymphoma

- Pembroke Welsh Corgi had an OR of 2.1 (95% CI 1.6–2.8) compared with a US referral population
(Villamil *et al.*, 2009)

Neurological conditions**Degenerative myelopathy**

- Cardigan Welsh Corgi had 1.51% prevalence and Pembroke Welsh Corgi had 0.58% prevalence, compared with 0.15% in referral cross-breeds in the USA
- Homozygosity for the A allele of the canine *SOD1* gene associated with degenerative myelopathy in Pembroke Welsh Corgi
- Prevalence of the mutant A allele was 69.7% in Pembroke Welsh Corgis in Japan, but only a small percentage may become clinically affected
(Coates *et al.*, 2007; Awano *et al.*, 2009; March *et al.*, 2009; Shelton *et al.*, 2012; Chang *et al.*, 2013)

Intervertebral disc disease (IVDD)

- Pembroke Welsh Corgi had 15.11% prevalence, compared with 4.43% for crossbreeds in a US referral caseload
- Welsh Corgi had RR 2.3 ($p < 0.01$) compared with an overall referral population in the USA
(Priester, 1976; Bellumori *et al.*, 2013)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 3.05% in Pembroke Welsh Corgi and 2.53% in Cardigan Welsh Corgi, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance: point mutation identified in Cardigan Welsh Corgi
- DNA-based test designed for detection of the codon 616 mutation in the alpha cyclic GMP phosphodiesterase gene in the Cardigan Welsh Corgi
(Petersen-Jones *et al.*, 1999; Petersen-Jones & Entz, 2002)

Renal and urinary conditions

Familial renal disease (familial nephropathy)

- Described in related Pembroke Welsh Corgis in the USA
(McKay *et al.*, 2004)

Telangiectasia

- Possibly congenital or hereditary
- Described in unrelated Pembroke Welsh Corgis in the USA
- May show marked haematuria and die from renal failure or anaemia
(Moore & Thornton, 1983)

Urolithiasis – cystine

- Small breeds and males are predisposed
- Males are strongly predisposed
- Welsh Corgi had OR 5.0 (95% CI 2.0–12.7) compared with crossbred dogs in the USA
(Low *et al.*, 2010)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Small breeds and females are predisposed
- Welsh Corgi comprised 16.2% of a referral caseload in New Zealand
(Jones *et al.*, 1998)

Reproductive conditions

Dystocia

- 35.7% of Pembroke Welsh Corgi litters born by caesarean section in the UK
(Evans & Adams, 2010a)

High perinatal mortality

- Older dams and increasing litter size associated with increased perinatal mortality
- Overall perinatal mortality risk in Norway was 8.0%
- Pembroke Welsh Corgi had the fifth-highest perinatal mortality (16.9%) among registered purebred litters in Norway
(Tønnessen *et al.*, 2012)

WELSH SPRINGER SPANIEL

See *Springer Spaniel*

WELSH TERRIER

Dermatological conditions

Atopic dermatitis (atopy)

- Welsh Terrier had the second-highest incidence in insured dogs in Sweden: 13.4 cases per 1000 DYAR (95% CI 9.4–17.4)
(Nødtvedt *et al.*, 2006)

Musculoskeletal conditions

Pectus excavatum (funnel chest)

- Brachycephalic breeds are over-represented
- Atypical form that resolves spontaneously described in Welsh Terriers in the USA
(Ellison & Halling, 2004)

Neoplastic conditions

Mammary neoplasia

- 37% of insured Welsh Terrier bitches in Sweden develop the condition by 10 years of age, compared with 13% for bitches overall
(Jitpean *et al.*, 2012)

Neoplasia – overall

- Welsh Terrier had the ninth-highest proportional mortality from cancer among pedigree breeds in the UK: 43.5% (95% CI 23.3–63.7)
(Adams *et al.*, 2010)

Ocular conditions**Cataract**

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 5.14%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005;
Donzel *et al.*, 2016)

Lens luxation – primary

- *ADAMTS17* mutation has been identified: 35.9% of US Welsh Terriers were carriers
- Welsh Terrier had 5.74 (95% CI 2.76–11.93) RR compared with overall referral dogs in the USA

(Sargan *et al.*, 2007; Gould *et al.*, 2011)

WEST HIGHLAND WHITE TERRIER (WHWT)**Behavioural conditions****Aggression**

- A genetic basis for aggression has been described
- A survey of UK veterinarians classified WHWT as having high aggression (Bradshaw *et al.*, 1996; Bradshaw & Goodwin, 1999; Zapata *et al.*, 2016)

Victim of dog-to-dog aggression

- WHWT was over-represented as the victim in dog-dog conflict in Germany, representing 2.4% of victims and 1.5% of the general population

(Roll & Unshelm, 1997)

Cardiovascular conditions**Atrioventricular (AV) block**

- WHWT had the highest prevalence (37.5%) of second-degree atrioventricular blocks among referred cardiology cases in Poland (Noszczyk-Nowak *et al.*, 2017)

Congenital heart disease

- Includes a range of congenital heart disorders
- WHWT had an OR of 1.8 compared with all referral dogs in Italy

(Oliveira *et al.*, 2011)

Pulmonic stenosis

- WHWT comprised 12.3% of a UK referral caseload

(Johnson *et al.*, 2004)

Sick sinus syndrome

- Describes disturbed cardiac rhythm affecting the function of the sinus node
- Characterized in WHWTs in the UK
- Mean age of 10.5 years

(Moneva-Jordan *et al.*, 2001)

Dermatological conditions**Atopic dermatitis (atopy)**

- OR 10.3 (95% CI 3.6–29.3) compared with a referral population in Australia
- WHWT was over-represented in a worldwide study and also in Switzerland, Hungary and the USA
- Fourth-highest incidence in insured dogs in Sweden: 8.4 cases per 1000 DYAR (95% CI 7.2–9.6)
- 8.58% prevalence for WHWT, compared with 1.08% for crossbreds, in a US referral caseload
- WHWT comprised 3.8% (95% CI 1.8–6.8) of an atopic caseload, compared with 0.6% of the overall hospital population, in a US referral study

(Zur *et al.*, 2002; Nødtvedt *et al.*, 2006; Tarpataki *et al.*, 2006; Picco *et al.*, 2008; Favrot *et al.*, 2010; Jaeger *et al.*, 2010; Bellumori *et al.*, 2013)

Demodicosis

- WHWT was over-represented in the UK: 16.1% of laboratory diagnoses, versus 3.6% of normal clinic population
- Described as a predilected breed in a Swedish referral study
- Third most commonly affected breed (6.9%) in a pan-European study (Day, 1997a; Holm, 2003; Mueller *et al.*, 2009)

Malassezia dermatitis

- WHWT had the highest prevalence (16.4%) among a dermatology referral caseload in Romania
- WHWT had RR 7.82 compared to a general laboratory population in the USA

(Mauldin *et al.*, 1997;
Mircean *et al.*, 2010)

Superficial necrolytic dermatitis

- Males and older dogs may be predisposed
- WHWT suggested as predisposed and comprised 16.7% of a US referral caseload (Outerbridge *et al.*, 2002)

Drug reactions**Anaesthetic-related complications**

- WHWT had 8% incidence among general small animal practices in Canada (Dyson *et al.*, 1998)

Endocrine conditions**Diabetes mellitus**

- OR 1.99 (95% CI 0.89–4.42) compared with crossbreeds in the UK
- Eighth-highest breed incidence in Sweden: 33 cases per 10 000 DYAR (95% CI 20–46)
- WHWT comprised 6.6% of cases, compared with 4.3% of the overall population of insured dogs in the UK (Davison *et al.*, 2005; Fall *et al.*, 2007; Catchpole *et al.*, 2008; Mattin *et al.*, 2014)

Hypoadrenocorticism (Addison's disease)

- Complex inheritance pattern likely
- Females predisposed
- OR 11.42 (95% CI 4.69–26.97) compared with an overall US referral population
- WHWT comprised 19.5% of a US referral caseload (Melián & Peterson, 1996; Peterson *et al.*, 1996)

Gastrointestinal conditions**Chronic hepatitis**

- WHWT was 9 times over-represented ($p < 0.001$) in laboratory samples in Sweden
- Clinical liver disease in WHWTs caused by many etiologic agents in the USA
- Associated with hereditary copper toxicosis in WHWT in the USA (Thornburg *et al.*, 1986, 1996; Andersson & Sevelius, 1991)

Congenital portosystemic shunt

- Referred WHWT (0.31% prevalence) had an OR of 6.0 (95% CI 2.6–12.2) compared with an overall US referral population
- WHWT comprised 14.3% of referral cases in the UK and were over-represented in another UK referral caseload (Tobias & Rohrbach, 2003; Goodfellow *et al.*, 2008; Gow *et al.*, 2012)

Exocrine pancreatic insufficiency (EPI)

- Presumed inherited and autoimmune-mediated
- WHWT had 15.8% (95% CI 11.6–20.6) prevalence in a UK study (Batchelor *et al.*, 2007)

Oesophageal and gastric foreign bodies

- Median age affected was 4 years
- OR 8.93 (95% CI 5.25–15.17) compared with an overall referral population in Switzerland
- WHWT had 21.4% prevalence in a UK referral study (Sale & Williams, 2006; Gianella *et al.*, 2009)

Haematological/immunological conditions**Primary hepatitis**

- Multiple causes including microorganisms, toxins and drugs, immune-mediated reactions and breed-associated metabolic errors
- Females predisposed ($p < 0.01$)
- WHWT was over-represented in a referral population in the Netherlands (Poldervaart *et al.*, 2009)

Pyruvate kinase deficiency

- Affected dogs have abnormal red blood cells with a lifespan of about 20 days
- A 6 base-pair insertion in the C domain of the *R-PK* gene was identified in WHWT
- Genetic test developed (Skelly *et al.*, 1999)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- Inherited as an autosomal recessive trait
- OR 33.2 (95% CI 15.7–70.4) compared to crossbreeds in the USA
- 26.2% of WHWT were affected in the UK (Robinson, 1992; Brenig *et al.*, 1999; LaFond *et al.*, 2002)

Cranial cruciate ligament (CCL) disease

- Females and older dogs predisposed
- OR 2.6 (95% CI 1.4–4.8) compared with an overall UK referral population (Adams *et al.*, 2011)

Cranio-mandibular osteopathy (lion jaw)

- Inherited as an autosomal recessive trait
- Usually affects dogs aged 3–8 months

- OR 1313 (95% CI 219–7874) compared to crossbreeds in the USA
(LaFond *et al.*, 2002; Macedo *et al.*, 2015)

Chondrodysplasia (short-limbed or disproportional dwarfism)

- Part of the WHWT breed conformation description
- An *fgf4* retrogene identified as a cause of chondrodysplasia
(Israel *et al.*, 2009; Parker *et al.*, 2009)

Panosteitis (enostosis, eosinophilic panosteitis)

- Young males predisposed
- OR 1.7 (95% CI 1.2–2.4) compared to crossbreeds in the USA
(LaFond *et al.*, 2002; Trostel *et al.*, 2003)

Patellar luxation

- 2.5% of UK and 3.6% of US WHWTs affected
- Mainly medial luxation observed, often bilateral
- Compared to crossbreeds, WHWT had odds ratios of 2.0 (95% CI 1.4–2.8) in the UK and 1.8 (95% CI 1.1–2.9) in the USA
(LaFond *et al.*, 2002; Orthopedic Foundation for Animals, 2015; O'Neill *et al.*, 2016c)

Neoplastic conditions

Adenoma/adenocarcinoma

- WHWT had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland
(Grüntzig *et al.*, 2016)

Canine cutaneous histiocytoma

- OR 2.60 (95% CI 2.27–2.99) compared to crossbreeds in the USA
(Goldschmidt & Mcmanus, 2000; Fulmer & Mauldin, 2007)

Lower urinary tract neoplasia

- WHWT was over-represented in the UK, comprising 11.4% of a referral caseload
(Burnie & Weaver, 1983)

Neurological conditions

Globoid cell leucodystrophy (Krabbe disease)

- Autosomal recessive inheritance
- Deficiency of galactocerebrosidase (GALC) activity that is responsible for lysosomal catabolism of certain galactolipids

- Disease-causing mutation of GALC cDNA identified in the WHWT
- Genetic test developed
(Victoria *et al.*, 1996)

Acute canine polyradiculoneuritis

- The most common form of acute polyneuropathy in dogs
- Diagnosis more common in autumn and winter
- WHWT had an OR of 8.56 (95% CI 1.39–52.90) compared with other referral dogs in the UK
(Laws *et al.*, 2017)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.71%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)
(Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Corneal ulceration (ulcerative keratitis)

- OR 2.05 (95% CI 1.34–3.15) compared with crossbreeds in the UK
(O'Neill *et al.*, 2017b)

Keratoconjunctivitis sicca

- WHWT was the third most common breed in a 2007 UK referral study, and accounted for 35% of cases in a 1985 UK referral study
- WHWT was the most common breed in a European multicentre study, comprising 18% of the cases
(Sansom & Barnett, 1985; Sanchez *et al.*, 2007; Ofri *et al.*, 2009)

Physiological conditions

Abnormal mediolateral location of medial fabella

- Medial fabella abnormally placed in 70% of WHWTs in the UK
- Considered as an incidental finding, and these dogs are not clinically affected
(Störk *et al.*, 2009)

High allergen-specific serum immunoglobulin E levels

- Non-atopic WHWTs had statistically significantly higher positive ELISA results than atopic WHWTs for 44 of 48 allergens tested

that are commonly regarded as significant in canine atopic dermatitis in Australia
(Roque *et al.*, 2011)

Renal and urinary conditions

Ectopic ureter

- WHWT bitches had relative risks (RR) of 6.9 (95% CI 2.5–17.5) and 16.71 (95% CI 4.38–59.12) compared with overall referral populations in two US studies
(Hayes, 1974b, 1984)

Urolithiasis

- WHWT was the fifth most common breed for uroliths overall submitted for laboratory testing in the UK (4.4%)
(Rogers *et al.*, 2011)

Urolithiasis – calcium oxalate

- OR 3.28 compared with crossbreeds in the USA
(Lekcharoensuk *et al.*, 2000a)

Reproductive conditions

Dystocia

- WHWT had an incidence of 15.2 per 1000 DYAR, compared with 5.7 per 1000 DYAR in insured bitches overall in Sweden
- OR 2.5 (95% CI 1.3–4.9) compared with first-opinion emergency-care crossbred bitches in the UK

(Bergström *et al.*, 2006;
O'Neill *et al.*, 2017c)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- RR 2.5 (95% CI 1.4–3.5) compared with all bitches insured by Agria for veterinary care in Sweden
(Egenvall *et al.*, 2001; Smith, 2006)

Respiratory conditions

Bronchiectasis

- OR 4.45 (95% CI 2.20–8.99) compared with an overall US referral population
(Hawkins *et al.*, 2003)

Chronic pulmonary disease (CPD)

- May be analogous to idiopathic pulmonary fibrosis in humans
- Associated with pulmonary hypertension in WHWT
- CPD in WHWTs results from aberrant collagen regulation

- Mean age 10.9 years
- Characterized in WHWTs in the USA, the UK and Finland

(Norris *et al.*, 2005a;
Schober & Baade, 2006;
Corcoran *et al.*, 2011; Syrjä *et al.*, 2013)

WETTERHOUN (DUTCH SPANIEL, FRISIAN WATER DOG)

Haematological/immunological conditions

Severe combined immunodeficiency

- Unexplained pup mortality reported in 10% of all litters of Wetterhoun
- Nonsense mutation in the gene coding for V(D)J recombination factor RAG1 identified
(Verfurden *et al.*, 2011)

Metabolic conditions

Overweight/obesity

- Wetterhoun had significantly higher body condition score (5.38) than other show dogs in Holland
(Courcier *et al.*, 2010; Corbee, 2013;
Raffan *et al.*, 2016)

WHIPPET

Cardiovascular conditions

Mitral valve disease

- OR 5.3 (95% CI 2.2–12.5) compared with crossbreeds in a UK primary-care veterinary population
(Parker & Kilroy-Glynn, 2012;
Mattin *et al.*, 2015a, 2015b)

Drug reactions

Multiple drug sensitivity

- High doses of many drugs (e.g. ivermectin or milbemycin) cause neurological signs
- Associated with the nt230(del4) *MDR1* mutation
- Long-haired Whippet had a 24.3% allelic frequency with 0.7% recessive mutant allele frequency in Europe, and had a 45% allelic frequency in Germany
- 58.3% of referral Long-haired Whippets in the USA were carriers for the mutated *ABCB1-1* allele

- Whippet had a 0% allelic frequency in Brazil (Dowling, 2006; Mealey & Meurs, 2008; Gramer *et al.*, 2011; Monobe *et al.*, 2015; Firdova *et al.*, 2016)

Haematological/immunological conditions

Eccentrocytosis

- Young dogs predisposed
- Whippet was the most strongly over-represented breed among a referral population in Italy, with 30.0% of Whippets affected (Caldin *et al.*, 2005)

Phosphofructokinase deficiency

- Inherited as an autosomal recessive trait
- Present at 1 year of age
- Described in Whippets in the UK (Gerber *et al.*, 2009)

Musculoskeletal conditions

Double muscling

- Heavily muscled dogs, often called 'bully' whippets
- Associated with muscle cramping and a distinctive overbite
- US 'bully' whippets are homozygous for a deletion mutation in the myostatin gene (*MSTN*)
- The 'bully' phenotype displays a simple autosomal recessive mode of inheritance (Mosher *et al.*, 2007; Stinckens *et al.*, 2011)

Neoplastic conditions

Canine cutaneous histiocytoma

- OR 1.94 (95% CI 1.14–3.32) compared with crossbreeds in the USA (Goldschmidt & Mcmanus, 2000; Fulmer & Mauldin, 2007)

Haemangiosarcoma

- Two US studies reported that Whippet had odds ratios of 31.76 (95% CI 7.86–128.33) and 7.3 (95% CI 3.0–17.9) compared with crossbreeds (Hargis *et al.*, 1992; Goldschmidt & Mcmanus, 2000)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- Prevalence of primary cataract 2.01%,

compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003) (Gelatt & MacKay, 2005; Donzel *et al.*, 2016)

Lens luxation – primary

- Whippet had relative risks (RR) of 3.28 (95% CI 1.65–6.54) and 4.57 (95% CI 2.57–8.12) compared with overall referral dogs in the USA (Sargan *et al.*, 2007)

Progressive retinal atrophy (PRA)

- Autosomal recessive inheritance suspected
- Whippet had a 31.37% prevalence in a primary-care veterinary caseload in Brazil (Somma *et al.*, 2017)

Physiological conditions

Innocent systolic heart murmur

- Associated with athletic training
- 58.1% of Whippets in Belgium had systolic murmurs in the absence of structural abnormalities (Bavegems *et al.*, 2011)

Large heart size

- Whippets in Belgium had larger echocardiographic heart size and radiographic vertebral heart size than other breeds of comparable body weight (Bavegems *et al.*, 2005, 2007)

Reduced thyroxine levels

- Healthy Whippets had significantly lower mean values for total T_4 (20.74 ± 1.35 nmol/l) than healthy control dogs of a variety of general breeds (mean total T_4 of 29.00 ± 1.91 nmol/l) ($p=0.0007$) in Belgium (van Geffen *et al.*, 2006)

WHITE SWISS SHEPHERD DOG (BERGER BLANC SUISSE)

Drug reactions

Multiple drug sensitivity

- High doses of many drugs (e.g. ivermectin or milbemycin) cause neurological signs
- Associated with the nt230(del4) *MDR1* mutation

- White Swiss Shepherd had a 16.2% allelic frequency with 0.8% recessive mutant allele frequency in Europe, and had a 14% allelic frequency in Germany

(Dowling, 2006; Geyer *et al.*, 2007; Gramer *et al.*, 2011; Firdova *et al.*, 2016)

Musculoskeletal conditions

Degenerative lumbosacral stenosis

- Involves soft and bony tissue alterations of the spine coupled with suspected instability of the L7–S1 intervertebral disk
- White Swiss Shepherd comprised 15% of a referral caseload in Germany and Switzerland (Gödde & Steffen, 2007)

Lumbosacral transitional vertebrae

- Large breeds predisposed
- White Swiss Shepherd had 13.5% prevalence, compared with an overall 10.0% prevalence among dogs screened for pelvic problems in the Czech Republic

(Fialová *et al.*, 2014)

WOLFHOUND – IRISH

See *Irish Wolfhound*

WOLFSPITZ

See *Keeshond*

YORKSHIRE TERRIER

Behavioural conditions

Aggression

- A genetic basis for aggression has been described
- Yorkshire Terrier was reported to have a high tendency to aggression in Italy

(Notari & Goodwin, 2007; Zapata *et al.*, 2016)

Victim of dog-to-dog aggression

- Yorkshire Terrier was over-represented as the victim in dog–dog conflict in Germany, representing 2.9% of victims and 2.1% of the general population

(Roll & Unshelm, 1997)

Cardiovascular conditions

Arrhythmia

- Yorkshire Terrier had the highest prevalence (23.0%) of sinus tachycardia among referred cardiology cases in Poland

(Noszczyk-Nowak *et al.*, 2017)

Mitral valve disease

- OR 3.1 (95% CI 1.9–5.0) compared with crossbreeds in a UK primary-care veterinary population
- Yorkshire Terrier comprised 10.4% of a referral caseload in Italy
- Yorkshire Terrier comprised 12% of an affected caseload, compared with 8% of healthy controls from a general population in France

(Borgarelli *et al.*, 2008; Serres *et al.*, 2008; Parker & Kilroy-Glynn, 2012; Mattin *et al.*, 2015a, 2015b)

Patent ductus arteriosus

- Median age at presentation was 4 months
- Small breeds and females appear predisposed
- Yorkshire Terrier comprised 8.6% of a referral caseload in the USA

(Selmic *et al.*, 2013)

Ruptured chordae tendineae

- Recorded among dogs diagnosed with mitral valve disease
- Yorkshire Terrier comprised 8.8% of a referral caseload in France

(Serres *et al.*, 2007)

Dental conditions

Periodontal disease

- Yorkshire Terrier had prevalence of 25.2% (95% CI 18.6–33.4), compared with crossbreeds at 9.2% (95% CI 7.4–11.0) in the UK
- Yorkshire Terrier comprised 22.7% of a dental referral caseload of small breeds in South Korea

(Kim *et al.*, 2013; O'Neill *et al.*, 2014b)

Dermatological conditions

Atopic dermatitis (atopy)

- OR 8.85 ($p < 0.0001$) compared with a referral population in Greece

(Saridomichelakis *et al.*, 1999)

Colour dilution alopecia

- Described in a series of 10 Yorkshire Terriers

(Roperto *et al.*, 1995)

Dermatophytosis (ringworm)

- Animals aged < 1 year predisposed
- Zoonosis
- Yorkshire Terriers were over-represented for positive cultures in the UK and Brazil
- Yorkshire Terrier had the highest prevalence (46.4%) of breeds tested in Italy (Sparkes *et al.*, 1993; Brillhante *et al.*, 2003; Cafarchia *et al.*, 2004; Cerundolo, 2004)

Drug reactions**Vaccine-associated adverse effect**

- Smaller breeds predisposed
- Yorkshire Terrier had the tenth-highest incidence (47.3 per 10 000 dogs within 3 days of vaccine administration; 95% CI 40.3–55.3), which was significantly higher than in an overall US primary-care population (38.2 per 10 000; 95% CI 37.1–39.3) (Moore *et al.*, 2005)

Endocrine conditions**Diabetes mellitus**

- Familial and inherited
- Peak incidence age of 7–9 years and older entire females predisposed
- Yorkshire Terrier comprised 4.0% of cases, compared with 1.7% of the overall population of insured dogs in the UK
- OR 2.62 ($p < 0.001$) compared with a referral population in Italy (Fracassi *et al.*, 2004; Davison *et al.*, 2005)

Diabetic ketoacidosis

- Yorkshire Terrier comprised 16.1% of a referral caseload in Belgium (De Causmaecker *et al.*, 2009)

Hyperadrenocorticism (Cushing's syndrome)

- Yorkshire Terrier was the most common breed, comprising 17.5% of a referral caseload in Austria
- Yorkshire Terrier comprised 13.6% of a referral caseload of pituitary-dependent hyperadrenocorticism in Spain (Alenza *et al.*, 2006; Zeugswetter *et al.*, 2008)

Gastrointestinal conditions**Chronic hepatitis**

- Familial
- OR 4.6 (95% CI 1.9–11.2) compared with overall unselected dogs from first-opinion practices in the UK

- RR 4.6 (95% CI 1.9–11.2) among referral post-mortem cases in the UK (Watson *et al.*, 2007, 2010)

Cleft lip and/or palate

- Birth defect, small breeds predisposed
- RR 6.2 compared with all referral dogs in the USA (Mulvihill *et al.*, 1980)

Congenital portosystemic shunt

- Small breeds predisposed
- Suspected to be hereditary in Yorkshire Terriers
- Yorkshire Terrier had 10.86% prevalence for portosystemic shunt, compared with 0.35% for crossbreeds in a US referral study
- Referred Yorkshire Terriers (2.9% prevalence) had OR 58.7 (95% CI 42.9–80.2) compared with an overall US referral population
- Yorkshire Terriers comprised 20.0% and 21.9% of two US referral caseloads and were over-represented in a UK referral caseload (Tobias, 2003; Tobias & Rohrbach, 2003; Winkler *et al.*, 2003; Toulza *et al.*, 2006; Goodfellow *et al.*, 2008; Bellumori *et al.*, 2013)

Inflammatory bowel disease

- Mode of inheritance is incompletely understood and may differ between breeds
- Yorkshire Terrier comprised 19.2% of a referral caseload in Spain (Luckschander *et al.*, 2006; Kathrani *et al.*, 2011)

Lymphangiectasia (resulting in protein-losing enteropathy)

- Mean age 7–8 years, females over-represented
- Associated with hypobalaminemia in Yorkshire Terrier
- OR 10.1 (95% CI 5.2–18.8) compared with an overall US referral population
- Yorkshire Terrier comprised 17.6% of a US referral caseload (Kimmel *et al.*, 2000; Kull *et al.*, 2001; Grützner *et al.*, 2013; Simmerson *et al.*, 2014)

Oesophageal and gastric foreign bodies

- Median age affected was 4 years
- OR 3.13 (95% CI 1.63–6.03) compared with an overall referral population in Switzerland (Gianella *et al.*, 2009)

Pancreatitis

- Middle- to older-aged, overweight and dogs with hyperlipidaemia and diabetes mellitus predisposed
- OR 4.3 (95% CI 1.2–15.3) compared with an overall referral population in the USA (Hess *et al.*, 1999; Lem *et al.*, 2008)

Portal vein hypoplasia

- May occur in conjunction with portosystemic shunt (PSS)
- Cases present from 6 months to 6 years
- Yorkshire Terrier was the most commonly affected breed, comprising 28.6% of a US referral caseload (Christiansen *et al.*, 2000)

Musculoskeletal conditions**Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)**

- Inherited as an autosomal recessive trait
- A study showed 60% of cases were Yorkshire Terriers in Holland
- OR 35.8 (95% CI 20.0–63.9) compared to crossbreeds in the USA
- 34.0% of Yorkshire Terriers were affected in the UK (Robinson, 1992; Piek *et al.*, 1996; Brenig *et al.*, 1999; LaFond *et al.*, 2002)

Cranial cruciate ligament (CCL) disease

- Females and older dogs predisposed
- OR 2.84 (95% CI 1.46–5.55) compared with an overall UK referral population (Adams *et al.*, 2011)

Patellar luxation

- 5.4% of UK Yorkshire Terriers affected
- Mainly medial luxation observed, often bilateral
- Yorkshire Terrier had odds ratios of 5.5 (95% CI 4.3–7.1) in the UK and 8.3 (95% CI 6.4–10.8) in the USA compared to crossbreeds, and was over-represented in Korea
- Yorkshire Terrier comprised 10.0% of a US caseload and 10.9% of a caseload in Thailand (LaFond *et al.*, 2002; Alam *et al.*, 2007; Campbell *et al.*, 2010; Nganvongpanit & Yano, 2011; Orthopedic Foundation for Animals, 2015; O'Neill *et al.*, 2016c)

Neoplastic conditions**Adenoma/adenocarcinoma**

- Yorkshire Terrier had an OR significantly above 1.0 ($p < 0.05$) compared with crossbreeds from a cancer registry in Switzerland (Grüntzig *et al.*, 2016)

Mammary neoplasia

- Yorkshire Terrier had an incidence of 188 cases per 10 000 DYAR, compared with 116 per 10 000 DYAR in crossbreeds among insured entire bitches in Sweden
- 25% of insured Yorkshire Terrier bitches in Sweden develop the condition by 10 years of age, compared with 13% for bitches overall (Egenvall *et al.*, 2005; Jitpean *et al.*, 2012)

Neurological conditions**Atlantoaxial subluxation/instability**

- Congenital or developmental condition
- Young small-breed dogs affected
- Yorkshire Terrier comprised 21.3% of a Japanese case study, 46.7% of a UK case study, and 21.1%, 32.6% and 41.7% of three US case study populations (Denny *et al.*, 1988; Beaver *et al.*, 2000; Platt *et al.*, 2004; Sanders *et al.*, 2004; Aikawa *et al.*, 2013)

Idiopathic epilepsy

- Hereditary basis established in several breeds
- Yorkshire Terrier had an incidence of 25.4 per 10 000 DYAR (95% CI 18.6–32.1) among insured dogs in Sweden, compared with 16.1 (95% CI 14.7–17.4) for crossbreeds
- Yorkshire Terrier comprised 6.7% of epilepsy cases and 8.1% of idiopathic epilepsy cases among referred dogs in Japan (Heske *et al.*, 2014; Hamamoto *et al.*, 2016)

Intervertebral disc disease (IVDD)

- Yorkshire Terrier comprised 6.1% of a cervical IVDD caseload and 0.6% of a thoracolumbar IVDD caseload at a Japanese referral hospital (Itoh *et al.*, 2008)

Necrotizing meningoencephalitis

- Young adult dogs affected in Switzerland
- Yorkshire Terrier predisposed in Switzerland (Tipold *et al.*, 1993; Tipold, 1995; Kuwamura *et al.*, 2002)

Ocular conditions

Cataract

- 28% of cataracts reported as breed-related in France
- OR 1.6 for all cataracts in France
- Prevalence of primary cataract 4.33%, compared to 1.61% in mixed-breed dogs, in a retrospective study of dogs presenting at North American teaching hospitals (VMDB, 1964–2003)

(Gelatt & MacKay, 2005;
Donzel *et al.*, 2016)

Congenital alacrima (congenital keratoconjunctivitis sicca)

- RR 22.7 compared with an overall US referral population
- Yorkshire Terrier was suggested as predisposed in a study from Argentina and Spain (Herrera, 2007; Westermeyer *et al.*, 2009)

Corneal ulceration (ulcerative keratitis)

- Yorkshire Terrier was the third commonest breed, comprising 16% of a referral caseload in South Korea
- OR 1.78 (95% CI 1.13–2.80) compared with crossbreeds in the UK (Kim *et al.*, 2009; O'Neill *et al.*, 2017b)

Glaucoma

- 4.5% of Yorkshire Terriers referred for eye or neurological conditions in Japan were affected (Kato *et al.*, 2006)

Physiological conditions

Litter size

- Smaller litters associated with older bitches and smaller breeds
- Yorkshire Terrier had the twelfth smallest mean litter size (3.5 puppies) among registered breeds in Norway (Borge *et al.*, 2011)

Renal and urinary conditions

Kidney disease

- Yorkshire Terrier had an incidence of 34 (95% CI 26–42) cases per 10 000 DYAR, compared with 15.8 per 10 000 DYAR overall among insured dogs in Sweden (Pelander *et al.*, 2015)

Renal calculi

- May be associated with urinary tract infections
- Yorkshire Terrier had odds ratios of 4.29 ($p < 0.001$) in females and 11.39 ($p < 0.001$) in males compared with crossbreeds in a US referral study (Ling *et al.*, 1998c)

Urolithiasis

- Yorkshire Terrier was the most common breed submitted for urolith analysis in both the Benelux countries (8.5%) and Spain and Portugal (17.5%) and was the second most common in both the UK (7.4%) and Germany (9.4%) (Hesse, 1990; Picavet *et al.*, 2007; Rogers *et al.*, 2011; Vrabelova *et al.*, 2011)

Urolithiasis – calcium oxalate

- Small breeds and males are predisposed
- Yorkshire Terrier was the fifth most common breed with submissions for calcium oxalate uroliths in Canada
- Yorkshire Terrier had odds ratios of 7.2 (95% CI 6.3–8.2) and 9.85 compared with crossbred dogs in two US studies, and 13.0 (95% CI 11.7–14.4) compared with all dogs in the UK (Lekcharoensuk *et al.*, 2000a; Houston & Moore, 2009; Low *et al.*, 2010; Roe *et al.*, 2012; Okafor *et al.*, 2014)

(Lekcharoensuk *et al.*, 2000a;
Houston & Moore, 2009;
Low *et al.*, 2010; Roe *et al.*, 2012;
Okafor *et al.*, 2014)

Urolithiasis – silica

- OR 2.7 ($p < 0.001$) compared with all dogs in a US study (Aldrich *et al.*, 1997)

Urolithiasis – struvite (magnesium ammonium phosphate)

- OR 3.6 (95% CI 3.2–4.0) compared with all dogs in the UK
- Small breeds and females are predisposed (Houston & Moore, 2009; Roe *et al.*, 2012)

Urolithiasis – urate

- Yorkshire Terrier had 4.5 OR (95% CI 3.6–5.6) compared with all dogs in the UK
- Younger and female animals predisposed (Roe *et al.*, 2012)

Reproductive conditions

Cryptorchidism

- Yorkshire Terrier had relative risks (RR) of 2.9 (95% CI 2.2–4.0) and 3.0 (95% CI 1.76–5.06) compared with overall referral populations in two US studies
- Yorkshire Terrier had a 7.6% prevalence in the UK, which was significantly higher than for crossbreeds ($p=0.0018$)
(Pendergrass & Hayes, 1975; Hayes *et al.*, 1985; Yates *et al.*, 2003)

Dystocia

- OR 2.7 (95% CI 1.6–4.6) compared with first-opinion emergency-care crossbred bitches in the UK
(O'Neill *et al.*, 2017c)

Respiratory conditions

Tracheal collapse

- Common cause of cough in mature, small-breed dogs
- Yorkshire Terrier comprised 16% and 30% of two US referral caseloads, 50% and 65% of two UK referral caseloads, and 17.6% of a referral caseload in Korea
(White & Williams, 1994; Macready *et al.*, 2007; Marolf *et al.*, 2007; Eom *et al.*, 2008; Fonfara *et al.*, 2011)

Soft-tissue conditions

Perineal herniation

- Yorkshire Terrier comprised 13.6% of a UK caseload
(Raffan, 1993)

PART II

CAT BREEDS

ABYSSINIAN

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Abyssinians were found to have increased scores for sociability with people and cat aggression, but lower scores for restraint resistance and vocalization and lower likelihood of fear of noises

(Wilhelmy *et al.*, 2016)

Cardiovascular conditions

Aortic thromboembolism

- Breed at increased risk in retrospective series of 127 cases in the USA: OR 6.03 ($p=0.0019$) compared to the general hospital population
- Males at increased risk: OR 1.75 ($p=0.003$)

(Smith *et al.*, 2003)

Dermatological conditions

Atopic dermatitis (atopy)

- Breed at increased risk in case series
- In a retrospective study based at an Australian dermatology referral practice the Abyssinian

was over-represented compared to the general hospital population

- Cases diagnosed with flea-bite hypersensitivity and adverse food reactions were excluded from the study

(Ravens *et al.*, 2014)

Endocrine conditions

Diabetes mellitus

- Breed at risk in a retrospective case series based on Swedish insurance data 2009–2013. Abyssinians had an increased incidence rate ratio (IRR) of 1.8 (95% CI 1.0–3.2) compared to all other breeds
- Males at increased risk: IRR 2.0 (95% CI 1.8–2.3) across all breeds
- Mean age at diagnosis: 10.7 years across all breeds

(Öhlund *et al.*, 2015)

Haematological/immunological conditions

Increased osmotic fragility of erythrocytes

- Results in haemolytic anaemia
- Inheritance suspected
- Affected cats have chronic intermittent severe anaemia and splenomegaly

(Kohn *et al.*, 2000; Tritschler *et al.*, 2016)

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats

(Grahn *et al.*, 2012)

Infectious conditions**Feline infectious peritonitis (FIP)**

- Breed at increased risk in a retrospective series of 60 cases from the patient records of a North American veterinary medicine teaching hospital 1986–2002: OR 8.98 (95% CI 2.71–29.77) compared to all cats
- Also found to be at increased risk in a study of 382 cases in Australia. Observed frequency in the FIP cohort was 4.4%, compared to an expected frequency 1.5%
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in a multi-cat environment at increased risk (all breeds)

(Pesteanu-Somogyi *et al.*, 2006;
Worthing *et al.*, 2012)

Mycobacterium avium complex infection

- Very rare
- 10/12 cases were Abyssinians in a case series in Australia and the USA
- It is believed that some lines may suffer from familial immunodeficiency which predisposes them to infection

(Baral *et al.*, 2006)

Urinary tract infections (UTI)

See under *Renal and urinary conditions*

Musculoskeletal conditions**Acquired myasthenia gravis**

See under *Neurological conditions*

Patellar luxation

- In a study of cases in the USA and Europe, 26/69 Abyssinian cats (38%) were found to be affected, compared to 1/84 cats of other breeds (1.2%)
- Dominant, possibly polygenic inheritance suspected

(Engvall & Bushnell, 1990)

Neurological conditions**Acquired myasthenia gravis**

- Breed at increased risk in a retrospective study of 235 cases in the USA 2001–2012: OR 4.97 compared to all other breeds
- Bimodal age of presentation: 2–3 years and 9–10 years

(Hague *et al.*, 2015)

Ocular conditions**Progressive retinal atrophy (PRA) – early-onset**

- Early-onset rod–cone dysplasia (PRA-Rdy), which is inherited dominantly
- Rarely seen
- Presents with retinal changes at 8–12 weeks; progresses rapidly to blindness at 1 year
- Mutation identified

(Curtis *et al.*, 1987; Bedford, 1989)

Progressive retinal atrophy (PRA) – late-onset

- Late-onset rod–cone degeneration (PRA-rdAc), which is inherited recessively
- More common than PRA-Rdy in the Abyssinian
- Clinical onset 1.5–2 years; progresses to blindness over 2–4 years
- Mutation identified

(Narfström, 1985; Bedford, 1989;
Menotti-Raymond *et al.*, 2010)

Physiological conditions**Blood group**

- In an Australian study, 89% of this breed were type A, 11% were type B and none were type AB
- In an American study of 230 Abyssinians, 13.5% were found to be type B

(Giger *et al.*, 1991b; Malik *et al.*, 2005)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Abyssinians produced larger kittens (mean 100.1 g) than the mean of all breeds (93.5 g)

(Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Abyssinians produced smaller litters (mean 3.9 kittens) than the mean of all breeds (4.6 kittens)

(Sparkes *et al.*, 2006)

Renal and urinary conditions

Renal amyloidosis

- Usually seen as part of reactive systemic amyloidosis
- Familial
- Clinical presentation is variable – only those cats with moderate or severe disease develop symptoms
- Renal amyloid deposits are found principally in the medulla; glomerular involvement is less common. Amyloid deposits may also be found in adrenal and thyroid glands, spleen, stomach, small intestine, heart, liver, pancreas and colon. These deposits often do not contribute to the clinical signs, which are principally due to chronic renal failure

(Boyce *et al.*, 1984;
DiBartola *et al.*, 1986a, 1986b)

Urinary tract infections (UTI)

- Breed at increased risk in a retrospective case series of 784 cases from the VMDB (1980–1997): OR 3.4 compared to all breeds
 - Increased risk was seen for spayed female cats and those > 10 years of age (all breeds)
 - In a more recent study of 155 cases, no breed predilection was found
- (Lekcharoensuk *et al.*, 2001;
Martinez-Ruzafa *et al.*, 2012)

Reproductive conditions

Dystocia

- Breed at increased risk in a study based on Swedish insurance data, 1999–2006
- Incidence rates (IR) were 22 cases per 10 000 CYAR for all cats, and 100 cases per 10 000 CYAR for the Abyssinian group of cats (which included Somalis)
- Incidence rate ratio (IRR) in the Abyssinian group was 1.5 compared to all other purebred cats

(Holst *et al.*, 2017)

Pyometra (cystic endometrial hyperplasia-pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Abyssinian group of cats (which included Somalis) was 27 cases per 10 000 CYAR, compared to 17 cases for all breeds combined

(Hagman *et al.*, 2014)

AMERICAN SHORTHAIR

Physiological conditions

Blood group

- In a US study, all cats of this breed were type A (Giger *et al.*, 1991a)

ASIAN

Physiological conditions

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Asian cats produced smaller kittens (mean 84.7 g) than the mean of all breeds (93.5 g) (Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Asian cats produced larger litters (mean 6.5 kittens) than the mean of all breeds (4.6 kittens)
- (Sparkes *et al.*, 2006)

AUSTRALIAN MIST

Infectious conditions

Feline infectious peritonitis (FIP)

- In an Australian series of 42 cases (1990–2002) Australian Mist cats were considered over-represented, numbering 5 of the 42 cases
- Increased risk 3 months to 3 years (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in a multi-cat environment were at increased risk (all breeds)
- This breed is predisposed to the effusive form (Norris *et al.*, 2005b)

BALINESE

Ocular conditions

Progressive retinal atrophy (PRA)

See under *Siamese*

Reproductive conditions

Dystocia

See under *Siamese*

Pyometra (cystic endometrial hyperplasia-pyometra complex)

See under *Siamese*

BENGAL

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Bengals were found to have increased scores for predatory behaviour and inappropriate elimination

(Wilhelmy *et al.*, 2016)

Gastrointestinal conditions

Tritrichomonas foetus infection

See under *Infectious conditions*

Haematological/immunological conditions

Blood group

- In a UK study of 100 Bengal cats, all were found to be type A

(Gunn-Moore *et al.*, 2009)

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

Infectious conditions

Feline infectious peritonitis (FIP)

- Breed at increased risk in a retrospective series of 60 cases from the patient records of a North American veterinary medicine teaching hospital 1986–2002: OR 41.03 (95% CI 4.91–342.85) compared to all cats
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in a multi-cat environment at increased risk (all breeds)

(Pesteanu-Somogyi *et al.*, 2006)

Tritrichomonas foetus infection

- Breed at increased risk in a UK study
- Cats < 1 year old predisposed

(Gunn-Moore *et al.*, 2007)

Musculoskeletal conditions

Thoracic wall abnormalities

- In a study of thoracic wall defects in kittens presented for vaccination at a single UK practice, 12/244 Bengal kitten were affected, compared to none of 1748 domestic short-haired kittens
- Defects included pectus excavatum, unilateral thoracic wall concavity and scoliosis
- Familial basis suspected

(Charlesworth & Strugess, 2012)

Neurological conditions

Polyneuropathy

- Characterized by recurrent myelination and demyelination
- 37 cases reported; all were Bengal cats
- 65% were male
- Mean age at onset 10.6 ± 7.9 months

(Bensfield *et al.*, 2011)

Ocular conditions

Cataract

- Cataracts were diagnosed in 45% of 51 Bengal cats recruited in an observational study to assess the national prevalence of ocular disease in France, October 2014 to November 2016
- Pedigree analysis suggested inheritance
- No visual impairment was reported

(Bourguet *et al.*, 2017)

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- Bengal cats had an OR of 5.6 (95% CI 2.83–10.98)
- Mean age 6.2 years (all breeds)
- Males were at increased risk: OR 1.1 (all breeds)

(Albasan *et al.*, 2012)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006

- IR for the Bengal was 64 cases per 10 000, CYAR compared to 17 cases for all breeds combined (Hagman *et al.*, 2014)

BIRMAN

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Birmans were found to have increased scores for fear-related aggression to familiar people and inappropriate elimination, but lower scores for activity and playfulness, vocalization, trainability and predatory behaviour (Wilhelmy *et al.*, 2016)

Wool-sucking (compulsive)

- Breed at increased risk
- Early weaning and small litter size were found to be additional risk factors in Birman cats in one study
- Affected cats had an abnormally intense appetite (Borns-Weil *et al.*, 2015)

Cardiovascular conditions

Aortic thromboembolism

- Breed at increased risk in a retrospective series of 127 cases in the USA: OR 10.52 ($p=0.0001$) compared to the general hospital population
- Males at increased risk: OR 1.75 ($p=0.003$) (Smith *et al.*, 2003)

Dermatological conditions

Congenital hypotrichosis

- Autosomal recessive inheritance, mutation identified
- May be associated with thymic aplasia and short life expectancy (Casal *et al.*, 1994; Abitbol *et al.*, 2015)

Gastrointestinal conditions

Tritrichomonas foetus infection

See under *Infectious conditions*

Haematological/immunological conditions

Atypical granulation of neutrophils

- Inherited as an autosomal recessive trait
- Common, with 46% of Birman cats studied affected

- Neutrophils have prominent eosinophilic granules
- No abnormalities of neutrophil function detected

(Hirsch & Cunningham, 1984)

Infectious conditions

Chlamydophilosis

- Breed at increased risk in a survey of cats in Britain
- Prevalence highest from 5 weeks to 9 months of age (all breeds)
- Males predisposed (all breeds) (Wills *et al.*, 1988)

Cryptococcosis

- Breed at significantly increased risk ($p=0.011$) in a retrospective series of 155 feline cases in Australia 1981–2001
- Across all breeds median age 6 years (range 1–16) with a peak at 2–3 years (O'Brien *et al.*, 2004)

Feline infectious peritonitis (FIP)

- Breed at increased risk in retrospective series of 60 cases from the patient records of a North American Veterinary medicine teaching hospital 1986–2002: OR 82.06 (95% CI 26.66–262.44) compared to all cats
- Inheritance of susceptibility is suspected to be polygenic, and heritability has been found to be high in Birmans
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in a multi-cat environment are at increased risk (all breeds) (Foley & Pedersen, 1996; Pesteanu-Somogyi *et al.*, 2006)

Tritrichomonas foetus infection

- Breed at increased risk in a study in Germany (Kuehner *et al.*, 2011)

Neurological conditions

Audiogenic reflex seizures

- Birmans were strongly represented (31% of cases) in a study of 96 cases in the UK recruited by questionnaire, September 2013 to March 2014
- Mean age of all cases was 15 years (10–19 years) (Lowrie *et al.*, 2015a)

Ocular conditions

Chlamydophila psittaci conjunctivitis

See under *Infectious conditions*

Physiological conditions

Azotaemia

- One study showed that 82% of healthy Birman cats <6 months of age had creatinine levels above the reference range. The significance of this was unknown
- In a separate study of 91 healthy Birman cats, new higher reference intervals of 97.2–221.0 mmol/l were established for Birman cats (compared to a general feline reference range of 70.7–159.1 mmol/l)
(Gunn-Moore *et al.*, 2002; Paltrinieri *et al.*, 2014)

Blood group

- In a US study, 82% of Birman cats were type A and 18% type B
- In a UK study (1995–1998) in a population of 24 Birmans 15 (62.5%) were type A, 7 (29.2%) were type B and 2 (8.3%) were type AB
(Giger *et al.*, 1991a; Knottenbelt *et al.*, 1999)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Birman cats produced larger kittens (mean 101 g) than the mean of all breeds (93.5 g)
(Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Birman cats produced smaller litters (mean 3.6 kittens) compared to the mean of all breeds (4.6 kittens)
(Sparkes *et al.*, 2006)

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- Birman cats had an OR of 6.77 (95% CI 5.01–9.15)
- Mean age (all breeds) 6.2 years
- Males were at increased risk (all breeds): OR 1.1

- In a separate study of 21 426 bladder uroliths, Birmans had an OR of 9.45 (95% CI 3.91–20.8) compared to domestic short-haired cats
(Albasan *et al.*, 2012; Houston *et al.*, 2016)

Reproductive conditions

Dystocia

- Breed at increased risk in a study based on Swedish insurance data, 1999–2006
- Incidence rates (IR) were 22 cases per 10 000 CYAR for all purebred cats combined and 101 cases per 10 000 CYAR for Birman cats
- Incidence rate ratio (IRR) in the Birman was 1.7 compared to all other purebred cats
(Holst *et al.*, 2017)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Birman was 35 cases per 10 000 CYAR compared to 17 cases for all breeds combined
(Hagman *et al.*, 2014)

BRITISH SHORTHAIR (BSH)

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM)

- Familial
- In a Danish study of 329 BSH cats screened by echocardiography for HCM, 8.5% were positive and 4.3% equivocal for the condition
- Median age at diagnosis 2.7 years
- Male cats at increased risk: OR 7.89 (95% CI 2.54–28.08)
(Granström *et al.*, 2011)

Gastrointestinal conditions

Trichomonas foetus infection

See under *Infectious conditions*

Haematological/immunological conditions

Haemophilia B

- Uncommon condition, occurs in this breed
(Littlewood, 1989)

Lymphoproliferative disease

- Novel disease recognized in multiple closely related BSH kittens in New Zealand

- Inheritance suspected
- Non-neoplastic proliferation of T cells (Aberdein *et al.*, 2015)

Infectious conditions

Feline infectious peritonitis (FIP)

- Breed at increased risk in a retrospective Australian series of 382 cases
- Observed frequency of BSH cats in the FIP cohort was 15.5%, compared to an expected frequency of 2.4%
- Susceptibility to FIP may be partially inherited polygenically, as suggested in one study of purebred catteries in the USA
- Increased risk 3 months to 3 years (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in a multi-cat environment at increased risk (all breeds) (Foley & Pedersen, 1996; Norris *et al.*, 2005b; Worthing *et al.*, 2012)

Tritrichomonas foetus infection

- Breed at increased risk in a study in Germany (Kuehner *et al.*, 2011)

Neurological conditions

Intervertebral disc disease (IVDD)

- In a UK retrospective study of the medical records of cats diagnosed with thoracolumbar intervertebral disc disease, 2008–2014, BSH cats were found to be at significantly increased risk ($p < 0.0001$) compared with the general hospital population (De Decker *et al.*, 2017)

Physiological conditions

Blood group

- In an Australian study including 8 BSH cats, 3 (38%) were type A, 5 (62%) were type B and none were type AB
- In a UK study (1995–1998), in a population of 121 BSH cats 48 (39.7%) were type A, 71 (58.7%) were type B and 2 (1.6%) were type AB
- In a small study including 5 BSH cats in England, 2 (40%) were type A and 3 (60%) were type B
- In a US study, 41% were type A and 59% type B (Giger *et al.*, 1991a; Knottenbelt *et al.*, 1999; Malik *et al.*, 2005; Forcada *et al.*, 2007)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, BSH cats produced larger kittens (mean 104.4 g) compared to the mean of all breeds (93.5 g) (Sparkes *et al.*, 2006)

Renal and urinary conditions

Polycystic kidney disease (PKD)

- Most common inherited disease in cats
- Some BSH cats with the condition share the same genetic mutation as Persians
- Autosomal dominant inheritance (Nivy *et al.*, 2015)

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, BSH cats had an OR of 7.8 (95% CI 4.87–34.82) compared to a control population of cats from the VMDB over the same period
- Males at increased risk: OR 1.5 (all breeds)
- Mean age 90 ± 41 months across all breeds (Lekcharoensuk *et al.*, 2000b)

Reproductive conditions

Dystocia

- Breed at increased risk in a study based on Swedish insurance data, 1999–2006
- Incidence rates (IR) were 22 cases per 10 000 CYAR for all cats and 157 cases per 10 000 CYAR for BSH cats
- Incidence rate ratio (IRR) in the BSH was 2.5 compared to all other purebred cats (Holst *et al.*, 2017)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- Incidence rate (IR) for the British Shorthair was 32 cases per 10 000 CYAR compared to 17 cases for all breeds combined (Hagman *et al.*, 2014)

BURMESE

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires,

Burmese were found to have increased scores for dog aggression, prey interest, sleeping in elevated/warm/hidden locations and separation anxiety

(Wilhelmy *et al.*, 2016)

Cardiovascular conditions

Endocardial fibroelastosis

- Inherited congenital anomaly
- Age of onset 3 weeks to 4 months
- Short illness in many cases, sudden death possible
- Cats with mild forms survive to adulthood (Zook & Paasch, 1982; Darke, 1989)

Dermatological conditions

Cutaneous asthenia (Ehlers–Danlos-like syndrome)

- Inherited (Hansen *et al.*, 2015)

Cutaneous food allergy

- Burmese cats were over-represented in a study of 48 cases (Scott & Miller, 2013)

Cutaneous neoplasia

See under *Neoplastic conditions*

Endocrine conditions

Diabetes mellitus

- Breed at increased risk in many studies (UK, Australia, Sweden)
- In a cohort of 193 563 cats in primary-care practice in England, 2009–2014 (VetCompass data), there were 1128 cases of diabetes mellitus (period prevalence 0.58%). Burmese were at increased risk: OR 3.0 (95% CI 2.0–4.4) compared to the overall population. Both being >4 kg body weight and >6 years of age were risk factors across all breeds ($p < 0.001$)
- Breed at increased risk in a retrospective series based on Swedish insurance data 2009–2013. Burmese had an increased incidence rate ratio (IRR) of 4.3 (95% CI 2.9–6.0) compared to all other breeds. Males at increased risk: IRR 2.0 (95% CI 1.8–2.3) across all breeds. Mean age at diagnosis 10.7 years across all breeds
- Inheritance suspected (O'Leary *et al.*, 2013; Öhlund *et al.*, 2015; O'Neill *et al.*, 2016b)

Infectious conditions

Feline infectious peritonitis (FIP)

- Breed at increased risk in Australian series of 42 cases. In this study male Burmese cats were over-represented, and 7/10 cases in Burmese cats were the non-effusive form
- In another study in the USA, Burmese were not found to be at increased risk compared to mixed-breed cats (Norris *et al.*, 2005b; Pesteanu-Somogyi, 2006)

Musculoskeletal conditions

Flat-chested kittens

- Reported to affect 3–4% of Burmese kittens in the UK (Sturges *et al.*, 1997)

Frontonasal dysplasia (Burmese head defect)

- Multiple congenital abnormalities of the head. Most kittens do not survive
- Inherited as autosomal recessive
- Created by selective breeding from a lineage with a more brachycephalic head shape (known as the Contemporary Burmese)
- Cats with shorter faces may be carriers
- Seen mainly in the USA (Sponenberg & Graf-Webster, 1986; Lyons *et al.*, 2016a)

Hypokalaemic polymyopathy

- Autosomal recessive inheritance. Mutation identified
- Seen in Burmese cats and closely related breeds
- Signs are often episodic and seen as muscle weakness and pain in the first year of life (Malik *et al.*, 2015)

Neoplastic conditions

Mast cell tumour (MCT) – cutaneous

- Breed at increased risk
- In a UK study of 287 records of feline MCTs, Burmese cats represented 2.4% of the study population, but only 1.1% of the control population
- Median age 11 years (range 5 months to 19 years) across all breeds (Melville *et al.*, 2015)

Neurological conditions

Feline orofacial pain syndrome

- Predominantly seen in Burmese cats. In a UK study of 113 cases, 100 were Burmese

- Suspected to be a neuropathic condition, may be triggered by oral/dental disease
- Symptoms of acute oral pain and self-mutilation are initially episodic but may become unremitting with time

(Rusbridge *et al.*, 2010)

Hereditary meningoencephalocele

- Seen as part of frontonasal dysplasia (see under *Musculoskeletal conditions*)

Lysosomal storage disease – GM₂ gangliosidosis

- Inherited, mutation identified
- Signs from 1–2 months of age include intention tremors, progressing to ataxia, hindlimb weakness and inability to stand by 4–5 months of age

(Bradbury *et al.*, 2009, 2105)

Ocular conditions

Corneal sequestrum

- In a retrospective review of 97 cases, 17 cats were Burmese
- Mean age 6.8 years

(Graham *et al.*, 2016)

Glaucoma – primary

- A retrospective study (1996–2001) of 6 affected Burmese cats led to a suggestion of a predisposition to primary narrow-angle glaucoma in this breed
- Affected cats were aged 7–10.5 years

(Hampson *et al.*, 2002)

Lipaemia of the aqueous humour

- Young cats (5 months to 1 year) affected
- Seen in the UK, Australia and New Zealand

(Gunn-Moore & Crispin, 1998)

Prolapse of the gland of the nictitating membrane ('cherry eye')

- Uncommon condition in the cat, reported most frequently in this breed

(Chahory *et al.*, 2004)

Uveal cysts

- Breed at increased risk
- In a retrospective study conducted in the UK and Australia, 36/5017 cats had uveal cysts (prevalence 0.72%), and 21 of these were Burmese (total of 516 Burmese cats; incidence 4.1%)

- Only 2 cats had concurrent intraocular disease

(Blacklock *et al.*, 2016)

Physiological conditions

Blood group

- In a UK study (1995–1998), 9/10 Burmese cats (90%) were type A and 1 (10%) was type B
- In an Australian study (1992–2003) of 30 Burmese cats, 28 (93%) were type A, 1 (3%) was type B and 1 (3%) was type AB
- In a UK study of 5 Burmese, all were found to be type A

(Knottenbelt *et al.*, 1999;

Malik *et al.*, 2005; Forcada *et al.*, 2007)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Burmese cats produced smaller kittens (mean 86.2g) than the mean of all breeds (93.5g)

(Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Burmese cats produced larger litters (mean 5.7 kittens) than the mean of all breeds (4.6 kittens)

(Sparkes *et al.*, 2006)

Triglyceridaemia

- A proportion of Burmese cats in Sydney, Australia, were found to have a more exaggerated and long-lasting triglyceridaemia than other cats, following a high-fat meal
- It has been suggested that this may result in increased insulin resistance and therefore risk of diabetes mellitus

(Kluger *et al.*, 2009)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Burmese cats had an OR of 3.74 (95% CI 2.08–7.22) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.73 (95% CI 1.63–1.82)
- Also found to be at risk in a US study

(Thumchai *et al.*, 1996; Houston *et al.*, 2016)

Respiratory conditions

Agensis of the nares

- Seen as part of frontonasal dysplasia (see under *Musculoskeletal conditions*)
(Noden & Evans, 1986)

CHARTREUX

Renal and urinary conditions

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in case series
- In a study of 7456 uroliths submitted to the Minnesota Urolith Center, 1981–1997, Chartreux cats had an OR of 4.51 (95% CI 1.66–21.51) compared to a control population of cats from the VMDB over the same period
- Males reported to be at reduced risk: OR 0.7 (all breeds)
- Mean age 69 ± 38 months (all breeds)
(Lekcharoensuk *et al.*, 2000b)

CHINCHILLA

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Persian*

COLOURPOINT PERSIAN

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Persian*

CORNISH REX

Infectious conditions

Feline infectious peritonitis (FIP)

- Cornish and Devon Rex combined were found to be at increased risk in a retrospective series of 60 cases from the patient records of a North American veterinary medicine teaching hospital, 1986–2002: OR 38.29 (95% CI 8.42–174.15) compared to all other cats
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)

- Cats in a multi-cat environment at increased risk (all breeds)
(Foley & Pedersen, 1996; Norris *et al.*, 2005b; Pesteanu-Somogyi *et al.*, 2006)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Cornish Rex was 50 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined
(Hagman *et al.*, 2014)

DEVON REX

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Devon Rex were found to have increased scores for playfulness, sociability with people and sleeping in elevated/warm/hidden locations
(Wilhelmy *et al.*, 2016)

Dermatological conditions

Malassezia dermatitis

- Increased numbers of *Malassezia* organisms are isolated in this breed from both seborrhoeic and healthy cats, compared to domestic short-haired and Cornish Rex cats
(Ahman *et al.*, 2007; Bond *et al.*, 2008)

Urticaria pigmentosa-like dermatosis

- Reported in Devon Rex cats
(Noli *et al.*, 2004; Colombo *et al.*, 2012)

Haematological/immunological conditions

Vitamin-K-dependent coagulopathy

- There are a number of reports suggesting the presence of this condition in Devon Rex cats in the UK and Australia
(Soute *et al.*, 1992)

Infectious conditions

Feline infectious peritonitis (FIP)

See under *Cornish Rex*

Infectious skin diseases

See under *Dermatological conditions*

Musculoskeletal conditions

Hereditary myopathy of Sphynx and Devon Rex cats

- Autosomal recessive inheritance
- Variably progressive
- Seen in the Devon Rex and the related Sphynx
- May represent a form of congenital myasthenia gravis

(Martin *et al.*, 2008; Gandolfi *et al.*, 2015)

Physiological conditions

Blood group

- In an Australian study including 71 Devon Rex cats, 32 cats (45%) were type A, 38 (54%) were type B and 1 (1.4%) was type AB

(Malik *et al.*, 2005)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Devon Rex cats had an OR of 2.4 (95% CI 1.21–5.09) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.73 (95% CI 1.63–1.82)

(Houston *et al.*, 2016)

Reproductive conditions

Dystocia

- Breed at increased risk
- In a survey of cat breeders reporting on 2928 litters, dystocia occurred in 5.8% of all litters, but in 18.2% of Devon Rex litters

(Gunn-Moore & Thrusfield, 1995)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Devon Rex was 59 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined

(Hagman *et al.*, 2014)

EGYPTIAN MAU**Haematological/immunological conditions**

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- Egyptian Mau cats had an OR of 44.41 (95% CI 32.95–59.86)
- Mean age was 6.2 years (all breeds)
- Males were at increased risk: OR 1.1 (all breeds)
- In a separate study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014, Egyptian Maus had an OR of 94.5 (95% CI 38.2–285)

(Albasan *et al.*, 2012; Houston *et al.*, 2016)

EUROPEAN SHORTHAIR**Renal and urinary conditions**

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- European Shorthair cats had an OR of 61.28 (95% CI 43.17–86.98)
- Mean age 6.2 years (all breeds)
- Males were at increased risk: OR 1.1 (all breeds)

(Albasan *et al.*, 2012)

EXOTIC SHORTHAIR

Ocular conditions

Epiphora

- Brachycephalia affects the course of the nasolacrimal duct and the efficiency of tear drainage. All 4 Exotic Shorthair cats were classified as severely brachycephalic in one study

(Schlueter *et al.*, 2009)

Renal and urinary conditions

Polycystic kidney disease (PKD)

- Autosomal dominant inheritance, mutation identified
- Affected cases may present with renal failure from 2 to 3 years of age
- In a French study of 64 Exotic Shorthair cats, 39.1% were found to be affected when screened with ultrasound
- Studies have demonstrated similar prevalences in USA, Australia and Italy

(Barrs *et al.*, 2001; Barthez *et al.*, 2003; Bonazzi *et al.*, 2009)

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Exotic Shorthair cats had an OR of 2.9 (95% CI 1.68–6.81) compared to a control population of cats from the VMDB over the same period
- Males reported to be at increased risk: OR 1.5 (all breeds)
- Mean age 90 ± 41 months across all breeds (Lekcharoensuk *et al.*, 2000b)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Persian*

Stillborn kittens

- In a UK questionnaire-based study of 1056 litters, Exotic Shorthair cats had a significantly higher prevalence of stillbirths (12.5%) than that of all breeds combined (7.2%)

(Sparkes *et al.*, 2006)

FOREIGN SHORTHAIR

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Foreign Shorthair cats had an OR of 7.15 (95% CI 3.75–11.77) compared to a control population of cats from the VMDB over the same period
- Males reported to be at increased risk: OR 1.5 across all breeds
- Mean age 90 ± 41 months (all breeds) (Lekcharoensuk *et al.*, 2000b)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in case series
- In a study of 7456 uroliths submitted to the Minnesota Urolith Center, 1981–1997, Foreign Shorthair cats had an OR of 18.46 (95% CI 11.12–25.74) compared to a control population of cats from the VMDB over the same period
- Males reported to be at reduced risk: OR 0.7 across all breeds
- Mean age 69 ± 38 months (all breeds) (Lekcharoensuk *et al.*, 2000b)

FOREIGN WHITE

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Siamese*

HAVANA BROWN

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Havana Brown cats had an OR of 5.47 (95% CI 2.49–18.38) compared to a control population of cats from the VMDB over the same period

- Males reported to be at increased risk: OR 1.5 across all breeds
- Mean age 90 ± 41 months (all breeds) (Lekcharoensuk *et al.*, 2000b)

HIMALAYAN

Cardiovascular conditions

Peritoneopericardial diaphragmatic hernia (PPDH)

- In a USA study of 67 cases 1987–2002, Himalayans were found to be significantly over-represented compared to the hospital population (prevalence of 1.45% in the Himalayan compared to 0.25% in the hospital population)
- Median age at presentation: 54 months (range 1.5–204 months) (Reimer *et al.*, 2004)

Dermatological conditions

Cheyletiellosis

- Himalayan cats were over-represented in a Canadian study of 111 cats with skin disease, accounting for 50% of the cases of cheyletiellosis, but only 6.9% of the hospital population (Scott & Paradis, 1990)

Cutaneous food allergy

- Himalayan cats were over-represented in a study of 48 cases (Scott & Miller, 2013)

Cutaneous neoplasia

See under *Neoplastic conditions*

Idiopathic facial dermatitis in Persians and Himalayans

- Breed at increased risk (Chung *et al.*, 2009)

Gastrointestinal conditions

Congenital portosystemic shunt

- In an Australian retrospective study of 9 cats, Himalayan cats were over-represented, with an OR 30.6 ($p < 0.001$) compared to the general hospital population
- Males may be predisposed

- High rate of cryptorchidism in affected males (24%)
- Both intra- and extrahepatic shunts can occur; extrahepatic are more common (Tillson & Winkler, 2002; Hunt, 2004)

Periodontal disease

- In a US study of 16 374 cats >5 years of age with a diagnosis of periodontal disease, Himalayans had an RR of 1.6 (95% CI 1.3–2.0) of being represented in the periodontal disease group compared to being represented in a hospital population of cats of similar age but without periodontal disease (Lund, 2012)

Infectious conditions

Aspergillosis

- Persian and Himalayan cats seem to be over-represented in a number of case reports (Barrs & Talbot, 2014)

Feline infectious peritonitis (FIP)

- Breed at increased risk in a retrospective series of 60 cases from the patient records of a North American veterinary medicine teaching hospital 1986–2002: OR 3.19 (95% CI 1.12–9.06) compared to all cats
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in multi-cat environment at increased risk (all breeds) (Foley & Pedersen, 1996; Pesteanu-Somogyi *et al.*, 2006)

Neoplastic conditions

Basal cell tumour

- Breed at increased risk in a series of 124 cases
- Risk increases with increasing age (Diters & Walsh, 1984)

Ocular conditions

Corneal sequestrum

- In a retrospective review of 97 cases, 18 cats were Himalayan
- Mean age 6.8 years (all breeds combined)
- Predisposition suggested in other studies (Morgan, 1994; Graham *et al.*, 2016)

Physiological conditions

Blood group

- In a US study, 80% were type A and 20% type B

(Giger *et al.*, 1991a)

Renal and urinary conditions

Polycystic kidney disease (PKD)

- Autosomal dominant inheritance
- In an Australian study of 48 Himalayan cats which were screened by renal ultrasound, 42% were found to be affected

(Barrs *et al.*, 2001)

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Himalayan cats had an OR of 2.7 (95% CI 2.02–2.81) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.73 (95% CI 1.63–1.82)
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Himalayan cats had an OR of 7.86 (95% CI 7.15–8.65) compared to a control population of cats from the VMDB over the same period

(Lekcharoensuk *et al.*, 2000b;

Houston *et al.*, 2016)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in a case series
- In a study of 7456 uroliths submitted to the Minnesota Urolith Center, 1981–1997, Himalayan cats had an OR of 2.01 (95% CI 1.85–2.49) compared to a control population of cats from the VMDB over the same period
- Males reported to be at reduced risk: OR 0.7, across all breeds
- Mean age 69 ± 38 months across all breeds

(Lekcharoensuk *et al.*, 2000b)

Respiratory conditions

Aspergillosis

- Persian and Himalayan cats seem to be over-represented in a number of case reports

(Barrs & Talbot, 2014)

Brachycephalic obstructive airway syndrome (BOAS)

- Likely a consequence of selective breeding for certain facial characteristics

(Ginn *et al.*, 2008; Meola, 2013)

JAPANESE BOBTAIL

Musculoskeletal conditions

Vertebral anomalies

- Japanese Bobtail cats feature variations of tail morphology caused by transitional vertebrae, changes in the numbers of vertebrae and hemivertebrae
- Simple autosomal dominant inheritance (with variable expression of tail length and kink placement)

(Pollard *et al.*, 2015)

JAVANESE

Ocular conditions

Progressive retinal atrophy (PRA)

See under *Siamese*

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Siamese*

KORAT

Neurological conditions

Lysosomal storage disease – GM₁ gangliosidosis

- Autosomal recessive inheritance, mutation identified
- Rare
- Signs seen at 3–6 months

(De Maria *et al.*, 1998)

Lysosomal storage disease – GM₂ gangliosidosis

- Autosomal recessive inheritance, mutation identified
- Rare
- Signs seen at 3–6 months

(Muldoon *et al.*, 1994)

Physiological conditions

Gestation length

- In a UK questionnaire-based study of 1056 litters, Korats had shorter gestation (mean 63 days) than the mean of all breeds (65.1 days)

(Sparkes *et al.*, 2006)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Korats produced smaller kittens (mean 72.7 g) than the mean of all breeds (93.5 g)

(Sparkes *et al.*, 2006)

LA PERM

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

MAINE COON

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Maine Coons were found to have increased scores for owner directed aggression and prey interest, but lower scores for attention-seeking behaviours, separation anxiety, sleeping in elevated/warm/hidden positions and inappropriate elimination

(Wilhelmy *et al.*, 2016)

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM)

- Inherited as an autosomal dominant trait, mutation identified
- Homozygosity for the mutation has been found to increase the risk of developing HCM compared to unaffected Maine Coon cats: OR

21.6 (95% CI 7.01–66.2). Heterozygosity did not increase the risk in this study

- In a UK-based study including 742 Maine Coon cats, the prevalence of the mutation was 39.4%
- Prevalence of the mutation has been described as 22% in Germany, 46% in Australia/New Zealand, and 34% worldwide
- Most affected cats developed HCM by 6 years of age, and males are at more risk of developing the disease

(Godiksen *et al.*, 2011;

Casamián-Sorrosal *et al.*, 2014)

Peritoneopericardial diaphragmatic hernia (PPDH)

- Congenital defect
- Breed at increased risk in a study of 31 feline cases, 2000–2007
- Maine Coons represented 4 of the 31 cases (12.9%), but only 2.2% of the hospital population

(Banz & Gottfried, 2010)

Dermatological conditions

Cutaneous food allergy

- Maine Coon cats were over-represented in a study of 48 cases

(Scott & Miller, 2013)

Gastrointestinal conditions

Intussusception

- Breed at increased risk in case series. Maine Coon cats represented 19/25 cases in a study, 2000–2012

(Vershoof *et al.*, 2015)

Trichomonas foetus infection

See under *Infectious conditions*

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

Infectious conditions

Tritrichomonas foetus infection

- Breed at increased risk in a study in Germany (Kuehner *et al.*, 2011)

Musculoskeletal conditions

Hip dysplasia

- In a retrospective US study of data recorded at the Orthopedic Foundation for Animals the prevalence of hip dysplasia was found to be 24.9% in Maine Coon cats
- There was a slightly higher prevalence in males (27.3%) than females (23.3%)
- 56% of cases were bilateral (Loder & Todhunter, 2017)

Peritoneopericardial diaphragmatic hernia (PPDH)

See under *Cardiovascular conditions*

Polydactyly

- Inherited in some lines of Maine Coons
- Variability in digit number and conformation
- Conformation of tarsus and carpus are also affected in some cases (Hamelin *et al.*, 2017)

Slipped capital femoral epiphysis (SCFE)

- In a retrospective study of cases presented to a first-opinion and referral practice in Austria, 2009–2015, 17/208 (8.17%) Maine Coon cats were diagnosed with SCFE, compared to 29/4348 (0.67%) cats of all breeds (Borak *et al.*, 2017)

Neoplastic conditions

Middle ear polyps

- In a study of 62 cases (2004–2014) Maine Coons were over-represented, with 23 cases (37%)
- Mean age (all breeds): 3.9 years (range 0.5–14 years) (Janssens *et al.*, 2017)

Physiological conditions

Blood group

- In a study of 357 blood donor cats in Italy, 21% were Maine Coon cats and all were type A (Spada *et al.*, 2014)

Haematological and biochemical differences

- In a retrospective study of 81 healthy Maine Coon cats presented as blood donors in Italy, 2011–2014, significant variations (from

published reference intervals in the general cat population) were found in a number of haematological and biochemical parameters. Breed-specific reference intervals have been created for Maine Coon cats

(Spada *et al.*, 2015)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Maine Coon cats produced larger kittens (mean 116.1 g) than the mean of all breeds (93.5 g)

(Sparkes *et al.*, 2006)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Maine Coon was 68 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined

(Hagman *et al.*, 2014)

MANX

Gastrointestinal conditions

Faecal incontinence (part of Manx syndrome)

See under *Musculoskeletal conditions*

Megacolon and constipation (part of Manx syndrome)

See under *Musculoskeletal conditions*

Rectal prolapse (part of Manx syndrome)

See under *Musculoskeletal conditions*

Musculoskeletal conditions

Manx syndrome

- Autosomal dominant inheritance
- The mutation causes a shortened or absent tail
- May also affect the spine and spinal cord, resulting in spina bifida; symptoms can include problems with the control of urination and defecation, rectal prolapse and paresis of the hindlegs
- Homozygosity for the mutation is lethal
- Variation in penetrance of the mutation results in variation of both tail length and severity of the spinal signs (Leipold *et al.*, 1974; Deforest & Basrur, 1979; Song *et al.*, 2016)

Neurological conditions

Spina bifida (part of Manx syndrome)

See under *Musculoskeletal conditions*

Renal and urinary conditions

Urinary incontinence (part of Manx syndrome)

See under *Musculoskeletal conditions*

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in case series
- Across all breeds, females reported to be more at risk than males, and neutered cats at greater risk than entire

(Cannon *et al.*, 2007)

NORWEGIAN FOREST CAT (NFC)

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM) and restrictive cardiomyopathy (RCM)

- Familial
- In a UK-based prospective study of 53 Norwegian Forest Cats screened for cardiomyopathy by echocardiogram, 13 were found to have signs of HCM
- As part of the same study, post-mortem results for 8 NFCs which died of cardiac-related problems were reviewed retrospectively. Pathological findings of both HCM and RCM were found in 7 cases

(März *et al.*, 2015)

Endocrine conditions

Diabetes mellitus

- Breed at increased risk in several case series (UK, Sweden)
- In a cohort of 193 563 cats in primary-care practice in England, 2009–2014 (VetCompass data), there were 1128 cases of diabetes mellitus (period prevalence 0.58%). Norwegian Forest Cats were at increased risk: OR 3.5 (95% CI 1.3–9.6) compared to the overall population. Both being >4kg body weight and >6 years of age were risk factors across all breeds ($p < 0.001$)
- Breed at risk in retrospective series based on Swedish insurance data 2009–2013. NFCs had an increased incidence rate ratio (IRR) of 1.9 (95% CI 1.5–2.3) compared to all other breeds
- Males at increased risk: IRR 2.0 (95% CI 1.8–2.3) across all breeds

- Mean age at diagnosis: 10.7 years across all breeds

(Öhlund *et al.*, 2015; O'Neill *et al.*, 2016b)

Gastrointestinal conditions

Tritrichomonas foetus infection

See under *Infectious conditions*

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

Infectious conditions

Tritrichomonas foetus infection

- Breed at increased risk in a study in Germany (Kuehner *et al.*, 2011)

Neurological conditions

Lysosomal storage disease – glycogen storage disease type IV

- Autosomal recessive inheritance
- Many affected kittens die perinatally (from hypoglycaemia). Survivors develop neuromuscular degeneration around 5 months of age
- In a US study screening 402 Norwegian Forest Cats, 58 were carriers and 4 were affected (Fyfe *et al.*, 2007)

Physiological conditions

Blood group

- In a US study, 100% of Norwegian Forest Cats were type A (Giger *et al.*, 1991a)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Norwegian Forest Cat was 39 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined (Hagman *et al.*, 2014)

OCICAT

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
 - Ocicats had an OR of 16.80 (95% CI 11.37–24.84)
 - Mean age (all breeds) 6.2 years
 - Males were at increased risk (all breeds): OR 1.1
 - In a separate study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014, Ocicats had an OR of 18.4 (95% CI 6.55–49.5)
- (Albasan *et al.*, 2012; Houston *et al.*, 2016)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
 - IR for the Ocicat was 105 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined
- (Hagman *et al.*, 2014)

ORIENTAL SHORTHAIR

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Orientals were found to have increased scores for compulsive (non-grooming) behaviours and cat aggression
- (Wilhelmy *et al.*, 2016)

Gastrointestinal conditions

Systemic amyloidosis

- Familial
 - At increased risk
 - Primarily involves the liver. May result in hepatic dysfunction or spontaneous hepatic haemorrhage
- (Piirsalu *et al.*, 1994; van der Linde-Sipman *et al.*, 1997)

Ocular conditions

Progressive retinal atrophy (PRA)

See under *Siamese*

Physiological conditions

Blood group

- In an Australian study, 100% of the Oriental/Siamese breed were type A
- (Malik *et al.*, 2005)

Gestation length

- In a UK questionnaire-based study of 1056 litters, Oriental Shorthair cats had longer gestation (mean 66.2 days) than the mean of all breeds (65.1 days)
- (Sparkes *et al.*, 2006)

Renal and urinary conditions

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in case series
 - In a study of 7456 uroliths submitted to the Minnesota Urolith Center, 1981–1997, Oriental Shorthair cats had an OR of 3.22 (1.21–10.18) compared to a control population of cats from the VMDB over the same period
 - Males reported to be at reduced risk: OR 0.7 across all breeds
 - Mean age 69 ± 38 months across all breeds
- (Lekcharoensuk *et al.*, 2000b)

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
 - Oriental cats had an OR of 3.81 (95% CI 1.68–8.63)
 - Mean age (all breeds): 6.2 years
 - Males were at increased risk (all breeds): OR 1.1
- (Albasan *et al.*, 2012)

Reproductive conditions

Dystocia

See under *Siamese*

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Siamese*

PERSIAN

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Persians were found to have lower scores for playfulness, predatory behaviour, prey interest and fear-related aggression (Wilhelmy *et al.*, 2016)

House-soiling

- Over-represented for soiling outside the litter box in a US retrospective series of 736 behaviour cases, 1991–2001 (Bamberger & Houpt, 2006)

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM)

- Breed at increased risk
- In a French retrospective study of 344 cats with HCM (2001–2011), there were 41 Persians
- Male Persians were significantly predisposed: 25/41 (61%)
- Median age was 7 years (0.5–19 years) across all breeds (Trehou-Sechi *et al.*, 2012)

Dermatological conditions

Dermatophytosis (ringworm)

- Breed at increased risk. In a French retrospective study Persians represented 75% of diagnosed cases but only 7.9% of the hospital population
- This breed may develop dermatophytic pseudomycetoma, characterized by one or more ulcerated and discharging nodules over the dorsal trunk or tail base (Scott & Paradis, 1990; Nuttall *et al.*, 2008)

Idiopathic facial dermatitis in Persians and Himalayans

- Breed at increased risk
- Unknown cause; possibly genetic basis (Bond *et al.*, 2000; Chung *et al.*, 2009)

Gastrointestinal conditions

Periodontal disease

- In a US study of 16 374 cats >5 years of age with a diagnosis of periodontal disease, Persians had an RR of 1.3 (95% CI 1.1–1.6) of

being represented in the periodontal disease group compared to being represented in a hospital population of cats of similar age but without periodontal disease

(Lund, 2012)

Polycystic liver disease

- Autosomal dominant inheritance, mutation identified
- Associated with polycystic kidney disease (see under *Renal and urinary conditions*)
- In one study of 27 cases of polycystic liver and/or kidney disease in cats, 21 were Persians or their crosses (Eaton *et al.*, 1997; Bosie *et al.*, 1998)

Haematological/immunological conditions

Chédiak–Higashi syndrome

- Seen in blue-smoke Persians with yellow eyes
- Autosomal recessive inheritance
- Multi-systemic condition

See also under *Ocular conditions*

(Kramer *et al.*, 1977)

Infectious conditions

Aspergillosis

- Persian and Himalayan cats seem to be over-represented in a number of case reports (Barrs & Talbot, 2014)

Dermatophytosis (ringworm)

See under *Dermatological conditions*

Neurological conditions

Intervertebral disc disease (IVDD)

- In a UK retrospective study of the medical records of cats diagnosed with thoracolumbar intervertebral disc disease, 2008–2014, Persian cats were found to be at significantly increased risk ($p < 0.0006$) compared to the general hospital population

(De Decker *et al.*, 2017)

Lysosomal storage disease – α -mannosidosis

- Inherited, mutation identified
- Signs, seen at 2–4 months, include cerebellar ataxia, corneal and lenticular opacities, skeletal abnormalities, hepatomegaly, thymic aplasia, gingival hyperplasia and polycystic kidneys
- Most cases are euthanized around 4–6 months (Berg *et al.*, 1997; Bradbury *et al.*, 2015)

Ocular conditions

Cataract

- Seen with Chédiak–Higashi syndrome
- Autosomal recessive inheritance
(Collier *et al.*, 1979)

Chédiak–Higashi syndrome

- Seen in blue-smoke Persians with yellow eyes
- Ocular signs include hypopigmentation of the fundus and iris, lack of a visible tapetum, nystagmus and cataracts

See also under *Haematological/immunological conditions*

(Collier *et al.*, 1979)

Corneal sequestrum

- In one retrospective review of 97 cases, 20 cats were Persians. In another (UK-based) study 31/64 cases were Persians
- Most seen at 2–7 years of age
- May be bilateral
(Featherstone & Sansom, 2004; Graham *et al.*, 2016)

Entropion

- Breed at increased risk of primary entropion
- Inheritance suspected
(Narfström, 1999b)

Epiphora

- Brachycephalia affects the course of the nasolacrimal duct and the efficiency of tear drainage. 23/31 Persians were classified as moderately or severely brachycephalic in one study, predisposing them to epiphora
(Schlueter *et al.*, 2009)

Lysosomal storage disease – α -mannosidosis

- Inherited, mutation identified
- Ocular symptoms include corneal and lenticular opacities

See also under *Neurological conditions*

(Berg *et al.*, 1997; Bradbury *et al.*, 2015)

Progressive retinal atrophy (PRA)

- Early-onset retinal degeneration
- Autosomal recessive inheritance, mutation identified
- Clinical signs at 2–3 weeks, blindness at 16 weeks
(Rah *et al.*, 2005; Alhaddad *et al.*, 2014)

Physiological conditions

Blood group

- In an Australian study, 67% of this breed were type A, 22% were type B and 11% were type AB
- In an English study, 88.2% of this breed were type A, 11.8% were type B and none were type AB
(Knottenbelt *et al.*, 1999; Malik *et al.*, 2005)

Litter size

- In a UK questionnaire-based study of 1056 litters, Persian cats produced smaller litters (mean 3.8 kittens) than the mean of all breeds (4.6 kittens)

(Sparkes *et al.*, 2006)

Renal and urinary conditions

Feline idiopathic cystitis

- Breed at increased risk in case series. In a prospective study of 55 cases, Persians represented 13% of affected cats but only 2% of the control population
- Neutered males at greater risk
- Mean age 4 years 11 months
(Gunn-Moore, 2003; Cameron *et al.*, 2004)

Polycystic kidney disease (PKD)

- Autosomal dominant inheritance
- It is estimated that 38% of Persian cats are affected worldwide
- In an Australian study of 230 Persian cats screened by renal ultrasound, 47% were found to be affected
- In a French study of 220 Persian cats screened by renal ultrasound, 41.8% were found to be affected
- Affected cases may present with renal failure from 2–3 years of age
- Some cases have liver cysts
(Barrs *et al.*, 2001; Barthez *et al.*, 2003; Young *et al.*, 2005)

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Persian cats had an OR of 2.21 (95% CI 1.82–2.69) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.73 (95% CI 1.63–1.82)

- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Persian cats had an OR of 3.30 compared to a control population of cats from the VMDB over the same period. Mean age 90 ± 41 months across all breeds

(Lekcharoensuk *et al.*, 2000b;
Houston *et al.*, 2016)

Reproductive conditions

Cryptorchidism

- Polygenic inheritance suspected
(Millis *et al.*, 1992)

Dystocia

- In a study based on Swedish insurance data, 1999–2006, incidence rates (IR) were 22 cases per 10 000 CYAR for all cats, and 38 cases per 10 000 CYAR for Persian cats. However, the Persian cat was found to be at reduced risk compared to all purebred cats (67 cases per 10 000 CYAR)
- In a separate Swedish retrospective study, 1986–1990, the Persian cat represented 37.4% of cases, by far the highest of all purebred cats
- In a study of data based on questionnaires completed by cat breeders, Persians were found to have high levels of dystocia
(Ekstrand & Linde-Forsburg, 1994;
Gunn-Moore & Thrusfield, 1995;
Holst *et al.*, 2017)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Persian group (which included Persians, Chinchillas, Colourpoint Persians, Exotic Shorthairs) was 34 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined
(Hagman *et al.*, 2014)

Stillborn kittens

- In a UK questionnaire-based study of 1056 litters, Persian cats had a significantly higher prevalence of stillbirths (10.8%) than that of all breeds combined (7.2%)
- Persians also have a higher number of kitten deaths in the first 8 weeks of life
(Sparkes *et al.*, 2006)

Respiratory conditions

Aspergillosis

- Persian and Himalayan cats seem to be over-represented in a number of case reports
(Barrs & Talbot, 2014)

Pulmonary carcinoma

- Breed at increased risk in case series
- In a USA-based retrospective study of 39 feline cases, Persians were represented at least 4 times more frequently than any other breed ($p < 0.0001$)
- Average age 12.3 years (range 6–18 years)
(D'Costa *et al.*, 2012)

RAGDOLL

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Ragdolls were found to have lower scores for attention-seeking, trainability and sleeping in elevated/warm/hidden locations
(Wilhelmy *et al.*, 2016)

Cardiovascular conditions

Aortic thromboembolism

- Breed at increased risk in a retrospective series of 127 cases in the USA: OR 14.40 ($p = 0.0016$) compared to the general hospital population
- Males at increased risk: OR 1.75 ($p = 0.003$)
(Smith *et al.*, 2003)

Hypertrophic cardiomyopathy (HCM)

- Inherited as an autosomal dominant trait, mutation identified
- In a UK-based study including 2018 Ragdoll cats, the prevalence of the mutation was 27%
- Prevalence has been described as 17% in Italy and 23% in the USA
- It is suspected that cats homozygous for the mutation have a shorter lifespan and are more likely to suffer cardiac death
- In a UK study of 127 cats with HCM, 1997–2005, Ragdolls with HCM were found to be younger and have a shorter survival time than other HCM cats
- Males predisposed
(Payne *et al.*, 2010; Borgeat *et al.*, 2014;
Casamián-Sorrosal *et al.*, 2014)

Dermatological conditions

Cutaneous neoplasia

See under *Neoplastic conditions*

Gastrointestinal conditions

Feline gastrointestinal eosinophilic sclerosing fibroplasia

- Breed at increased risk. In a retrospective study of 13 cases (from Australia and UK) Ragdolls (7/13 cases) and males (9/13 cases) were over-represented

(Linton *et al.*, 2015)

Infectious conditions

Cryptococcosis

- Breed at significantly increased risk ($p < 0.001$) in a retrospective series of 155 feline cases in Australia, 1981–2001
- Across all breeds, median age was 6 years (range 1–16) with a peak at 2–3 years

(O'Brien *et al.*, 2004)

Feline infectious peritonitis (FIP)

- Breed at increased risk in a retrospective series of 60 cases from the patient records of a North American veterinary medicine teaching hospital, 1986–2002: OR 52.22 (95% CI 11.14–244.79) compared to all cats
- Increased risk from 3 months to 3 years of age (all breeds)
- Males and sexually intact cats predisposed (all breeds)
- Cats in multi-cat environment at increased risk (all breeds)

(Foley & Pedersen, 1996;
Pesteanu-Somogyi *et al.*, 2006)

Neoplastic conditions

Mast cell tumour (MCT) – cutaneous

- Breed at increased risk
- In a UK study of 287 records of feline MCTs, Ragdoll cats represented 2.1% of the study population, compared to 0.7% of the control population
- Median age 11 years (range 5 months to 19 years)

(Melville *et al.*, 2015)

Physiological conditions

Blood group

- In an Italian study of 61 Ragdoll cats, 77.1% were type A, 4.9% type B, and 18% type AB
- (Proverbio *et al.*, 2013)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
 - In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–1997, Ragdoll cats had an OR 7.85 (95% CI 4.74–26.56) compared to a control population of cats from the VMDB over the same period
 - Males reported to be at increased risk: OR 1.5 across all breeds
 - Mean age 90 ± 41 months across all breeds
- (Lekcharoensuk *et al.*, 2000b)

Urolithiasis – struvite (magnesium ammonium phosphate)

- Breed at increased risk in case series
 - In a study of 7456 uroliths submitted to the Minnesota Urolith Center, 1981–97, Ragdoll cats had an OR 4.9 (95% CI 1.98–14.35) compared to a control population of cats from the VMDB over the same period
 - Males reported to be at reduced risk: OR 0.7 across all breeds
 - Mean age 69 ± 38 months across all breeds
- (Lekcharoensuk *et al.*, 2000b)

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- Ragdoll cats had an OR of 5.14 (95% CI 3.15–8.40)
- Mean age (all breeds): 6.2 years
- Males were at increased risk (all breeds): OR 1.1

(Albasan *et al.*, 2012)

Reproductive conditions

Dystocia

- Breed at increased risk in a study based on Swedish insurance data, 1999–2006
- Incidence rates (IR) were 22 cases per 10 000 CYAR for all cats, and 102 cases per 10 000 CYAR for Ragdoll cats
- Incidence rate ratio (IRR) in the Ragdoll cat was 1.5 compared to all other purebred cats

(Holst *et al.*, 2017)

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Ragdoll was 80 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined

(Hagman *et al.*, 2014)

RUSSIAN BLUE

Dermatological conditions

Cutaneous neoplasia

See under *Neoplastic conditions*

Endocrine conditions

Diabetes mellitus

- Breed at increased risk in a retrospective series based on Swedish insurance data 2009–2013. Russian Blues had an increased incidence rate ratio (IRR) of 3.8 (95% CI 2.1–6.4) compared to all other breeds
- Males at increased risk: IRR 2.0 (95% CI 1.8–2.3) across all breeds
- Mean age at diagnosis: 10.7 years across all breeds

(Öhlund *et al.*, 2015)

Neoplastic conditions

Mast cell tumour (MCT) – cutaneous

- Breed at increased risk
- In a UK study of 287 records of feline MCTs, Russian Blue cats represented 2.1% of the study population, compared to 0.1% of the control population
- Median age 11 years (range 5 months to 19 years)

(Melville *et al.*, 2015)

Renal and urinary conditions

Urolithiasis

- Breed at risk in case series
- Type of urolith not specified

(Lekcharoensuk *et al.*, 2000b)

SAVANNAH

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutation identified

- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Grahn *et al.*, 2012)

SCOTTISH FOLD

Musculoskeletal conditions

Osteochondrodysplasia

- Autosomal dominant inheritance with incomplete penetrance

(Takanosu *et al.*, 2008)

Physiological conditions

Blood group

- In a US study, 85% reported as type A, 15% as type B

(Giger *et al.*, 1991a)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in case series
- In a study of 7934 calcium oxalate uroliths submitted to the Minnesota Urolith Center, 1981–97, Scottish fold cats had an OR 5.21 (95% CI 3.53–10.52) compared to a control population of cats from the VMDB over the same period
- Males reported to be at increased risk: OR 1.5 across all breeds
- Mean age 90 ± 41 months across all breeds

(Lekcharoensuk *et al.*, 2000b)

SEYCHELLOIS

Reproductive conditions

Dystocia

See under *Siamese*

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Siamese*

SIAMESE

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires,

Siamese were found to have increased scores for stranger-directed aggression, but lower scores for sociability and fear-related aggression

(Wilhelmy *et al.*, 2016)

Wool-sucking (compulsive)

- Breed at increased risk
- The presence of a medical condition was found to be an increased risk factor for wool-sucking in Siamese cats in one study
- Affected cats had an abnormally intense appetite

(Bornes-Weil *et al.*, 2015)

Cardiovascular conditions

Endocardial fibroelastosis

- Inherited congenital anomaly
- Age of onset 3 weeks to 4 months
- Short illness in many cases, sudden death possible
- Cats with mild forms survive to adulthood

(Zook & Paasch, 1982)

Mitral stenosis

- Breed at increased risk in US retrospective case series
- Males over-represented across all breeds
- May present as an asymptomatic murmur, or as respiratory distress due to congestive heart failure and/or hindlimb paralysis due to aortic thromboembolism

(Campbell & Thomas, 2012)

Dermatological conditions

Blastomycosis

See under *Infectious conditions*

Cryptococcosis

See under *Infectious conditions*

Cutaneous neoplasia

See under *Neoplastic conditions*

Endocrine conditions

Diabetic ketoacidosis

- Breed at increased risk

(Cooper *et al.*, 2015)

Primary hyperparathyroidism

- Breed at increased risk in retrospective case series

- Siamese represented 5/7 cats with primary hyperparathyroidism
- 5 of the cats were female
- Ages ranged from 8 to 15 years, mean 12.9 years

(Richter *et al.*, 1990)

Gastrointestinal conditions

Intestinal neoplasia

- Breed at increased risk in case series
- In a retrospective study of VMDB records of 1129 cases of feline intestinal tumours, 1964–2004, Siamese cats had an OR of 1.79 compared to other breeds
- Cats of all breeds >7 years old were at significantly increased risk
- Siamese were found to be at increased risk of both intestinal adenocarcinoma and lymphoma
- In a separate study, 6/11 cases of intestinal adenocarcinoma were found to be in the Siamese. Males were at increased risk

(Cribb, 1988; Risetto *et al.*, 2011)

Megaoesophagus and pyloric dysfunction

- Breed at increased risk in case series. In a study of 13 cases of pyloric dysfunction, 12 cats were Siamese; 8 cases had concurrent oesophageal dilatation
- Inheritance suspected
- Vomiting started soon after weaning

(Pearson *et al.*, 1974)

Pancreatitis

- Breed at increased risk in a retrospective review of 40 cases, 1976–1989

(Hill & Van Winkle, 1993)

Periodontal disease

- In a US study of 16 374 cats >5 years of age with a diagnosis of periodontal disease, Siamese had an RR of 1.3 (95% CI 1.1–1.5) of being represented in the periodontal disease group compared to being represented in a hospital population of cats of similar age but without periodontal disease
- In a Japanese study of 323 cats, 7.1% were affected by gingivostomatitis, but among the Siamese the incidence was 40%

(Fujitsu & Sakai, 1999; Lund, 2012)

Pyloric dysfunction

See *Megaoesophagus and pyloric dysfunction*

Salivary gland neoplasia

- Siamese cats were over-represented in a study of 30 cats
- Most were adenocarcinomas (Hammer *et al.*, 2001)

Systemic amyloidosis

- Familial
- Primarily involves the liver. May result in hepatic dysfunction or spontaneous hepatic haemorrhage (van der Linde-Sipman *et al.*, 1997; Niewold *et al.*, 1999)

***Tritrichomonas foetus* infection**

See under *Infectious conditions*

Haematological/immunological conditions**Haemophilia B**

- Factor IX deficiency occurs in this breed (Littlewood, 1989)

Infectious conditions**Blastomycosis**

- In a literature review of cases, 1961–1988, 9/23 cases were in the Siamese cat
- 70% of cases across all breeds were male
- 75% of cases across all breeds were <4 years old (Miller *et al.*, 1990)

Cryptococcosis

- Breed at significantly increased risk ($p=0.013$) in a retrospective series of 155 feline cases in Australia, 1981–2001
- Across all breeds, the median age was 6 years (range 1–16) with a peak at 2–3 years (O'Brien *et al.*, 2004)

Mycobacterial infections

- In a UK study of 339 cases of feline mycobacterial infections, 39 cases were seen in pedigree cats, of which 18 were Siamese
- Cutaneous lesions were the most common presenting signs
- Siamese may also be predisposed to disseminated *Mycobacterium avium* complex infection (Baral *et al.*, 2006; Gunn-Moore *et al.*, 2011)

Sporotrichosis

- Breed at increased risk in a US case series
- Young males predisposed (Davies & Troy, 1996)

***Tritrichomonas foetus* infection**

- Breed at increased risk in a UK study
- Cats <1 year old predisposed (Gunn-Moore *et al.*, 2007)

Musculoskeletal conditions**Mucopolysaccharidosis VI**

- Autosomal recessive inheritance
- Causes dwarfism, facial dysmorphism, multiple skeletal, neurological and retinal deficits
- Two mutations identified: D520N (which causes a very mild phenotype) and L476P (which causes a more severe phenotype) (Crawley *et al.*, 2003; Lyons *et al.*, 2016b)

Slipped capital femoral epiphysis (SCFE)

- Siamese were over-represented (23% of cases) in a study of 13 cases
- Across all breeds, males were over-represented (85%)
- 90% of the male cases were overweight
- Age range 4.5–24 months across all breeds (Craig, 2001)

Neoplastic conditions**Mast cell tumour (MCT) – cutaneous**

- Breed at increased risk in some case series
- In one study the head was the most commonly affected site
- In a UK study of 287 records of feline MCTs, Siamese cats represented 5.2% of the study population, compared to 1.8% of the control population
- Median age 11 years (range 5 months to 19 years)
- Histiocytic mast cell tumours, which are multicentric and regress spontaneously, are found in young Siamese cats (<1 year old) (Chastain *et al.*, 1988; Miller *et al.*, 1991; Melville *et al.*, 2015)

Intestinal neoplasia

See under *Gastrointestinal conditions*

Lymphoma

- Breed at increased risk in several case series
- Males over-represented (3.2:1 in one study)

- Siamese cats are at increased risk of mediastinal lymphoma, tending to present at a younger age than other breeds and being feline leukaemia virus-negative (Louwerens *et al.*, 2005; Fabrizio *et al.*, 2014)

Mammary neoplasia

- Breed at increased risk
- Generally seen in older cats, but may occur at an earlier age in this breed
- Primarily a disease of female cats, but occasionally seen in males (Hayes *et al.*, 1981)

Salivary gland neoplasia

See under *Gastrointestinal conditions*

Neurological conditions

Feline hyperaesthesia syndrome

- Unknown aetiology
- May be due to a myopathy (Tuttle, 1980)

Lysosomal storage disease

- Autosomal recessive inheritance
- Types seen in the Siamese are GM₁ and GM₂ gangliosidosis, ceroid lipofuscinosis, mucopolysaccharidosis type VI, sphingomyelinosis (Niemann–Pick disease)

See also under *Musculoskeletal conditions* and *Ocular conditions*

(Skelly & Franklin, 2002)

Ocular conditions

Congenital glaucoma and microphakia

- Autosomal recessive inheritance
- Additionally, Siamese cats may be predisposed to primary open-angle glaucoma (McLellan *et al.*, 2004; McLellan & Teixeira, 2015)

Convergent strabismus and nystagmus

- Congenital; breed at increased risk (Johnson, 1991)

Lens luxation

- Breed at increased risk in series of 345 cases
- Most common age at presentation 7–9 years
- Males were over-represented (Olivera *et al.*, 1991)

Lysosomal storage disease

- Autosomal recessive inheritance

- Ocular signs include visual deficits and corneal opacities

See also under *Neurological conditions*

(Skelly & Franklin, 2002)

Progressive retinal atrophy (PRA)

- Late-onset rod–cone degeneration (PRA-rdAc), which is inherited recessively
- Siamese group (including, Oriental Shorthair, Balinese and Javanese) are at increased risk
- Clinical onset 1.5–2 years, progressing to blindness over 2–4 years
- Mutation identified (Menotti-Raymond *et al.*, 2010)

Physiological conditions

Blood group

- In Portuguese, English, Australian and American studies, 100% of purebred Siamese cats were type A (Giger *et al.*, 1991a; Silvestre-Ferreira *et al.*, 2004; Malik *et al.*, 2005; Forcada *et al.*, 2007)

Gestation length

- In a UK questionnaire-based study of 1056 litters, Siamese cats had longer gestation (mean 66.1 days) than the mean of all breeds (65.1 days) (Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Siamese cats produced larger litters (mean 4.9 kittens) than the mean of all breeds (4.6 kittens) (Sparkes *et al.*, 2006)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Siamese cats had an OR of 1.53 (95% CI 1.25–1.89) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.73 (95% CI 1.63–1.82) (Houston *et al.*, 2016)

Urolithiasis – urate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre 1998–2014

- Siamese cats had an OR of 4.52 (95% CI 3.37–5.96) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined: OR 1.3 (95% CI 1.16–1.54) (Houston *et al.*, 2016)

Reproductive conditions

Dystocia

- Breed at increased risk in a study based on Swedish insurance data, 1999–2006
- Incidence rates (IR) were 22 cases per 10 000 CYAR for all cats, and 135 cases per 10 000 CYAR for the Oriental group (including the Siamese, Balinese, Seychellois and Oriental Shorthair)
- Incidence rate ratio (IRR) in the Oriental group was 2.2 compared to all other pure-bred cats
- In a study of data based on questionnaires completed by cat breeders, Siamese-type were found to have high levels of dystocia (Gunn-Moore & Thrusfield, 1995; Holst *et al.*, 2017)

Mammary neoplasia

See under *Neoplastic conditions*

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Siamese group (which included Balinese, Foreign White, Javanese, Oriental Shorthair, Seychellois) was 84 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined (Hagman *et al.*, 2014)

Respiratory conditions

Feline asthma

- Breed at increased risk in some (but not all) case series
- Siamese represented 12/22 cases in a Greek study
- Siamese were also over-represented in a study of 65 cases of feline bronchial disease, 1980–1986 (Moise *et al.*, 1989; Adamama-Moraitou *et al.*, 2004; Trzil & Reinero, 2014)

Mediastinal lymphoma

See under *Neoplastic conditions*

SIBERIAN FOREST CAT

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM)

- Believed to be inherited in Siberian Forest Cats (Meurs *et al.*, 2009)

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms (Grahm *et al.*, 2012)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Siberian cat was 125 cases per 10 000 CYAR, compared to 17 cases for all breeds combined (Hagman *et al.*, 2014)

SINGAPURA

Haematological/immunological conditions

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms (Grahm *et al.*, 2012)

SNOWSHOE

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control

population of cats without urinary tract disease derived from the VMDB over the same period

- Snowshoe cats had an OR of 16.91 (95% CI 4.93–58.06)
- Mean age (all breeds): 6.2 years
- Males were at increased risk (all breeds): OR 1.1

(Albasan *et al.*, 2012)

SOMALI

Haematological/immunological conditions

Increased osmotic fragility of erythrocytes

- Results in haemolytic anaemia
- Inheritance suspected
- Affected cats have chronic intermittent severe anaemia and splenomegaly

(Kohn *et al.*, 2000; Tritschler *et al.*, 2016)

Pyruvate kinase deficiency

- Results in haemolytic anaemia
- Autosomal recessive inheritance, mutations identified
- Breed at increased risk in a study of DNA samples from 14 179 cats
- Carriers are asymptomatic
- Variable onset and severity of symptoms

(Kohn, 2000; Grahn *et al.*, 2012)

Musculoskeletal conditions

Acquired myasthenia gravis

See under *Neurological conditions*

Neurological conditions

Acquired myasthenia gravis

- Breed at increased risk in a retrospective study of 235 cases in the USA, 2001–2012: Somali cats had an OR of 11.60 compared to all other breeds
- Bimodal age of presentation, at 2–3 years and 9–10 years

(Hague *et al.*, 2015)

Ocular conditions

Progressive retinal atrophy (PRA)

- Late-onset rod–cone degeneration (PRA-rdAc), which is inherited recessively
- Clinical onset 1.5–2 years, progressing to blindness over 2–4 years
- Mutation identified

(Narfström, 1985; Bedford, 1989; Menotti-Raymond *et al.*, 2010)

Physiological conditions

Blood group

- In the USA, 78% reported as type A, 22% as type B

(Giger *et al.*, 1991a)

Litter size

- In a UK questionnaire-based study of 1056 litters, Somali cats produced smaller litters (mean 3.6 kittens) than the mean of all breeds (4.6 kittens)

(Sparkes *et al.*, 2006)

Renal and urinary conditions

Renal amyloidosis

- Breed at increased risk

(van Rossum *et al.*, 2004)

Reproductive conditions

Dystocia

See under *Abyssinian*

Pyometra (cystic endometrial hyperplasia–pyometra complex)

See under *Abyssinian*

SPHYNX

Cardiovascular conditions

Hypertrophic cardiomyopathy (HCM)

- Breed at increased risk. In a French prospective study of 114 healthy Sphynx cats examined by echocardiography, 2004–2011, 23 (20.2%) were found to have HCM
- Males and females were equally affected
- Autosomal dominant inheritance with incomplete penetrance proposed

(Chetboul *et al.*, 2012)

Mitral valve dysplasia

- Breed at increased risk. In a French prospective study of 114 healthy Sphynx cats examined by echocardiography, 2004–2011, 16 (14%) were found to have congenital heart disease. In most cases this was mitral valve dysplasia

(Chetboul *et al.*, 2012)

Dermatological conditions

Malassezia skin colonization

- Studies have demonstrated a high rate of *Malassezia* carriage in normal Sphynx cats

(Volk *et al.*, 2010)

Musculoskeletal conditions

Hereditary myopathy of Sphynx and Devon Rex cats

- Autosomal recessive inheritance
- Variably progressive
- Seen in Sphynx and the related Devon Rex
- May represent a congenital myasthenic condition

(Martin *et al.*, 2008; Gandolfi *et al.*, 2015)

Renal and urinary conditions

Urolithiasis – urate

- Breed at increased risk in a retrospective study of uroliths examined at the Minnesota Urolith Center, 1981–2008, compared to a control population of cats without urinary tract disease derived from the VMDB over the same period
- Sphynx cats had an OR of 12.88 (95% CI 4.52–36.75)
- Mean age (all breeds): 6.2 years
- Males were at increased risk (all breeds): OR 1.1 (Albasan *et al.*, 2012)

Reproductive conditions

Pyometra (cystic endometrial hyperplasia–pyometra complex)

- Breed at increased risk based on Swedish insurance data, 1999–2006
- IR for the Sphynx was 433 cases per 10 000 CYAR, compared to 17 cases per 10 000 CYAR for all breeds combined

(Hagman *et al.*, 2014)

TONKINESE

Behavioural conditions

General behavioural traits

- In a study of 574 single-breed registered cats, based on owner-completed questionnaires, Tonkinese were found to have increased scores for sociability with people, playfulness, vocalization, attention-seeking behaviour, separation anxiety and trainability, but lower scores for restraint resistance, owner-directed aggression, fear-related aggression and cat aggression

(Wilhelmy *et al.*, 2016)

Endocrine conditions

Diabetes mellitus

- Breed at increased risk in case series
- In a cohort of 193 563 cats in primary-care practice in England, 2009–2014 (VetCompass data), there were 1128 cases of diabetes mellitus (period prevalence 0.58%). Tonkinese were at

increased risk, with an OR of 4.1 (95% CI 1.8–9.6) compared to the overall population. Both being >4 kg body weight and >6 years of age were risk factors across all breeds ($p < 0.001$)

(O'Neill *et al.*, 2016b)

Physiological conditions

Blood group

- In a study in the USA, 100% of Tonkinese cats tested were type A

(Giger *et al.*, 1991a)

Kitten birth weight

- In a UK questionnaire-based study of 1056 litters, Tonkinese cats produced smaller kittens (mean 84.1 g) than the mean of all breeds (93.5 g) (Sparkes *et al.*, 2006)

Litter size

- In a UK questionnaire-based study of 1056 litters, Tonkinese cats produced larger litters (mean 5.3 kittens) than the mean of all breeds (4.6 kittens)

(Sparkes *et al.*, 2006)

Renal and urinary conditions

Urolithiasis – calcium oxalate

- Breed at increased risk in a retrospective study of 21 426 bladder uroliths submitted to the Canadian Veterinary Urolith Centre, 1998–2014
- Tonkinese cats had an OR 5.02 (95% CI 1.89–17.3) compared to domestic short-haired cats
- Males were found to be at increased risk in all breeds combined, OR 1.73 (95% CI 1.63–1.82)

(Houston *et al.*, 2016)

TURKISH ANGORA

Physiological conditions

Blood group

- In a study of 28 Turkish Angora cats in Turkey, 53.6% were type A and 46.4% were type B

(Arikan *et al.*, 2003)

TURKISH VAN

Physiological conditions

Blood group

- In a study of 85 Turkish Van cats in Turkey, 40% were type A and 60% were type B

(Arikan *et al.*, 2003)

PART III

DISEASE DESCRIPTIONS

Abnormal dentition

Dentition (teeth) may develop abnormally in several ways in dogs and cats. Possible abnormalities include persistent deciduous teeth, unerupted teeth, malformed or maldirected teeth, malocclusion, missing or supernumerary teeth (polyodontia). For example, missing incisors are seen in some Havanese dogs as part of a wider syndrome of multiple developmental abnormalities.

Abnormal heart sounds

These include murmurs, gallop rhythms and adventitious sounds.

Abnormal mediiodistal location of medial fabella

The lateral and medial fabellae are sesamoid bones in the tendon of origin of each head of the gastrocnemius muscle, associated with the stifle (knee) joint.

Accessory pathway arrhythmia

Accessory pathways are abnormal cardiac muscle fibres which can conduct the electrical impulse that causes the heart to contract, bypassing

the atrioventricular node which helps control heart rate. This can lead to arrhythmias such as Wolf–Parkinson–White syndrome.

Achilles tendon rupture

The Achilles tendon attaches to the calcaneus of the hock joint. Rupture will lead to lameness due to a loss of ability to extend the hock.

Acne

Also known as folliculitis, acne involves skin inflammation, often around the chin or face, with signs including redness, swelling, papules and pustules. It can be idiopathic or bacterial in origin.

Acquired megaesophagus

A motility disorder resulting in flaccidity of the oesophagus. Affected animals suffer from regurgitation and are at increased risk of aspiration pneumonia.

Acquired myasthenia gravis

An autoimmune disorder affecting the receptors of the neurotransmitter acetylcholine resulting in muscular weakness. Signs may include fatigue

on exercise and megaesophagus. Myasthenia gravis may be primary, or secondary to another condition such as thymoma.

Acral mutilation and analgesia

A neurological or skin condition which results in mutilation of the distal extremities. Pups are often affected from between 3 and 5 months of age. Temperature and pain sensation of the toes is lost. The prognosis is often poor. See also *Sensory neuropathy*.

Acrodermatitis

Inherited condition of Bull Terriers causing coat, skin and footpad lesions. There may be respiratory and gastrointestinal signs, and the prognosis is poor. A defect in zinc metabolism may be involved in the pathogenesis.

Acromegaly

Acromegaly results from overproduction of growth hormone by the pituitary gland. In the cat, this most commonly results from a tumour of the pituitary gland, while in the dog it results from excess progesterone either in older intact bitches or following exogenous progesterone administration. Signs of acromegaly include enlargement of the paws, mandible and internal organs. Some cases present with diabetes mellitus.

Actinic (solar) keratosis

An inflammatory skin condition caused by prolonged exposure to the sun. It can lead to skin cancer.

Acute canine polyradiculoneuritis

An autoimmune disease of the peripheral nerves leading to paralysis. It is sometimes known as *coonhound paralysis*, although this more correctly only relates to the condition when it is caused by toxicity from a racoon bite.

Acute febrile neutrophilic vasculitis

An immune-mediated skin condition of unknown cause. Leads to multifocal areas of skin discoloration and oedema.

Acute non-compressive nucleus pulposus extrusion (type III disc extrusion)

This occurs when a small piece of disc material breaks off and impacts the spinal cord causing bruising. This can lead to paresis or paralysis,

which may be reversible over time. This condition presents clinically in a similar way to a fibrocartilaginous embolism, although acute non-compressive nucleus pulposus extrusion is more commonly associated with spinal hyperaesthesia.

Adaptation to hypoxia

Certain breeds which have traditionally been kept and bred at high altitudes, such as Tibetan Mastiffs, show genetic adaptation to the low oxygen levels found at altitude.

Adenoma/adenocarcinoma

Adenomas are benign tumours of glandular tissue. Adenocarcinomas are malignant tumours of glandular tissue.

Adult-onset deafness

There can be various causes of deafness in later life, including infection, trauma, iatrogenic injury and degenerative processes.

Adult-onset hair loss

There are several causes of hair loss in adult dogs, such as hormonal disease, parasitism and self-trauma. Some cases are unexplained, but as they occur more commonly in certain breeds, there may be a genetic component.

Afghan myelopathy

A progressive disease of the white matter of the spinal cord. Signs include pelvic limb ataxia and paresis, progressing to thoracic limb involvement, tetraplegia and eventually death from respiratory paralysis.

Agenesis of the nares

This congenital condition can predispose to laryngeal collapse. Dyspnoea, mouth breathing and snoring are seen.

Aggression

Some dog breeds are noted in studies to be more prone to aggression than others. However, different types of aggression are recognized, including fear-related, aggression to owners, aggression to strangers, predatory aggression, protective aggression and aggression to other dogs. Aggression is a common reason for owners to seek euthanasia, and so is an important cause of mortality, as well as being a public health risk.

Alanine aminotransferase (ALT) levels

This enzyme is routinely tested in blood samples as a marker of liver damage.

Alaskan Malamute chondrodysplasia

This condition leads to short limbs with bowed front legs and laterally deviated paws. Haemolytic anaemia is often also seen.

Alexander disease

This rare neurodegenerative disorder causes neurological signs including ataxia, tremors and weakness, progressing over a few weeks to paralysis.

Alopecia X

Also known as adrenal hyperplasia-like syndrome, biopsy-responsive alopecia, castration-responsive alopecia, castration-responsive dermatosis, growth-hormone-responsive alopecia, pseudo-Cushing's syndrome. This is a poorly understood cosmetic condition affecting coat quality and colour in early adulthood. It is characterized by partial to complete alopecia that spares the head and forelimbs and the alopecic skin may become hyperpigmented. It is progressive and symmetrical but non-pruritic.

Amblyopia and quadriplegia

This is a lethal inherited condition of Irish Setters. Puppies are unable to walk, and progression to visual impairment, nystagmus and seizures occurs.

Anaesthetic-related complications

All general anaesthesia carries a degree of risk. Some breeds are more at risk of complications. This may be due to anatomical factors such as airway anatomy, or to individual tolerance of certain anaesthetics.

Anal furunculosis (perianal fistula)

Chronically infected and often deep tracts in the soft tissues around the anus. An immune-mediated basis is suspected.

Anal sac adenocarcinoma

Anal sac adenocarcinoma is a malignant tumour which is palpable as a discrete or infiltrative mass in the anal sac. These tumours are often associated with hypercalcaemia and metastasize early to the sublumbar lymph nodes, spleen and lung. They are rare in the cat.

Angiostrongylosis

Angiostrongylus vasorum, also known as lungworm or French heartworm, can cause respiratory disease, neurological disease, clotting disorders and hypercalcaemia. It has been increasing in geographical spread in the UK in recent years.

Antibiotic-responsive diarrhoea (small intestinal bacterial overgrowth)

This condition was previously known as small intestinal bacterial overgrowth. It is believed to result from abnormal interaction between the intestinal immune system and antigens derived from gastrointestinal bacteria.

Antral pyloric hypertrophy (pyloric stenosis, chronic hypertrophic pyloric gastropathy)

In this syndrome, gastric outlet obstruction is caused by hypertrophy of the pyloric muscle, hyperplasia of the antral mucosa or both, and results in persistent vomiting. Congenital hypertrophy of the pyloric muscle is seen in young Boxers and Boston Terriers and may be referred to as congenital pyloric stenosis.

Adult-onset antral pyloric hypertrophy syndrome is seen in older (> 6 years) small oriental canine breeds. In these cases, there may be antropyloric mucosal hyperplasia only or a combination of mucosal and muscular hypertrophy.

Aortic or cardiac mineralization

An incidental finding noted in some dogs during radiography or computed tomography.

Aortic stenosis – subaortic stenosis (SAS)

This condition accounts for up to a third of reported cases of congenital canine heart disease. It appears to be inherited, but not in a simple way, with more than one gene thought to be involved. The condition develops in the first 3–8 weeks of life, and may be first detected as a heart murmur at this age. If the condition is mild, clinical signs are minimal, but more severely affected animals may suffer from weakness, collapse and sudden death. Screening programmes allowing selective breeding may reduce the incidence of this defect.

Aortic thromboembolism

A blood clot that lodges in the caudal aorta, especially in cats but occasionally in dogs, causing signs of hindlimb paresis or paralysis, cold extremities and pain. It is often associated with

cardiac disease in cats, and many cases may present with concurrent heart failure.

Arachnoid cyst – spinal

Arachnoid cysts are a rare cause of focal spinal cord compression in young dogs. Neurological deficits depend on the site of the lesion.

Arrhythmogenic right ventricular cardiomyopathy (Boxer cardiomyopathy)

The condition is also seen in cats. Affected animals can present with syncope, weakness or congestive heart failure.

Arthrosis of cervical articular facet joints

Abnormalities of the facet joints in the neck resulting in severe pain.

Aspergillosis

Aspergillosis is an opportunist fungal infection caused by *Aspergillus* spp. In the dog, it generally presents as a nasal and frontal sinus infection, and dolichocephalic (long-nosed) breeds seem predisposed. Disseminated aspergillosis is a more serious and often fatal form presenting as a systemic disease with multiple organ involvement.

Aspiration pneumonia

Inhalation of foreign material leading to lung inflammation. This is a serious and often fatal disease.

Atherosclerosis

Thickening of arterial walls.

Atlantoaxial subluxation/instability

This is seen primarily in young dogs of toy breeds which present with neck pain and neurological deficits in all four limbs due to cervical spinal cord compression. A variety of congenital defects, including a lack of or hypoplasia of the dens and shortening of the axis, lead to instability of the atlantoaxial articulation. The condition may also be acquired in any breed as a result of fracture of the dens or damage to the ligamentous support.

Atopic dermatitis (atopy)

This is a common condition involving hypersensitivity reactions to environmental allergens, affecting around 10% of the canine population. It is thought to be inherited, but the exact mode of inheritance has not yet been determined. Signs

include pruritus, erythema and self-trauma with secondary bacterial infection.

Atresia ani

Congenital lack of a patent anus. Atresia ani is a rare disease (0.007% cases in a review of 1.6 million dogs from the VMDDB).

Atrial fibrillation (AF)

This arrhythmia is usually seen in conjunction with severe heart disease such as dilated cardiomyopathy. However, some large-breed dogs can show an asymptomatic atrial fibrillation in the absence of structural heart disease.

Atrial septal defect

This uncommon congenital condition involves a defect in the septum between the atria. Small defects may be asymptomatic, but larger lesions can lead to congestive heart failure.

Atrioventricular (AV) block

Arrhythmia due to a disruption of the conduction of the electrical impulse from the pacemaker in the atrium to the ventricles.

Atypical granulation of neutrophils

Fine eosinophilic granules are noted in the neutrophils. This does not appear to affect the function of the cells.

Audiogenic reflex seizures

Reflex seizures are fits consistently triggered by environmental factors. Audiogenic reflex seizures are triggered by high-pitched sounds.

Avascular necrosis of the femoral head (Legg–Calvé–Perthes disease)

This disorder of small-breed dogs involves an aseptic, avascular necrosis of the femoral head. Signs are usually seen from 5 months of age. Ischaemia of the femoral head leads to degeneration of the bone which presents as a progressive uni- or bilateral hindlimb lameness.

Avulsion of the tibial tuberosity

This is a fracture of the growth plate of the tibial tuberosity in young dogs, which is seen more commonly in dogs with large quadriceps muscles.

Azotaemia

Higher than normal levels of urea, creatinine and other nitrogen-containing compounds in the blood.

Babesiosis

A tickborne protozoal disease caused by *Babesia* species. Infection results primarily in haemolytic anaemia; however, complications in multiple organs may develop. Infection may be peracute, acute, chronic or subclinical. It has been proposed that the disease may also be transmitted between dogs by biting.

Bacterial cholecystitis and bactibilia

Bacterial infection of the gallbladder and biliary tree.

Bacterial folliculitis/furunculosis

Inflammation of the skin associated with bacteria.

Bacterial keratitis

Bacterial infection and resulting inflammation of the cornea.

Basal cell tumour

These common skin tumours arise from the basal epithelial cells which give rise to the epidermis. They are usually well circumscribed, firm, freely mobile masses found in the dermis and subcutis around the head and neck. They are generally slow growing and benign in behaviour, rarely metastasizing. Basal cell carcinoma has low-grade malignancy and is a more invasive form that is reasonably common in cats but uncommon in dogs and rarely metastasizes in either species. Some dermatopathologists now consider most basal cell tumours to be more accurately classified as trichoblastomas.

Benign familial hyperphosphatasemia

A hereditary condition causing elevations in serum alkaline phosphatase.

Black hair follicular dysplasia

A rare disorder causing alopecia in areas of black hair. Thought to be inherited as an autosomal recessive trait.

Blastomycosis

Blastomycosis is a systemic fungal infection, caused by *Blastomyces dermatitidis* and seen primarily in North America. The lungs are the site of initial infection, and from there it may spread to the lymphatics, skin, eyes and bones. Dogs are more commonly infected than cats, young, male, large sporting-breed dogs living close to water being at greatest risk.

Blood group

Canine blood groups are classified according to the dog erythrocyte antigen (DEA). There are six main groups, DEA 1.1, 1.2, 3, 4, 5 and 7. DEA 1.1 and 1.2 are alleles so it is not possible to be positive for both. Since there are no naturally occurring autoantibodies to these two groups, an initial blood transfusion does not usually cause an acute transfusion reaction. However, if the donor and recipient are incompatible, an immune response will be mounted after the first transfusion, so subsequent incompatible transfusions are likely to cause a reaction. DEA 1.1 is inherited as an autosomal dominant. A DEA 1.1-negative female bred with a DEA 1.1-positive male may therefore have DEA 1.1-positive pups, and this can lead to neonatal isoerythrolysis.

Feline blood groups are classified into A, B and AB. A is dominant over B, but the inheritance of the rare AB blood group is determined by a different gene. There are high levels of naturally occurring antibodies against the other groups, so there are no universal donors, and crossmatching or typing is vital before transfusing.

Bone marrow necrosis

Necrosis of the bone marrow can be associated with toxins, neoplasia and infections. This leads to cytopenias such as anaemia, thrombocytopenia and neutropenia.

Bone tumour

Common bone tumours include osteosarcomas and chondrosarcomas. Signs of bone tumours include pain, swelling and pathological fractures.

Border Collie collapse

This condition occurs after strenuous exercise and can lead to ataxia, altered mentation and collapse. It is thought to be an episodic diffuse central nervous system disorder.

Borreliosis (Lyme disease)

This is caused by the tick-borne organism *Borrelia burgdorferi*, and causes signs including pyrexia and polyarthritis.

Brachycephalic obstructive airway syndrome (BOAS)

This term is used to describe a group of anatomical deformities which lead to respiratory compromise in brachycephalic breeds. An intronic, transposable element within the *SMOC2* gene is

associated with disruption of facial skeleton development in a dose-dependent manner. The deformities include stenotic nares, laryngeal deformities and hypoplastic trachea. The clinical signs are of upper airway obstruction and secondary complications; concurrent conditions such as laryngeal oedema and bronchopneumonia can occur in severely affected dogs.

Bradyarrhythmia

An abnormally slow heart rate, often associated with an irregular rhythm.

Bronchiectasis

Dilatation of the bronchi, often occurring as a complication of chronic bronchitis or bronchopneumonia. The changes are irreversible once present.

Calcinosis circumscripta

An uncommon condition, usually of unknown cause, although it may be associated with hyperadrenocorticism, or with tissue damage from mechanical, chemical, infectious or other factors. Calcinosis circumscripta involves localized deposition of mineral, causing a tumour-like nodule.

Calcinosis cutis

Calcinosis cutis involves localized or widespread mineralized deposits in the skin.

Calcium phosphate deposition

Calcium phosphate deposition can cause paraplegia in very young puppies, resulting from narrowing of the cervical vertebral canal. Soft-tissue deposition of mineral also occurs.

Cane toad toxicity

Exposure to the cane toad (*Bufo marinus*) commonly leads to hypersalivation and erythema of the mucous membranes. Seizures and arrhythmias are also seen.

Canine acanthomatous ameloblastoma (acanthomatous epulis)

An oral tumour which, while technically benign, can be locally invasive.

Canine autoimmune inflammatory disease (AID)

This disease involves widespread systemic inflammation, leading to signs including fever, arthritis, dermatitis, otitis and amyloidosis.

Canine cutaneous histiocytoma

These skin tumours are commonly seen in young dogs (< 4 years old), and appear as solitary, firm, well-circumscribed intradermal nodules on the head, limbs or trunk. Occasionally the surface will ulcerate. They are benign, and most will regress spontaneously over a period of months. See also *Histiocytic diseases*.

Canine distemper

This disease is caused by the canine distemper virus, and causes gastrointestinal signs, respiratory signs and footpad hyperkeratosis. Neurological signs can also occur. The disease can be fatal but is reliably prevented by vaccination.

Canine ectodermal dysplasia (hairlessness)

A rare genetic fragile skin condition seen in puppies.

Canine epileptoid cramping syndrome (Spike's disease)

This is a paroxysmal dyskinesia or movement disorder which can resemble epilepsy. Most cases begin before the age of 3 years, with signs of tremor, dystonia and difficulty walking lasting from 2 to 30 minutes. Some cases are associated with gastrointestinal signs, and in some cases diet changes may improve the severity of signs.

Canine follicular dysplasia (seasonal flank alopecia)

This is a localized cyclic follicular dysplasia causing a symmetrical alopecia. Hair loss and hyperpigmentation are noted especially on the flanks. Some dogs lose their hair in spring and regrow it spontaneously in autumn, while for others the reverse occurs.

Canine juvenile cellulitis

This condition is also known as juvenile pyoderma, puppy strangles or juvenile sterile granulomatous dermatitis and lymphadenitis. It is an uncommon disorder causing pustules on the face and pinnae of puppies. The submandibular lymph nodes are often greatly enlarged.

Canine leproid granuloma

A mycobacterial disease causing nodules and papules, usually around the head, which may ulcerate.

Canine leucocyte adhesion deficiency (CLAD)

An inherited defect in the ability of neutrophils to bind to endothelial cells causes severe and recurrent bacterial infections. A molecular diagnostic test is available to diagnose this condition.

Canine multiple system degeneration (progressive neuronal abiotrophy)

A hereditary neurodegenerative disease causing degeneration of the cerebellum, caudate nucleus and substantia nigra. Signs start from 3 months of age and start with classic cerebellar signs, but progress to difficulty initiating movement and balance problems.

Canine odontogenic parakeratinized cyst

A number of subtypes have been recognized, including radicular (periapical) cysts, dentigerous (follicular) cysts and lateral periodontal cysts. Odontogenic cysts are pathological, epithelial-lined cavities containing fluid or semi-solid material that are derived from odontogenic epithelium and can be developmental or the result of an inflammatory process. Odontogenic keratocysts are developmental cysts with an unusual propensity for recurrence.

Canine pigmented epidermal naevus

Some cases are considered to be an inherited disorder similar to that in humans, and some may be associated with canine papillomavirus infection.

Canine spongiform leucoencephalomyelopathy

A progressive neurological disease causing signs from 2–9 weeks of progressive neurological dysfunction including ataxia, paresis, paralysis and spasticity.

Cardiac hypertrophy

Thickening of the heart muscles can be present as a normal variation in athletic breeds, as a response to obstruction of outflow such as aortic stenosis, or as a primary myocardial disorder such as hypertrophic cardiomyopathy.

Cardiac tumour

Various tumours can affect the heart, including haemangiosarcomas, which are often found attached to the right atrium, and chemodectomas, which are often attached to the aorta.

Cardiomyopathy – X-linked muscular dystrophy (Golden Retriever muscular dystrophy)

In this rare inherited condition, skeletal muscle signs predominate, starting at about 8 weeks of age, but severe cardiac involvement may develop later. Few dogs affected with this condition survive past 5 years of age.

Carpal laxity syndrome

This condition involves either carpal (wrist) hyperextension or carpal hyperflexion. It occurs in puppies and is usually self-limiting.

Cataract

An opacity which may affect all or part of the lens or lens capsule, unilaterally or bilaterally. Cataracts may be primary (where a hereditary basis is suspected) or secondary, for example, to ocular inflammation, metabolic disease or congenital anomalies such as persistent pupillary membranes or persistent hyaloid artery.

Cataracts may be detected first in a variety of different areas of the lens and may progress at different rates. A complete cataract involves the whole lens and obscures the fundus, resulting in blindness in the affected eye.

Catechol-O-methyltransferase gene polymorphism

Alterations in this gene are associated with psychological problems in humans, and may be linked to behavioural abnormalities, including in territorial defence behaviours, in animals.

Central axonopathy

Several breeds have had central axonopathies described. These often have a very early age of onset, are progressive, and show signs of ataxia and tremor.

Central tarsal bone fracture

Fracture of one of the bones of the hock.

Central vestibular disease

Vestibular disease leads to signs of head tilt, nystagmus and ataxia. Central vestibular disease results from lesions in the brainstem and/or cerebellum.

Cerebellar ataxia

Ataxia related to the cerebellum is usually characterized by hypermetria, tremor, incoordination, but retention of conscious proprioception.

Cerebellar degeneration

Also known as cerebellar cortical abiotrophy, cerebellar cortical degeneration, hereditary cerebellar degeneration. Cerebellar cells can undergo premature ageing, degeneration and death (termed abiotrophy), leading to signs of cerebellar dysfunction (intention tremor, ataxia, hypermetria and menace deficits). In most cases the condition is believed to be hereditary.

Cerebellar hypoplasia

Congenital malformations of the cerebellum include hypoplasia and aplasia of the whole or part of the cerebellum. Some may have a genetic basis, while others result from a teratogen. Clinical signs are seen as soon as the animal becomes mobile and are non-progressive. They include hypermetria, head tremor and a wide-based stance. There is no treatment, but animals may make suitable pets if not severely affected.

Cerebellar Purkinje cell degeneration and coat colour dilution

Signs include growth retardation, progressive ataxia and dilution of the coat colour.

Cervical spondylomyelopathy (cervical vertebral malformation, wobbler syndrome)

A collection of abnormalities of the cervical vertebrae, including spinal canal stenosis, hypertrophy of the ligamentum flavosum and dorsal annulus, and disc herniation. Causes progressive signs of ataxia, starting in the hindlimbs and progressing to the forelimbs.

Chédiak-Higashi syndrome

A condition seen in the Persian cat. Affected cats have abnormal neutrophils and/or neutropenia, and may have increased susceptibility to disease. Platelet storage pool deficiency results in increased bleeding tendencies with normal coagulation profiles. Signs may include ocular and cutaneous albinism, cataracts, susceptibility to infection and bleeding tendencies.

Chemodectomas (aortic body and carotid body)

Chemodectomas are derived from the chemoreceptor cells of the aortic and carotid bodies which detect changes in the blood pH, oxygen and carbon dioxide levels. Chemodectoma of the aortic body arises at the heart base and is

reported more frequently than carotid body tumours, which arise at the bifurcation of the carotid artery and present as a cervical mass. Both are relatively uncommon in the dog and cat, but brachycephalic dogs appear predisposed. Chemodectomas may be locally invasive and have the potential to metastasize.

Cheyletiellosis

A non-suppurative dermatitis caused by *Cheyletiella* spp. mites.

Chiari malformation/syringomyelia

This condition involves a fluid-filled dilatation in the cervical spinal cord, which may lead to clinical signs of neck pain, persistent scratching around the head and neck, and sometimes ataxia. It has become increasingly recognized because of the increased availability of MRI, which is necessary for the diagnosis. Syringomyelia is often associated with a malformation of the occipital bone of the skull, which is known as Chiari malformation, or caudal occipital malformation syndrome.

Chlamydophilosis

This zoonotic disease caused by *Chlamydia psittaci* is more commonly seen in birds.

Chondrodysplasia (short-limbed or disproportional dwarfism)

Disproportionate growth in which the legs are shorter than expected for the head and body. This is considered normal in some breeds (e.g. Dachshunds), and abnormal in others.

Chondrosarcoma (skeletal)

This is the second most common primary bone tumour of dogs, accounting for 5–10% of cases. It generally grows slowly and metastasizes less frequently than the osteosarcomas.

Chronic atrophic gastritis

Chronic inflammation of the stomach leading to loss of glandular tissue.

Chronic hepatitis

Inflammatory liver disease which usually progresses to cirrhosis. There are many types of hepatitis, and classification remains controversial; however, certain breeds of dog may be predisposed, and individual breeds may demonstrate particular patterns of inflammation.

Chronic hepatitis (copper-associated)

Copper storage hepatopathy in the Bedlington Terrier results from a defect in biliary copper excretion leading to accumulation of copper in hepatocytes and hepatic necrosis. Copper-associated hepatopathy occurs in other breeds, but other mechanisms may be involved in the copper accumulation.

Chronic hypertrophic pyloric gastropathy

See *Antral pyloric hypertrophy*.

Chronic kidney disease (chronic renal failure, chronic renal insufficiency)

Many cases are idiopathic, but the disease may also be secondary to conditions such as amyloidosis, glomerulonephritis, neoplasia, renal dysplasia and pyelonephritis. The International Renal Interest Society (IRIS) grades severity on a scale of 1 to 4 based on serum creatinine, with subcategories based on proteinuria and hypertension. Chronic kidney disease causes signs such as polyuria–polydipsia, weight loss, inappetence, vomiting and anaemia. See also *Familial renal disease* and *Kidney disease*.

Chronic pulmonary disease (CPD)

A chronic respiratory illness, typified by inspiratory crackles and an increased interstitial pattern on thoracic radiography that is particularly prevalent in West Highland White Terriers. It has been suggested that this condition may be analogous to idiopathic pulmonary fibrosis (IPF) in humans.

Chronic superficial keratitis (pannus)

A bilateral progressive inflammatory disease of the cornea. A fleshy, vascular lesion spreads towards the central cornea from the temporal limbus. Corneal pigmentation follows and, if severe, vision loss occurs. It is suspected to have an immune-mediated basis and is influenced by ultraviolet radiation. The condition is more severe in dogs living at high altitude. May be seen with plasma cell infiltration of the nictitating membrane (plasmoma).

Chylothorax

An accumulation of chyle in the pleural space. This is often idiopathic, although it can be secondary to conditions such as heart failure and neoplasia. Signs are attributable to the pleural effusion, i.e. respiratory signs.

Cleft lip and/or palate

A congenital defect in the hard or soft palate allowing abnormal communication between the oral cavity and the nasal cavity/nasopharynx. Signs including failure to thrive and reflux from the nose during suckling are present from an early age.

Colitis

Inflammation of the lower intestine causing signs such as watery diarrhoea, fresh blood in the faeces and tenesmus.

Collie eye anomaly

A bilateral congenital condition characterized by abnormal development of the eye. The severity and effect on vision is variable. Mild cases may have only choroidal hypoplasia (inadequate development of the choroid). More severe cases may also have optic nerve colobomas, retinal detachment and intraocular haemorrhage. The condition is best diagnosed at 6–7 weeks, and it is advisable to remove affected dogs from a breeding programme.

Colorectal polyps

Polyps are more commonly found in the rectum than in the colon. They may be single or multiple, and while often benign, may have malignant characteristics.

Colour dilution alopecia

Hypotrichosis and recurrent bacterial folliculitis occur in colour dilute areas.

Cone degeneration (hemeralopia or day blindness)

Day blindness with no ophthalmoscopically visible abnormality. The condition results from selective degeneration of the cone photoreceptors of the retina. Dogs show severe loss of vision in daylight from 8–12 weeks but are able to see in dim light.

Congenital achromatopsia (rod monochromacy, day blindness)

Achromatopsia refers to lack of colour vision, which can be detected from 8 weeks using electroretinography.

Congenital alacrima (congenital keratoconjunctivitis sicca)

A lack of lacrimal tissue and tears causing dry eye.

Congenital deafness

This has been observed in numerous breeds (especially Dalmatians and blue-eyed white cats) and usually results from a partial or complete failure of development of the organ of Corti.

Congenital glaucoma and microphakia

High intraocular pressure with a small lens.

Congenital heart disease

A number of heart conditions can be caused by developmental defects, including septal defects, stenosis of major vessels, valve malformations and patent ductus arteriosus.

Congenital hypothyroidism

Can lead to disproportionate dwarfism, mental dullness and coat changes.

Congenital hypotrichosis

Animals affected with this condition are born without normal coats or lose their coats in the first few months of life.

Congenital megaesophagus

A motility disorder which when congenital shows signs of regurgitation around the time of weaning.

Congenital myasthenia gravis

Clinical signs in dogs include muscle weakness on exercise which improves with rest, and megaesophagus. The onset may be chronic or acute and the condition can be generalized or focal. Signs in cats include drooling, ventroflexion of the neck, regurgitation, weakness and lameness.

Congenital portosystemic shunt

Failure of fetal venous shunts to close after birth leads to persistent shunting of blood from the gastrointestinal tract to the systemic circulation without hepatic metabolism. Shunts may be single or multiple, intrahepatic or extrahepatic. Large breeds of dog are more likely to have intrahepatic shunts, small breeds and cats are more likely to have extrahepatic shunts.

Congenital umbilical hernia

A defect in the body wall at the umbilicus.

Conjunctival haemangiosarcoma

Malignant tumour of the conjunctiva.

Conjunctivitis

Inflammation of the conjunctiva.

Convergent strabismus and nystagmus

Results from abnormal visual pathway development associated with an absence of ocular pigment. Convergent strabismus is an abnormal deviation of the eyeballs medially. Nystagmus is a repetitive drift or flick of the eye. Other symptoms include decreased visual activity and loss of binocular vision.

Corneal and anterior segment foreign body trauma

Damage to the cornea from foreign bodies leading to penetration can cause glaucoma and blindness.

Corneal dystrophy

A primary, non-inflammatory bilateral (though not necessarily symmetrical) opacity of the cornea. The term *dystrophy* implies a hereditary condition. However, in many cases of corneal dystrophy, firm evidence of inheritance is lacking, although no underlying disease can be found. Different layers of the cornea may be affected, giving *epithelial*, *endothelial* and *stromal* dystrophies. The appearance, age of onset and rate of progression vary with the breed. Visual disturbance may occur if the lesion becomes extensive.

Corneal pigmentation (pigmentary keratopathy)

Occurs secondary to local irritation, e.g. from entropion or distichiasis.

Corneal sequestrum

A disease of the cornea seen in the cat, characterized by the development of a pigmented lesion in the centre of the cornea. In some cases, it is secondary to feline herpes virus keratitis, or chronic corneal irritation from entropion or trichiasis.

Corneal ulceration (ulcerative keratitis)

May occur secondary to dry eye, foreign bodies, chronic irritation such as entropion, or infection.

Corns (on the footpad)

Well-circumscribed regions of hyperkeratosis on the footpad.

Cranial cruciate ligament (CCL) disease

This common injury often presents as a severe, acute onset lameness. Diagnosis is by ascertaining the presence of a 'cranial drawer' motion in the stifle, by radiography, and by arthroscopy or arthrotomy.

Craniomandibular osteopathy (lion jaw)

This condition is inherited in some breeds, but other factors may be important in other breeds. Clinical signs include mandibular swelling, drooling and pain on opening the mouth. Irregular bone proliferation is seen on radiographs. The disorder is usually self-limiting, although in some cases hemimandibulectomy may be necessary.

Cricopharyngeal dysfunction

Difficulty in swallowing due to failure of the cricopharyngeal sphincter (the upper oesophageal sphincter) to relax.

Cryptococcosis

Cryptococcosis is a systemic fungal infection, caused by *Cryptococcus neoformans*, which is found worldwide and may be spread by pigeons. It infects a wide range of mammalian species but is most commonly seen in the cat. Clinical signs may reflect nasal, respiratory, central nervous system, ocular or cutaneous involvement.

Cryptorchidism

A failure of one or both testes to descend into the scrotum. The undescended testis may be found in the inguinal canal or abdomen. Cryptorchidism may be unilateral (in which case the right testis is more commonly affected in the dog) or bilateral. Smaller breeds of dog seem to have a higher risk. The risk of testicular neoplasia is higher in retained testes.

Cutaneous and renal glomerular vasculopathy (Alabama rot)

A disease of unknown cause, leading to skin ulceration and acute kidney injury. This was originally a disease of Greyhounds seen mainly in the USA; more recently a similar disease has been recognized in the UK, but without obvious breed predispositions.

Cutaneous asthenia (Ehlers–Danlos-like syndrome)

Causes skin hyperextensibility and alopecia.

Cutaneous food allergy

Allergic reaction to allergens within food causing skin signs.

Cutaneous haemangioma

Haemangiomas are benign tumours arising from the vascular endothelial cells of the dermis and subcutis. They are common in dogs but rare in cats. They appear as well-circumscribed blue/purple masses. It has been suggested that prolonged exposure to sunlight may be a predisposing factor, making light-skinned dogs with short coats more vulnerable.

Cutaneous haemangiosarcoma

Malignant tumours arising from the vascular endothelial cells of the dermis and subcutis. May be indistinguishable from haemangiomas or may be infiltrative with large areas of haemorrhage.

Cutaneous lupus erythematosus

Dogs typically develop clinical signs before 10 months of age, but age of onset has been reported as late as 2.75 years. Clinical signs consist of excessive scaling and crusting that first occur on the face, ears and back and then progress to a generalized form.

Cutaneous melanoma

Malignant skin tumour involving melanocytes.

Cutaneous mucinosis

Excessive mucin in the skin and subcutis, leading to signs of skin thickening and folding, with vesicles which may ooze fluid when squeezed.

Cutaneous soft-tissue sarcoma

A malignant skin tumour.

Dancing Dobermann disease

This is believed to be a neuromuscular disease of the gastrocnemius muscle; the underlying cause is not known. It has only been reported in Dobermann Pinschers. Affected dogs initially flex one pelvic limb while standing. As progression occurs to involve the other pelvic limb, the dog is seen to alternately flex and extend each pelvic limb in a dancing motion.

Deafness and vestibular disease

Hearing and balance are sensed in the ear and communicated to the brain via cranial nerve VIII. Lesions in any of these areas can lead to simultaneous deafness and balance disorders.

Deficiency of third component of complement

An immunodeficiency syndrome related to deficiencies in the complement system.

Degenerative encephalopathy

There are several neurodegenerative diseases reported in animals, including one affecting Nova Scotia Duck Tolling Retrievers which is associated with marked movements during sleep, anxiety and noise phobia, and gait abnormalities. It is seen from a young age and is progressive.

Degenerative lumbosacral stenosis

A disease of the lumbosacral space caused by conditions such as disc protrusion, instability of the vertebrae and foraminal stenosis. Causes signs of pain and lameness.

Degenerative myelopathy

A degenerative disease primarily seen in older dogs. Diffuse degeneration of the white matter of the thoracolumbar spinal cord results in progressive pelvic limb ataxia, paresis and loss of conscious proprioception. A DNA mutation that is a major risk factor for this condition has been identified.

Delivery by caesarean section

Some breeds have an increased frequency of delivery by the caesarean section. This may be due to dystocia, or may be elective.

Demodicosis

Generalized demodicosis is a severe skin disease caused by *Demodex* spp. mite. It can lead to pyoderma and deep folliculitis.

Dentigerous cyst

A cyst of the oral epithelium. May be incidental, or may be associated with gingival swelling, pain and anorexia.

Dermal arteritis of the nasal philtrum

Linear, well-circumscribed ulcers of nasal philtrum caused by arteritis. Mild arterial bleeding can result.

Dermatophyte pseudomycetoma

Subcutaneous fungal infections that cause nodules, ulcers and draining tracts.

Dermatophytosis (ringworm)

This fungal skin condition is particularly prevalent in long-haired cats.

Dermoid sinus

A dermoid sinus is a developmental defect arising from the incomplete separation of the skin and neural tube. It may be found midline in the cervical, cranial thoracic or sacrococcygeal regions. In cases where the sinus communicates with the dura mater, neurological signs may be seen. The condition is most commonly found in the Rhodesian Ridgeback, and it is believed to be hereditary in this breed.

Diabetes mellitus

Diabetes mellitus occurs where there is hyperglycaemia resulting from an absolute or relative lack of insulin. Where the blood glucose level exceeds the renal threshold, glycosuria results.

Diabetic ketoacidosis

A serious metabolic state caused by uncontrolled diabetes mellitus, in which ketones build up in the blood, and acidosis and usually hyperkalaemia are present. Older animals are predisposed.

Diffuse idiopathic skeletal hyperostosis (DISH)

Calcification and ossification of the ligaments. Can lead to spinal hyperaesthesia and stiffness, due to compression of the nerve roots.

Dilated cardiomyopathy (DCM)

This condition involves a dilatation of the heart causing larger chamber size, thinner heart walls and reduced strength of the heartbeat. It is thought that the majority of cases are genetic or familial, but it is not certain that all cases are genetic in origin. Nutritional abnormalities may also contribute, and it is possible that viral and immune-mediated causes may be involved in some cases.

Dirofilariasis (heartworm)

Infection by *Dirofilaria immitis* leading to a number of clinical signs, from asymptomatic infection to weight loss, dyspnoea and heart murmur. Some dogs develop pulmonary hypertension or pulmonary thromboembolism.

Discoid lupus erythematosus

Uncommon in dogs and very rare in cats. This condition is an immune-mediated dermatosis with no systemic involvement. Clinical signs include depigmentation, scaling, erosion and ulceration. The nose is primarily affected, but the pinnae, limbs and genitalia can be involved.

Discospondylitis

Infection of the intervertebral disc with osteomyelitis of adjoining vertebral bodies. Infection occurs secondarily to spinal surgery, foreign body migration or septic emboli from the skin or urinary/genital tract, or from a concurrent endocarditis. Clinical signs may include pyrexia, anorexia, spinal pain and paresis.

Disseminated aspergillosis

See *Aspergillosis*.

Disseminated histiocytoma

See *Histiocytic diseases*.

Distal limb fractures and dislocations

These fractures are common in racing Greyhounds, due to the stresses of racing.

Distal tibial valgus deformity

Angular limb deformity of the hindlimb.

Distichiasis

Abnormally positioned cilia (eyelashes) which emerge through or close to meibomian gland orifices. They are often of no clinical significance but in some cases may cause ocular irritation. Inheritance is suspected, given the high incidence in certain breeds, though the exact mode of transmission is unknown.

Dorsal ridge

The dorsal hair ridge is considered normal for Rhodesian Ridgebacks but can be associated with dermoid sinus.

Double muscling

Mutations in the gene for the myostatin protein augment muscle growth.

Dysmyelination of the central nervous system

A severe myelin deficiency of the central nervous system can lead to signs of tremors from a young age.

Dysplasia of the cerebellar cortex

An inherited malformation of the central nervous system characterized by hydrocephalus and an unusual dysplasia of the cerebellar cortex.

Dystocia

Dystocia can be defined as a difficulty or an inability in giving birth. It may result from a wide range of maternal or fetal factors. In dogs, brachycephalic

breeds are predisposed because of a combination of a narrow maternal pelvis and a large fetal head and shoulders. Small nervous breeds may be predisposed owing to a tendency to psychological inhibition and primary uterine inertia. Purebred cats are at higher risk than mixed-breed cats, with dolicocephalic and brachycephalic types at greater risk than mesocephalic.

Eccentrocytosis

A red blood cell abnormality associated with oxidative stress. Eccentrocytes are red blood cells which when observed under a microscope look as if they have most of their haemoglobin on one side only. They have been associated with various conditions including onion/garlic ingestion and vitamin K antagonism.

Eclampsia (puerperal tetany)

Depletion of extracellular calcium resulting in hypocalcaemia. Signs include nervousness, panting, restlessness, tremors and finally seizures. It is most common within 6 weeks of parturition when demand for milk is great. It can affect any breed but is most common in small breeds.

Ectopic cilia

Cilia (eyelashes) which emerge directly through the palpebral conjunctiva to cause corneal irritation, ulceration and pain.

Ectopic ureter

A congenital anomaly involving one or both ureters. There is failure of the affected ureter to terminate in the trigone region of the bladder, opening instead into the urethra, vagina or uterus. Continuous or intermittent urinary incontinence may be seen as a result, usually in the juvenile bitch. Urinary incontinence is less commonly associated with ectopic ureters in the male because of the longer urethra and stronger urethral sphincter.

Ectropion

Eversion (rolling out) of all or part of the eyelid margin leading to exposure and subsequent irritation of the ocular tissues. In some cases ectropion is seen as part of *macroblepharon*.

Ehrlichiosis

Disease caused by the tick-borne parasite *Ehrlichia canis*. This can lead to ocular, haematological and central nervous system signs, such as uveitis, thrombocytopenia and seizures.

Elbow dysplasia

Genetics and rapid growth predispose to this disease complex, which includes an ununited anconeal process, medial coronoid process disease and osteochondritis dissecans of the medial humeral condyle. It may be that elbow dysplasia is caused by osteochondrosis, possibly relating to incongruities of the trochlear notch. In the UK and USA, there are screening schemes for elbow dysplasia in operation.

Electrocardiographic abnormalities

Abnormalities in the electrocardiogram include changes in the height and width of complexes, presence or absence of some of the components of the PQRS complex, and changes in the heart rhythm.

Enamel hypoplasia

Lack of tooth enamel.

Endocardial fibroelastosis

Severe thickening of the endocardium, sometimes involving the mitral valve leaflets. Heart murmur, failure to thrive and congestive heart failure are seen.

Endocrine alopecia

Loss of hair due to hormonal causes.

Entropion

An inward rolling of all or part of the eyelid margin resulting in irritation of the conjunctival and corneal surfaces. Entropion may occur alone, or as part of *macroblepharon*, where it may occur in the upper eyelid and/or lateral and medial to central ectropion in the lower eyelid. Where entropion occurs in the upper eyelid, trichiasis often results. In many cases, entropion arises as a result of conformation of the skull and orbit, and the amount of skin on the head. Most cases present in the first year; some severely affected cases present as early as 2–6 weeks of age. Some mild cases disappear as the dog matures.

Eosinophilia

Increased blood levels of the type of white blood cell known as an eosinophil.

Eosinophilic dermatitis and oedema

This rare condition presents as acute-onset erythematous macules which progress to form plaques. Variable oedema is seen.

Epidermal hyperkeratosis

A mild epidermolytic ichthyosis. Adult dogs with the disease show generalized, pigmented hyperkeratosis with epidermal fragility.

Epidermolysis bullosa acquisita

This is a group of inherited bullous diseases involving abnormal keratin production. Junctional epidermolysis bullosa causes bullae, vesicles and erosions in various locations. Dystrophic epidermolysis bullosa shows erosions in the mucocutaneous junctions. See also *Junctional epidermolysis bullosa*.

Epilepsy

Recurrent seizures caused by functional disorders of the brain. If no structural or metabolic cause can be found, the condition is termed idiopathic epilepsy, and this tends to have an age of onset in dogs of 6 months to 6 years. The high incidence in certain breeds of dog suggests an inherited basis.

Epiphora

Overflow of tears down the face.

Episodic falling (hypertonicity, collapsing Cavalier King Charles Spaniels)

This condition is poorly understood, but it is speculated that it is due to a disorder in GABA transmission. Extensor rigidity induced by exercise is seen, but mentation remains normal.

Erythema multiforme

An acute skin disorder resulting from a hypersensitivity reaction, causing erythematous macules, pustules and plaques.

Eversion of the cartilage of the nictitating membrane

Scrolling of the cartilage of the third eyelid which may result in chronic conjunctivitis. Seen most commonly in young large-breed dogs.

Exercise-induced collapse

Labradors are prone to a condition of collapse on exercise; extensive investigations have as yet failed to reveal a cause.

Exfoliative cutaneous lupus erythematosus (ECLE, lupoid dermatosis of the German Short-haired Pointer)

Hyperkeratosis and alopecia, mainly of the head, from 5–7 months of age.

Exocrine pancreatic insufficiency (EPI)

Lack of digestive enzymes that are produced by the pancreas, causing signs of diarrhoea and weight loss due to maldigestion.

Extraskelatal soft-tissue osteosarcomas

Most osteosarcomas affect the bone, but they sometimes arise from non-bony soft tissue.

Factor I deficiency

Deficiencies in factor I (fibrinogen) prolong the prothrombin time in tests and can cause bleeding disorders.

Factor II deficiency

Factor II deficiencies prolong prothrombin time and can cause bleeding disorders.

Factor VII deficiency

This condition leads to a mild clotting disorder. Prothrombin time (PT) is usually prolonged, but activated partial thromboplastin time (aPTT) is usually normal, as is consistent with a disorder of the extrinsic pathway.

Factor X deficiency

Factor X is part of the common pathway, so PT and aPTT are both prolonged. A specific factor X assay is used to confirm the diagnosis. The severity of bleeding is variable, and some affected dogs may survive into adulthood.

Factor XI deficiency

Factor XI is part of the intrinsic pathway of secondary haemostasis. Deficiency leads to prolongation of PT and aPTT. Affected animals can have prolonged bleeding after surgery or trauma but do not exhibit spontaneous haemorrhage.

Faecal incontinence (part of Manx syndrome)

See *Manx syndrome*.

Familial canine dermatomyositis

This is a hereditary disease causing skin lesions around the face, ear tips and digits.

Familial cutaneous lupus erythematosus (CLE)

An autoimmune skin disorder.

Familial renal disease (familial nephropathy)

Familial diseases include those that occur in related individuals with a greater frequency than chance alone would allow. Familial renal disease should be suspected whenever chronic renal failure occurs in an immature or young animal. If chronic renal failure develops before physical maturity, stunting will develop. Familial renal diseases vary in clinical signs and pathology depending on the breed.

- **Glomerular basement membrane disorder.**

The main lesion in this condition is a thickening and splitting of the glomerular basement membrane, usually resulting in early-onset proteinuria and leading to renal failure.

- **Membranoproliferative glomerulonephritis.**

Glomerulonephritis results from the presence of immune complexes in the glomerular capillary walls, leading to glomerular damage. In most cases, there is significant proteinuria leading to hypoalbuminaemia which, if severe, is manifest as peripheral oedema or ascites (nephrotic syndrome). The condition may progress to renal failure.

- **Periglomerular fibrosis.** Periglomerular fibrosis progresses to generalized interstitial fibrosis and results in renal failure.

- **Polycystic kidney disease:** see under separate heading.

- **Renal amyloidosis:** see under separate heading.

- **Renal cystadenocarcinoma:** see under separate heading.

- **Renal dysplasia:** see under separate heading.

Familial Shar Pei fever

A familial renal amyloidosis, which can also affect the liver. Females are predisposed, and the mean age of onset is 4 years. Signs include fever, joint swelling, vomiting, anorexia and jaundice.

Familial vasculopathy

This is a rare group of conditions which can cause dermatological and systemic signs. In some breeds the disease is inherited.

Familial vasculopathy of the nasal planum, nostrils and nasal mucosa

A presumably hereditary pyogranulomatous and vasculitic disorder of the nasal planum, nostrils and nasal mucosa. Clinical signs include bilateral nasal discharge or a bilateral ulcerative and

destructive process of the nasal planum, nostrils and nasal mucosa.

Fanconi syndrome

Fanconi syndrome results from generalized proximal renal tubular dysfunction resulting in abnormal reabsorption of many solutes, including glucose, amino acids and phosphate. Low blood levels of the solutes involved may result. Affected dogs are polyuric and polydipsic. The condition may progress to acute renal failure or pyelonephritis.

Feline asthma (feline bronchitis, allergic bronchitis)

This condition may be chronic or acute, and can present with mild or severe signs. Coughing and dyspnoea are seen.

Feline gastrointestinal eosinophilic sclerosing fibroplasia

A non-neoplastic gastrointestinal mass. The cause is unknown.

Feline hyperaesthesia syndrome

Cats with this condition show a combination of dermatological, neurological and behavioural signs including self-trauma, vocalization and apparent hallucinations. The cause is not known, but there can be a variety of trigger factors such as skin irritants and stress.

Feline idiopathic cystitis

Signs of cystitis with no obvious underlying cause. Cases present with dysuria, pollakiuria, haematuria and sometimes urethral obstruction. Predisposing factors are believed to include being male, neutered, overweight, using an indoor litter box, limited outside access, dry diet and living in a stressful relationship with one or more other cats.

Feline infectious peritonitis (FIP)

Caused by feline coronavirus. While many cats are seropositive for feline coronavirus, only 10% will go on to develop FIP. FIP can develop in two forms: 'dry' or non-effusive FIP where pyogranulomas develop in various organs throughout the body, and 'wet' or effusive FIP where ascites and pleural effusions develop.

Feline orofacial pain syndrome

In this condition, chewing motions and pawing at the face are seen. This can be episodic or continuous. Oral disease is often present.

Femoral artery occlusion

Femoral pulses are found to be weak or absent in these cases.

Fibroadnexal hamartoma

Non-inflammatory, non-neoplastic skin nodule.

Fibrocartilaginous embolic myelopathy (FCEM, fibrocartilaginous embolism (FCE), ischaemic myelopathy)

Fibrocartilage causes a focal obstruction of blood flow to the spinal cord. The exact mechanism is unknown, but signs often occur acutely at exercise, and lead to lateralized, usually non-painful signs of myelopathy such as paresis and paralysis. Large and giant breeds are more likely to be predisposed.

Fibroma/fibrosarcoma

A fibrosarcoma is a malignant tumour derived from fibrous tissues and may be found in many sites, including the bone, skin, spleen and oral cavity. Fibroma is the benign form. Tumour behaviour varies with the site and histological grade. In general, fibrosarcomas are locally invasive but have a relatively low rate of metastasis. Oral fibrosarcomas tend to be histologically low grade but biologically high grade with a tendency to aggressive local infiltration and a higher potential for local and distant metastasis. Cutaneous fibrosarcomas are more common in cats than in dogs. Tumours vary from slow-growing, well-circumscribed lesions which are more benign in behaviour to rapidly growing, poorly circumscribed lesions which may be more malignant.

Flank-sucking

A compulsive behaviour seen in some dogs, often just before they fall asleep. It may be associated with blanket-sucking and kneading of the paws.

Flat-chested kittens

A congenital thoracic wall defect.

Flea-bite hypersensitivity

Severe inflammatory reactions may occur due to hypersensitivity reactions to allergens in flea saliva.

Focal metatarsal fistulation (sterile idiopathic pedal panniculitis)

Causes sinus tracts in the metatarsi that can discharge a sterile exudate.

Folate – low serum level

Folate (vitamin B₉) levels in the blood may be low because of disease in the small intestine.

Follicular dysplasia

This is a group of conditions having alopecia and coat changes in common. They are suspected to be inherited in many breeds.

Folliculitis

Inflammation of the hair follicles.

Footpad hyperkeratosis

This condition is familial in some breeds. Signs of severe hyperkeratosis are usually seen by 6 months of age.

Frontonasal dysplasia (Burmese head defect)

This congenital defect is inherited and has been described as autosomal recessive or autosomal dominant with incomplete penetrance. Heterozygotes have a range of facial dimensions. The condition is fatal to homozygotes. It is seen in the eastern, 'new look' or contemporary strain of Burmese. The upper maxillary region is duplicated.

Gallbladder mucocoele

Distension of the gallbladder with mucus. The cause is unknown, but it can be associated with hypothyroidism and hyperadrenocorticism.

Gastric carcinoma

A primary, malignant stomach tumour derived from epithelial cells.

Gastric dilatation/volvulus (bloat, GDV)

Gastric distension due to the rapid accumulation of food, fluid or gas resulting in torsion of the stomach. Usually an acute and severe condition which is rapidly fatal without treatment. Large deep-chested dogs are predisposed. Mesenteric volvulus involves a twist of the small intestines, which is also acutely life-threatening.

Gastric neoplasia

Tumours of the stomach include adenocarcinoma, lymphoma, leiomyomas and leiomyosarcomas.

Gastrocnemius musculotendinopathy

A chronic strain injury which can lead to rupture of the gastrocnemius tendon.

Gastroduodenal perforation

Rupture of the upper gastrointestinal tract, secondary for example to ulceration, will cause life-threatening peritonitis.

Gastrointestinal changes seen with BOAS

There is a high prevalence of gastrointestinal disorders in dogs with brachycephalic obstructive airway syndrome (BOAS), diagnosed clinically, endoscopically and histologically. These changes improve following surgical management of the upper airway disease.

Gastrointestinal foreign bodies

Foreign bodies in the gastrointestinal tract can cause vomiting, and may cause obstruction, leading to dehydration, shock, rupture and peritonitis. See also *Oesophageal and gastric foreign bodies*.

Generalized sebaceous gland hyperplasia

Rare idiopathic disease causing greasy coat and hair clumping.

German Shepherd Dog pyoderma

This is an idiopathic deep pyoderma affecting German Shepherd Dogs. Probably an inherited immunodeficiency causes susceptibility to infection.

Gestation length

Gestation is generally shorter with larger litters, but breed seems to exert an independent effect.

Giant axonal neuropathy

A progressive peripheral neuropathy causing paresis, hyporeflexia and hypotonia and progressing to faecal incontinence and sensory deficits.

Giant platelets and thrombocytopenia

Some breeds have fewer but larger platelets, without a reduction in clotting function.

Giardiasis

Caused by *Giardia* spp. parasites that can cause signs of diarrhoea.

Gingival and oropharyngeal neoplasia

Cancer of the gums, oral cavity and throat.

Glaucoma

A group of diseases characterized by degeneration of the retinal ganglion cells and optic nerve resulting in progressive loss of vision. The condition is

associated with an increase in intraocular pressure. Primary glaucoma develops without the presence of other intraocular disease and may be hereditary, with potential for bilateral involvement. Primary glaucomas may be divided by the appearance of the iridocorneal filtration angle into *open-angle* and *closed-angle* glaucomas. Causes of secondary glaucomas include lens luxation, uveitis, neoplasia and cataracts.

Globoid cell leucodystrophy (Krabbe disease)

An enzyme deficiency that causes accumulation of galactocerebroside in oligodendrocytes and Schwann cells. Leads to paralysis hyper/dysmetria, tremors, blindness and death.

Gluten-sensitive enteropathy

An intolerance to gluten-containing foods, most commonly seen in Irish Setters. Weight loss and chronic diarrhoea due to malabsorption are seen.

Glycogen storage disease type III

Type III glycogen storage disease (Cori's disease) leads to poor growth, weakness and liver disease. This disease is rare and also has a poor prognosis.

Gracilis contracture (fibrotic myopathy)

This is usually seen in athletic dogs and leads to a gait alteration. This condition may be part of a complex with *fibrotic myopathy of semitendinosus muscle*. Surgery can help, but the condition can recur.

Granulomatous meningoencephalitis

See *Meningoencephalitis*.

Granulomatous sebaceous adenitis

See *Sebaceous adenitis*.

Granulosa-theca cell ovarian tumour

A type of sex cord stromal tumour of the ovary. This tumour is potentially malignant.

Grass awn migration

Grass awns are reported to comprise 61% of foreign bodies. Predilection sites include ears, eyes, interdigital web and nose.

Grey eosinophils

Eosinophils stain abnormally, which may lead to underestimating their numbers on automated counts.

Haemangioendothelioma

A group of vascular neoplasms.

Haemangioma

See *Cutaneous haemangioma*.

Haemangiosarcoma

This is a highly malignant tumour arising from vascular endothelial cells. Primary sites include the spleen (most common site in the dog), the right atrium of the heart, liver, skin, bone, nervous system, kidney, bladder and oral cavity. Metastasis to a wide variety of sites is common, and in many cases micrometastasis has occurred by the time of diagnosis.

Haematological differences

Some breeds differ in the normal range of their haematological parameters, possibly related to the purposes they have been bred for, such as racing Greyhounds.

Haemophilia A

This deficiency of factor VIII can cause moderate to severe bleeding. aPTT is prolonged, but PT is normal. Many cases may arise from new mutations. The condition is inherited as a sex-linked recessive.

Haemophilia B (factor IX deficiency, Christmas disease)

This clotting deficiency is caused by a deficiency in factor IX. This is a sex-linked condition, but because the gene for factor IX is smaller than for factor VIII, spontaneous mutations are less common for haemophilia B than for haemophilia A. aPTT is prolonged, but PT is normal.

Haemorrhagic gastroenteritis

An acute haemorrhagic vomiting and diarrhoea syndrome of unknown cause. Most commonly seen in middle-aged small-breed dogs.

Heart block

A slowing or failure of electrical conduction in the heart. First-degree heart block involves a prolongation of the PR interval, second-degree involves intermittent block of the conduction from sinoatrial node to ventricles, and third-degree involves a complete block. Can lead to signs of weakness, collapse and syncope, depending on severity.

Heart disease

Heart disease can be acquired, such as cardiomyopathies and valve degenerations, or congenital, such as stenosis or valve malformations.

Heart murmur

An abnormal heart sound. A murmur may be associated with heart disease or may be innocent.

Hemivertebrae (wedge-shaped vertebrae)

Among the most frequent vertebral malformations in dogs, hemivertebrae are congenitally malformed vertebrae most commonly seen at the level of thoracic vertebrae 7–9. Neurological signs, e.g., pelvic limb ataxia, paresis, faecal and urinary incontinence, may result from spinal cord compression.

Haemophagocytic syndrome

Proliferation of haemophagocytic histiocytes causes cytopenias. May be idiopathic or secondary to other disorders such as infectious diseases, immune-mediated disorders and neoplasia. Familial and acquired forms have been described in humans.

Hepatocerebellar degeneration

A syndrome causing progressive cerebellar and hepatic disease, leading to neurological signs from the age of 4–6 weeks.

Hereditary ataxia

Progressive ataxia results from degeneration of the white matter of the cervical and thoracic spinal cord in young Smooth-haired Fox Terriers and Jack Russell Terriers.

Hereditary lupoid dermatosis (exfoliative cutaneous lupus erythematosus)

This is a familial disease of unknown cause. Scaling begins on the head and progresses to become generalized. There may be systemic signs. No consistently successful treatment has been described.

Hereditary meningoencephalocoele

Herniation of the brain through the skull which is fatal. Is also associated with craniofacial abnormalities.

Hereditary myopathy of Sphynx and Devon Rex cats

A variably progressive myopathy causing signs of muscle weakness, megaoesophagus and fatigue.

Hereditary nasal parakeratosis in Labrador Retrievers

Hyperkeratosis and depigmentation of the nasal planum are seen in this condition. The dogs are otherwise healthy.

Hereditary necrotizing myelopathy

This disease causes hindlimb paresis in young dogs. It appears similar to Afghan myelopathy.

Hereditary polioencephalomyelopathy of the Australian Cattle dog

This is a degenerative condition primarily affecting the grey matter, causing focal malacia in the cerebellum, brainstem and spinal cord.

Hereditary retinal dystrophy of Briards (congenital stationary night blindness)

A retinal dystrophy causing congenital night blindness with a variable effect on day vision. The disease is slowly progressive, with no fundoscopic changes until 2–3 years of age. Nystagmus may be present. A defect in retinal polyunsaturated fatty acid metabolism may be involved.

Hereditary spectrin deficiency

Spectrin deficiency with increased red blood cell osmotic fragility is associated with hereditary spherocytosis and haemolytic anaemia in humans. Although spectrin deficiency was found to be common in a population of Golden Retrievers, it was unclear whether this was associated with haemolytic anaemia in these animals.

Hereditary thrombopathy (Glanzmann's thrombasthenia)

This condition affects platelets, leading to decreased platelet retention and absent platelet aggregation. This causes severe mucosal bleeding. Platelet count is usually normal or possibly slightly decreased. The buccal mucosal bleeding time is increased and clot retraction is abnormal. It is inherited as an autosomal recessive trait.

Hiatal hernia

Protrusion of part of the oesophagus and/or part of the stomach through the oesophageal hiatus into the thoracic space. Signs include vomiting, regurgitation and hypersalivation.

High allergen-specific serum immunoglobulin E levels

High levels of allergen-specific IgE are associated with atopy, but can be found in dogs of some breeds which do not have atopy.

Hip dysplasia

This very common condition occurs in a wide range of breeds, with large and giant breeds particularly affected. Various deformities in the hip lead to joint instability with the development of degenerative joint disease. Many affected dogs show no or minimal clinical signs. Genetics undoubtedly play a role, but environmental factors such as nutrition and exercise are also important. Various screening programmes are used around the world in an attempt to reduce the incidence of this condition. Estimates of heritability vary from 20% to 60%, depending on breed, population examined and methods applied.

Histiocytic diseases

Histiocytes are white cells that occur in the tissues and include macrophages and dendritic cells. There are several well-defined conditions, but the terminology of these diseases is somewhat confused.

- **Reactive histiocytoses** are considered non-neoplastic, but occur due to a degree of immune dysregulation. *Cutaneous histiocytosis* is confined to the skin and draining lymph nodes; *systemic histiocytosis* may involve other body sites.
- **Histiocytic sarcoma complex** is a malignant disease. *Histiocytic sarcomas* are localised lesions which may occur in many tissues including spleen, lymph nodes, skin, and brain tissue. Histiocytic sarcomas have also been termed 'localised histiocytosis'. *Disseminated histiocytoma* occurs when histiocytic sarcomas occur in multiple organs. This is also sometimes known as 'malignant histiocytosis' and is rapidly progressive and fatal.
- **Canine cutaneous histiocytoma** is a solitary lesion in a young dog which often regresses spontaneously: see under separate heading.

Histoplasmosis

The causal organism, *Histoplasma capsulatum*, is a saprophytic soil fungus. Histoplasmosis is uncommon and occurs mainly in the central USA, India and southeastern Asia. Clinical signs include fever, anorexia, weight loss, cough,

dyspnoea and ocular and skin lesions; however, many infections are subclinical.

Horner's syndrome

Neurological disease characterized by miosis (constricted pupil), ptosis (drooping eyelid), enophthalmos (shrunken eye), and prolapsed nictitans. Causes in dogs include trauma and ocular disease, but 50% of cases are idiopathic (cause unknown).

Hydrocephalus

Hydrocephalus occurs where there is dilatation of all or part of the ventricular system of the brain, and may be congenital or acquired (usually secondary to neoplasia or inflammatory disease). Signs include a domed cranium, seizures and altered mental status.

Hyperadrenocorticism (Cushing's syndrome)

One of the most commonly diagnosed endocrinopathies in the dog, but rare in the cat. Hyperadrenocorticism occurs where there is a sustained and inappropriately elevated secretion of cortisol from the adrenal cortex. Hyperadrenocorticism may be pituitary-dependent, where there is excessive adrenocorticotrophic hormone (ACTH) secretion leading to adrenal cortical hyperplasia and increased cortisol secretion, or it may be adrenal-dependent, where there is a functional adrenocortical tumour.

Hypercholesterolaemia

High levels of serum cholesterol. Associated with atherosclerosis and coronary heart disease in humans. May be primary or secondary, e.g. to endocrine diseases, nephrotic syndrome or cholestasis.

Hyperhomocysteinaemia

This multifactorial condition has been reported in Greyhounds with suspected gastrointestinal disease.

Hypertrophic cardiomyopathy (HCM)

In this condition a hypertrophied ventricle causes congestive heart failure and dysrhythmias. The incidence is reported as being 1.6–5.2% in cats, but it is rare in dogs. It is likely that the condition is inherited, although modifying factors may cause variable expression of the condition.

Hypertrophic neuropathy

An inherited neuropathy reported in the Tibetan Mastiff which results in generalized weakness, hyporeflexia and dysphonia from 7–10 weeks of age. There is no treatment, and the prognosis is guarded.

Hyperuricosuria

Excessive excretion of uric acid in the urine. Predisposes to urate urolithiasis. See also *Urolithiasis*.

Hypoadrenocorticism (Addison's disease)

A condition in which there is inadequate adrenocortical hormone production, leading to a deficiency in mineralocorticoids and/or glucocorticoids. Primary hypoadrenocorticism (Addison's disease) results most commonly from immune-mediated destruction of the adrenal cortices leading to deficiencies of all adrenocortical hormones. A hereditary factor has been suggested in some breeds. It is very rare in the cat.

Hypocobalaminaemia

Low blood cobalamin (vitamin B₁₂) levels.

Hypodontia

Missing teeth due to a failure of the teeth to develop.

Hypokalaemic polymyopathy

Clinical signs of this condition include ventroflexion of the neck and transient weakness. There may also be a tremor. Serum potassium levels of less than 3 mmol/l are seen.

Hypomagnesaemia

Low levels of magnesium can occur for a wide number of reasons such as reduced intake, reduced gastrointestinal absorption (with gastrointestinal disease) and increased renal loss (e.g. renal disease, diabetes mellitus, hyperadrenocorticism). It can lead to a variety of signs including weakness, arrhythmia and seizures.

Hypomyelination syndrome

In this condition there is a reduction or absence of myelin in the axons of the central nervous system. The myelin is biochemically normal but present in reduced amounts, leading to nervous signs such as tremors, ataxia and hypermetria. The myelin deficiency may be permanent or

simply delayed, and in some cases the clinical signs disappear with age.

Hypospadias

Hypospadias results from incomplete fusion of the urethral folds during the formation of the male urethra such that the urethra opens abnormally on the underside of the penis, proximal to the normal urethral opening. It may be seen with other congenital defects such as cryptorchidism and intersexuality. It has also been described in the female dog, where it is mostly seen in intersex states.

Hypothyroidism

A common endocrine disease in the dog. There is a deficiency in the secretion of thyroid hormone either as a result of thyroid gland destruction (primary hypothyroidism), inadequate pituitary production of thyroid-stimulating hormone (TSH) (secondary hypothyroidism) or inadequate hypothalamic secretion of thyrotropin-releasing hormone (TRH) (tertiary hypothyroidism). Many breeds seem predisposed to primary hypothyroidism, most notably Doberman Pinschers and Golden Retrievers.

Ichthyosis (fish scale disease)

Extreme hyperkeratosis is seen, and affected dogs are abnormal at birth. Scaling, erythema and severe seborrhoea are seen.

Idiopathic epilepsy

See *Epilepsy*.

Idiopathic facial dermatitis in Persians and Himalayans

This is an uncommon disease demonstrating crusting, erythema and self-trauma to the head and neck. Secondary bacteria and yeasts are common.

Idiopathic head tremor syndrome

The head tremors seen in this condition are of unknown aetiology, but may be seizures or dyskinesia. Dogs remain fully conscious during the event, which last for a variable amount of time.

Idiopathic nasodigital hyperkeratosis

This rare keratinization disorder causes hyperkeratosis of the dorsal nasal planum.

Idiopathic polyneuropathy in Alaskan Malamutes

A progressive polyneuropathy of young adults causing neurological signs such as paraparesis and muscle atrophy.

Idiopathic sterile granuloma and pyogranuloma

Thought to be an immune-mediated disease, causing non-painful lesions, typically around the face and feet.

Idiopathic ulcerative dermatosis in Shetland Sheepdogs and Collies

This condition may be related to dermatomyositis. Vesicles and bullae progress to ulcers, especially in the inguinal and axillar regions and mucocutaneous junctions.

Immune-mediated haemolytic anaemia (IMHA)

Mild to severe anaemia, chronic or acute, can be seen with this condition. Haematology shows anaemia (with a regenerative response if the anaemia is not too acute) and spherocytosis. Although many cases respond well to treatment, the potential for serious complications means that the prognosis for this condition is guarded.

Immune-mediated inflammatory myopathy with severe tongue muscular atrophy

Thought to be a generalized myopathy which particularly targets the muscles of the head and tongue to cause atrophy.

Immune-mediated polyarthritis

Inflammation affecting multiple joints. This may be primary or secondary to other conditions.

Immune-mediated rheumatic disease (IMRD)

Chronic stiffness and joint pain characterize this disorder, which may be related to systemic lupus erythematosus.

Immune-mediated thrombocytopenia (IMTP)

This disorder is characterized by immune-mediated destruction of platelets. Epistaxis, haematochezia and mucosal haemorrhage are often seen.

Immunodeficiency syndrome

Several different breeds have primary immunodeficiencies, predisposing to a variety of infections. Respiratory conditions in related Irish Wolfhounds have been attributed to an underlying immunodeficiency, possibly in cell-mediated immunity or in IgA.

Immunoglobulin deficiency

See *Pneumocystis carinii* infection.

Immunoproliferative enteropathy (of Basenjis)

A specific disease of Basenjis of unknown cause. Several forms of pathology may be identified in the gastrointestinal tract including hypertrophic gastritis, lymphocytic plasmacytic enteritis and villous clubbing and fusion resulting in chronic diarrhoea, anorexia and weight loss.

Incomplete ossification of the humeral condyle (IOHC)

This condition may present with a history of mild intermittent lameness which is unresponsive to anti-inflammatory drugs. Acute severe lameness may follow exercise or mild trauma, corresponding with a humeral condylar fracture.

Increased C-reactive protein (CRP)

CRP is an acute-phase protein that is a marker for inflammation.

Increased osmotic fragility of erythrocytes (resulting in haemolytic anaemia)

Extreme fragility of the erythrocytes (red blood cells) can lead to recurrent and severe anaemia, with splenomegaly and weight loss. Prednisolone and blood transfusions may be helpful.

Inflammatory bowel disease

Immune-mediated intestinal disease. This may be a reaction to food allergens, but other mechanisms are possible. Causes signs of malabsorption such as diarrhoea, vomiting and weight loss.

Inflammatory myopathy

Generalized inflammation of the muscles may be due to immune-mediated polymyositis, infectious causes and preneoplastic syndromes. Focal inflammation may be due to diseases such as masticatory myositis and dermatomyositis.

Inherited focal retinal degeneration

This condition is distinct from generalized progressive retinal degeneration. Lesions can progress to blindness and cataract formation.

Inguinal/scrotal herniation

A weakness in the body wall in the inguinal region.

Inherited myopathy of Great Danes

Previously known as central core myopathy, this very rare condition leads to signs of muscle weakness, which becomes progressively more severe. The reported cases were euthanized before 2 years of age.

Inherited polyneuropathy

An idiopathic polyneuropathy of Leonbergers which is thought to have a sex-linked inheritance. Similar to Charcot–Marie–Tooth disease in humans.

Innocent systolic heart murmur

Abnormal heart noises may indicate cardiac disease, or may be incidental.

Insulinoma

Insulinomas are functional insulin-secreting tumours of the pancreatic beta cells. Insulin secretion is independent of the normal negative feedback control, resulting in hypoglycaemia. These tumours are generally slow growing but malignant with a high metastatic potential.

Intervertebral disc disease (IVDD)

Degeneration of the intervertebral discs, resulting in extrusion or protrusion of the nucleus pulposus, may result in spinal cord compression and pain/paresis. Nuclear extrusion occurs early in chondrodystrophoid breeds, e.g., Pekingese, Dachshunds, Beagles, Welsh Corgis, French Bulldogs, some Spaniels and Basset Hounds, giving rise to signs in younger dogs.

Intestinal adenocarcinoma

A malignant tumour found most frequently in the colon of dogs and the jejunum and ileum of cats. Most are locally invasive and metastasize early to the lymph nodes and liver.

Intestinal adenoma

A benign intestinal tumour.

Intracutaneous cornifying epithelioma (keratoacanthoma)

Benign cutaneous neoplasms which can be solitary or multiple (generalized).

Intussusception

A condition where one part of the intestines telescopes inside another, which can cause a partial or complete obstruction.

Iridociliary cysts

These cysts of the eye occur spontaneously in older dogs. They are often incidental findings.

Iris hypoplasia

Failure of the iris to develop properly.

Irish Setter hypochondroplasia

The limbs are slightly shortened in this condition. The ulna and radius may be bowed and carpal valgus is seen.

Ischaemic stroke

Blockage of a blood vessel, usually by a thrombus, causing damage to part of the brain.

Ivermectin and milbemycin

Toxicosis from the use of these parasiticides causes signs including drooling, vomiting and seizures.

Joint flexibility

Unique flexibility of the joints (especially the shoulder and neck) seen in the Norwegian Lundehund.

Junctional epidermolysis bullosa (JEB)

Separation of the dermis from the epidermis, leading to bullae and ulcers over prominences and claw dystrophy. See also *Epidermolysis bullosa acquisita*.

Juvenile-onset distal myopathy

This condition has been reported in several pups and has been described as a muscular dystrophy. Clinical signs include decreased activity and various postural abnormalities.

Juvenile renal disease

Chronic kidney disease can occur at a young age, causing a number of signs including polyuria, polydipsia, vomiting and weight loss.

Keratoconjunctivitis sicca (KCS, dry eye)

A common disease characterized by reduced aqueous tear production resulting in drying and inflammation of the conjunctiva and cornea. The condition may be congenital (rarely), or result from infectious, drug-induced, neurological or immune-mediated causes. A genetic influence is suggested by the high incidence in a number of breeds.

Kidney disease

A number of diseases can affect the kidneys, including inherited disorders such as dysplasia, neoplastic disorders, and idiopathic conditions. Signs include vomiting, weight loss and polyuria/polydipsia. See also *Chronic kidney disease* and *Familial renal disease*.

Kitten birth weight

The average weight of newborn kittens can vary by breed. Kitten birth weight tends to be lower with larger litters and shorter gestation, but breed seems to exert an independent effect

L-2-Hydroxyglutaric aciduria (organic aciduria)

This metabolic disorder causes neurological signs with onset from 6 months of age to 7 years. Typical changes are observed on MRI scans.

Lafora's disease

Progressive myoclonic epilepsy is seen in this condition. Diagnosis is by histopathological examination of affected tissue.

Laryngeal paralysis–polyneuropathy syndrome

Usually idiopathic but may be related to generalized myopathies or neuropathies. Stridor, aggravated by excitement and exercise, is the main clinical sign, although severe cases may progress to cyanosis and collapse.

Lateral luxation of the superficial digital flexor tendon

A tear in the retinaculum of the superficial digital flexor tendon can cause mild to moderate lameness with luxation of the tendon.

Left atrial rupture

This is a rare cause of pericardial effusion, and is usually secondary to mitral valve disease.

Left systolic apical murmur

Heart murmurs in this region are often related to mitral valve disease.

Leishmaniasis

A disease caused by the protozoa *Leishmania* spp. The visceral form causes systemic signs including lymphadenopathy, weight loss, epistaxis and melaenia. The cutaneous form causes ulcerative nodules. The sandfly is the vector of transmission.

Lens luxation

Displacement of the lens from its normal position. Lens luxation may be primary (not associated with other ocular conditions) and considered inherited, or secondary to trauma, cataract formation, glaucoma, neoplasia or uveitis. Lens luxation is a potentially serious condition and may result in raised intraocular pressure, glaucoma and loss of vision. Where both lens luxation and glaucoma occur, it is not always clear which condition is primary. Primary lens luxation (PLL) is an inherited disease in which the lens comes free from its attachments and moves anteriorly or posteriorly in the eye. It is usually bilateral, though both lenses do not usually luxate simultaneously. Terrier breeds are predisposed, and it is usually first diagnosed when dogs are 2–6 years old. The lens in the eye can also luxate secondary to conditions such as glaucoma, cataract and trauma.

Leptospirosis

A bacterial infection caused by *Leptospira* spp., of which there are many subtypes (serovars). Causes signs of renal and liver failure.

Lethal astrocytosis

A neurological disorder of Gordon Setters manifesting as early onset of gait and postural abnormalities. The condition is progressive, and affected puppies are recumbent at 5–6 weeks of age.

Leucopenia

Low white blood cell count.

Leucodystrophy

Several leucodystrophies have been reported in different breeds, generally presenting with signs of ataxia and paresis leading to paralysis.

Leucoencephalomyelopathy

This is believed to be an inherited condition. Degeneration of the myelin of the spinal cord, brainstem, cerebellum and sometimes optic tracts results in ataxia, tetraparesis and loss of conscious proprioception, with increased spinal reflexes and muscle tone. Vision is usually unaffected. The condition progresses over 6–12 months.

Limber tail

A sudden development of a flaccid tail. Thought to be caused by a strain of the tail muscles.

Lingual haemangiosarcoma

A malignant tumour of the tongue.

Lingual plasma cell tumour

A tumour of the tongue.

Lipaemia of the aqueous humour

A transient milky appearance in the eyes of young Burmese cats which may be related to abnormal lipid metabolism.

Lipoid dystrophy (amaurotic idiocy)

Amaurotic idiocy is one of several lipoid dystrophies involving the central nervous system.

Lipoma

A benign tumour of fat cells, generally found in the subcutaneous tissues. Lipomas are common, affecting up to 16% of dogs. Infiltrative lipomas are locally invasive, making surgical excision more difficult, but they do not metastasize.

Lissencephaly

A developmental anomaly where the cerebral cortex has reduced or absent gyri or sulci, resulting in a smooth appearance. Clinical signs are usually seen from a few months of age and may include behavioural abnormalities, lack of training, aggressive behaviour, visual deficits and seizures.

Localized parakeratotic hyperkeratosis

A scaling disorder which may be responsive to zinc supplementation.

Low fertility

Various medical and environmental conditions can contribute to poor fertility.

Low immunological response to vaccination

The immune response is a complex continuous trait with multiple genetic, environmental and lifestyle factors. Various factors including breed can affect the strength and duration of response to vaccination, emphasizing the need for widespread vaccination to provide protection by herd immunity. Larger breeds and dogs aged < 1 year are predisposed to lower titre responses.

Low Schirmer tear test (STT)

The Schirmer tear test measures tear production.

Low serum cobalamin and cTLI concentrations

Low levels of vitamin B₁₂ and trypsin-like immunoreactivity, a marker for exocrine pancreatic function.

Low thiopurine methyltransferase (TPMT) activity in red blood cells

This enzyme is important in the metabolism of certain drugs such as azathioprine, and its activity is lower in some breeds than others.

Low thrombocyte count (physiological thrombocytopenia)

Low platelet levels.

Lower urinary tract neoplasia

Neoplasia of the lower urinary tract can cause signs of haematuria, pollakiuria, dysuria and urinary obstruction. Transitional cell carcinoma is one of the most common types of lower urinary tract neoplasia.

Lumbosacral disease

Stenosis (narrowing) of the lumbosacral vertebral canal and/or intervertebral foramina causes compression of the lumbosacral nerve roots. Clinical signs may include pain on palpation of the area, pelvic limb paresis or lameness, tail paralysis, hypotonia of the anal sphincter and bladder atonicity ('lumbosacral syndrome').

Lumbosacral transitional vertebrae

Abnormally formed vertebra occur congenitally between the last normal lumbar vertebra and the first normal sacral vertebra. This vertebral anomaly may predispose to cauda equina syndrome.

Lung lobe torsion

Rotation of a lung lobe along its long axis. This rare and life-threatening condition is more common in large, deep-chested breeds. Presenting signs include dyspnoea and pleural effusion. There may be an accompanying chylothorax.

Lyme nephritis

This condition is seen in 1–2% of cases that have recovered from borreliosis, and involves a protein-losing nephropathy.

Lymphangiectasia

Distension of the lymphatic vessels in the intestinal mucosa. It may be a primary congenital disease, or it may be secondary to other disorders including inflammatory bowel disorders, neoplasia, thoracic duct occlusion or right-sided heart failure. The condition results in *protein-losing enteropathy* leading to weight loss, diarrhoea and in some cases oedema and ascites.

Lymphoma

Lymphoma is a malignant lymphoproliferative disease also known as malignant lymphoma or lymphosarcoma. Lymphoma is the most common haematopoietic tumour in the dog and cat. Lymphoma may be classified anatomically by the location of the disease (multicentric, mediastinal, alimentary, cutaneous or extranodal), histologically, or immunophenotypically as B-cell or T-cell.

Lymphoproliferative disease

Excessive production of lymphocytes, usually due to neoplastic disease.

Lysosomal storage disease

Lysosomal storage diseases include ceroid lipofuscinosis, fucosidosis, glycogen storage disease type II (Pompe disease), glycogen storage disease type IV, GM₁ gangliosidosis, GM₂ gangliosidosis and α -mannosidosis.

These rare diseases result from a failure of normal metabolic processes due to a deficiency of an enzyme within the lysosomes of neuronal tissues. As a result, substrate accumulates, causing cellular dysfunction and eventually death. One of a variety of lysosomal enzymes may be affected. Signs usually occur before 1 year of age and may include ataxia, tremors, seizures, dementia and blindness. Most lysosomal storage diseases are believed to be inherited as an autosomal recessive trait.

Macroblepharon (diamond eye)

An abnormally large palpebral fissure. In some cases (e.g. brachycephalic breeds), this occurs as a result of exophthalmos (protrusion of the globe) leading to exposure keratopathy. In others it results from overlong eyelid margins and laxity of the canthal structures leading to both upper lid entropion and lower lid ectropion, resulting in 'diamond eye' in severe cases. The precorneal tear film is often disturbed, and corneal and conjunctival disease may occur secondarily.

Malassezia dermatitis

This yeast infection by *Malassezia pachydermatis* commonly causes pruritus and a greasy, scaly skin disease.

Malassezia skin colonization

Some breeds are prone to increased carriage of *Malassezia pachydermatis* despite being clinically healthy.

Male pseudohermaphroditism (persistent Müllerian duct syndrome)

A pseudohermaphrodite is an individual in whom the chromosomal and gonadal sex agree, but the phenotypic sex is reversed. Therefore, a male pseudohermaphrodite has both an X and a Y chromosome, testes (usually undescended) and female genitalia. Persistent Müllerian duct syndrome is an inherited form seen in Miniature Schnauzers.

Mammary neoplasia

This is common in both the dog and the cat. Mammary tumours are derived from the epithelial and sometimes myoepithelial tissues of the mammary glands. In dogs approximately 50% are benign, in cats over 80% are malignant. Entire animals or those spayed after several seasons are predisposed. Behaviour varies depending on the histological grade, but malignant mammary tumours may be very aggressive, metastasizing to the local lymph nodes, the lungs, and occasionally the abdominal organs and bone.

Manx syndrome

The Manx breed has a mutation resulting in shortening or absence of the tail. This mutation may also result in spina bifida, a developmental abnormality of the spine which may result in loss of control of urination and defecation

(urinary and faecal incontinence), and rectal prolapse. Some cases also have reduction in the control of the hindlegs (paresis).

Mast cell tumour (MCT)

Mast cell tumours are relatively common in dogs, representing up to 20% of skin tumours. Up to 40% of cases have mutations in a proto-oncogene, *c-kit*. MCTs may present in a wide variety of forms, and therefore they need to be included in the differential of all skin masses. Clinical behaviour varies from benign to highly aggressive malignant tumours which have the potential to metastasize (usually to the liver, spleen or kidney).

Cutaneous mast cell tumours are less common in the cat. 'Mastocytic' mast cell tumours are similar to the canine mast cell tumours and occur mostly as solitary nodules in older cats. A variant classified histologically as a 'histiocytic' mast cell tumour has been seen in young Siamese cats. These tumours are multicentric and regress spontaneously over a number of months. Systemic and intestinal forms of mast cell tumour may also be seen in this species.

Medial displacement of the biceps brachii tendon

A gradual onset in lameness which is exacerbated by exercise is seen with this unusual condition. Palpation and manipulation of the shoulder can reveal pain, crepitus and sometimes a palpable popping of the tendon out of the intertubercular groove.

Mediastinal lymphoma

See *Lymphoma*.

Megacolon and constipation (part of Manx syndrome)

Dilatation of the colon which results in constipation.

Megaesophagus and pyloric dysfunction

Oesophageal dilatation and reduced motility resulting in regurgitation. Megaesophagus can be classified as congenital idiopathic megaesophagus, which presents shortly after weaning, or acquired megaesophagus, which occurs later in life and may be idiopathic or secondary to another condition such as myasthenia gravis.

Melanoma

Cutaneous melanomas represent 4–6% of canine skin tumours and 1–2% of feline skin tumours. They present as firm, pigmented dermal masses and are more common in dark-skinned animals. Those found on the digits and close to mucocutaneous junctions tend to be more malignant and may metastasize to local lymph nodes, lungs and other more distant sites.

Oral melanomas are the most common oropharyngeal malignancy in the dog. They are highly metastatic: up to 20% may have metastatic disease at the time of diagnosis.

Melanomas are the most common primary tumour of the globe. They are most commonly found in the anterior uvea (canine anterior uveal melanoma – CAUM) but may be found in other areas, for example the limbus. In the dog they have low metastatic potential but expansion may lead to destruction of the eye. In the cat they may be malignant.

Meningioma

See *Primary brain tumour*.

Meningoencephalitis

Also known as meningoencephalitis of unknown origin (MUO) or meningoencephalitis of unknown aetiology. This is a group of inflammatory diseases including necrotizing encephalitis and granulomatous encephalitis of unknown cause. The disease may be focal or diffuse and may affect any part of the central nervous system, leading to a wide range of clinical signs, including seizures, ataxia, nystagmus and visual deficits. The disease is often chronic and progressive. Small-breed dogs are most commonly affected.

Metaphyseal osteopathy (hypertrophic osteodystrophy)

This uncommon idiopathic condition affects the metaphyses of the long bones of young rapidly growing large-breed dogs, causing pain, swelling, lameness and pyrexia. It affects dogs aged 2–6 months, and males may be predisposed.

Methylmalonic aciduria

This organic aciduria is a metabolic disease, often associated with cobalamin deficiency, which can lead to neurological signs.

Microvascular portal dysplasia

Congenital malformation of the intrahepatic portal circulation resulting in vascular shunting and hepatic dysfunction. Cases are often sub-clinical; signs, where they occur, are similar to those of portosystemic shunts but occur later and are generally milder.

Middle ear polyps

Fleshy lumps originating from the tympanic epithelium. The cause is unknown but chronic inflammation may be a factor.

Mitochondrial encephalopathy

Inbuilt errors of mitochondrial metabolism can lead to neurological disorders.

Mitochondrial myopathy

A defect in mitochondrial function leads to decreased exercise tolerance with tachycardia and tachypnoea resulting from severe acidosis. This condition can occasionally cause sudden death.

Mitral stenosis

Narrowing of the mitral valve opening restricts blood flow from the atrium to the ventricle. This condition is usually congenital. Left atrial enlargement and congestive heart failure are common sequelae. There may be other concurrent congenital cardiac disorders present.

Mitral valve disease

Also known as chronic degenerative valvular disease, endocardiosis. This is the most common cause of heart disease in the dog, with up to 75% of dogs with congestive heart failure suffering from this condition. The heart valves become deformed by a myxomatous degeneration leading to regurgitation and congestive heart failure. This condition is likely to be inherited, although other proposed causes include stress, hypertension, hypoxia, infection and endocrine abnormalities. Males, older dogs and smaller breeds are predisposed.

Mitral valve dysplasia

This is the commonest congenital heart disease of cats but uncommon in dogs (in which it represents 8–10% of all congenital heart disease). It is thought to be inherited in some breeds of dog. The lesion consists of a malformation of the mitral valve, the normal function of which is to prevent blood flowing back from the left ventricle to the left atrium. Many

animals with this condition do not show signs, and those that do usually demonstrate exercise intolerance and congestive heart failure.

Mucocutaneous pyoderma

A dermatological condition causing crusting of the mucocutaneous junctions.

Mucopolysaccharidosis (including types IIIA, IIIB, IV and VI)

This group of conditions results from inherited chromosomal abnormalities and leads to a metabolic bone disease. There are a number of sub-categories. Type I disease causes a large broad head which may be associated with ocular and cardiac abnormalities. Pectus excavatum, fusion of the cervical vertebrae and hip subluxations also occur. Type III, divided into subtypes A and B, causes neurological signs. Type IV disease causes gross physical abnormalities of the skeleton, including a severe general laxity and luxation of the joints. Type VI disease causes dwarfism and skeletal, neurological and retinal abnormalities. There is no treatment, although some affected animals may have an acceptable quality of life.

Multifocal chorioretinitis (Borzoi chorioretinopathy)

A clinically inapparent, non-progressive chorioretinitis described in the Borzoi. Lesions may be uni- or bilateral and consist of focal retinal oedema which becomes pigmented and hyper-reflective later.

Multifocal retinopathy

Multifocal grey lesions develop initially in the outer retina, followed by the development of areas of serous retinal detachment more centrally. No significant progression with time and no loss of vision.

Multiple autoimmune diseases syndrome

A disorder similar to autoimmune polyendocrine syndrome has been reported in Italian Greyhounds.

Multiple collagenous naevi (nodular dermatofibrosis)

Cutaneous hamartomas, non-neoplastic lumps formed by normal parts of an organ that are arranged erroneously.

Multiple drug sensitivity

A mutation of the gene *MDR1* (*ABCB1*) affects the P-glycoprotein drug transport pump. Dogs with this mutation can be prone to drug toxicity including antiparasitic and chemotherapy drugs.

Multiple ocular defects

Several congenital defects present in the same eye.

Multisystem neuronal degeneration

A slowly progressive degenerative disease. Diffuse neuronal loss throughout the subcortical, brainstem and cerebellar nuclei results in signs, including loss of recognition of the owner, apathy, hyperactivity, hypersexuality and aggression.

Mural folliculitis and parakeratosis

Seen in Labrador Retrievers, this is a skin condition in which cracks and fissures develop in the nasal planum.

Muscle cramping ('Scottie cramp')

An inherited disorder of Scottish Terriers. Affected dogs are normal at rest, but exercise may provoke muscle spasms which in the mildest form of the disease appear as pelvic limb stiffness. Severe attacks cause rigidity of all muscles, including facial muscles, causing the dog to fall over into a tightly curled ball. Consciousness is maintained and the animal makes a spontaneous recovery. The cause is unknown but it is believed to be a disorder of central nervous system neurotransmitters. A similar condition has been reported in Dalmatians and Norwich Terriers.

Muscular dystrophy – X-linked

A sex-linked muscular dystrophy of Golden Retriever found in the USA has been found to be similar to Duchenne muscular dystrophy of humans. Clinical signs include exercise intolerance, gait abnormalities, trismus and occasionally cardiac involvement. Creatinine kinase (CK) is massively elevated on biochemistry.

Muscular dystrophy of pharyngeal and oesophageal muscles causing dysphagia

Atrophy of oesophageal and pharyngeal muscles resulting in difficulty swallowing.

Muscular stiffness of Labrador Retrievers

A movement disorder of young Labradors causing extreme muscle stiffness. An X-linked hereditary disease was suspected.

Mycobacterial infections

Mycobacteria, including *M. tuberculosis* and *M. avium*, are aerobic, acid-fast bacteria, with each species having a variation in host affinity and disease potential. The bacterium causes progressive systemic signs, with granulomatous masses affecting organs such as lymph nodes, lungs and liver.

***Mycobacterium avium* complex infection**

Mycobacterium avium complex has a worldwide distribution in soil and water, but disease is rare. The organisms cause extensive granuloma formation, and the site of formation determines the signs. The bowel, spleen, liver and mesenteric lymph nodes are usually involved in dogs, while regional lymph nodes, skin and gastrointestinal tract are commonly involved in cats.

Myocardial infarction (MI)

A clot affecting the blood supply to the heart causing cardiac damage.

Myokymia and neuromyotonia

Myokymia involves a worm-like writhing of muscles, while neuromyotonia involves spontaneous muscle activity from hyperexcitable peripheral nerves.

Myopathy

Muscle disease characterized by muscle weakness and atrophy (wasting). Usually progressive in nature.

Myopia (short-sightedness)

Loss of ability to focus eyesight on long distance. Vision is blurred because images are focused in front of the retina.

Myositis

Inflammatory muscle disease can be generalized, such as immune-mediated polymyositis, or localized, such as with masticatory myositis. Affected muscles may swell in the early stages and atrophy in later stages, and signs include pain and joint stiffness.

Myotonia

Clinical signs of this condition include excess muscle mass, stiff gait after rest and collapse. Dyspnoea may be seen if the respiratory muscles are involved.

Narcolepsy-cataplexy

Narcolepsy is characterized by excessive sleepiness at inappropriate times, while cataplexy is acute flaccid paralysis from which the animal makes a complete recovery after a few seconds to several minutes. In dogs, cataplexy seems to be the more common, often associated with excitement, e.g., eating or playing.

Nasal cavity tumours

The most common nasal cavity tumours diagnosed in the dog are carcinomas (in particular adenocarcinomas). Other types include sarcomas (fibrosarcoma, chondrosarcoma or osteosarcoma), lymphoma and melanoma. Most are malignant, causing local invasion and progressive destruction, but are slow to metastasize. Doliocephalic dogs, particularly of large and medium size, are reported to be at increased risk; disease in these breeds may be associated with pollutants, e.g., passive smoking or living in urban environments. In the cat, lymphoma is the most common nasal cavity tumour, followed by adenocarcinoma and squamous cell carcinoma.

Nasopharyngeal turbinates

The presence of nasal turbinates protruding caudally from the choanae into the nasopharynx of dogs and cats. May be a component contributor to brachycephalic obstructive airway syndrome.

Necrotizing meningoencephalitis

Usually fatal, mean survival of 3 months. See also *Meningoencephalitis*.

Neonatal cerebellar ataxia

Causes neurological signs such as ataxia and tremor from the age of 2 weeks.

Neonatal encephalopathy with seizures (NEWS)

Puppies affected with this condition are small and weak and often die in the first week of life. Those surviving longer show tremors, ataxia and seizures. None have been reported to survive past 7 weeks of age.

Neuroaxonal dystrophy

A degenerative central nervous system disorder of unknown cause, seen primarily in Rottweilers. Pathological findings include swellings of the distal axons within the central nervous system and cerebellar atrophy. Signs include ataxia,

hypermetria and intention tremors which may be slowly progressive over several years.

Neuronal ceroid lipofuscinosis

This lysosomal storage disease causes signs including reduction in vision, ataxia, tremors, seizures and death. Clinical signs may not appear until adulthood.

Nodular granulomatous episclerokeratitis (proliferative keratoconjunctivitis)

Single or multiple, raised fleshy masses originating at the limbus and invading the cornea. Involvement of the nictitating membrane may occur. Usually a bilateral condition. The lesions are inflammatory and respond to prednisolone and azathioprine.

Noise sensitivity

Some breeds are more prone to hypersensitivity to loud noises.

Non-accidental injury (NAI, battered pet)

This refers to deliberate injury and/or abuse, which may be inflicted by a caregiver, someone else in the household, or someone outside the household.

Non-spherocytic haemolytic anaemia

This condition is due to a defect in the calcium pump system. Chronic haemolysis may lead to myelofibrosis.

Norwegian Elkhound chondrodysplasia

Dwarfism occurring as a result of a disturbance of endochondral ossification.

NT-ProBNP measurement

This peptide is increased with ventricular stretch and is a marker for heart disease.

Ocular melanosis (abnormal pigment deposition)

Large numbers of melanocytes infiltrate the iridocorneal angle, episclera, choroid and iris, predisposing to glaucoma (melanocytic glaucoma).

Oculocutaneous albinism

Albinism of the eyes and skin. May be associated with other ocular abnormalities.

Oculoskeletal dysplasia

This condition is inherited as an autosomal recessive, and causes clinical signs of shortened

limbs and deviated joints, associated with ocular signs such as cataracts and retinal detachment.

Oesophageal and gastric foreign bodies

Some breeds are predisposed to obstruction of the oesophagus or stomach caused by ingestion of foreign bodies, such as toys, household items or bones, leading to signs of vomiting and regurgitation, and possibly resulting in rupture and mediastinitis or peritonitis. See also *Gastrointestinal foreign bodies*.

Olecranon fracture

A fracture of the proximal part of the ulna.

Optic nerve hypoplasia

A congenitally small optic disc with reduced numbers of optic nerve axons and visual impairment.

Oral epulis

A swelling on the gingival (gum) margin, which may be neoplastic. Classified into four types: fibromatous, ossifying, acanthomatous and giant cell.

Oral fibrosarcoma

A malignant tumour of the mouth.

Oral neoplasia

A number of cancers can affect the mouth, including squamous cell carcinoma, melanoma and lymphoma.

Oropharyngeal neoplasia

The most common canine oropharyngeal tumours are squamous cell carcinomas, malignant melanomas, epulides and fibrosarcomas. The most common feline oropharyngeal tumours are squamous cell carcinomas and fibrosarcomas. See also *Melanoma*.

Oropharyngeal penetrating injury

Penetrations of the oropharynx can occur because of foreign body ingestion or 'stick injuries' involving penetration caused by a thrown or retrieved stick. These can lead to severe and difficult-to-treat infections.

Osteochondrodysplasia

A disorder of development of the bone and cartilage.

Osteochondrosis

Including osteochondrosis of the elbow, hock, sacrum, shoulder and stifle. Osteochondrosis is characterized by abnormal development of the cartilage in the physal and epiphysal sites. Common regions for osteochondrosis are the caudal humeral head, the medial condyle of the humerus, the medial coronoid process of the ulna, the anconeal process of the elbow, the lateral and medial condyles of the stifle and the medial ridge of the talus. Lumbosacral osteochondrosis causes signs of cauda equina syndrome.

Osteogenesis imperfecta

This group of inherited diseases causes osteopenia and increased bone fragility. The underlying defect is probably in collagen formation. The condition is rare, and the exact mode of inheritance is unknown. Cases may present with a history of multiple fractures following little or no trauma.

Osteosarcoma

Osteosarcoma is the most common of the malignant primary bone tumours in the dog, representing 80–90% of bone tumours in large dogs. It is rapid in growth and highly invasive and destructive. Osteosarcoma of the appendicular skeleton of dogs is highly malignant and metastasizes early (commonly to the lungs). In common with other primary bone tumours, appendicular osteosarcoma is more common in large- and giant-breed dogs. They generally occur in older dogs, but those affecting giant breeds may be seen at an earlier age. Osteosarcoma of the axial skeleton (including the skull) is generally considered less malignant. Osteosarcoma in the cat is also less aggressive.

Otitis externa

Inflammation/infection of the external ear canal, causing signs of pain, discharge and irritation. Causes include underlying skin disease and foreign bodies.

Otitis media

Inflammation/infection of the middle ear, which can lead to vestibular signs and pain. May occur due to haematogenous spread of bacteria or local extension from otitis externa.

Overweight/obesity

Some breeds are prone to developing an increased body condition, often related to an abnormally large appetite. Overall, 20–50% of dogs are obese, but classifying obesity is subjective. A genetic basis for obesity has been described.

Pain perception and fear memory resilience

An ability to overcome pain to maintain sufficient focus and continue to work is a strong asset for a working dog. Dogs that are resilient have an enhanced chance of working success especially in challenging environments such as the Australian outback.

Palate agenesis, anotia and polydactyly

A familial condition in which the external ear and palate fail to form, associated with extra toes.

Palpebral neoplasia

Cancer of the eyelids.

Pancreatitis

Inflammation of the pancreas which may be acute or chronic and results in lethargy, abdominal pain and vomiting. The inciting causes are often unknown, but it can be associated with hyperlipoproteinaemia, certain drugs, trauma or pancreatic ischaemia, and it may follow a high-fat meal.

Panosteitis (enostosis, eosinophilic panosteitis)

This condition is fairly common, affecting young dogs aged 6–18 months. Acute intermittent lameness affecting one or more limbs is seen, often associated with pyrexia. A viral aetiology is suspected. The condition is usually self-limiting.

Paroxysmal dyskinesia

Intermittent movement disorders characterized by disturbed movement, with the animal being normal between episodes.

Parvovirus enteritis

An infectious viral disease which most commonly causes a severe gastroenteritis; however, it may also manifest as acute myocarditis or neonatal mortality.

Patellar luxation

Usually presents as an intermittent lameness, although in bilateral cases it may present as a hindlimb gait abnormality. It is usually seen from 6 months of age, although in some cases it may not cause clinical signs until the animal is older.

Patent ductus arteriosus

The ductus arteriosus carries blood from the pulmonary artery to the aorta in the fetus to bypass the lung, which is not in use in utero. The ductus arteriosus normally closes within the first week after birth. Failure of the vessel to close leads to a patent ductus arteriosus. Females are at a higher risk of developing the condition than males. Signs range from none, to congestive heart failure and poor body condition, to weakness, collapse and seizures. Congenital heart disease is uncommon in dogs, but patent ductus arteriosus is one of the more common types (third most common in one study, representing 20.9% of diagnosed cases). Previous studies have suggested that patent ductus arteriosus is the most common form of congenital heart disease.

Pattern baldness (follicular dysplasia, bald thigh syndrome)

A disease causing baldness of the lateral and caudal thighs. An endocrinopathy may underlie this condition.

Pectinate ligament dysplasia (PLD)

Abnormal development of the iridocorneal filtration angle which may predispose to closed-angle glaucoma later in life.

Pectus excavatum (funnel chest)

Ventral thoracic wall deformity.

Pelger–Huet anomaly

This anomaly involves a decreased segmentation of granulocyte nuclei. There does not seem to be an increased predisposition to infection in these cases.

Pemphigus foliaceus

This is probably the most common autoimmune skin condition of dogs and cats. Clinical signs of crusting and pustules usually start on the face and ears and progress to become multifocal or generalized. Secondary bacterial infection is common.

Perianal (hepatoid) gland adenomas

These are benign tumours that arise from the modified sebaceous glands of the perianal area. They appear as well-circumscribed raised lesions which may ulcerate.

Perianal adenocarcinomas

These are malignant and difficult to distinguish from their benign counterparts in gross appearance. They are less common than perianal gland adenomas.

Pericardial effusion

This condition involves a build-up of fluid between the heart wall and the pericardium. It can be caused by tumours of the heart, but often has no known single cause ('benign' or 'idiopathic' pericardial effusion). The idiopathic form is poorly understood, and it is not known whether it is inherited. However, it does tend to affect large- and giant-breed dogs. Signs are precipitated by the inability of the heart to fill properly because of restriction caused by the fluid-filled pericardial sac. Chronic cases exhibit weight loss, ascites and dyspnoea resulting from pleural effusion. Acute cases may show rapidly progressing weakness, collapse and death.

Perineal herniation

This relatively common condition presents as a swelling in the perineal region or as a defect palpable per rectum. Urinary bladder retroflexion with associated metabolic complications is seen in 20% of affected dogs.

Perinuclear antineutrophilic cytoplasmic autoantibodies (pANCA)

These autoantibodies are primarily associated with intestinal inflammation. Levels are often high in healthy Soft-Coated Wheaten Terriers, especially if they are related to individuals with protein-losing nephropathy or enteropathy.

Periodontal disease

Disease of the gingiva and periodontal tissue which can lead to infection and tooth loss.

Peritoneopericardial diaphragmatic hernia (PPDH)

This uncommon congenital disease is seen in only about 0.5% of cases of congenital heart disease. It involves a continuation of the pericardium with the peritoneum, and often abdominal

viscera are found within the pericardial sac. It may be asymptomatic, or may cause signs of respiratory distress, vomiting and colic.

Persistent hyaloid remnants

Remains of the hyaloid artery may persist into adulthood. This may not cause clinical problems, but can lead to cataract formation.

Persistent hyperplastic primary vitreous (PHPV)

A congenital condition in which there is abnormal development and regression of the hyaloid system and primary vitreous. It is often associated with persistent hyperplastic tunica vasculosa lentis (PHTVL), in which there is persistence of an embryonic vascular system attached to the posterior lens capsule. The condition is rare, but is seen more frequently in Doberman Pinschers and Staffordshire Bull Terriers. The condition varies in severity. In its most severe form, it is associated with microphthalmia and other ocular defects.

Persistent hyperplastic tunica vasculosa lentis

A congenital ocular disease.

Persistent pupillary membranes (PPM, membrana pupillaris persistens)

Persistent pupillary membranes are uveal remnants which fail to regress normally in the first 6 weeks of life and persist in the anterior chamber (either unilaterally or bilaterally). Strands which bridge from iris to iris are generally of no clinical significance; however, iris-to-cornea or iris-to-lens strands may cause focal corneal and lenticular opacities, respectively. PPMs are a common finding, but severe visual impairment is rare.

Pes varus

This limb deformity is seen at 5–6 months of age, and may be related to an underlying tibial dysplasia.

Phosphofructokinase deficiency

Haemolytic crises and exertional myopathy are seen with this relatively common enzyme deficiency. Exercise intolerance is often seen.

Photoreceptor dysplasia

One of the diseases that make up progressive retinal atrophy.

Pigmentary uveitis

Inflammation of the iris and ciliary body associated with abnormal pigment deposition. Seen most commonly in the Golden Retriever and often seen in conjunction with iris cysts. Cataract and glaucoma are common sequelae.

Pilomatrixoma

An uncommon benign tumour of the hair follicle. Pilomatrixoma presents as a solitary, firm mass in the dermis or subcutis, with ulceration of the overlying epidermis. They usually occur over the neck, back and tail of dogs. They are rare in cats.

Pituitary dwarfism (hyposomatotropism)

Pituitary dwarfism results from a failure of growth hormone secretion in an immature animal. The most striking abnormality is a failure to grow, animals remaining of small stature. There may be a wide range of other clinical signs such as hair and coat changes and changes in dentition, or concurrent failure of other pituitary hormones (panhypopituitarism). The condition is most commonly seen in the German Shepherd Dog but has been identified in several other breeds.

Plasma cell infiltration of the nictitating membrane (plasmoma)

Bilateral plasma cell infiltration of the nictitating membrane resulting in follicle formation and depigmentation. Often associated with chronic superficial keratitis (pannus).

Platelet disorder

Abnormality of platelet function which is associated with blood clotting.

Pneumocystis carinii infection

A predisposition to respiratory disease caused by *Pneumocystis carinii* is suspected to be promoted by an underlying immunodeficiency, but the exact nature of this, and inheritance of the condition, is not known. However, IgG levels are lower in affected dogs than in control animals.

Polyarthritis

Canine idiopathic polyarthritis is the most common form of immune-mediated arthropathy. Approximately 25% of cases are associated with chronic infection remote from the joint, 15% are associated with gastrointestinal disease, and another subset is associated with

neoplasia remote from the joints. In the other cases, which account for about 50%, there is no other pathology or underlying aetiology detected.

Polycystic kidney disease (PKD)

In this disorder, large portions of the renal parenchyma are replaced by multiple cysts. Both kidneys are generally involved and in some cases cysts are also found in the liver. Kidneys may be palpably enlarged and irregular, and the diagnosis can be confirmed by ultrasound. The condition progresses to renal failure at a variable rate. Renal cysts may also be seen in cases of renal dysplasia or neoplasia.

Polycystic liver disease

Fluid-filled cysts in the liver parenchyma. Liver function may or may not be affected. Often seen in conjunction with polycystic kidney disease.

Polydactyly

Several breeds show a tendency to have more than the typical number of digits (polydactyly) or fused digits (syndactyly). These abnormalities do not generally cause severe clinical problems. In some breeds, an extra digit or claw is considered 'normal'.

Polymyopathy

Generalized muscle disease.

Polyneuropathy

Polyneuropathies can cause signs of weakness, ataxia and sensory deficits.

Polyneuropathy with ocular abnormalities and neuronal vacuolation (POANV)

A juvenile-onset polyneuropathy which is also associated with microphthalmia and cataracts.

Polyodontia

Additional teeth, which may be seen in temporary or permanent dentition. Incisors and premolars are most commonly affected.

Portal vein hypoplasia

Previously known as hepatic microvascular dysplasia. Describes a microscopic histopathological vascular anomaly of dogs and cats that is often present in conjunction with a macroscopic portosystemic shunt (PSS). Clinical signs that may be seen with portal vein hypoplasia are

suggestive of a PSS, although some dogs may be asymptomatic. Signs include hepatoenkephalopathy, vomiting, diarrhoea, urinary tract signs associated with ammonium biurate urolithiasis, stunted growth, and prolonged recovery from anaesthesia.

Postoperative bleeding

It has been noted that retired Greyhounds bleed postoperatively more frequently than expected, and a clotting disorder is suspected. However, a disorder of primary or secondary haemostasis has not been identified.

Primary brain tumour

Primary brain tumours are derived from tissues of the nervous system, including nerve cells, glial cells, meninges and neuroepithelial cells. They are generally solitary, and most cases will present with signs of a space-occupying lesion in the brain, the specific signs varying with the location. Meningiomas and gliomas are the most common primary brain tumours in dogs. Meningiomas are the most common primary brain tumours in the cat and may be single or multiple in this species. Meningiomas are generally histologically benign, but can be locally invasive, especially in dogs. Choroid plexus neoplasms are tumours derived from the choroid plexus (the plexus of cells producing cerebrospinal fluid in the brain).

Primary ciliary dyskinesia

In this condition, the mechanism for removing mucus from the airways is defective, leading to respiratory infections. Other conditions associated with defective ciliary function include loss of hearing and loss of sperm motility, with consequent infertility.

Primary hepatitis

Primary hepatitis covers the most frequently occurring group of liver diseases in dogs. It comprises all inflammatory hepatic diseases that are not characterized by the non-specific changes that are described as non-specific reactive hepatitis. The diagnosis of hepatitis in dogs is mainly based on histologic morphology. Primary hepatitis spans a wide range of aetiologies including disease caused by microorganisms, toxins and drugs, immune-mediated reactions and breed-associated metabolic errors.

Primary hyperlipidaemia (primary hypertriglyceridaemia, idiopathic hypertriglyceridaemia)

This familial condition can cause multisystemic signs such as abdominal pain, seizures and pancreatitis.

Primary hyperoxaluria

An autosomal recessive disorder of glyoxylate metabolism which predisposes to oxalate urolithiasis.

Primary hyperparathyroidism

Primary hyperparathyroidism is usually the result of a functional parathyroid adenoma. The excess of parathyroid hormone results in hypercalcaemia. The Keeshond seems particularly susceptible to the condition.

Primary hypoparathyroidism

This is an uncommon condition in which lymphocytic plasmacytic destruction of the parathyroid glands results in a deficiency of parathyroid hormone and hypocalcaemia.

Primary lens luxation (PLL)

See *Lens luxation*.

Primary orthostatic tremor

This disease is characterized by a severe tremor present when standing, i.e. abolished by walking or sitting.

Primary seborrhoea

This is an inherited disorder of keratinization and cornification which has an early age of onset. Clinical signs include flaking, scaling, crusting and greasy, smelly skin. The claws may also be affected. Secondary bacterial and fungal infections are common.

Primary secretory otitis media

Cavalier King Charles Spaniels are prone to getting a build-up of secretions in the middle ear, which can lead to signs of middle ear disease.

Primary splenic torsion

Twisting of the spleen. Can be acute, with signs including collapse and shock, or may be vague and intermittent in the chronic form.

Progressive axonopathy

An inherited neuropathy causing progressive pelvic limb ataxia from 2–3 months of age.

Progressive degenerative polyneuropathy

A progressive degenerative polyneuropathy with a distal axonal distribution.

Progressive retinal atrophy (PRA)

Degeneration of the retinal cells. An autosomal recessive inheritance is suspected in most breeds. Different breeds are affected at different ages by different types of PRA. However, all cases are bilateral and progress to blindness. In most cases, the earliest clinical sign is night blindness, with day vision being lost at a variable time later. Ophthalmoscopically there is attenuation of retinal vessels and tapetal hyper-reflectivity. In the later stages, the condition is often accompanied by cataracts.

Progressive rod–cone degeneration (PRCD)

Progressive rod–cone degeneration is an autosomal recessive photoreceptor degeneration of late onset in dogs. It results in complete blindness in almost every case, and is one of several inherited diseases designated as progressive retinal atrophy (PRA). The disease is homologous to human retinitis pigmentosa (RP).

Prolapse of the gland of the nictitating membrane ('cherry eye')

Prolapse of the tear gland normally located behind the nictitating membrane results in exposure and irritation of the gland. It is usually first seen in young dogs, less than 2 years of age.

Prolonged activated partial thromboplastin time (aPTT)

aPTT is a test to assess the clotting ability of the blood, and prolonged test results are associated with reduced ability to clot.

Pronounced eosinophilic response

Immune-mediated, allergic, parasitic and neoplastic diseases can cause an increase in eosinophils.

Proptosis

Expulsion of the globe from the orbit with entrapment of the eyelids behind the eyeball.

Prostate disease

Common diseases of the prostate include benign prostatic hyperplasia, prostatitis and neoplasia.

Prostate neoplasia – including prostatic carcinoma

Prostatic carcinoma is very invasive locally and metastasizes early to the sublumbar lymph nodes, pelvic bones, lumbar vertebrae, and further afield to the lungs. In contrast to the disease in humans, canine prostatic carcinoma is not androgen-dependent, and several studies have suggested an increased risk in castrated dogs.

Protothecosis

The causal organisms of this condition, *Prototheca* spp., are ubiquitous algae. Infection is rare. Gastrointestinal, ocular and nervous signs are more common than dermatological signs.

Pseudohyperkalaemia

False readings of potassium levels, especially in certain breeds with intra-erythrocyte potassium.

Pulmonary artery dissection

Dissection of major elastic arteries such as the aorta or pulmonary artery involves a tear in the tunica intima and subsequent infiltration of blood into the vessel wall. Dissection in animals may be associated with systemic hypertension, elastin dysplasia, aortic aneurysm, with uncorrected patent ductus arteriosus or secondary to tumour infiltration.

Pulmonary carcinoma

A malignant lung tumour.

Pulmonic stenosis

This is a congenital condition, affecting 20% of dogs diagnosed with congenital heart disease. The condition may be asymptomatic, or may cause signs of syncope and right-sided congestive heart failure.

Pyometra (cystic endometrial hyperplasia–pyometra complex)

Accumulation of purulent fluid in the uterus, usually in dioestrus. Clinical signs include inappetence, polyuria/polydipsia, vomiting and a vulval discharge if the cervix is open.

Pyothorax

Purulent infection in the thoracic cavity. This can be caused by haematogenous spread of bacteria or penetrating foreign bodies, though in many cases a cause is not identified. Can cause fever, dyspnoea and sepsis.

Pyotraumatic folliculitis (acute moist dermatitis, hot spot, wet eczema, summer sores)

Acute inflammation of the skin, which may be triggered by a stimulus such as a flea bite, but is perpetuated by self-trauma. This can be acutely painful.

Pyruvate dehydrogenase phosphatase 1 deficiency

This rare inherited condition causes exercise intolerance, and may shorten life expectancy owing to heart and lung problems.

Pyruvate kinase deficiency

In this condition of red blood cells, the affected cells lose the ability to retain their normal shape and have a reduced affinity for oxygen and a shortened lifespan. Genetic carriers are asymptomatic. Onset and severity of symptoms are variable.

Radial carpal bone fracture

Fracture of one of the bones of the carpus.

Raised serum alkaline phosphatase

Alkaline phosphatase (ALP) is an enzyme found in many tissues, including bone and liver, as well as in corticosteroid-induced isoenzymes. ALP is a marker for liver disease, but can also be elevated in young animals, bone disease and neoplasia.

Rectal prolapse (part of Manx syndrome)

See *Manx syndrome*.

Red cell microcytosis

Small red blood cells may be normal in some breeds.

Renal amyloidosis

Amyloidosis results from the deposition of an insoluble fibrillar protein (amyloid) in a variety of organs, resulting in their dysfunction. Reactive systemic amyloidosis is a systemic syndrome in which amyloid deposition can be found in many organs. Chronic inflammatory or neoplastic disease may predispose to systemic amyloidosis in other breeds. Amyloid deposits in the kidney lead to progressive renal dysfunction. In most cases there is glomerular involvement, resulting in moderate to severe proteinuria and sometimes nephrotic syndrome. In most breeds, renal amyloidosis is seen in older dogs; however, in

the Shar Pei dog and Abyssinian cat it is seen at an earlier age.

Renal arteriosclerosis

Endothelial lesions are found in the renal arteries of young greyhounds.

Renal calculi

Kidney stones – see *Urolithiasis*.

Renal cystadenocarcinomas

This is a condition of bilateral and multifocal renal neoplasia which is seen primarily in the German Shepherd Dog. Cases present at 5–11 years of age with anorexia, weight loss and polydipsia. The renal condition is associated with generalized cutaneous nodules and, in females, with multiple uterine leiomyomas.

Renal dysplasia

The term renal dysplasia refers to conditions where there is disorganized development of the renal parenchyma resulting in the persistence of structures inappropriate to the stage of development, e.g., immature glomeruli.

Respiratory distress syndrome (acute respiratory distress syndrome, ARDS)

This condition has been reported in related Dalmatians. Progressive pulmonary failure occurred, leading to death in 3 weeks. No known risk factors for ARDS could be identified.

Restrictive cardiomyopathy (RCM)

A form of heart muscle disease seen most commonly in cats. There is a loss of ability of the myocardium to relax, leading to diastolic dysfunction. Sequelae include heart failure and thromboembolism.

Retinal degeneration

Hereditary retinal degenerations are blinding disorders characterized by dysfunction and death of rod and cone photoreceptor cells of the retina. They are genetically and phenotypically heterogeneous, with differing causative mutations and clinical presentations.

Retinal detachment

Separation of the retina from the underlying tissues resulting in loss of function of the detached portion. May be partial or complete. Usually secondary to other ocular or systemic disease.

Retinal dysplasia

Retinal dysplasia is characterized by disorganized retinal development. There are three categories: multifocal, geographic and complete. More severe forms are associated with significant vision loss. Has various causes but is commonly genetic in origin.

Retinal pigment epithelial dystrophy (RPED, central progressive retinal atrophy)

Abnormal accumulation of pigment within the retina resulting in a progressive retinal degeneration and visual deficiencies. The condition was previously reported most commonly in England, but is now infrequently seen. Similar retinal lesions have been seen in dogs fed vitamin-E deficient diets, and in some breeds RPED lesions have been associated with a deficiency of vitamin E due to a metabolic abnormality.

Retinopathy (rod–cone dystrophy)

Causes day blindness.

Rhinitis/bronchopneumonia syndrome

Leads to transient or chronic cough, dyspnoea and nasal discharge. Some affected dogs have decreased IgA levels.

Rhipicephalus sanguineus

The brown dog tick, which can be a vector for diseases such as ehrlichiosis.

Ruptured chordae tendineae

The chordae tendineae are major components of the atrioventricular valve apparatus. They anchor the mitral valve, and rupture can lead to severe and acute congestive heart failure.

Sacrocaudal fusion

May result from breeding for speed, and is often an incidental finding.

Sacroiliac joint biomechanics

The sacroiliac joint (SIJ) is formed between the surfaces of the sacrum and ilium bones. Its function is weight-bearing and transmission of the propulsion from the pelvic limbs to the spine during locomotion. In a four-legged stance, about 30–40% of body weight is distributed to the pelvic limbs when standing but the hip joint is subjected to forces approximately three times body weight during locomotion.

Salivary gland neoplasia

Tumours of the salivary gland include adenocarcinoma, squamous cell carcinoma and sarcoma.

Salmonellosis

Salmonella spp. are zoonotic bacteria that lead to gastrointestinal signs such as vomiting and diarrhoea as well as weight loss, fever and abdominal pain.

Sarcoma – soft tissue

A malignant tumour arising in non-epithelial tissue.

Sarcoptic mange (scabies, fox mange)

A contagious skin disease caused by the *Sarcoptes* spp. mite, which can lead to severe itching and hair loss.

Seasonal flank alopecia

See *Canine follicular dysplasia*.

Sebaceous adenitis

An autoimmune skin condition that is uncommon in dogs and rare in cats. Different breeds show different manifestations, including scaling, alopecia, hyperkeratosis and occasional systemic signs. Generally the disease is non-pruritic unless there is secondary bacterial infection.

Sebaceous gland tumours

One of the most common skin tumours of the dog, but less common in the cat. They may be single or multiple. There are various histologic types: *sebaceous hyperplasia* (which is not a neoplastic condition but may be the precursor of sebaceous adenomas or epitheliomas), *sebaceous epitheliomas*, *sebaceous adenomas* and *sebaceous adenocarcinomas*. All types appear as wart-like lesions which may be ulcerated or melanocytic, regardless of histopathological type. With the exception of adenocarcinomas, which may (rarely) metastasize, sebaceous gland tumours are generally benign in behaviour.

Secondary lens luxation

See *Lens luxation*.

Selective IgA deficiency

This condition causes different clinical signs in different breeds, including dermatological disease, respiratory disease such as rhinitis, and gastrointestinal disease such as antibiotic-responsive diarrhoea and anal furunculosis.

Selective malabsorption of cobalamin (vitamin B₁₂)

Malabsorption of vitamin B₁₂ (cobalamin) may occur as a result of an absence of the appropriate receptors in the ileum. Signs include inappetence, failure to grow, non-regenerative anaemia and neutropenia. The condition resembles Imerslund–Gräsbeck syndrome in humans.

Sensory ataxic neuropathy

A slowly progressive neurological disorder of Golden Retrievers which leads to ataxia and dysmetria from the age of 2–8 months.

Sensory neuropathy (acral mutilation syndrome)

Sensory neuropathies have been seen in a number of breeds. In Pointers, signs of self-mutilation associated with loss of pain sensation predominate, whereas in Dachshunds loss of proprioception and ataxia may be seen. In Boxers, the condition is termed *progressive axonopathy* and is characterized by pelvic limb hyporeflexia, hypotonia and proprioceptive loss. In some cases acral mutilation will result. See also *Acral mutilation and analgesia*.

Sesamoid disease

Diseases of these small bones of the metacarpal/metatarso-phalangeal joints include degeneration and fracture leading to pain and crepitus.

Severe combined immunodeficiency

Severe combined immunodeficiency leads to a marked susceptibility to infection and frequently results in early mortality.

Severe subacute necrotizing encephalopathy (Leigh-like syndrome)

This condition causes rapidly progressive vestibular signs at 6–8 weeks of age.

Sex reversal

See *XX sex reversal*.

Shaker dog disease

This condition has been most commonly observed in dogs with white hair coats, particularly Maltese and West Highland White Terriers. Dogs develop a fine whole-body tremor which may worsen with excitement and stress. Other signs may include nystagmus, menace deficits,

proprioceptive deficits and seizures. There may be an underlying mild lymphocytic encephalitis, and affected animals are usually responsive to immunosuppressive doses of corticosteroids with benzodiazepines.

Sialocoele

A swelling of the salivary glands or their ducts causing a subcutaneous or submucosal cavity containing mucinous saliva. It may be idiopathic or secondary to salivary calculi, infections and foreign bodies.

Short tail (bobtail)

Tail length is often a breed-specific trait. It depends on the number of the caudal vertebrae, which can vary significantly between individual dogs and breeds. A C189G mutation in exon 1 of the T-box transcription factor T gene has been identified as a genetic cause of short-tail phenotype. This mutation is thought to cause embryonic lethality in homozygotes and is associated with reduced litter size.

Sialorrhoea with spirocercosis

Infection with the nematode *Spirocerca* spp., which is mainly transmitted by dung beetles, can affect the salivary glands. See also *Spirocercosis*.

Sick sinus syndrome (bradycardia–tachycardia syndrome)

This dysrhythmia often involves periods of bradycardia and tachycardia, leading to syncope.

Sinonasal neoplasia

Cancer of the nose and associated nasal sinuses.

Slipped capital femoral epiphysis (SCFE)

A Salter–Harris fracture of the femur which can occur with minimal trauma.

Small intestinal volvulus

A rare condition where the small intestine rotates about its mesenteric axis, resulting in intestinal obstruction and ischaemia. The condition is rapidly fatal. Large-breed dogs are predisposed.

Solid intraocular xanthogranuloma

A solid intraocular mass comprised of macrophages. Hyperlipidaemia may be a predisposing factor.

Spiculosis

In this condition, multiple bone spicules are found, particularly over the lateral hock.

Spina bifida

This is a developmental defect resulting from the failure of the two halves of the dorsal spinous processes to fuse, most commonly in the lumbar spine. Protrusion of the spinal cord or meninges may result in signs including pelvic limb ataxia, paresis and urinary or faecal incontinence. If no protrusion occurs the condition is termed *spina bifida occulta*.

Spinal arachnoid pseudocysts

Developmental lesions of the spine leading to ataxia and hypermetria.

Spinal dysraphism

This is a congenital malformation of the spinal cord resulting in a wide-based stance and bunny-hopping gait of the hindlimbs. It may be associated with hemivertebrae or spina bifida. The condition is non-progressive.

Spinal muscular atrophy

A condition similar to motor neurone disease in humans, where premature degeneration of motor neurone cell bodies in the spinal cord and brainstem result in generalized weakness which may progress to muscular atrophy and tetraparesis/tetraplegia.

Spinocerebellar ataxia (hereditary ataxia)

Early-onset degeneration of the area of the spinal cord which carries messages to the cerebellum results in the development of cerebellar ataxia as early as 2–6 months of age and progresses. Myokymia (involuntary twitching of muscles) develops and may progress to muscle spasms. Some cases develop epileptic seizures. Most dogs are euthanized young due to poor quality of life.

Spinocerebellar degeneration (late onset)

This is a late-onset degeneration of the spinocerebellar tracts, resulting in ataxia in older dogs.

Spirocercosis

Infection with *Spirocerca lupi*. *S. lupi* is carried by dung beetles and is most prevalent in warm climates. Infection causes nodules to develop in the oesophagus, stomach and aorta resulting in signs including regurgitation, vomiting and breathing

difficulties. Neoplastic transformation of the nodules to osteosarcoma or fibrosarcoma may occur. See also *Sialorrhoea with spirocercosis*.

Splenic disease

Diseases of the spleen include masses, torsion and inflammation.

Splenic masses

Masses of the spleen include lymphosarcoma, fibrosarcoma and haemangiosarcoma.

Splenic torsion

Splenic torsion results from the twisting of the spleen on itself, preventing normal blood drainage and resulting in congestion and swelling. It is mostly seen in large- and giant-breed dogs.

Spondylocostal dysostosis

A rare inherited growth disorder characterized by severe abnormalities of the spine and ribs.

Spondylosis deformans

Also known as ankylosing spondylitis. Bony spurs (osteophytes) form around the margins of the vertebral endplates. They become more common with age, and are often apparent radiographically, but are rarely of clinical significance.

Spontaneous chronic corneal epithelial defects (refractory corneal ulceration, indolent ulcers)

Slow-healing, superficial corneal ulcers which may represent a form of corneal epithelial dystrophy. Originally described in the Boxer, but also occurs in other breeds. Usually seen in middle-aged dogs.

Spontaneous pneumothorax

A closed pneumothorax resulting from air leakage from the lungs within the chest cavity, with no apparent history of trauma. It may occur secondary to many diseases, including neoplasia, abscess, pneumonia, parasites, asthma or migrating foreign body (e.g. a grass awn).

Sporotrichosis

A fungal infection caused by *Sporothrix schenckii*. The organism is found worldwide in soils rich in decaying matter. Infection results in dermal and subcutaneous nodules, and may result in disseminated disease with multiple organ involvement, more commonly in cats than in dogs.

Squamous cell carcinoma

One of the more common malignant skin tumours in the dog, and the most common in the cat. Dogs are commonly affected on the head, trunk and perineum, whereas cats are commonly affected on the pinnae, nasal planum and eyelids. Squamous cell carcinoma is locally invasive but the incidence of metastasis is variable. Exposure to ultraviolet (UV) light may predispose to the development of actinic solar dermatitis, which may progress to squamous cell carcinoma.

Squamous cell carcinoma of the cornea is rare but may occur more commonly in brachycephalic breeds and those predisposed to keratoconjunctivitis sicca.

Squamous cell carcinoma is the most common cutaneous tumour of the digit in dogs. It is locally invasive, resulting in bone lysis, and metastasizes more frequently than squamous cell carcinomas found in other cutaneous sites. It occurs most commonly in large-breed dogs with black coats. Some dogs have multiple tumours.

Sterile nodular panniculitis

Inflammation of the subcutaneous fat presenting as deep nodules which may or may not have draining tracts.

Steroid-responsive meningitis–arteritis (SRMA)

An immune-mediated disorder seen in young dogs. Signs include neck pain and fever.

Stillborn kittens

Kittens may be born dead for many reasons including infectious or congenital disorders.

Stomatocytosis

Stomatocytes are bowl-shaped red blood cells. They have increased osmotic fragility and may predispose to haemolytic anaemia.

Stomatocytosis–hypertrophic gastritis

Abnormally shaped red blood cells which have increased osmotic fragility (stomatocytosis) seen with gastric mucosal hypertrophy.

Sudden acquired retinal degeneration (SARD, amaurosis)

This is an idiopathic irreversible loss of vision that occurs over days to months, without an apparent lesion affecting the eye.

Sulfonamide-associated hypersensitivity

Sulfonamides are associated with idiosyncratic drug reactions in some dogs. These reactions occur at normal dose rates and can include skin eruptions, uveitis, keratoconjunctivitis sicca, polyarthropathy, hepatotoxicity, thrombocytopenia, neutropenia and haemolytic anaemia. The widespread use of sulfonamides is therefore limited in dogs, as in humans.

Superficial necrolytic dermatitis

This is a drug reaction to certain shampoos. Cutaneous and systemic signs usually occur within 3 days. Cutaneous signs include papules, plaques and pustules; systemic signs include pyrexia and depression.

Sweat gland tumour

These may be adenomas (of glandular or ductal origin) or adenocarcinomas. They are uncommon in the dog and cat. They may present as small solitary nodules in the dermis and subcutis with or without ulceration. An inflammatory form of adenocarcinoma is poorly circumscribed and more infiltrative. Adenocarcinomas are highly invasive and may metastasize to local and regional lymph nodes, and occasionally to more distant sites, e.g. the lungs.

Swimmer puppy syndrome

Developmental abnormality of neonatal puppies. Affected puppies have splayed legs and are unable to walk. The condition may also be seen in kittens.

Symmetric lupoid onychodystrophy

All claws are shed. They may regrow but are usually brittle and deformed. An immune-mediated pathogenesis is suspected.

Synovial myxoma

Joint tumour in which the joint is filled with myxomatous nodules. Large-breed, middle-aged dogs are predisposed, and the stifle and digit are the most commonly affected joints.

Systemic amyloidosis

Deposition of amyloid (a type of protein) in multiple organs of the body, disrupting function.

Systemic histiocytosis

See *Histiocytic diseases*.

Systemic lupus erythematosus (SLE)

Diagnosis of this uncommon disease is made on the basis of the presence of at least two signs of autoimmune disease, together with high levels of antinuclear antibodies (ANA) (although 10% of cases are ANA-negative). Clinical manifestations include polyarthritis, mucocutaneous lesions, glomerular disease, autoimmune haemolytic anaemia, autoimmune thrombocytopenia and neurological signs.

Tail-chasing

Tail-chasing is a repetitive behaviour in which a dog spins slowly or rapidly in tight circles with or without biting its tail. The potential causes of this behaviour include an acute conflict behaviour only manifested in conflict situations, attention-seeking behaviour, neurological diseases (e.g. seizures), dermatological disease (e.g. flea allergy dermatitis), and compulsive/obsessive-compulsive disorder.

Tapetal degeneration

An inherited condition seen in Beagles, in which there is progressive degeneration of tapetal cells without any effect on vision.

Telangiectasia

Multiple vascular lesions across several organs including the kidneys, subcutaneous tissue, spleen and brain.

Temporomandibular joint dysplasia

Abnormal development of the articular surfaces leads to temporomandibular dysplasia with laxity of the joint. The condition can be seen from 6 months of age. Clinical signs include open mouth locking often associated with yawning, and luxation, which may be chronic. Degenerative joint disease and masticatory muscle wastage may be seen in chronic cases.

Testicular neoplasia

Testicular neoplasia is common in the dog. There are three main tumour types: Sertoli cell tumour, seminoma and interstitial (Leydig) cell tumour. They may be unilateral or bilateral, and two or more tumour types may occur in one or both testes. Neoplasia occurs more commonly in retained than descended testes: an increased risk of 13.6 times has been reported.

Tetralogy of Fallot

Pulmonic stenosis, ventricular septal defect, a dextrapositioned or over-riding aorta and a secondary right ventricular hypertrophy comprise the four parts of this congenital cardiac abnormality.

Thiobarbiturates

Anaesthetic drugs used less commonly nowadays.

Thoracic spinal stenosis

Narrowing of the thoracic spinal canal which results in neurological signs.

Thoracic vertebral malformations

Congenital vertebral malformations are common, especially in small brachycephalic dogs. Hemivertebrae are frequently reported in screw-tailed brachycephalic breeds such as the French bulldog and are assumed to be hereditary. These vertebral malformations are most frequently found in the thoracic vertebral column and can affect single or multiple vertebrae. Despite the potential to cause clinical signs of spinal cord dysfunction, vertebral malformations are frequently not associated with clinical disease.

Thoracic wall abnormalities

Congenital thoracic wall defects which may include pectus excavatum, unilateral thoracic wall concavity and scoliosis.

Thymoma

Thymoma is a tumour of the epithelial cells of the thymus gland, which is situated in the cranial mediastinum. It is uncommon in both dogs and cats. Thymomas are generally benign and slow-growing. Signs relate to the presence of a cranial mediastinal mass and may vary: cough, dyspnoea, regurgitation and occasionally obstruction of the cranial vena cava, leading to facial and forelimb oedema ('precaval syndrome'). Autoimmune conditions such as myasthenia gravis may be associated with thymoma.

Thyroid hormones

Levels of hormones produced by the thyroid gland below the normal reference range may be seen in healthy dogs of certain breeds (notably sight hounds).

Thyroid neoplasia

Most thyroid tumours in the dog are invasive and malignant carcinomas presenting as readily detectable masses in the neck. Adenomas do occur but are usually small and rarely detected during life. Thyroid tumours in the dog are generally non-functional, only 5–20% being functional, producing clinical signs of hyperthyroidism. Up to 30% of cases become hypothyroid as normal thyroid tissue is destroyed by the tumour. The remainder are euthyroid.

Tracheal and laryngeal tumours

Cartilaginous tumours of the larynx and trachea are uncommon in the dog. Arctic breeds seem to be the most commonly affected.

Tracheal collapse

This condition can occur in young dogs with a severe form of the condition, or later in life in those less severely affected. Clinical signs include coughing and inspiratory stridor. A characteristic 'goose-honk' cough may be heard.

Tracheal hypoplasia

Underdevelopment of the trachea can lead to marked respiratory complications.

Trance-like syndrome

During episodes, affected dogs may have a slow gait, aimless repetitive pacing and yet be aware of their surroundings and responsive. It is unclear whether this is a compulsive behaviour or a neurological condition. Dogs are normal between episodes.

Transmissible venereal tumour

Contagious venereal tumour which most commonly affects the external genitalia. Common in tropical and subtropical areas in sexually mature dogs. May also affect the oral and nasal mucosa, rectum and intra-abdominal organs.

Trapped neutrophil syndrome (cyclic haematopoiesis)

In this condition, neutrophil numbers decrease every 12 days. There may also be cyclic decreases in platelet, monocyte and reticulocyte numbers.

Traumatic proptosis

Anterior displacement of the eye from the orbit as a result of blunt trauma. Brachycephalic breeds most commonly affected.

Trichoepithelioma

An uncommon benign tumour of the hair follicle. Trichoepitheliomas present as solitary, firm masses in the dermis or subcutis, often with ulceration of the overlying epidermis. They usually occur over the back, trunk and limbs of dogs.

Tricuspid valve dysplasia

This is a relatively uncommon malformation of the tricuspid valve (3.1% of all congenital heart defects in one study), the consequence of which is to allow blood to flow back from the right ventricle into the right atrium. Affected dogs may have other congenital heart lesions. In some cases affected animals are asymptomatic for many years, while others show progressive heart failure leading to death.

Trigeminal neuropathy

Disease of the trigeminal nerve (cranial nerve V) which causes signs including masticatory muscle atrophy and loss of the palpebral reflex. Trigeminal neuritis is an idiopathic and self-limiting condition.

Triglyceridaemia

Increased levels of circulating triglycerides which, while considered normal after a fatty meal, would be considered abnormal if seen in the fasting state.

Tritrichomonas foetus infection

Tritrichomonas foetus is a parasite that infects the gastrointestinal system of cats, causing diarrhoea which may be accompanied by blood and mucus and persist for many months.

Ulcerative colitis (histiocytic ulcerative colitis)

Chronic inflammatory disease of the colon characterized histologically by an inflammatory infiltrate with large numbers of histiocytes. Ulceration of the mucosa may be a prominent feature, and the disease tends to be refractory to treatment.

Urethral prolapse

Prolapse of the urethral mucosa through the external urethral orifice may occur in young male dogs. Brachycephalic breeds may be predisposed.

Urethral sphincter mechanism incompetence

A weak urinary sphincter allows urine leakage, usually when the animal is relaxed and lying down. The condition is the most common cause of urinary incontinence in the adult dog and is most commonly diagnosed in neutered females of medium to large breeds (> 20 kg). In many cases, the condition is responsive to reproductive hormone and sympathomimetic administration.

Urinary incontinence (part of Manx syndrome)

See *Manx syndrome*.

Urinary tract infections (UTI)

Bacterial infection of the urinary tract.

Urolithiasis

The formation of stones (uroliths) anywhere within the urinary tract.

- **Apatite urolithiasis.** These are relatively uncommon uroliths composed of calcium phosphate. They are most commonly seen in patients with primary hyperparathyroidism, hyperadrenocorticism and hypercalciuria.
- **Calcium oxalate urolithiasis.** Calcium oxalate uroliths are commonly found in both dogs and cats. In dogs, hypercalciuria, hypercalcaemia and hyperadrenocorticism predispose to calcium oxalate urolith formation. Dietary risk factors include excessive calcium, excessive oxalic acid, high protein, high sodium, restricted phosphorus, restricted potassium and dry diet formulation. Calcium oxalate uroliths tend to be radiodense, often rough and round or oval in shape. The risk is higher in males and in older dogs. In cats the risk is higher in males and neutered cats.
- **Cystine urolithiasis.** Cystinuria occurs as a result of an (autosomal recessive) inherited defect in cystine transport in the renal tubules. Cystinuria predisposes to cystine urolithiasis. Most cystine uroliths are relatively radiolucent, smooth and oval, and they are more likely to form in acid urine. They represent 1–3% of canine uroliths in the USA. They are uncommon in cats. With the exception of the Newfoundland, almost all cases recorded have been in male dogs.
- **Silica urolithiasis.** Silica uroliths are uncommon. They have a characteristic jackstone appearance, are relatively radiodense and are

less soluble in acid urine. There may be a link between dietary ingredients (notably corn gluten feed and soybean hulls) and silica urolith formation. Most affected dogs are male.

- **Struvite (magnesium ammonium phosphate) urolithiasis.** Struvite uroliths are relatively common in cats and dogs. Previously considered the most common type of urolith, several reports have suggested that calcium oxalate is now more common. They are generally radiodense, smooth, round or faceted stones. Most struvite uroliths are infection-induced in the dog, and since females are predisposed to urinary tract infections this may explain the strong female predisposition seen. Alkaline urine favours their formation.
- **Urate urolithiasis.** Urate uroliths are the third most common urolith in the dog after calcium oxalate and struvite, accounting for approximately 8% of canine uroliths. They are relatively radiolucent and are usually found as multiple, small, smooth, round or oval stones of brown-green colour. They tend to form more in acid urine. Dalmatians are predisposed to the development of urate uroliths because of a reduced capacity to convert uric acid to allantoin in the liver, leading to high levels of urinary uric acid excretion. Animals with hepatic portal vascular anomalies are predisposed to urate uroliths, owing to the reduced hepatic ability to convert ammonia to urea and uric acid to allantoin, leading to increased urinary excretion of these substances.

Urticaria

Sudden development of wheals (hives) as part of an anaphylactic allergic reaction.

Urticaria pigmentosa-like dermatosis

A condition seen in Devon Rex cats. Erythematous papules, some with crusting and pigmentation, develop in a linear pattern on the head, neck, chest and abdomen. May represent an atypical presentation of dermatophytosis or a reaction to allergic skin disease.

Uterine leiomyoma

A form of benign tumour affecting the smooth muscle of the uterus.

Uveal cysts

Cysts of the iris and ciliary body of the eye are usually benign and may be single or multiple, unilateral or bilateral. They may be congenital or acquired, most occurring spontaneously in adult dogs. They are generally benign and of no significance unless large or multiple.

Uveitis

Inflammation of the iris, ciliary body and/or choroid of the eye. Provoking causes include neoplasia, infection, excessive pigment and immune-mediated disease.

Uveodermatologic syndrome (Vogt–Koyanagi–Harada-like syndrome)

Believed to be an immune-mediated disorder similar to Vogt–Koyanagi–Harada syndrome in humans. Melanocytes are targeted by the immune system. Ocular signs include anterior uveitis, uveal depigmentation and retinal damage. Dermatological signs may include vitiligo (depigmentation) of the eyelids, nasal planum and lips.

Vaccine-associated adverse effect

Adverse reactions to vaccines are uncommon, but can range from mild to severe with signs including facial swelling, nausea, shock and anaphylaxis. In Weimaraners, routine vaccinations occasionally cause an acute systemic vasculitis, with gastrointestinal signs, metaphyseal osteopathy and lameness.

Vaccine-associated vasculitis with hypertrophic osteopathy

This vasculitis, which is associated with routine vaccinations, leads to initial gastrointestinal signs, which later develop into lameness and metaphyseal osteopathy.

Vaginal hyperplasia/vaginal prolapse

Vaginal hyperplasia is an exaggerated response of the vaginal mucosa to normal circulating oestrogen during pro-oestrus or oestrus. Vaginal oedema and thickening occurs and may result in a degree of vaginal prolapse. Young large-breed and brachycephalic bitches are most commonly affected.

Vascular ring anomaly – persistent right aortic arch (PRAA)

Vascular ring anomalies are uncommon congenital abnormalities of the major thoracic arteries leading to entrapment of the oesophagus and

clinical signs of regurgitation at or after weaning. Persistent right aortic arch is the most common vascular ring anomaly.

Vascular tumours

Cancers of components of the wall of blood vessels.

Vasculitis

This uncommon condition causes various skin lesions such as wheals, papules, oedema, ulceration and scarring. Various cell types may be involved histologically, but the classification is usually of a type III hypersensitivity reaction. Various triggering factors may be involved, such as vaccinations, arthropod bites and mast cell tumours.

Ventral comedone syndrome

In this condition, comedones form because of pressure and friction on the sternum. Secondary bacterial infection, which may respond to topical antibiotic preparations, is common.

Ventricular arrhythmia and sudden death

Abnormal heart rhythm originating in the ventricles which can result in sudden death without prior signs of heart disease.

Ventricular septal defect

Congenital heart disease that is uncommon; patent ventricular septal defect was the fourth most common type in one study, representing 7.5% of diagnosed cases. This condition involves a hole in the interventricular septum. Defects vary in size but can be very large. Small defects may be asymptomatic, but larger defects can cause congestive heart failure, pulmonary vascular disease and pulmonary hypertension.

Vertebral anomalies

See *Thoracic vertebral malformations*.

Vertebral heart score (VHS)

VHS is a radiographic measurement used to assess heart size independently of the size of the dog. However, some breeds have a higher than normal range of VHS.

Vertebral tumours

Cancers of the vertebral column and its contents.

Vesicular cutaneous lupus erythematosus

A cutaneous ulcerative disease seen in Collie breeds. Aggressive immunosuppressive treatment is indicated.

Vincristine-associated myelosuppression

Vincristine is a vinca-alkaloid antineoplastic drug which may cause bone marrow suppression and reduction of blood cell counts. Border Collies seem to be unusually sensitive to this effect.

Vitamin-A-responsive dermatosis

This condition is characterized by a relatively refractory seborrhoea. There is often a waxy otitis externa. Vitamin A supplementation can lead to complete resolution, but therapy may need to be lifelong.

Vitamin-K-dependent coagulopathy

Clinical signs of this condition include prolonged bleeding after surgery or trauma. It is thought that an autosomal recessive defect in hepatic vitamin K metabolism leads to reduced levels of factors II, VII, IX and X. Affected animals can be detected by demonstrating a prolonged PT and aPTT.

Vitiligo

This is thought to be an autoimmune disease causing depigmentation of the face. The condition is usually seen in young adults.

Vitreous syneresis

Degeneration of the vitreous results in liquefaction. This is a common age-related finding and usually of no clinical significance; however, in some breeds it may occur at an earlier age. Extension of the abnormal vitreous into the anterior chamber may predispose to glaucoma. Rarely, it may predispose to retinal detachment.

Vitreoretinopathy

Liquefaction of the vitreous leads to retinal detachment and retinal tears, predisposing to glaucoma.

von Willebrand's disease (vWD)

This is the commonest inherited disorder of haemostasis in dogs, and is caused by a deficiency in von Willebrand factor (vWF), which is vital for platelet function. There are three recognized types. Type I involves a quantitative reduction in vWF, and the severity of the disease varies with breed; some individuals respond to desmopressin (DDAVP) treatment. Type II disease also involves a quantitative reduction in vWF, but the condition is more severe than type I, and there is no response to DDAVP. Type III disease is characterized by a complete absence of vWF leading to the most severe clinical disease, again with no response to DDAVP. Genetic tests are also available to identify carriers of the gene to allow appropriate selection of breeding stock.

Webbed feet

Webbed feet consist of toes that are connected by a membrane. These are characteristic of breeds that may spend time in aquatic environments.

Wool-sucking (compulsive)

This abnormal behaviour appears to be partly genetic, but environmental factors may also play a role, such as early weaning and small litter size. All cats affected in one study had abnormally intense appetites.

XX sex reversal

Sex reversal refers to the situation where the chromosomal and phenotypic sex do not agree. In XX sex reversal a phenotypic male is chromosomally a female.

Zinc-responsive dermatosis

Two syndromes of this condition are recognized. Syndrome I occurs in diets with sufficient zinc. Erythema, alopecia, crusting, scaling and suppuration are seen around the mouth, chin, eyes and ears. In syndrome II, zinc-deficient diets can lead to hyperkeratotic plaques on the footpads and nasal planum.

References

- Aberdein, D., Munday, J.S., Fairley, R.A. *et al.* (2015) A novel and likely inherited lymphoproliferative disease in British Shorthair Kittens. *Veterinary Pathology*, **52** (6), 1176–1182.
- Abitbol, M., Bossé, P., Thomas, A. *et al.* (2015) A deletion in FOXN1 is associated with a syndrome characterized by congenital hypotrichosis and short life expectancy in Birman cats. *PLoS One*, **10** (3), e0120668.
- Abramson, C.J., Platt, S.R., Jakobs, C. *et al.* (2003) L-2-Hydroxyglutaric aciduria in Staffordshire Bull Terriers. *Journal of Veterinary Internal Medicine*, **17** (4), 551–556.
- Abramson, C.J., Platt, S.R. & Shelton, G.D. (2004) Pyruvate dehydrogenase deficiency in a Sussex Spaniel. *Journal of Small Animal Practice*, **45** (3), 162–165.
- Acland, G.M. & Aguirre, G.D. (1987) Retinal degenerations in the dog: IV. Early retinal degeneration (erd) in Norwegian Elkhounds. *Experimental Eye Research*, **44** (4), 491–521.
- Acland, G.M., Blanton, S.H., Hershfield, B. & Aguirre, G.D. (1994) XLPRA: A canine retinal degeneration inherited as an X-linked trait. *American Journal of Medical Genetics*, **52** (1), 27–33.
- Adam, F.H., German, A.J., McConnell, J.F. *et al.* (2012) Clinical and clinicopathologic abnormalities in young dogs with acquired and congenital portosystemic shunts: 93 cases (2003–2008). *Journal of the American Veterinary Medical Association*, **241** (6), 760–765.
- Adamama-Moraitou, K.K., Patsikas, M.N. & Koutinas, A.F. (2004) Feline lower airway disease: a retrospective study of 22 naturally occurring cases from Greece. *Journal of Feline Medicine and Surgery*, **6**, 227–233.
- Adams, P., Bolus, R., Middleton, S., Moores, A.P. & Grierson, J. (2011) Influence of signalment on developing cranial cruciate rupture in dogs in the UK. *Journal of Small Animal Practice*, **52** (7), 347–352.
- Adams, V.J., Evans, K.M., Sampson, J. & Wood, J.L.N. (2010) Methods and mortality results of a health survey of purebred dogs in the UK. *Journal of Small Animal Practice*, **51** (10), 512–524.
- Adaszek, Ł., Martinez, A.C. & Winiarczyk, S. (2011) The factors affecting the distribution of babesiosis in dogs in Poland. *Veterinary Parasitology*, **181** (2–4), 160–165.
- Adkins, E.A. & Hendrix, D.V.H. (2005) Outcomes of dogs presented for cataract evaluation: a retrospective study. *Journal of the American Animal Hospital Association*, **41** (4), 235–240.
- Adler, J.A., Drobatz, K.J. & Hess, R.S. (2007) Abnormalities of serum electrolyte concentrations in dogs with hypoadrenocorticism. *Journal of Veterinary Internal Medicine*, **21** (6), 1168–1173.
- Affolter, V.K. & Moore, P.F. (2002) Localized and disseminated histiocytic sarcoma of dendritic cell origin in dogs. *Veterinary Pathology*, **39** (1), 74–83.
- Agler, C., Nielsen, D.M., Urkasemsin, G. *et al.* (2014) Canine hereditary ataxia in Old English Sheepdogs and Gordon Setters is associated with a defect in the autophagy gene encoding RAB24. *PLoS Genetics*, **10** (2), e1003991.
- Aguirre, A.L., Center, S.A., Randolph, J.F. *et al.* (2007) Gallbladder disease in Shetland Sheepdogs: 38 cases (1995–2005). *Journal of the American Veterinary Medical Association*, **231** (1), 79–88.
- Aguirre, G.D. & Rubin, L.F. (1975) Rod-cone dysplasia (progressive retinal atrophy) in Irish Setters. *Journal of the American Veterinary Medical Association*, **166** (2), 157–164.
- Aguirre-Hernández, J., Wickström, K. & Sargan, D.R. (2007) The Finnish Lapphund retinal atrophy locus to the centromeric region of CFA9. *BMC Veterinary Research*, **10** (3), 14.
- Aherne, M. & Beijerink, N.J. (2013) Congenital heart disease: a case series of 135 dogs. *Australian Veterinary Practice*, **43** (3), 466–470.
- Ahlgren, J. & Uimari, P. (2016) Heritability of hypothyroidism in the Finnish Hovawart population. *Acta Veterinaria Scandinavica*, **58** (1), 39.

- Ahman, S., Perrins, N. & Bond, R. (2007) Carriage of *Malassezia* spp. yeasts in healthy and seborrhoeic Devon Rex cats. *Medical Mycology*, **45**, 449–455.
- Ahonen, S.J., Arumilli, M. & Lohi, H. (2013) A CNGB1 frameshift mutation in Papillon and Phalène Dogs with progressive retinal atrophy. *PLoS One*, **8** (8), e72122.
- Ahonen, S.J., Kaukonen, M., Nussdorfer, F.D. *et al.* (2014) A novel missense mutation in ADAMTS10 in Norwegian Elkhound primary glaucoma. *PLOS One*, **9** (11), e111941.
- Aikawa, T., Shibata, M. & Fujita, H. (2013) Modified ventral stabilization using positively threaded profile pins and polymethylmethacrylate for atlantoaxial instability in 49 dogs. *Veterinary Surgery*, **42** (6), 683–692.
- Aikawa, T., Shibata, M., Asano, M. *et al.* (2014) A comparison of thoracolumbar intervertebral disc extrusion in French Bulldogs and Dachshunds and association with congenital vertebral anomalies. *Veterinary Surgery*, **43** (3) 301–307.
- Aksenovich, T.I., Zorkal'tsov, I.V. Aul'chenko, Iu. S. *et al.* (2004) Inheritance of hypodontia in Kerry Blue Terrier dogs *Genetika*, **40** (5) 658–666.
- Alam, M.R., Lee, J.I., Kang, H.S. *et al.* (2007) Frequency and distribution of patellar luxation in dogs: 134 cases (2000 to 2005). *Veterinary and Comparative Orthopaedics and Traumatology*, **20** (1), 59–64.
- Albasan, H., Lulich, J.P., Osborne, C.A. *et al.* (2005) Evaluation of the association between sex and risk of forming urate uroliths in Dalmatians. *Journal of the American Veterinary Medical Association*, **227** (4), 565–569.
- Albasan, H., Osborne, C.A., Lulich, J.P. *et al.* (2012) Risk factors for urate uroliths in cats *Journal of the American Veterinary Medical Association*, **240** (7), 842–847.
- Alberch, P. (1985) Developmental constraints: why St. Bernards often have an extra digit and Poodles never do. *American Naturalist*, **126** (3), 430–433.
- Alderton, D. & Morgan, T. (1993). *Dogs*. Dorling Kindersley, London.
- Aldrich, J., Ling, G.V., Ruby, A.L., Johnson, D.L. & Franti, C.E. (1997) Silica-containing urinary calculi in dogs (1981–1993). *Journal of Veterinary Internal Medicine*, **11** (5), 288–295.
- Alenza, D.P., Arenas, C., Lopez, M.L. & Melian, C. (2006) Long-term efficacy of trilostane administered twice daily in dogs with pituitary-dependent hyperadrenocorticism. *Journal of the American Animal Hospital Association*, **42** (4), 269–276.
- Alhaddad, H., Gandolfi, B., Grah, R.A. *et al.* (2014) Genome-wide association and linkage analyses localize a progressive retinal atrophy locus in Persian cats. *Mammalian Genome*, **25** (7–8), 354–362.
- Allan, F.J., Thompson, K.G., Jones, B.R. *et al.* (1996) Neutropenia with a probable hereditary basis in Border Collies. *New Zealand Veterinary Journal*, **44**, 67–72.
- Allbaugh, R.A. & Stuhr, C.M. (2013) Thermal cautery of the canine third eyelid for treatment of cartilage eversion. *Veterinary Ophthalmology*, **16** (5) 392–395.
- Alroy, J., Rush, J.E., Freeman, L. *et al.* (2000) Inherited infantile dilated cardiomyopathy in dogs: genetic, clinical, biochemical, and morphologic findings. *American Journal of Medical Genetics Part A*, **95** (1), 57–66.
- Alroy, J., Rush, J.E. & Sarkar, S. (2005) Infantile dilated cardiomyopathy in Portuguese water dogs: Correlation of the autosomal recessive trait with low plasma taurine at infancy. *Amino Acids*, **28** (1), 51–56.
- Alves, L., Hülsmeier, V., Jaggy, A. *et al.* (2011) Polymorphisms in the ABCB1 Gene in phenobarbital responsive and resistant idiopathic epileptic Border Collies. *Journal of Veterinary Internal Medicine*, **25**, 484–489.
- Amat, M., Manteca, X., Mariotti, V.M., Ruiz de la Torre, J.L. & Fatjó, J. (2009) Aggressive behavior in the English Cocker Spaniel. *Journal of Veterinary Behavior: Clinical Applications and Research*, **4**, 111–117.
- Amberger, C., Hagen, A., Hauser, K. *et al.* (1996) [ECG abnormalities in Airedale Terriers]. *Tierärztliche Praxis*, **24**, 278–283.
- Amis, T.C. (1974) Tracheal collapse in the dog. *Australian Veterinary Journal*, **50** (7), 285–289.
- Amort, K.H., Ondreka, N., Rudorf, H. *et al.* (2012) MR-imaging of lumbosacral intervertebral disc degeneration in clinically sound German Shepherd Dogs compared to other breeds. *Veterinary Radiology and Ultrasound*, **53** (3) 289–295.

- Anderson, M.A., Constantinescu, G.H., Dee, L.G. *et al.* (1995a) Fractures and dislocations of the Racing Greyhound. Part I. *Compendium on Continuing Education for the Practicing Veterinarian*, **17**, 779–786.
- Anderson, M.A., Constantinescu, G.H., Dee, L.G. *et al.* (1995b) Fractures and dislocations of the Racing Greyhound. Part II. *Compendium on Continuing Education for the Practicing Veterinarian*, **17**, 899–909.
- Andersson, M. & Sevelius, E. (1991) Breed, sex and age distribution in dogs with chronic liver disease: a demographic study. *Journal of Small Animal Practice*, **32** (1), 1–5.
- Anfinsen, K.P., Grotmol, T., Bruland, O.S. & Jonasdottir, T.J. (2011) Breed-specific incidence rates of canine primary bone tumors: a population based survey of dogs in Norway. *Canadian Journal of Veterinary Research*, **75** (3), 209–215.
- Angles, J.M., Famula, T.R. & Pedersen, N.C. (2005) Uveodermatologic (VKH-like) syndrome in American Akita dogs is associated with an increased frequency of DQA1*00201. *Tissue Antigens*, **66** (6), 656–665.
- Angus, J.C., Lichtensteiger, C., Campbell, K.L. & Schaeffer, D.J. (2002) Breed variations in histopathologic features of chronic severe otitis externa in dogs: 80 cases (1995–2001). *Journal of the American Veterinary Medical Association*, **221**, 1000–1006.
- Animal Health Trust (2017). *Canine DNA Testing*. http://www.aht.org.uk/cms-display/genetics_canine.html (accessed October 2017).
- Anis, E.A., Frank, L.A., Francisco, R. *et al.* (2016) Identification of canine papillomavirus by PCR in Greyhound dogs. *Peer J*, **4**, e2744.
- Aoki, H., Mishina, M., Watanabe, T. (2012) Canine transitional cell carcinoma of the bladder and urethra: a retrospective study of 82 cases (2004–2009). *Journal of the Japan Veterinary Medical Association*, **65**, 289–292.
- Arceneaux, K.A., Taboada, J. & Hosgood, G. (1998) Blastomycosis in dogs: 115 cases (1980–1995). *Journal of the American Veterinary Medical Association*, **213** (5), 658–664.
- Arikan, S., Duru, S.Y., Gurkan, M. *et al.* (2003) Blood type A and B frequencies in Turkish Van and Angora cats in Turkey *Journal of Veterinary Medicine*, **50** (6) 303–306.
- Arikan, S., Guzel, M., Mamak, N. & Ograk, Y.Z. (2009) Frequency of blood types DEA 1.1, 3, 4, 5, and 7 in Kangal dog. *Revue de Médecine Vétérinaire*, **160** (4), 180–183.
- Arnold, S. (1997) Urinary incontinence in castrated bitches. Part I: Significance, clinical aspects and etiopathogenesis. *Schweizer Archiv für Tierheilkunde*, **139** (6), 271–276.
- Arnott, E.R., Peek, L., Early, J.B. *et al.* (2015) Strong selection for behavioural resilience in Australian stock working dogs identified by selective sweep analysis. *Canine Genetics and Epidemiology*, **2** (1), 6.
- Aronsohn, M. (1985) Canine thymoma. *Veterinary Clinics of North America: Small Animal Practice*, **15** (4), 755–767.
- Arrol, L., Penderis, J., Garosi, L. *et al.* (2012) Aetiology and long-term outcome of juvenile epilepsy in 136 dogs. *Veterinary Record*, **170**, 335.
- Arthurs, G.I. & Langley-Hobbs, S.J. (2006) Complications associated with corrective surgery for patellar luxation in 109 dogs. *Veterinary Surgery*, **35**, 559–566.
- Arvelius, P., Asp, H.E., Fikse, W.F., Strandberg, E. & Nilsson, K. (2014) Genetic analysis of a temperament test as a tool to select against everyday life fearfulness in Rough Collie. *Journal of Animal Science*, **92**, 4843–4855.
- Aslanian, M.E., Sharp, C.R., Rozanski, E.A. *et al.* (2014) Clinical outcome after diagnosis of hemophilia A in dogs. *Journal of the American Veterinary Medical Association*, **245** (6) 677–683.
- Auger, J.M. & Riley, S.M. (1997) Combined hiatal and pleuroperitoneal hernia in a shar-pei. *Canadian Veterinary Journal*, **38** (10), 640–642.
- Averill, D.R. & Bronson, R.T. (1977) Inherited necrotizing myelopathy of Afghan Hounds. *Journal of Neuropathology and Experimental Neurology*, **36**, 734–747.
- Awano, T., Johnson, G.S., Wade, C.M. *et al.* (2009) Genome-wide association analysis reveals a SOD1 mutation in canine degenerative myelopathy that resembles amyotrophic lateral sclerosis. *Proceedings of the National Academy of Sciences of the United States of America*, **106** (8), 2794–2799.

- Awano, T., Katz, M.L., O'Brien, D.P. *et al.* (2006) A mutation in the cathepsin D gene (CTSD) in American Bulldogs with neuronal ceroid lipofuscinosis. *Molecular Genetics and Metabolism*, **87** (4), 341–348.
- Babo, V. von, Eberle, N., Mischke, R. *et al.* (2012) Canine non-hematopoietic gastric neoplasia. Epidemiologic and diagnostic characteristics in 38 dogs with post-surgical outcome of five cases. *Tierärztliche Praxis, Ausgabe K, Kleintiere/Heimtiere*, **40**, 243–249.
- Bacon, N.J., Oni, O. & White, R.A.S. (2002) Treatment of urethral sphincter mechanism incompetence in 11 bitches with a sustained-release formulation of phenylpropanolamine hydrochloride. *Veterinary Record*, **151** (13), 373–376.
- Bagley, R.S., Kornegay, J.N. & Wheeler, S.J. (1993) Generalized tremors in Maltese: clinical findings in seven cases. *Journal of the American Animal Hospital Association*, **29** (2), 141–145.
- Baker, L.A., Kirkpatrick, B., Rosa, G.J.M. *et al.* (2017) Genome-wide association analysis in dogs implicates 99 loci as risk variants for anterior cruciate ligament rupture. *PLoS One*, **12** (4), e0173810.
- Baker-Gabb, M., Hunt, G.B. & France, M.P. (2003) Soft tissue sarcomas and mast cell tumours in dogs; clinical behaviour and response to surgery. *Australian Veterinary Journal*, **81** (12), 732–738.
- Bamberger, M. & Houpt, K.A. (2006) Signalment factors, comorbidity, and trends in behavior diagnoses in dogs: 1,644 cases (1991–2001). *Journal of the American Veterinary Medical Association*, **229** (10), 1591–1601.
- Bannasch, D.L., Ling, G.V., Bea, J. *et al.* (2004) Inheritance of urinary calculi in the Dalmatian. *Journal of Veterinary Internal Medicine*, **18** (4), 483–487.
- Bannasch, D., Safra, N., Young, A. *et al.* (2008) Mutations in the SLC2A9 gene cause hyperuricosuria and hyperuricemia in the dog. *PLoS Genetics*, **4** (11), e1000246.
- Banz, A.C. & Gottfried, S.D. (2010) Peritoneopericardial diaphragmatic hernia: a retrospective study of 31 cats and eight dogs. *Journal of the American Animal Hospital Association*, **46** (6), 398–404.
- Baral, R.M., Metcalfe, S.S., Krockenberger, M.B. *et al.* (2006). Disseminated Mycobacterium avium infection in young cats: overrepresentation of Abyssinian cats. *Journal of Feline Medicine and Surgery*, **8** (1), 23–44.
- Baranowska, I., Jäderlund, K.H., Nennesmo, I. *et al.* (2009) Sensory ataxic neuropathy in golden retriever dogs is caused by a deletion in the mitochondrial tRNA^{Tyr} gene. *PLoS Genetics*, **5** (5), e1000499.
- Barker, E.N., Dawson, L.J., Rose, J.H. *et al.* (2016) Degenerative encephalopathy in Nova Scotia Duck Tolling Retrievers presenting with a rapid eye movement sleep behaviour disorder. *Journal of Veterinary Internal Medicine*, **30** (5), 1681–1689.
- Barnett, K.C. (1980) Hereditary cataract in the Welsh Springer Spaniel. *Journal of Small Animal Practice*, **21** (11), 621–625.
- Barnett, K.C. (1988) Inherited eye disease in the dog and cat. *Journal of Small Animal Practice*, **29**, 462–475.
- Barnett, K.C. (2006) Congenital keratoconjunctivitis sicca and ichthyosiform dermatosis in the Cavalier King Charles Spaniel. *Journal of Small Animal Practice*, **47**, 524–528.
- Barnhart, K.F., Credille, K.M., Ambrus, A. *et al.* (2004). A heritable keratinization defect of the superficial epidermis in Norfolk Terriers. *Journal of Comparative Pathology*, **130**, 246–254.
- Barrientos, L.S., Crespi, J.A., It, V. *et al.* (2013) Prevalence of canine juvenile generalized demodicosis in the Buenos Aires region, Argentina. *Japanese Journal of Veterinary Dermatology*, **19**, 57–61.
- Barrs, V.R., Gunew, M., Foster, S.F., Beatty, J.A. & Malik, R. (2001) Prevalence of autosomal dominant polycystic kidney disease in Persian cats and related breeds in Sydney and Brisbane. *Australian Veterinary Journal*, **79** (4), 257–259.
- Barrs, V.R. & Talbot, J.J. (2014) Feline aspergillosis. *Veterinary Clinics of North America: Small Animal Practice*, **44** (1), 51–73.
- Bartels, K.E., Higbee, R.G., Bahr, R.J. *et al.* (2003) Outcome of and complications associated with prophylactic percutaneous laser disk ablation in dogs with thoracolumbar disk disease: 277 cases (1992–2001). *Journal of the American Veterinary Medical Association*, **222** (12), 1733–1739.
- Bartges, J.W., Osborne, C.A., Lulich, J.P. *et al.* (1994) Prevalence of cystine and urate uroliths in Bulldogs and urate uroliths in Dalmatians. *Journal of the American Veterinary Medical Association*, **204** (12), 1914–1918.

- Barthez, P.Y., Rivier, P. & Begon, D. (2003) Prevalence of polycystic kidney disease in Persian and Persian-related cats in France. *Journal of Feline Medicine and Surgery*, **5** (6), 345–347.
- Bassett, R.J., Burton, G.G. & Robson, D.C. (2004) Antibiotic responsive ulcerative dermatoses in German Shepherd Dogs with mucocutaneous pyoderma. *Australian Veterinary Journal*, **82**, 485–489.
- Basso, C., Fox, P.R., Meurs, K.M. *et al.* (2004) Arrhythmogenic right ventricular cardiomyopathy causing sudden cardiac death in Boxer dogs. *Circulation*, **109**, 1180–1185.
- Batchelor, D.J., Noble, P.-J.M., Cripps, P.J. *et al.* (2007) Breed associations for canine exocrine pancreatic insufficiency. *Journal of Veterinary Internal Medicine*, **21** (2), 207–214.
- Bateson, P. (2010). *Independent Inquiry into Dog Breeding*. University of Cambridge, Cambridge.
- Batt, R.M., Barnes, A., Rutgers, H.C. *et al.* (1991) Relative IgA deficiency and small intestinal bacterial overgrowth in German Shepherd Dogs. *Research in Veterinary Science*, **50** (1), 106–111.
- Batt, R.M., Carter, M.W. & McClean, L. (1984) Morphological and biochemical studies of a naturally occurring enteropathy in the Irish Setter dog: a comparison with coeliac disease in man. *Research in Veterinary Science*, **37**, 339–346.
- Battersby, I.A., Giger, U. & Hall, E.J. (2005) Hyperammonaemic encephalopathy secondary to selective cobalamin deficiency in a juvenile Border Collie. *Journal of Small Animal Practice*, **46** (7), 339–344.
- Bauer, B.S., Sandmeyer, L.S. & Grahn, B.H. (2015) Diagnostic ophthalmology. Microphthalmos and multiple ocular anomalies (MOA) OU consistent with merle ocular dysgenesis (MOD). *Canadian Veterinary Journal*, **56** (7), 767–768.
- Baumworcel, N., Soares, A.M.B., Helms, G., Rei, P.R.L. & Castro, M.C.N. (2009) Three hundred and three dogs with cataracts seen in Rio de Janeiro, Brazil. *Veterinary Ophthalmology*, **12** (5), 299–301.
- Bavegems, V., Duchateau, L.U.C., Sys, S.U. & De Rick, A. (2007) Echocardiographic reference values in Whippets. *Veterinary Radiology and Ultrasound*, **48** (3), 230–238.
- Bavegems, V., Van Caelenberg, A., Duchateau, L. *et al.* (2005) Vertebral heart size ranges specific for Whippets. *Veterinary Radiology and Ultrasound*, **46** (5), 400–403.
- Bavegems, V.C., Duchateau, L., Polis, I.E. *et al.* (2011) Detection of innocent systolic murmurs by auscultation and their relation to hematologic and echocardiographic findings in clinically normal Whippets. *Journal of the American Veterinary Medical Association*, **238** (4), 468–471.
- Bearss, J.J., Schulman, F.Y. & Carter, D. (2012) Histologic, Immunohistochemical, and clinical features of 27 mammary tumors in 18 male dogs. *Veterinary Pathology*, **49**, 602–607.
- Beaver, D.P., Ellison, G.W., Lewis, D.D. *et al.* (2000) Risk factors affecting the outcome of surgery for atlantoaxial subluxation in dogs: 46 cases (1978–1998). *Journal of the American Veterinary Medical Association*, **216** (7), 1104–1109.
- Beck, J.A., Lloyd, S., Hafezparast, M. *et al.* (2000) Genealogies of mouse inbred strains. *Nature Genetics*, **24** (1), 23–25.
- Beck, J.J., Staatz, A.J., Pelsue, D.H. *et al.* (2006) Risk factors associated with short-term outcome and development of perioperative complications in dogs undergoing surgery because of gastric dilatation-volvulus: 166 cases (1992–2003). *Journal of the American Veterinary Medical Association*, **229** (12), 1934–1939.
- Beckers, E., Poucke, M. van, Ronsyn, L. & Peelman, L. (2016) Frequency estimation of disease-causing mutations in the Belgian population of some dog breeds. Part 2: Retrievers and other breed types. *Vlaams Diergeneeskundig Tijdschrift*, **85**, 185–196.
- Beco, L., Fontaine, J., Gross, T.L. & Charlier, G. (1996) Colour dilution in alopecia in seven Dachshunds. A clinical study and the hereditary, microscopical and ultrastructural aspect of the disease. *Veterinary Record*, **7** (2), 91–97.
- Bedford, P.G. (1982a) Collie eye anomaly in the Border Collie. *Veterinary Record*, **111** (2), 34–35.
- Bedford, P.G. (1982b) Collie eye anomaly in the United Kingdom. *Veterinary Record*, **111** (12), 263–270.
- Bedford, P.G. (1984) Retinal pigment epithelial dystrophy (CPRA): a study of the disease in the Briard. *Journal of Small Animal Practice*, **25**, 129–138.
- Bedford, P.G. (1989) Control of inherited retinal degeneration in dogs and cats in the United Kingdom. *Journal of Small Animal Practice*, **30** (3), 172–177.
- Bedford, P.G. (1998) Collie eye anomaly in the Lancashire Heeler. *Veterinary Record*, **143** (13), 354–356.

- Bedford, P.G. (2009) Retinal pigment epithelial dystrophy in the Briard. *Veterinary Record*, **164**, 377.
- Bedford, P.G. (2017) Open-angle glaucoma in the Petit Basset Griffon Vendeen. *Veterinary Ophthalmology*, **20** (2), 98–102.
- Bedu, A.S., Labruyère, J.J., Thibaud, J.-L. *et al.* (2012) Age-related thoracic radiographic changes in Golden and Labrador Retriever muscular dystrophy. *Veterinary Radiology and Ultrasound*, **53** (5), 492–500.
- Beer, K.A.S., Syring, R.S. & Drobatz, K.J. (2012) Evaluation of plasma lactate concentration and base excess at the time of hospital admission as predictors of gastric necrosis and outcome and correlation between those variables in dogs with gastric dilatation-volvulus: 78 cases (2004–2009). *Journal of the American Veterinary Medical Association*, **242** (1), 54–58.
- Behr, S. & Cauzinille, L. (2006) Aseptic suppurative meningitis in Juvenile Boxer Dogs: Retrospective study of 12 cases. *Journal of the American Animal Hospital Association*, **42**, 277–282.
- Belanger, J.M., Bellumori, T.P., Bannasch, D.L., Famula, T.R. & Oberbauer, A.M. (2017) Correlation of neuter status and expression of heritable disorders. *Canine Genetics and Epidemiology*, **4** (1), 6.
- Bell, T.G., Butler, K.L., Sill, H.B. (2002) Autosomal recessive severe combined immunodeficiency of Jack Russell Terriers. *Journal of Veterinary Diagnostic Investigation*, **14** (3), 194–204.
- Bellenger, C.R. (1980) Perineal hernia in dogs. *Australian Veterinary Journal*, **56**, 434–438.
- Bellenger, C.R. (1996) Inguinal and scrotal herniation in 61 dogs. *Australian Veterinary Practitioner*, **26**, 58–59.
- Bellenger, C.R. & Simpson, D.J. (1992) Canine sialocoeles: 60 clinical cases. *Journal of Small Animal Practice*, **33** (8), 376–380.
- Bellenger, C.R., Maddison, J.E., Macpherson, G.C. & Ilkiw, J.E. (1990) Chronic hypertrophic pyloric gastropathy in 14 dogs. *Australian Veterinary Journal*, **67** (9), 317–320.
- Belluco, S., Brisebard, E., Watrelot, D. *et al.* (2013) Digital squamous cell carcinoma in dogs. *Veterinary Pathology*, **50**, 1078–1082.
- Bellumori, T.P., Famula, T.R., Bannasch, D.L., Belanger, J.M. & Oberbauer, A.M. (2013) Prevalence of inherited disorders among mixed-breed and purebred dogs: 27,254 cases (1995–2010). *Journal of the American Veterinary Medical Association*, **242** (11), 1549–1555.
- Bende, B., Kovács, K. B., Solymosi, N. & Németh, T. (2015) Characteristics of urolithiasis in the dog population of Hungary from 2001 to 2012. *Acta Veterinaria Hungarica*, **63**, 323–336.
- Bennett, D., Tennant, B., Lewis, D.G. *et al.* (1988) A reappraisal of anterior cruciate ligament disease in the dog. *Journal of Small Animal Practice*, **29** (5), 275–297.
- Bennett, P.F., DeNicola, D.B., Bonney, P. *et al.* (2002) Canine anal sac adenocarcinomas: clinical presentation and response to therapy. *Journal of Veterinary Internal Medicine*, **16** (1), 100–104.
- Bensfield, A.C., Evans, J., Pesayco, J.P. *et al.* (2011) Recurrent demyelination and remyelination in 37 young Bengal cats with polyneuropathy. *Journal of Veterinary Internal Medicine*, **25** (4) 882–889.
- Bensignor, E. & Guaguere, E. (2012) Sebaceous adenitis in dogs: a review. *Pratique Medicale et Chirurgicale de l'Animal de Compagnie*, **43** (7), 65–71.
- Bentley, E. (2005) Spontaneous chronic corneal epithelial defects in dogs: a review. *Journal of the American Animal Hospital Association*, **41** (3), 158–165.
- Berdoulay, A., English, R.V. & Nelson, B. (2005) Effect of topical 0.02% tacrolimus aqueous suspension on tear production in dogs with keratoconjunctivitis sicca. *Veterinary Ophthalmology*, **8** (4), 225–232.
- Berendt, M., Gredal, H. & Pedersen, L.G. (2002) A cross-sectional study of epilepsy in Danish Labrador Retrievers: prevalence and selected risk factors. *Journal of Veterinary Internal Medicine*, **16** (3), 262–268.
- Berendt, M., Gullov, C.H., Christensen, S.L. *et al.* (2008) Prevalence and characteristics of epilepsy in the Belgian shepherd variants Groenendael and Tervueren born in Denmark 1995–2004. *Acta Veterinaria Scandinavica*, **50**, 51.
- Berg, T., Tollersrud, O.K., Walkley, S.U. *et al.* (1997) Purification of feline lysosomal alpha-mannosidase, determination of its cDNA sequence and identification of a mutation causing alpha-mannosidosis in Persian cats. *Biochemistry Journal*, **328** (3), 863–870.
- Berger, B. & Feldman, E.C. (1987) Primary hyperparathyroidism in dogs: 21 cases (1976–1986). *Journal of the American Veterinary Medical Association*, **191** (3), 350–356.

- Bergknut, N., Egenvall, A., Hagman, R. *et al.* (2012) Incidence of intervertebral disk degeneration-related diseases and associated mortality rates in dogs. *Journal of the American Veterinary Medical Association*, **240** (11), 1300–1309.
- Bergman, P.J., Withrow, S.J., Straw, R.C. *et al.* (1994) Infiltrative lipoma in dogs: 16 cases (1981–1992). *Journal of the American Veterinary Medical Association*, **205** (2), 322–324.
- Bergsjø, T., Arnesen, K., Heim, P. *et al.* (1984) Congenital blindness with ocular developmental anomalies, including retinal dysplasia, in Doberman Pinscher dogs. *Journal of the American Veterinary Medical Association*, **184** (11), 1383–1386.
- Bergström, A., Nødtvedt, A., Lagerstedt, A.S. & Egenvall, A. (2006) Incidence and breed predilection for dystocia and risk factors for cesarean section in a Swedish population of insured dogs. *Veterinary Surgery*, **35** (8), 786–791.
- Bergvall, K.E. & Shokrai, A. (2014) Clinical and histological characterization of multifocal, spontaneous, noninfectious alopecia in Norwegian Puffin Dogs (Lundehunds). *Veterinary Dermatology*, **25** (2), 112–119.
- Bernard, M.A. & Valli, V.E. (1977) Familial renal disease in Samoyed dogs. *Canadian Veterinary Journal*, **18** (7), 181–189.
- Berthelin, C.F., Bailey, C.S., Kass, P.H. *et al.* (1994) Cryptococcosis of the nervous system of dogs, part 1: epidemiologic, clinical and neuropathologic features. *Progress in Veterinary Neurology*, **5** (3), 88–97.
- Besalti, O., Pekcan, Z., Sirin, Y.S. & Erbas, G. (2006) Magnetic resonance imaging findings in dogs with thoracolumbar intervertebral disk disease: 69 cases (1997–2005). *Journal of the American Veterinary Medical Association*, **228** (6), 902–908.
- Betschart, S., Hässig, M. & Spiess, B. (2014) [Lens luxation in dogs: a retrospective study of 134 dogs (2000–2011).] *Schweizer Archiv für Tierheilkunde*, **156**, 125–131.
- Beuing, R., Mues, C., Tellhelm, B. & Erhardt, G. (2000) Prevalence and inheritance of canine elbow dysplasia in German Rottweilers. *Journal of Animal Breeding and Genetics*, **117** (6), 375–383.
- Bexfield, N.H., Andres-Abdo, C., Scase, T.J., Constantino-Casas, F. & Watson, P.J. (2011) Chronic hepatitis in the English Springer Spaniel: clinical presentation, histological description and outcome. *Veterinary Record*, **169** (16), 415.
- Bexfield, N.H., Buxton, R.J., Vitek, T.J. *et al.* (2012a) Breed, age and gender distribution of dogs with chronic hepatitis in the United Kingdom. *Veterinary Journal*, **193** (1), 124–128.
- Bexfield, N.H., Watson, P.J., Aguirre-Hernández, J. *et al.* (2012b) DLA class II alleles and haplotypes are associated with risk for and protection from chronic hepatitis in the English Springer Spaniel. *PLoS One*, **7** (8), e42584.
- Bhalerao, D.P., Rajpurohit, Y., Vite, C.H. & Giger, U. (2002) Detection of a genetic mutation for myotonia congenita among Miniature Schnauzers and identification of a common carrier ancestor. *American Journal of Veterinary Research*, **63** (10), 1443–1447.
- Bhatti, S.F., Vanhaesebrouck, A.E., Van Soens, I. *et al.* (2011) Myokymia and neuromyotonia in 37 Jack Russell Terriers. *Veterinary Journal*, **189** (3), 284–288.
- Bianchi, M., Dahlgren, S., Massey, J. *et al.* (2015) A multi-breed genome-wide association analysis for canine hypothyroidism identifies a shared major risk locus on CFA12. *PLoS One*, **10** (8), e0134720.
- Bichsel, P., Vandevelde, M., Lang, J. & Kull-Hächler, S. (1983) Degenerative myelopathy in a family of Siberian Husky dogs. *Journal of the American Veterinary Medical Association*, **183** (9), 998–1000, 1965.
- Biel, M., Kramer, M., Forterre, F. *et al.* (2013) Outcome of ventriculoperitoneal shunt implantation for treatment of congenital internal hydrocephalus in dogs and cats: 36 cases (2001–2009). *Journal of the American Veterinary Medical Association*, **242**, 948–958.
- Bielfelt, S.W., Redman, H.C., McClellan, R.O. (1971) Sire- and sex-related differences in rates of epileptiform seizures in a purebred beagle dog colony. *American Journal of Veterinary Research*, **32**, 2039–2048.
- Biezyński, J., Skrzypczak, P., Piatek, A. *et al.* (2012) Assessment of treatment of osteochondrosis dissecans (OCD) of shoulder joint in dogs: the results of two years of experience. *Polish Journal of Veterinary Sciences*, **15** (2), 285–290.
- Bilek, A. & Hirt, R. A. (2007) Breed-associated increased occurrence of gastric carcinoma in Chow-chows. *Wiener Tierärztliche Monatsschrift*, **94**, 71–79.

- Binder, D.R., Herring, I.P. & Gerhard, T. (2007) Outcomes of nonsurgical management and efficacy of demecarium bromide treatment for primary lens instability in dogs: 34 cases (1990–2004). *Journal of the American Veterinary Medical Association*, **231** (1), 89–93.
- Binder, H., Arnold, S., Scherring, C. *et al.* (2000) Palmoplantar hyperkeratosis in Irish Terriers: evidence of autosomal recessive inheritance. *Journal of Small Animal Practice*, **41**, 52–55.
- Bingel, S.A. & Sande, R.D. (1982) Chondrodysplasia in the Norwegian Elkhound. *American Journal of Pathology*, **107**, 219–229.
- Bingel, S.A. & Sande, R.D. (1994) Chondrodysplasia in five Great Pyrenees. *Journal of the American Veterinary Medical Association*, **205** (6), 845–845.
- Bingel, S.A., Sande, R.D. & Wight, T.N. (1985) Chondrodysplasia in the Alaskan Malamute: characterization of proteoglycans dissociatively extracted from dwarf growth plates. *Laboratory Investigations*, **53**, 479–485.
- Birkenheuer, A.J., Correa, M.T., Lew, M.G. *et al.* (2005) Geographic distribution of babesiosis among dogs in the United States and association with dog bites: 150 cases (2000–2003). *Journal of the American Veterinary Medical Association*, **227** (6), 942–947.
- Bishop, M.A., Xenoulis, P.G., Berghoff, N. *et al.* (2012) Partial characterization of cobalamin deficiency in Chinese Shar Peis. *Veterinary Journal*, **191** (1), 41–45.
- Bishop, M.A., Xenoulis, P.G., Levinski, M.D., Suchodolski, J.S. & Steiner, J.M. (2010) Identification of variants of the SPINK1 gene and their association with pancreatitis in Miniature Schnauzers. *American Journal of Veterinary Research*, **71** (5), 527–533.
- Bizikova, P., Linder, K.E., Wofford, J.A. *et al.* (2015) Canine epidermolysis bullosa acquisita: a retrospective study of 20 cases. *Veterinary Dermatology*, **26** (6), 441–450.
- Bjerkås, E. (1991) Hereditary cataract of dogs in Norway. *Norsk Veterinærtidsskrift*, **103**, 5–14.
- Bjerkås, E., Ekesten, B. & Farstad, W. (2002) Pectinate ligament dysplasia and narrowing of the irido-corneal angle associated with glaucoma in the English Springer Spaniel. *Veterinary Ophthalmology*, **5** (1), 49–54.
- Bjerkås, E. & Haaland, M.B. (1995) Pulverulent nuclear cataract in the Norwegian Buhund. *Journal of Small Animal Practice*, **36** (11), 471–474.
- Bjerkås, E. & Narfström, K. (1994) Progressive retinal atrophy in the Tibetan Spaniel in Norway and Sweden. *Veterinary Record*, **134** (15), 377–379.
- Bjotvedt, G., Hendricks, G.M. & Brandon, T.A. (1988) Hemodynamic basis of renal arteriosclerosis in young greyhounds. *Laboratory Animal Science*, **38** (1), 62–67.
- Black, V., Garosi, L., Lowrie, M., Harvey, R.J. & Gale, J. (2014) Phenotypic characterisation of canine epileptoid cramping syndrome in the Border Terrier. *Journal of Small Animal Practice*, **55**, 102–107.
- Blacklock, B.T., Grundon, R.A., Meehan, M., Tetas Pont, R. & Hartley, C. (2016) Uveal cysts in domestic cats: a retrospective evaluation of thirty-six cases. *Veterinary Ophthalmology*, **1** (suppl. 1) 56–60.
- Blackshaw, J.K. (1991) An overview of types of aggressive behaviour in dogs and methods of treatment. *Applied Animal Behaviour Science*, **30** (3), 351–361.
- Blackshaw, J.K., Sutton, R.H. & Boyhan, M.A. (1994) Tail chasing behaviour in dogs. *Canine Practice*, **19**, 7–10.
- Blake, S. & Lapinski, A. (1980) Hypothyroidism in different breeds. *Canine Practice*, **7** (2), 48, 51.
- Blehaut, T.R.W., Hardstaff, J.L., Chapman, P.S. *et al.* (2014) Spatial, demographic and clinical patterns of Angiostrongylus vasorum infection in the dog population of Southern England. *Veterinary Record*, **175**, 148.
- Blendinger, C., Blendinger, K. & Bostedt, H. (1995) Urinary incontinence following ovariectomy in bitches. I. Origin, frequency and predisposition. *Tierärztliche Praxis*, **23**, 291–299.
- Bley, T., Gaillard, C., Bilzer, T. *et al.* (2002) Genetic aspects of Labrador Retriever myopathy. *Research in Veterinary Science*, **73**, 231–236.
- Blum, J.R., Cork, L.C., Morris, J.M., Olson, J.L. & Winkelstein, J.A. (1985) The clinical manifestations of a genetically determined deficiency of the third component of complement in the dog, *Clinical Immunology and Immunopathology*, **34**, 304–315.
- Blum, M. & Distl, O. (2013) Congenital sensorineural deafness in the dog. *Praktische Tierarzt*, **94** (3), 678–688.

- Boerkamp, K.M., Teske, E., Boon, L.R. *et al.* (2014) Estimated incidence rate and distribution of tumours in 4,653 cases of archival submissions derived from the Dutch Golden Retriever population. *BMC Veterinary Research*, **10** (1), 34.
- Boerno, C.M. & Eaton-Wells, R.D. (1995) Medial displacement of the tendon of origin of the biceps brachii muscle in 10 Greyhounds. *Journal of Small Animal Practice*, **36**, 69–73.
- Boevé, M.H., van der Linde-Sipman, T. & Stades, F.C. (1988) Early morphogenesis of persistent hyperplastic tunica vasculosa lentis and primary vitreous. The dog as an ontogenetic model. *Investigative Ophthalmology and Visual Science*, **29** (7), 1076–1086.
- Böhm, J., Vasli, N., Maurer, M. *et al.* (2013) Altered splicing of the BIN1 muscle-specific exon in humans and dogs with highly progressive centronuclear myopathy. *PLoS Genetics*, **9** (6), e1003430.
- Boldizsar, H., Szenci, O., Muray, T. *et al.* (1992) Studies on mammary tumours 1. Age, seasonal and breed distribution. *Acta Veterinaria Hungarica*, **40** (1–2), 75–87.
- Bomassi, E., Misbach, C., Tissier, R. *et al.* (2015) Signalment, clinical features, echocardiographic findings, and outcome of dogs and cats with ventricular septal defects: 109 cases (1992–2013). *Journal of the American Veterinary Medical Association*, **247** (2), 166–175.
- Bomhard, D. von & Drejack, J. (1977) Statistical studies of the occurrence of mammary gland tumours in bitches. *Kleintierpraxis*, **22**, 205–209.
- Bonazzi, M., Volta, A., Gnudi, G. *et al.* (2009) Comparison between ultrasound and genetic testing for the early diagnosis of polycystic kidney disease in Persian and Exotic Shorthair cats. *Journal of Feline Medicine and Surgery*, **11** (6), 430–434.
- Bond, R., Curtis, C.F., Ferguson, E. *et al.* (2000) An idiopathic facial dermatitis of Persian Cats. *Veterinary Dermatology*, **11** (1), 35–41.
- Bond, R., Elwood, C.M., Littler, R.M., Pinter, L. & Lloyd, D.H. (1998) Humoral and cell-mediated responses to *Malassezia pachydermatis* in healthy dogs and dogs with *Malassezia* dermatitis. *Veterinary Record*, **143**, 381–384.
- Bond, R., Ferguson, E.A., Curtis, C.F., Craig, J.M. & Lloyd, D.H. (1996) Factors associated with elevated cutaneous *Malassezia pachydermatis* populations in dogs with pruritic skin disease. *Journal of Small Animal Practice*, **37**, 103–107.
- Bond, R., Stevens, K., Perrins, N. *et al.* (2008) Carriage of *Malassezia* spp. yeasts in Cornish Rex, Devon Rex and Domestic short-haired cats: a cross-sectional survey. *Veterinary Dermatology*, **19** (5), 299–304.
- BongKyeong, K., YoungMin, Y., JeKyung, S. & KangMoon, S. (2001) Spontaneous ophthalmic diseases of Beagles in Korea. *Korean Journal of Veterinary Research* **41**, 113–121.
- Bonnett, B.N., Egenvall, A., Hedhammar, Å. & Olson, P. (2005) Mortality in over 350,000 insured Swedish dogs from 1995–2000: I. breed-, gender-, age- and cause-specific rates. *Acta Veterinaria Scandinavica*, **46** (3), 105–120.
- Borak, D., Wunderlin, N., Brückner, M. *et al.* (2017) Slipped capital femoral epiphysis in 17 Maine Coon cats. *Journal of Feline Medicine and Surgery*, **19** (1), 13–20.
- Borgarelli, M., Santilli, R.A., Chiavegato, D. *et al.* (2006) Prognostic indicators for dogs with dilated cardiomyopathy. *Journal of Veterinary Internal Medicine*, **20** (1), 104–110.
- Borgarelli, M., Savarino, P., Crosara, S. *et al.* (2008) Survival characteristics and prognostic variables of dogs with mitral regurgitation attributable to myxomatous valve disease. *Journal of Veterinary Internal Medicine*, **22** (1), 120–128.
- Borge, K.S., Tønnessen, R., Nødtvedt, A. & Indrebø, A. (2011) Litter size at birth in purebred dogs: a retrospective study of 224 breeds. *Theriogenology*, **75** (5), 911–919.
- Borgeat, K., Casamián-Sorrosal, D., Helps, C. *et al.* (2014) Association of the myosin binding protein C3 mutation (MYBPC3 R820W) with cardiac death in a survey of 236 Ragdoll cats. *Journal of Veterinary Cardiology*, **16** (2), 73–80.
- Borissov, I., Sivrev, D. & Milev, N. (2004) Incidence of some teeth and occlusion abnormalities in dogs: a retrospective study (1995–2002). *Bulgarian Journal of Veterinary Medicine*, **7** (4), 245–250.
- Borns-Weil, S., Emmanuel, C., Longo, J. *et al.* (2015) A case–control study of compulsive wool-sucking in Siamese and Birman cats ($n = 204$). *Journal of Veterinary Behaviour*, **10** (6), 543–558.
- Bosie, J.T., van den Ingh, T.S. & van der Linde-Sipman, J.S. (1998) Polycystic liver and kidney disease in cats. *Veterinary Quarterly*, **20** (4), 136–139.

- Boston, S.E., Lu, X., Culp, W. *et al.* (2014) Efficacy of systemic adjuvant therapies administered to dogs after excision of oral malignant melanomas: 151 cases (2001–2012). *Journal of the American Veterinary Medical Association*, **245** (4), 401–407.
- Boudreaux, M.K., Kvam, K., Dillon, A.R. *et al.* (1996) Type I Glanzmann's thrombasthenia in a Great Pyrenees Dog. *Veterinary Pathology*, **33** (5), 503–511.
- Boudreaux, M.K. & Lipscomb, D.L. (2001) Clinical, biochemical, and molecular aspects of Glanzmann's thrombasthenia in humans and dogs. *Veterinary Pathology*, **38** (3), 249–260.
- Boudreaux, M.K. & Martin, M. (2011) P2Y12 receptor gene mutation associated with postoperative hemorrhage in a Greater Swiss Mountain dog. *Veterinary Clinical Pathology*, **40**, 202–206.
- Bound, N., Zakai, D., Butterworth, S.J. *et al.* (2009) The prevalence of canine patellar luxation in three centres. Clinical features and radiographic evidence of limb deviation. *Veterinary and Comparative Orthopaedics and Trauma*, **22** (1), 32–37.
- Bourguet, A., Chaudieu, G., Briatta, A. *et al.* (2017) Cataracts in a population of Bengal cats in France. *Veterinary Ophthalmology*, April 25. doi: 10.1111/vop.12470 [Epub ahead of print].
- Bovee, K.C., Joyce, T., Reynolds, R. *et al.* (1978) The Fanconi syndrome in Basenji dogs: a new model for renal tubular defects. *Science*, **201** (4361), 1129–1131.
- Bowles, C.A., Alsaker, R.D. & Wolfe, T.L. (1979) Studies of the Pelger–Huët anomaly in foxhounds. *American Journal of Pathology*, **96** (1), 237–247.
- Boyce, J.T., DiBartola, S.P., Chew, D.J. *et al.* (1984) Familial renal amyloidosis in Abyssinian cats. *Veterinary Pathology*, **21** (1), 33–38.
- Boydell, P. (2000) Idiopathic Horner syndrome in the Golden Retriever. *Journal of Neuro-ophthalmology*, **20** (4), 288–290.
- Bradbury, A.M., Gurda, B.L., Casal, M.L. *et al.* (2015) A review of gene therapy in canine and feline models of lysosomal storage disorders *Human Genetic Therapy Clinical Development*, **26** (1), 27–37.
- Bradbury, A.M., Morrison, N.E., Hwang, M. *et al.* (2009) Neurodegenerative lysosomal storage disease in European Burmese cats with hexosaminidase beta-subunit deficiency. *Molecular Genetics and Metabolism*, **97** (1), 53–59.
- Bradshaw, J.W. & Goodwin, D. (1999) Determination of behavioural traits of pure-bred dogs using factor analysis and cluster analysis; a comparison of studies in the USA and UK. *Research in Veterinary Science*, **66** (1), 73–76.
- Bradshaw, J.W., Goodwin, D., Lea, A.M. & Whitehead, S.L. (1996) A survey of the behavioural characteristics of pure-bred dogs in the United Kingdom. *Veterinary Record*, **138** (19), 465–468.
- Brandão, C.V.S., Borges, A.G., Ranzani, J. J.T. *et al.* (2002) Transmissible venereal tumour in dogs: a retrospective study of 127 cases (1998–2000). *Revista de Educação Continuada do CRMV-SP*, **5**, 25–31.
- Brandão, C.V.S., Ranzani, J.J.T., Marinho, L.F.L.P., Rodrigues, G.N. & Cremonini, D.N. (2005) Proptose em cães e gatos: análise retrospectiva de 64 casos. *Archives of Veterinary Science*, **10** (1).
- Braund, K.G., Shores, A., Cochran, S. *et al.* (1994a) Laryngeal paralysis–polyneuropathy complex in young Dalmatians. *American Journal of Veterinary Research*, **55**, 534–552.
- Braund, K.G., Shores, A., Lowrie, C.T. *et al.* (1997) Idiopathic polyneuropathy in Alaskan Malamutes. *Journal of Veterinary Internal Medicine*, **11**, 243–249.
- Braund, K.G., Toivio-Kinnucan, M., Vallat, J.M., Mehta, J.R. & Levesque, D.C. (1994b) Distal sensorimotor polyneuropathy in mature Rottweiler dogs. *Veterinary Pathology*, **31** (3), 316–326.
- Breit, S. & Künzel, W. (2001) On biomechanical properties of the sacroiliac joint in purebred dogs. *Annals of Anatomy–Anatomischer Anzeiger*, **183** (2), 145–150.
- Bremer, H.D., Vilson, Á., Bonnett, B.N. *et al.* (2015) Disease patterns and incidence of immune-mediated disease in insured Swedish Nova Scotia Duck Tolling Retrievers. *Veterinary Record*, **177** (3), 74.
- Brenig, B., Leeb, T., Jansen, S. & Kopp, T. (1999) Analysis of blood clotting factor activities in canine Legg–Calvé–Perthes' disease. *Journal of Veterinary Internal Medicine*, **13** (6), 570–573.
- Brennan, K.E. & Ihrke, P.J. (1983) Grass awn migration in dogs and cats: a retrospective study of 182 cases. *Journal of the American Veterinary Medical Association*, **182** (11), 1201–1204.
- Brenner, O., de Lahunta, A., Summers, B.A. *et al.* (1997) Hereditary polioencephalomyopathy of the Australian Cattle Dog. *Acta Neuropathologica*, **94**, 54–66.

- Breur, G.J., Zerbe, C.A., Slocombe, R.F., Padgett, G.A. & Braden, T.D. (1989) Clinical, radiographic, pathologic, and genetic features of osteochondrodysplasia in Scottish Deerhounds. *Journal of the American Veterinary Medical Association*, **195** (5), 606–612.
- Brilhante, R.S.N., Cavalcante, C.S.P., Soares-Junior, F.A. *et al.* (2003) High rate of *Microsporium canis* feline and canine dermatophytoses in Northeast Brazil: Epidemiological and diagnostic features. *Mycopathologia*, **156** (4), 303–308.
- Brockman, D.J., Washabau, R.J. & Drobatz, K.J. (1995) Canine gastric dilatation/volvulus syndrome in a veterinary critical care unit: 295 cases (1986–1992). *Journal of the American Veterinary Medical Association*, **207** (4), 460–464.
- Brodey, R. S., Misdorp, W., Riser, W. H. & Heul, R. O., van der (1974) Canine skeletal chondrosarcoma: a clinicopathologic study of 35 cases. *Journal of the American Veterinary Medical Association* **165**, 68–78.
- Brons, A.K., Henthorn, P.S., Raj, K *et al.* (2013) SLC3A1 and SLC7A9 mutations in autosomal recessive or dominant canine cystinuria: a new classification system. *Journal of Veterinary Internal Medicine*, **27** (6), 1400–1408.
- Brooks, M. (1999) A review of canine inherited bleeding disorders: biochemical and molecular strategies for disease characterization and carrier detection. *Journal of Heredity*, **90**, 112–118.
- Brooks, M., Dodds, W.J. & Raymond, S.L. (1992) Epidemiologic features of von Willebrand's disease in Doberman pinschers, Scottish terriers, and Shetland sheepdogs: 260 cases (1984–1988) *Journal of the American Veterinary Medical Association*, **200** (8), 1123–1127.
- Brooks, M.B., Barnas, J.L., Fremont, J. *et al.* (2005) Cosegregation of a factor VIII microsatellite marker with mild hemophilia A in Golden Retriever dogs. *Journal of Veterinary Internal Medicine*, **19** (2), 205–210.
- Broome, C., Burbidge, H.M. & Pfeiffer, D.U. (2000) Prevalence of laryngeal paresis in dogs undergoing general anaesthesia. *Australian Veterinary Journal*, **78** (11), 769–772.
- Brown, N.O., Parks, J.L. & Greene, R. W. (1977) Canine urolithiasis: retrospective analysis of 438 cases. *Journal of the American Veterinary Medical Association*, **170**, 414–418.
- Brown, N.O., Patnaik, A.K. & MacEwen, E.G. (1985) Canine hemangiosarcoma: retrospective analysis of 104 cases. *Journal of the American Veterinary Medical Association*, **186** (1), 56–58.
- Bruder, M.C., Shoieb, A.M., Shirai, N., Boucher, G.G. & Brodie, T.A. (2010) Renal dysplasia in Beagle Dogs, *Toxicologic Pathology*, **38**, 1051–1057.
- Brunnberg, L., Schebitz, H., Vollmerhaus, B. *et al.* (1983) Olecranon fracture in the dog: treatment and results. *Kleintierpraxis*, **28**, 17–22.
- Bruun, C.S., Jäderlund, K.H., Berendt, M. *et al.* (2013) A Gly98Val mutation in the N-Myc Downstream Regulated Gene 1 (NDRG1) in Alaskan Malamutes with polyneuropathy. *PLoS One*, **8**, e54547.
- Bruyette, D.S. & Feldman, E.C. (1988) Primary hypoparathyroidism in the dog: Report of 15 cases and review of 13 previously reported cases. *Journal of Veterinary Internal Medicine*, **2** (1), 7–14.
- Bryan, J.N., Keeler, M.R., Henry, C.J. *et al.* (2007) A population study of neutering status as a risk factor for canine prostate cancer. *The Prostate*, **67** (11), 1174–1181.
- Bryden, S.L., White, S.D., Dunston, S.M., Burrows, A.K. & Olivry, T. (2005) Clinical, histopathological and immunological characteristics of exfoliative cutaneous lupus erythematosus in 25 German Short-Haired Pointers. *Veterinary Dermatology*, **16** (4), 239–252.
- Buber, T., Saragusty, J., Ranen, E. *et al.* (2007) Evaluation of lidocaine treatment and risk factors for death associated with gastric dilatation and volvulus in dogs: 112 cases (1997–2005). *Journal of the American Veterinary Medical Association*, **230** (9), 1334–1339.
- Buchanan, J.W. (2001) Patent ductus arteriosus morphology, pathogenesis, types and treatment. *Journal of Veterinary Cardiology*, **3** (1), 7–16.
- Buchanan, J.W. (2004) Tracheal signs and associated vascular anomalies in dogs with persistent right aortic arch. *Journal of Veterinary Internal Medicine*, **18**, 510–514.
- Buchanan, J.W., Beardow, A.W. & Sammarco, C.D. (1997) Femoral artery occlusion in Cavalier King Charles Spaniels. *Journal of the American Veterinary Medical Association*, **211**, 872–874.
- Buchanan, J.W. & Bücheler, J. (1995) Vertebral scale system to measure canine heart size in radiographs. *Journal of the American Veterinary Medical Association*, **206** (2), 194–199.

- Burbidge, H.M. (1995) A review of laryngeal paralysis in dogs. *British Veterinary Journal*, **151**, 71–82.
- Bureau, S., Monnet, E. & Orton, E.C. (2005) Evaluation of survival rate and prognostic indicators for surgical treatment of left-to-right patent ductus arteriosus in dogs: 52 cases (1995–2003). *Journal of the American Veterinary Medical Association*, **227** (11), 1794–1799.
- Burk, R. L., Corley, E. A., Corwin, L. A. (1978) The radiographic appearance of pulmonary histoplasmosis in the dog and cat: a review of 37 case histories. *Journal of the American Veterinary Radiology Society*, **19**, 2–7.
- Burkert, B.A., Kerwin, S.C., Hosgood, G.L., Pechman, R.D. & Fontenelle, J.P. (2005) Signalment and clinical features of diskospondylitis in dogs: 513 cases (1980–2001). *Journal of the American Veterinary Medical Association*, **227** (2), 268–275.
- Burkett, G. (2000) Skin diseases in greyhounds. *Veterinary Medicine*, **95**, 115–124.
- Burnie, A.G. & Weaver, A.D. (1983) Urinary bladder neoplasia in the dog; a review of seventy cases. *Journal of Small Animal Practice*, **24** (3), 129–143.
- Burns, C.G., Bergh, M.S. & McLoughlin, M.A. (2013) Surgical and nonsurgical treatment of peritoneopericardial diaphragmatic hernia in dogs and cats: 58 cases (1999–2008). *Journal of the American Veterinary Medical Association*, **242** (5), 643–650.
- Burns, M.S., Bellhorn, R.W., Impellizzeri, C.W. *et al.* (1988) Development of hereditary tapetal degeneration in the beagle dog. *Current Eye Research*, **7** (2), 103–114.
- Burrows, C.F. & Ignaszewski, L.A. (1990) Canine gastric dilatation–volvulus. *Journal of Small Animal Practice*, **31** (10), 495–501.
- Bussadori, C., Pradelli, D., Borgarelli, M. *et al.* (2009) Congenital heart disease in boxer dogs: Results of 6 years of breed screening. *Veterinary Journal*, **181**, 187–192.
- Cafarchia, C., Romito, D., Sasanelli, M. *et al.* (2004) The epidemiology of canine and feline dermatophytoses in southern Italy. *Mycoses*, **47** (11–12), 508–513.
- Cairo, J., Font, J., Gorraiz, J. *et al.* (1999) Intestinal volvulus in dogs: a study of four clinical cases. *Journal of Small Animal Practice*, **40** (3), 136–140.
- Caldin, M., Carli, E., Furlanello, T. *et al.* (2005) A retrospective study of 60 cases of eccentrocytosis in the dog. *Veterinary Clinical Pathology*, **34** (3), 224–231.
- Callan, M.B., Washabau, R.J., Saunders, H.M. *et al.* (1993) Congenital esophageal hiatal hernia in the Chinese Shar-Pei Dog. *Journal of Veterinary Internal Medicine*, **7** (4), 210–215.
- Cameron, J.M., Maj, M.C., Levandovskiy, V. *et al.* (2007) Identification of a canine model of pyruvate dehydrogenase phosphatase 1 deficiency. *Molecular Genetics and Metabolism*, **90** (1), 15–23.
- Cameron, M.E., Casey, R.A., Bradshaw, J.W.S. *et al.* (2004) A study of environmental and behavioural factors that may be associated with feline idiopathic cystitis. *Journal of Small Animal Practice*, **45** (3), 144–147.
- Campbell, C.A., Horstman, C.L., Mason, D.R. & Evans, R.B. (2010) Severity of patellar luxation and frequency of concomitant cranial cruciate ligament rupture in dogs: 162 cases (2004–2007). *Journal of the American Veterinary Medical Association*, **236** (8), 887–891.
- Campbell, F.E. & Thomas, W.P. (2012) Congenital supra-auricular mitral stenosis in 14 cats. *Journal of Veterinary Cardiology*, **14** (1), 281–292.
- Campbell, K.A., Payne, J.T., Doornink, M.T. & Haggerty, J. (2016) Outcome of tibial closing wedge osteotomy in 55 cranial cruciate ligament-deficient stifles of small dogs (<15 kg). *Veterinary Surgery*, **45**, 1056–1062.
- Campora, C., Freeman, K.P., Lewis, F.I. *et al.* (2011) Determination of haematological reference intervals in healthy adult Greyhounds. *Journal of Small Animal Practice*, **52** (6), 301–309.
- Canier, P., Gallo, L., Sturaro, E. *et al.* (2004) Prevalence of spondylosis deformans and estimates of genetic parameters for the degree of osteophytes development in Italian Boxer Dogs. *Journal of Animal Science*, **82**, 85–92.
- Cannizzo, K.L., McLoughlin, M.A., Mattoon, J.S. *et al.* (2003) Evaluation of transurethral cystoscopy and excretory urography for diagnosis of ectopic ureters in female dogs: 25 cases (1992–2000). *Journal of the American Veterinary Medical Association*, **223** (4), 475–481.
- Cannon, A.B., Westropp, J.L., Ruby, A.L. *et al.* (2007) Evaluation of trends in urolith composition in cats: 5230 cases (1985–2004). *Journal of the American Veterinary Medical Association*, **231** (4), 570–576.

- Capak, D., Bali, R., Butkovic, V. *et al.* (2007) Appearance and frequency of digit neoplastic masses in dogs. *Veterinarska Stannica*, **38** (6), 321–335.
- Capt, A., Spirito, F., Guaguere, E. *et al.* (2005) Inherited junctional epidermolysis bullosa in the German Pointer: Establishment of a large animal model. *Journal of Investigative Dermatology*, **124** (3), 530–535.
- Cardy, T.J.A., Tzounos, C.E., Volk, H.A. & De Decker, S. (2016) Clinical characterization of thoracolumbar and lumbar intervertebral disk extrusions in English Cocker Spaniels. *Journal of the American Veterinary Medical Association*, **248**, 405–412.
- Carette, F., Ragetly, G., Cauzinille, L. (2014) Origin and diagnostic approach to thymoma in dogs and cats. *Pointe Vétérinaire*, **45** (347 part 1), 24–29.
- Cargill, E.J., Famula, T.R., Strain, G.M. *et al.* (2004) Heritability and segregation analysis of deafness in U.S. Dalmatians. *Genetics*, **166** (3), 1385–1393.
- Carmichael, K.P., Miller, M., Rawlings, C.A. *et al.* (1996) Clinical, hematologic, and biochemical features of a syndrome in Bernese Mountain Dogs characterized by hepatocerebellar degeneration. *Journal of the American Veterinary Medical Association*, **208**, 1277–1279.
- Caro-Vadillo, A., García-Guasch, L., Carretón, E., Montoya-Alonso, J.A. & Manubens, J. (2013) Arrhythmic right ventricular cardiomyopathy in Boxer Dogs: a retrospective study of survival. *Veterinary Record*, **172**, 268.
- Carpenter, J.L., Myers, A.M., Connor, M.W. *et al.* (1988) Tuberculosis in five Basset Hounds. *Journal of the American Veterinary Medical Association*, **192** (11), 1563–1568.
- Carr, A.P., Panciera, D.L. & Kidd, L. (2002) Prognostic factors for mortality and thromboembolism in canine immune-mediated hemolytic anemia: a retrospective study of 72 dogs. *Journal of Veterinary Internal Medicine*, **16**, 504–509.
- Carr, J.G., Tobias, K.M. & Smith, L. (2014) Urethral prolapse in dogs: a retrospective study. *Veterinary Surgery*, **43** (5), 574–580.
- Carrera, I., Hammond, G.J.C. & Sullivan, M. (2008) Computed tomographic features of incomplete ossification of the canine humeral condyle. *Veterinary Surgery*, **37** (3), 226–231.
- Carrig, C.B., Sponenberg, D.P., Schmidt, G.M. *et al.* (1988). Inheritance of associated ocular and skeletal dysplasia in Labrador Retrievers. *Journal of the American Veterinary Medical Association*, **193** (10), 1269–1272.
- Carter, W.J., Crispin, S.M., Gould, D.J. *et al.* (2005) An immunohistochemical study of uveodermatologic syndrome in two Japanese Akita dogs. *Veterinary Ophthalmology*, **8** (1), 17–24.
- Casal, M.L., Giger, U., Bovee, K.C. *et al.* (1995) Inheritance of cystinuria and renal defect in Newfoundlands. *Journal of the American Veterinary Medical Association*, **207** (12), 1585–1589.
- Casal, M.L., Munure, R.M., Janis, M.A., Werner, P. & Henthorn, P.S. (2006) Epilepsy in Irish Wolfhounds. *Journal of Veterinary Internal Medicine*, **20** (1), 131–135.
- Casal, M.L., Straumann, U. & Sigg, C. (1994) Congenital hypotrichosis with thymic aplasia in nine Birman kittens. *Journal of the American Animal Hospital Association*, **30**, 600–602.
- Casamián-Sorrosal, D., Chong, S.K., Fonfara, S. & Helps, C. (2014) Prevalence and demographics of the MYBPC3-mutations in Ragdolls and Maine Coons in the British Isles. *Journal of Small Animal Practice*, **55** (5), 269–273.
- Case, L.C., Ling, G.V., Franti, C.E. *et al.* (1992) Cystine-containing urinary calculi in dogs: 102 cases (1981–1989). *Journal of the American Veterinary Medical Association*, **201** (1), 129–133.
- Castellano, M.C. & Idiart, J. R. (2005) Multifocal renal cystadenocarcinoma and nodular dermatofibrosis in dogs. *Compendium on Continuing Education for the Practicing Veterinarian*, **27** (11), 846–853.
- Castellano, M.C., Idiart, J.R., Ruager, J. *et al.* (2000) Generalised nodular dermatofibrosis and cystic renal disease in five German Shepherd Dogs. *Canine Practice*, **25** (2), 18–21.
- Castro, N.B., Boos, G.S., Wurster, F. *et al.* (2013) Post-mortem diagnosis of gastric dilatation volvulus syndrome in dogs. *Acta Scientiae Veterinariae*, **41**, 1120.
- Catchpole, B., Kennedy, L.J., Davison, L.J. & Ollier, W.E.R. (2008) Canine diabetes mellitus: from phenotype to genotype. *Journal of Small Animal Practice*, **49** (1), 4–10.
- Catchpole, B., Ristic, J.M., Freeman, L.M. *et al.* (2005) Canine diabetes mellitus: can old dogs teach us new tricks? *Diabetologica*, **48** (10), 1948–1956.

- Cattanach, B.M., Dukes-McEwan, J., Wotton, P.R., Stephenson, H.M. & Hamilton, R.M. (2015) A pedigree-based genetic appraisal of Boxer ARVC and the role of the Striatin mutation. *Veterinary Record*, **176**, 492–492.
- Cauzinille, L. & Kornegay, J.N. (1996) Fibrocartilaginous embolism of the spinal cord in dogs: Review of 36 histologically confirmed cases and retrospective study of 26 suspected cases. *Journal of Veterinary Internal Medicine*, **10** (4), 241–245.
- Caywood, D.D., Klausnier, J.S., O'Leary, T.P. *et al.* (1988) Pancreatic insulin-secreting neoplasms: clinical diagnostic and prognostic features in 73 dogs. *Journal of the American Animal Hospital Association*, **24** (5), 577–584.
- Cerda-Gonzalez, S., Talarico, L. & Todhunter, R. (2016) Noninvasive assessment of neuromuscular disease in dogs: use of the 6-minute walk test to assess submaximal exercise tolerance in dogs with centronuclear myopathy. *Journal of Veterinary Internal Medicine*, **30** (3), 808–812.
- Cerundolo, R. (2004) Generalized *Microsporum canis* dermatophytosis in six Yorkshire Terrier dogs. *Veterinary Dermatology*, **15** (3), 181–187.
- Cerundolo, R., Lloyd, D.H., McNeil, P. *et al.* (2000) An analysis of factors underlying hypotrichosis and alopecia in Irish Water Spaniels in the United Kingdom. *Veterinary Dermatology*, **11**, 107–122.
- Cerundolo R., Mauldin E.A., Goldschmidt M.H. *et al.* (2005) Adult-onset hair loss in Chesapeake Bay Retrievers: a clinical and histological study. *Veterinary Dermatology*, **16**, 39–46.
- Cetinkava, M.A., Yardimki, C. & Saglam, M. (2007) Carpal laxity syndrome in forty-three puppies. *Veterinary and Comparative Orthopaedics and Traumatology*, **20**, 126–130.
- Chahory, S., Crasta, M., Trio, S. *et al.* (2004) Three cases of prolapse of the nictitans gland in cats. *Veterinary Ophthalmology*, **7** (6), 417–419.
- Chandler, H.L., Gemensky-Metzler, A. J., Bras, I. D. *et al.* (2010) In vivo effects of adjunctive tetracycline treatment on refractory corneal ulcers in dogs. *Journal of the American Veterinary Medical Association*, **237**, 378–386.
- Chandler, M.L., Elwood, C., Murphy, K.F., Gajanayake, I. and Syme, H.M. (2007) Juvenile nephropathy in 37 Boxer Dogs. *Journal of Small Animal Practice*, **48**, 690–694.
- Chang, H.-S., Kamishina, H., Mizukami, K. *et al.* (2013) Genotyping assays for the canine degenerative myelopathy-associated c.118G>A (p.E40K) mutation of the SOD1 gene using conventional and real-time PCR methods: a high prevalence in the Pembroke Welsh Corgi breed in Japan. *Journal of Veterinary Medical Science*, **75** (6), 795–798.
- Chapman, P.S., Boag, A.K., Guitian, J. & Boswood, A. (2004) *Angiostrongylus vasorum* infection in 23 dogs (1999–2002). *Journal of Small Animal Practice*, **45** (9), 435–440.
- Charlesworth, T.M. & Sturgess, C.P. (2012) Increased incidence of thoracic wall deformities in related Bengal kittens *Journal of Feline Medicine and Surgery*. **14** (6), 566–570.
- Chase, K. (2006) Understanding the genetics of autoimmune disease: two loci that regulate late onset Addison's disease in Portuguese Water Dogs. *International Journal of Immunogenetics*, **33** (3), 179–184.
- Chastain, C.B. & Panciera, D. (2002) Thyroid function testing in Greyhounds. *Small Animal Clinical Endocrinology*, **12** (1), 4.
- Chastain, C.B. & Panciera, D. (2004) Sex hormone concentrations in dogs with alopecia. *Small Animal Endocrinology*, **14** (1), 37–38.
- Chastain, C.B. & Panciera, D. (2007) Primary hyperparathyroidism in Keeshonden. *Small Animal Clinical Endocrinology*, **17** (2), 8.
- Chastain, C.B., Panciera, D. & Waters, C. (1999) Associations between age, parity, hormonal therapy and breed, and pyometra in Finnish dogs. *Small Animal Clinical Endocrinology*, **9** (2), 18.
- Chastain, C.B., Panciera, D. & Waters, C. (2001) Eclampsia in dogs: 31 cases (1995–1998). *Small Animal Clinical Endocrinology*, **11** (1), 9.
- Chastain, C.B., Turk, M.A. & O'Brien, D. (1988) Benign cutaneous mastocytomas in two litters of Siamese kittens. *Journal of the American Veterinary Medical Association*, **193** (8), 959–960.
- Chaudieu, G. (1995) Study of inherited focal degeneration (retinopathy) in the Borzoi: a report on 160 dogs. *Pratique Médicale & Chirurgicale de l'Animal de Compagnie*, **30**, 461–472.
- Chaudieu, G. (2001) Study of progressive retinal atrophy in Border Collie. Data provided by 213 examinations carried out on 161 dogs. *Pratique Médicale et Chirurgicale de l'Animal de Compagnie*, **36**, 55–73.

- Chaudieu, G. & Molon-Noblot, S. (2004) Early retinopathy in the Bernese Mountain Dog in France: preliminary observations. *Veterinary Ophthalmology*, **7** (3), 175–184.
- Chaudieu, G., Molon-Noblot, S. & Duprat, P. (1993) Primary lens luxation in the Brittany Spaniel: clinical aspects and aetiopathogenic study, *Pratique Médicale et Chirurgicale de l'Animal de Compagnie*, **28**, 37–47.
- Chaudieu, G., Olivier, A., Thomas, A. *et al.* (2014) Atrophie progressive de la rétine du Border Collie: étude rétrospective (1996–2012), *Revue Vétérinaire Clinique*, **49**, 93–101.
- Chaves, R.O., Beckmann, D.V., Copat, B. *et al.* (2014) Doença vestibular em cães: 81 casos (2006–2013). *Pesquisa Veterinária Brasileira*, **34**, 1231–1235.
- Chavkin, M., Roberts, S., Salman, M., Severin, G. & Scholten, N. (1994) Risk factors for development of chronic superficial keratitis in dogs. *Journal of the American Veterinary Medical Association*, **204** (10), 1630–1634.
- Chen, X., Johnson, G.S., Schnabel, R.D. *et al.* (2008) A neonatal encephalopathy with seizures in standard poodle dogs with a missense mutation in the canine ortholog of ATF2. *neurogenetics*, **9** (1), 41–49.
- Cheng, S., Wigney, D., Haase, B. (2016) Inheritance of chronic superficial keratitis in Australian Greyhounds. *Animal Genetics*, **47** (5), 629.
- Chetboul, V., Charles, V., Nicolle, A. *et al.* (2006a) Retrospective study of 156 atrial septal defects in dogs and cats (2001–2005). *Journal of Veterinary Medicine Series A*, **53**, 179–184.
- Chetboul, V., Petit, A., Gouni, V. *et al.* (2012) Prospective echocardiographic and tissue Doppler screening of a large Sphynx cat population: reference ranges, heart disease prevalence and genetic aspects. *Journal of Veterinary Cardiology*, **14** (4), 497–509.
- Chetboul, V., Trolle, J.M., Nicolle, A. *et al.* (2006b) Congenital heart diseases in the Boxer dog: a retrospective study of 105 cases (1998–2005). *Journal of Veterinary Medicine A, Physiology, Pathology and Clinical Medicine*, **53**, 346–351.
- Chew, D.J., DiBartola, S.P., Boyce, J.T. *et al.* (1983) Juvenile renal disease in Doberman Pinscher dogs. *Journal of the American Veterinary Medical Association*, **182** (5), 481–485.
- Chieffo, C., Stalis, I.H., Van Winkle, T.J., Haskins, M.E. & Patterson, D.F. (1994) Cerebellar Purkinje's cell degeneration and coat color dilution in a family of Rhodesian Ridgeback dogs. *Journal of Veterinary Internal Medicine*, **8** (2), 112–116.
- Choisunirachon, N., Lee Shuan, L., Tanaka, Y. *et al.* (2014) Retrospective study of computed tomographic characterization of canine oral malignant melanoma in 24 dogs. *Thai Journal of Veterinary Medicine*, **44**, 497–503.
- Chrisman, C.L. (1990) Dancing Doberman disease: clinical findings and prognosis. *Progress in Veterinary Neurology*, **1** (1), 83–90.
- Chrisman, C.L. (1992) Neurological diseases of Rottweilers: neuroaxonal dystrophy and leukoencephalomalacia. *Journal of Small Animal Practice*, **33** (10), 500–504.
- Christensen, N., Canfield, P., Martin, P. *et al.* (2009) Cytopathological and histopathological diagnosis of canine splenic disorders. *Australian Veterinary Journal*, **87**, 175–181.
- Christiansen, J.S., Hottinger, H.A., Allen, L., Phillips, L. & Aronson, L.R. (2000) Hepatic microvascular dysplasia in dogs: a retrospective study of 24 cases (1987–1995). *Journal of the American Animal Hospital Association*, **36** (5), 385–389.
- Chung, T.H., Ryu, M.-H., Kim, D.Y. *et al.* (2009) Topical tacrolimus (FK506) for the treatment of feline idiopathic facial dermatitis. *Australian Veterinary Journal*, **87** (10), 417–420.
- Churcher, R.K. & Watson, A.D. (1997) Canine histiocytic ulcerative colitis. *Australian Veterinary Journal*, **75** (10), 710–713.
- Ciekot, P.A., Powers, B.E., Withrow, S.J. *et al.* (1994) Histologically low grade yet biologically high grade fibrosarcomas of the mandible and maxilla in dogs: 25 cases (1982–1991). *Journal of the American Veterinary Medical Association*, **204** (4), 610–615.
- Clark, L.A., Credille, K.M., Murphy, K.E. & Rees, C.A. (2005) Linkage of dermatomyositis in the Shetland Sheepdog to chromosome 35. *Veterinary Dermatology*, **16** (6), 392–394.
- Clark, P. & Parry, B.W. (1995) Survey of Irish Wolfhounds in Australia for von Willebrand's disease. *Australian Veterinary Journal*, **72** (10), 393.

- Clemente, M., Marin, L., Iazbik, M.C. *et al.* (2010) Serum concentrations of IgG, IgA, and IgM in retired racing Greyhound dogs. *Veterinary Clinical Pathology*, **39** (4), 436–439.
- Clements, D.N., Fitzpatrick, N., Carter, S.D. & Day, P.J. (2007) Cartilage gene expression correlates with radiographic severity of canine elbow osteoarthritis. *Veterinary Journal*, **179** (2), 211–218.
- Clements, D.N., Gear, R.N.A., Tattersall, J. *et al.* (2004) Type I immune-mediated polyarthritis in dogs: 39 cases (1997–2002). *Journal of the American Veterinary Medical Association*, **224**, 1323–1327.
- Clercx, C., Reichler, I., Peeters, D. *et al.* (2003) Rhinitis/bronchopneumonia syndrome in Irish Wolfhounds. *Journal of Veterinary Internal Medicine*, **17** (6), 843–849.
- Clinkenbeard, K.D., Wolf, A.M., Cowell, R.L. *et al.* (1989) Canine disseminated histoplasmosis. *Compendium on Continuing Education for the Practicing Veterinarian*, **11** (11), 1347–1351, 1354–1357, 1360.
- Coates, J.R., March, P.A., Oglesbee, M. *et al.* (2007) Clinical characterization of a familial degenerative myelopathy in Pembroke Welsh Corgi dogs. *Journal of Veterinary Internal Medicine*, **21** (6), 1323–1331.
- Coates, J.R., O'Brien, D.P., Kline, K.L. *et al.* (2002) Neonatal cerebellar ataxia in Coton de Tulear dogs. *Journal of Veterinary Internal Medicine*, **16**, 680–689.
- Cochrane (2017). <http://www.cochrane.org> (accessed October 2017).
- Cockcroft, P.D. & Holmes, M.A. (2003). *Handbook of Evidence-Based Veterinary Medicine*. Blackwell, Oxford.
- Cockett, P.A. & Jones, D.G.C. (1985) The incidence of humeral condylar fractures in the dog: a survey of seventy-nine cases. *Journal of Small Animal Practice*, **26**, 437–444.
- Cohn-Bendit, F. (1995) Tumeurs malignes de la thyroïde chez le chien: étude rétrospective de 137 cas. *Pratique Médicale et Chirurgicale de l'Animal de Compagnie*, **30**, 85–92.
- Cole, L.K., Samii, V.F., Wagner, S.O. & Rajala-Schultz, P.J. (2015) Diagnosis of primary secretory otitis media in the Cavalier King Charles Spaniel. *Veterinary Dermatology*, **26**, 459–e107.
- Collier, L.L., Bryan, G.M. & Prieur, D.J. (1979) Ocular manifestations of the Chédiak-Higashi syndrome in four species of animals. *Journal of the American Veterinary Medical Association*, **175** (6), 587–590.
- Collins, B.K., Collier, L.L., Johnson, G.S. *et al.* (1992) Familial cataracts and concurrent ocular anomalies in Chow Chows. *Journal of the American Veterinary Medical Association*, **200** (10), 1485–1491.
- Collins, D., Angles, J.M., Christodoulou, J. *et al.* (2013) Severe subacute necrotizing encephalopathy (Leigh-like syndrome) in American Staffordshire Bull Terrier Dogs. *Journal of Comparative Pathology*, **148**, 345–353.
- Colombo, S., Scarampella, F., Ordeix, L. *et al.* (2012) Dermatophytosis and papular eosinophilic/mastocytic dermatitis (urticaria pigmentosa-like dermatitis) in three Devon Rex cats. *Journal of Feline Medicine and Surgery*, **14** (7), 498–502.
- Comerford, E.J., Tarlton, J.F., Wales, A. *et al.* (2006) Ultrastructural differences in cranial cruciate ligaments from dogs of two breeds with a differing predisposition to ligament degeneration and rupture. *Journal of Comparative Pathology*, **134** (1), 8–16.
- Comito, B., Knowles, K.E. & Strain, G.M. (2012) Congenital deafness in Jack Russell Terriers: prevalence and association with phenotype. *Veterinary Journal*, **193** (2), 404–407.
- Conceição, L.G., Acha, L.M.R., Borges, A.S. *et al.* (2011) Epidemiology, clinical signs, histopathology and molecular characterization of canine leproid granuloma: a retrospective study of cases from Brazil. *Veterinary Dermatology*, **22**, 249–256.
- Conrado, F.O., Oliveira, S.T., Lacerda, L.A. *et al.* (2014) Clinicopathologic and electrocardiographic features of Akita dogs with high and low erythrocyte potassium phenotypes. *Veterinary Clinical Pathology*, **43** (1), 50–54.
- Constantino-Casas, F., Mayhew, D., Hoather, T.M. & Dobson, J.M. (2011) The clinical presentation and histopathologic-immunohistochemical classification of histiocytic sarcomas in the Flat Coated Retriever. *Veterinary Pathology*, **48** (3), 764–771.
- Contreary, C.L., Outerbridge, C.A., Affolter, V.K. *et al.* (2015) Canine sterile nodular panniculitis: a retrospective study of 39 dogs. *Veterinary Dermatology* **26** (6) 451–458.
- Contreras, A.M., Li, O.M., Gavidia C.M., Diego, D.C. & Hoyos, L.A. (2009) Retrospective case control study of canine ehrlichiosis at the Veterinary Faculty of Universidad Nacional Mayor de San Marcos: 2002–2005 period. *Revisita de Investigaciones Veterinarias del Perú*, **20** (2), 270–276.

- Cooley, D.M., Beranek, B.C., Schlittler, D.L. *et al.* (2002) Endogenous gonadal hormone exposure and bone sarcoma risk. *Cancer Epidemiology Biomarkers and Prevention*, **11** (11), 1434.
- Cooper, A.E., Ahonen, S., Rowlan, J.S. *et al.* (2014a) A novel form of progressive retinal atrophy in Swedish Vallhund dogs. *PLoS One*, **9** (9), e106610.
- Cooper, B.J., Valentine, B.A., Wilson, S. *et al.* (1988) Canine muscular dystrophy: confirmation of X-linked inheritance. *Journal of Heredity*, **79**, 405–408.
- Cooper, J.J., Schatzberg, S.J., Vernau, K.M. *et al.* (2014b) Necrotizing meningoencephalitis in atypical dog breeds: a case series and literature review. *Journal of Veterinary Internal Medicine*, **28** (1), 198–203.
- Cooper, R.L., Drobatz, K.J., Lennon, E.M. *et al.* (2015) Retrospective evaluation of risk factors and outcome predictors in cats with diabetic ketoacidosis (1997–2007): 93 cases. *Journal of Veterinary Emergency and Critical Care*, **25** (2), 263–272.
- Coopman, F., Verhoeven, G., Saunders, J., Duchateau, L. & Bree, H.v. (2008) Prevalence of hip dysplasia, elbow dysplasia and humeral head osteochondrosis in dog breeds in Belgium. *Veterinary Record*, **163** (22), 654–658.
- Copping, J. (2014) Pedigree dogs ‘as healthy as mongrels’, say vets. *Sunday Telegraph*, 13 April 2014. <http://www.telegraph.co.uk/news/health/pets-health/10762988/Pedigree-dogs-as-healthy-as-mongrels-say-vets.html>.
- Corbee, R.J. (2013) Obesity in show dogs. *Journal of Animal Physiology and Animal Nutrition*, **97** (5), 904–910.
- Corcoran, B.M., King, L.G., Schwarz, T., Hammond, G. & Sullivan, M. (2011) Further characterisation of the clinical features of chronic pulmonary disease in West Highland White Terriers. *Veterinary Record*, **168** (13), 355–355.
- Corcoran, K.A. & Koch, S.A. (1993) Uveal cysts in dogs: 28 cases (1989–1991). *Journal of the American Veterinary Medical Association*, **203** (4), 545–546.
- Cork, L.C., Griffin, J.W., Choy, C., Padula, C.A. & Price, D. L. (1982) Pathology of motor neurons in accelerated hereditary canine spinal muscular atrophy. *Laboratory Investigation*, **46**, 89–99.
- Cork, L.C., Price, D.L., Griffin, J.W. *et al.* (1990) Hereditary canine spinal muscular atrophy: canine motor neuron disease. *Canadian Journal of Veterinary Research*, **54**, 77–82.
- Cornelissen, J.M.R. & Hopster, H. (2010) Dog bites in the Netherlands: a study of victims, injuries, circumstances and aggressors to support evaluation of breed specific legislation. *Veterinary Journal*, **186** (3), 292–298.
- Corr, S. A., Draffan, D., Kulendra, E., Carmichael, S. & Brodbelt, D. (2010) Retrospective study of Achilles mechanism disruption in 45 dogs. *Veterinary Record*, **167**, 407–411.
- Cosgrove, L., Hammond, G. & McLauchlan, G. (2015) Primary portal vein hypoplasia and SLC2A9 mutation associated with urate urolithiasis in a Spanish Water Dog. *Canadian Veterinary Journal*, **56** (11), 1153–1157.
- Cottrell, B.D. & Barnett, K.C. (1988) Primary glaucoma in the Welsh Springer Spaniel. *Journal of Small Animal Practice*, **29** (3), 185–199.
- Courcier, E.A., Thomson, R.M., Mellor, D.J. & Yam, P.S. (2010) An epidemiological study of environmental factors associated with canine obesity. *Journal of Small Animal Practice*, **51** (7), 362–367.
- Couto, C.G., Krakowka, S., Johnson, G. *et al.* (1989) In vitro immunologic features of Weimaraner dogs with neutrophil abnormalities and recurrent infections. *Veterinary Immunology and Immunopathology*, **23** (1), 103–112.
- Couto, K.M., Iazbik, M.C., Marin, L.M. *et al.* (2015) Plasma N-terminal pro-B-type natriuretic peptide concentration in healthy retired racing Greyhounds. *Veterinary Clinical Pathology*, **44** (3), 405–409.
- Cowan, L.A., Hertzke, D.M., Fenwick, B.W. *et al.* (1997) Clinical and clinicopathologic abnormalities in Greyhounds with cutaneous and renal glomerular vasculopathy: 18 cases (1992–1994). *Journal of the American Veterinary Medical Association*, **210**, 789–793.
- Cowan, S., Bartges, J.W. & Gompf, R.E. (2004) Giant platelet disorder in the Cavalier King Charles Spaniel. *Experimental Haematology*, **32**, 344–350.
- Cox, R.H., Peterson, L.H. & Detweiler, D.K. (1976) Comparison of arterial hemodynamics in the mongrel dog and the Racing Greyhound. *American Journal of Physiology*, **230**, 211–218.

- Cox, V.S., Wallace, L.J. & Jessen, C.R. (1978) An anatomic and genetic study of canine cryptorchidism. *Teratology*, **18** (2), 233–240.
- Cox, V.S., Wallace, L.J., Anderson, V.E. & Rushmer, R.A. (1980) Hereditary esophageal dysfunction in the Miniature Schnauzer dog. *American Journal of Veterinary Research*, **41** (3), 326–330.
- Coyle, K. & Steinberg, H. (2004) Characterization of lymphocytes in canine gastrointestinal lymphoma. *Veterinary Pathology*, **41** (2), 141–146.
- Coyne, B.E. & Fingland, R.B. (1992) Hypoplasia of the trachea in dogs: 103 cases (1974–1990). *Journal of the American Veterinary Medical Association*, **5**, 768–772.
- Craig, L.E., Julian, M.E., & Ferracone, J.D. (2002). The diagnosis and prognosis of synovial tumors in dogs: 35 cases. *Veterinary Pathology Online*, **39** (1), 66–73.
- Craig, L.E. (2001) Physeal dysplasia with slipped capital femoral epiphysis in 13 cats. *Veterinary Pathology*, **38** (1), 92–97.
- Craven, M., Simpson, J.W., Ridyad, A.E. *et al.* (2004) Canine inflammatory bowel disease: retrospective analysis of diagnosis and outcome in 80 cases (1995–2002). *Journal of Small Animal Practice*, **45** (7), 336–342.
- Crawley, A.C., Muntz, F.H., Haskins, M.E. *et al.* (2003) Prevalence of mucopolysaccharidosis type VI mutations in Siamese cats. *Journal of Veterinary Internal Medicine*, **17** (4), 495–498.
- Credille, K.M., Minor, J.S., Barnhart, K.F. *et al.* (2009) Transglutaminase 1-deficient recessive lamellar ichthyosis associated with a LINE-1 insertion in Jack Russell Terrier dogs. *British Journal of Dermatology*, **161** (2), 265–272.
- Crews, L.J., Feeney, D.A., Jessen, C.R. *et al.* (2008) Radiographic findings in dogs with pulmonary blastomycosis 125 cases (1989–2006). *Journal of the American Veterinary Medical Association*, **232** (2), 215–222.
- Crews, L.J., Feeney, D.A., Jessen, C.R., Rose, N.D. & Matise, I. (2009) Clinical, ultrasonographic, and laboratory findings associated with gallbladder disease and rupture in dogs: 45 cases (1997–2007). *Journal of the American Veterinary Medical Association*, **234** (3), 359–366.
- Crha, M., Konvalinová, J., Fichtel, T. & Pavlica, Z. (2003) Migration of grass awn in dogs: a retrospective study of 140 cases. *Kleintierpraxis* **48**, 427–433.
- Cribb, A.E. (1988) Feline gastrointestinal adenocarcinoma: a review and retrospective study. *Canadian Veterinary Journal*, **29** (9), 709–712.
- Crispin, S.M. (1988) Cystalline corneal dystrophy in the dog. Histochemical and ultrastructural study. *Cornea*, **7** (2), 149–161.
- Cruikshank, J., Quaas, R.L., Li, J. *et al.* (2009) Genetic analysis of ventricular arrhythmia in young German Shepherd Dogs. *Journal of Veterinary Internal Medicine*, **23** (2), 264–270.
- Crumlish, P.T., Sweeney, T., Jones, B. & Angles, J.M. (2006) Hypertrophic osteodystrophy in the Weimaraner dog: Lack of association between DQA1 alleles of the canine MHC and hypertrophic osteodystrophy. *Veterinary Journal*, **171** (2), 308–313.
- Cruz, F., Vilà, C. & Webster, M.T. (2008) The legacy of domestication: accumulation of deleterious mutations in the dog genome. *Molecular Biology and Evolution*, **25** (11), 2331–2336.
- Culp, W.T.N., Ehrhart, N., Withrow, S.J. *et al.* (2013) Results of surgical excision and evaluation of factors associated with survival time in dogs with lingual neoplasia: 97 cases (1995–2008). *Journal of the American Veterinary Medical Association*, **242** (10), 1392–1397.
- Cummings, J.F., Cooper, B.J., de Lahunta, A. & van Winkle, T.J. (1981) Canine inherited hypertrophic neuropathy. *Acta neuropathologica*, **53** (2), 137–143.
- Cummings, J.F. & de Lahunta, A. (1978) Hereditary myelopathy of Afghan hounds, a myelinolytic disease, *Acta neuropathologica*, **42**, 73–81.
- Cummings, J.F., de Lahunta, A., Braund, K.G. & Mitchell, W.J. (1983) Hereditary sensory neuropathy. Nociceptive loss and acral mutilation in Pointer dogs: canine hereditary sensory neuropathy. *American Journal of Pathology*, **112** (1), 136–138.
- Cummings, J.F., de Lahunta, A., Simpson, S.T. & McDonald, J.M. (1984) Reduced substance P-like immunoreactivity in hereditary sensory neuropathy of Pointer dogs. *Acta neuropathologica*, **63** (1), 33–40.

- Cummings, J.F., Summers, B.A., de Lahunta, A. & Lawson, C. (1986) Tremors in Samoyed pups with oligodendrocyte deficiencies and hypomyelination. *Acta neuropathologica*, **71** (3), 267–277.
- Cunningham, J.G. & Farnbach, G.C. (1988) Inheritance and idiopathic canine epilepsy. *Journal of the American Animal Hospital Association*, **24**, 421–424.
- Curtis, R. (1988) Retinal diseases in the dog and cat: an overview and update. *Journal of Small Animal Practice*, **29** (7), 397–415.
- Curtis, R. & Barnett, K.C. (1980) Primary lens luxation in the dog. *Journal of Small Animal Practice*, **21** (12), 657–668.
- Curtis, R. & Barnett, K.C. (1981) Canine adenovirus-induced ocular lesions in the Afghan hound. *Cornell veterinarian*, **71**, 85–95.
- Curtis, R. & Barnett, K.C. (1989) A survey of cataracts in Golden and Labrador Retrievers. *Journal of Small Animal Practice*, **30** (5), 277–286.
- Curtis, R., Barnett, K.C. & Leon, A. (1987) An early onset retinal dystrophy with dominant inheritance in the Abyssinian cat. Clinical and pathological findings. *Investigative Ophthalmology and Visual Science*, **28** (1), 131–139.
- Curtis, R., Barnett, K.C. & Startup, F.G. (1983) Primary lens luxation in the Miniature Bull Terrier. *Veterinary Record*, **112** (14), 328–330.
- Cuthbert, S., Morton, M. & Lowrie, M. (2016) Immunodeficiency syndrome in a Weimaraner involving the peripheral nervous system. *Veterinary Record Case Reports*, **4** (1).
- D'Anjou, Penninck, D., Cornejo, L. *et al.* (2004) Ultrasonographic diagnosis of portosystemic shunting in dogs and cats. *Veterinary Radiology and Ultrasound*, **45** (5), 424–437.
- da Costa, R.C. (2010) Cervical spondylomyelopathy (wobbler syndrome) in dogs. *Veterinary Clinics of North America: Small Animal Practice*, **40** (10), 881–913.
- Dahl, K., Moe, L., Indrebø, A. & Gamlem, H. (2002) Incidence of mammary tumours in related Boxers. *Norsk Veterinærtidsskrift*, **114**, 615–622.
- Dahlbom, M., Anderson, M., Juga, J. & Alanko, M. (1997) Fertility parameters in male Irish Wolfhounds: a two-year follow-up study. *Journal of Small Animal Practice*, **38** (12), 547–550.
- Dahlgren, S., Ziener, M.L. & Lingaas, F. (2016) A genome-wide association study identifies a region strongly associated with symmetrical onychomadesis on chromosome 12 in dogs. *Animal Genetics*, **47** (6), 708–716.
- Dambach, D.M., Lannon, A., Sleeper, M.M. & Buchanan, J. (1999) Familial dilated cardiomyopathy of young Portuguese Water Dogs. *Journal of Veterinary Internal Medicine*, **13** (1), 65–71.
- Dambach, D.M., Smith, C.A., Lewis, R.M. *et al.* (1997) Morphologic, immunohistochemical, and ultrastructural characterization of a distinctive renal lesion in dogs putatively associated with *Borrelia burgdorferi* infection: 49 cases (1987–1992). *Veterinary Pathology*, **34** (2), 85–96.
- Damur-Djuric, N., Steffen, F., Hassig, M. *et al.* (2006) Lumbosacral transitional vertebrae in dogs: classification, prevalence, and associated with sacroiliac morphology. *Veterinary Radiology and Ultrasound*, **47**, 32–38.
- Dandrieux, J.R.S., Noble, P.J.M., Halladay, L.J., McLean, L. & German, A.J. (2013) Canine breed predispositions for marked hypocobalaminaemia or decreased folate concentration assessed by a laboratory survey. *Journal of Small Animal Practice*, **54** (3), 143–148.
- Dang, Q.Q., Dung, T.H. & Dung, C.A. (2017) The relation of body score (body height/body length) and haplotype E on Phu Quoc Ridgeback dogs (*Canis familiaris*). *Journal of Entomology and Zoology Studies*, **5** (1), 388–394.
- Darke, P.G.G. (1989) Congenital heart disease in dogs and cats. *Journal of Small Animal Practice*, **30**, 599–607.
- Davidson, A.G., Bell, E.J., Lees, G.E. *et al.* (2007) Genetic causes of autosomal recessive hereditary nephropathy in the English Cocker Spaniel. *Journal of Veterinary Internal Medicine*, **21** (3), 394–401.
- Davidson, A.P., Pollard, R.E., Bannasch, D.L. *et al.* (2004) Inheritance of cricopharyngeal dysfunction in Golden Retrievers. *American Journal of Veterinary Research*, **65** (3), 344–349.
- Davies, C. & Troy, G.C. (1996) Deep mycotic infections in cats. *Journal of the American Animal Hospital Association*, **32**, 380–391.

- Davies, D.R. & Irwin, P.J. (2003) Degenerative neurological and neuromuscular disease in young Rottweilers. *Journal of Small Animal Practice*, **44** (9), 388–394.
- Davison, L.J., Herrtage, M.E. & Catchpole, B. (2005) Study of 253 dogs in the United Kingdom with diabetes mellitus. *Veterinary Record*, **156** (15), 467–471.
- Day, M.J. (1997a) An immunohistochemical study of the lesions of demodicosis in the dog. *Journal of Comparative Pathology*, **116** (2), 203–216.
- Day, M.J. (1997b) Review of thymic pathology in 30 cats and 36 dogs. *Journal of Small Animal Practice*, **38** (9), 393–403.
- Day, M.J. (1999) Possible immunodeficiency in related Rottweiler dogs. *Journal of Small Animal Practice*, **40** (12), 561–568.
- Day, M.J. & Weaver, B.M.Q. (1992) Pathology of surgically resected tissue from 305 cases of anal furunculosis in the dog. *Journal of Small Animal Practice*, **33** (12), 583–589.
- D'Costa, S., Yoon, B.-I., Kim, D.-Y. *et al.* (2012) Morphologic and molecular analysis of 39 spontaneous feline pulmonary carcinomas. *Veterinary Pathology*, **49** (6), 971–978.
- De Causmaecker, V., Daminet, S. & Paepe, D. (2009) Diabetes ketoacidosis and diabetes ketosis in 54 dogs: a retrospective study. *Vlaams Diergeneeskundig Tijdschrift*, **78**, 327–337.
- De Decker, S., Warner, A.-S. & Volk, H.A. (2017) Prevalence and breed predisposition for thoracolumbar intervertebral disc disease in cats. *Journal of Feline Medicine and Surgery*, **19** (4), 419–423.
- De Lorenzi, D., Bertonecello, D. & Drigo, M. (2009) Bronchial abnormalities found in a consecutive series of 40 brachycephalic dogs. *Journal of the American Veterinary Medical Association*, **235** (7), 835–840.
- De Maria, R., Divari, S., Bo, S. *et al.* (1998) Beta-galactosidase deficiency in a Korat cat: a new form of feline GM1-gangliosidosis. *Acta Neuropathologica*, **96** (3), 307–314.
- de Morais, H.S., DiBartola, S.P. & Chew, D.J. (1996) Juvenile renal disease in Golden Retrievers: 12 cases (1984–1994). *Journal of the American Veterinary Medical Association*, **209** (4), 792–797.
- De Risio, L., Forman, O.P., Mellersh, C.S. *et al.* (2016a) Paroxysmal dyskinesia in Norwich Terrier Dogs. *Movement Disorders in Clinical Practice*, **3** (6), 573–579.
- De Risio, L., Freeman, J. & Lewis, T. (2016b) Prevalence, heritability and genetic correlations of congenital sensorineural deafness and coat pigmentation phenotype in the English Bull Terrier. *BMC Veterinary Research*, **12**, 146.
- De Risio, L., Lewis, T., Freeman, J. *et al.* (2011) Prevalence, heritability and genetic correlations of congenital sensorineural deafness and pigmentation phenotypes in the Border Collie. *Veterinary Journal*, **188**, 286–290.
- De Risio, L., Newton, R., Freeman, J. & Shea, A. (2015) Idiopathic epilepsy in the Italian Spinone in the United Kingdom: prevalence, clinical characteristics, and predictors of survival and seizure remission. *Journal of Veterinary Internal Medicine*, **29** (3), 917–924.
- Debenham, S.L., Millington, A., Kijast, J. *et al.* (2002) Canine leucocyte adhesion deficiency in Irish Red and White Setters. *Journal of Small Animal Practice*, **43** (2), 74–75.
- Dedola, C., Ressel, L., Hill, P. B., van den Broek, A.H.M. & Thoday, K. L. (2010) Idiopathic generalized sebaceous gland hyperplasia of the Border Terrier: a morphometric study. *Veterinary Dermatology*, **21**, 494–502.
- Deehr, A.J. (1998) A histopathological study of iridociliary cysts and glaucoma in Golden Retrievers. *Progress in Veterinary and Comparative Ophthalmology*, **1** (2), 153–158.
- Dees, D.D., Fritz, K.J., MacLaren, N.E. *et al.* (2014) Efficacy of prophylactic antiglaucoma and anti-inflammatory medications in canine primary angle-closure glaucoma: a multicenter retrospective study (2004–2012). *Veterinary Ophthalmology*, **17**, 195–200.
- Deforest, M.E. & Basrur, P.K. (1979) Malformations and the Manx syndrome in cats. *Canadian Veterinary Journal*, **20** (11), 304–314.
- Degen, M. (1987) Pseudohyperkalemia in Akitas. *Journal of the American Veterinary Medical Association*, **190**, 541–543.
- DeGroot, W., Giuffrida, M.A., Rubin, J. *et al.* (2016) Primary splenic torsion in dogs: 102 cases (1992–2014). *Journal of the American Veterinary Medical Association*, **248** (6), 661–668.
- Dekomien, G., Runte, M., Gödde, R. & Epplen, J.T. (2000) Generalized progressive retinal atrophy of Sloughi dogs is due to an 8-bp insertion in exon 21 of the PDE6B gene. *Cytogenetic and Genome Research*, **90** (3–4), 261–267.

- Dekomien, G., Vollrath, C., Petrasch-Parwez, E. *et al.* (2010) Progressive retinal atrophy in Schapendoes dogs: mutation of the newly identified CCDC66 gene. *neurogenetics*, **11** (2), 163–174.
- de Lahunta, A. & Averill, D.R. (1976) Hereditary cerebellar cortical and extrapyramidal nuclear abiotrophy in Kerry Blue Terriers. *Journal of the American Veterinary Medical Association*, **168** (12), 1119–1124.
- de Lahunta, A., Fenner, W.R., Indrieri, R.J. *et al.* (1980) Hereditary cerebellar cortical abiotrophy in the Gordon Setter. *Journal of the American Veterinary Medical Association* **177** (6) 538–541.
- de Lahunta, A., Ingram, J.T., Cummings, J.F. *et al.* (1994) Labrador Retriever central axonopathy. *Progress in Veterinary Neurology*, **5**, 117–122.
- Delamaide Gasper, J.A., Rylander, H., Stenglein, J.L. *et al.* (2014) Osseous-associated cervical spondylomyelopathy in dogs: 27 cases (2000–2012). *Journal of the American Veterinary Medical Association*, **244** (11), 1309–1318.
- Delverdier, M., Guire, F. & Haverbeke, G. van (1991) Tumours of the buccal cavity in dogs: study of 117 cases. *Revue de Médecine Vétérinaire*, **142** 811–816.
- Denerolle, P., Bourdoiseau, G., Magnol, J.-P. *et al.* (1998) German Shepherd Dog pyoderma: a prospective study of 23 cases. *Veterinary Dermatology*, **9**, 243–248.
- Dennis, M.M., Ehrhart, N., Duncan, C.G., Barnes, A.B. & Ehrhart, E.J. (2006) Frequency of and risk factors associated with lingual lesions in dogs: 1,196 cases (1995–2004). *Journal of the American Veterinary Medical Association*, **228** (10), 1533–1537.
- Denny, H.R., Gibbs, C. & Waterman, A. (1988) Atlanto-axial subluxation in the dog: a review of thirty cases and an evaluation of treatment by lag screw fixation. *Journal of Small Animal Practice*, **29** (1), 37–47.
- Dervisis, N.G., Kiupel, M., Qin, Q. & Cesario, L. (2016) Clinical prognostic factors in canine histiocytic sarcoma. *Veterinary and Comparative Oncology*, June 23. doi: 10.1111/vco.12252 [Epub ahead of print].
- Diaz, J.V., Duque, C. & Geisel, R. (2007) Neuroaxonal dystrophy in dogs: case report in 2 litters of Papillon puppies. *Journal of Veterinary Internal Medicine*, **21** (3), 531–534.
- DiBartola, S.P., Hill, R.L., Fechheimer, N.S. *et al.* (1986a) Pedigree analysis of Abyssinian cats with familial amyloidosis. *American Journal of Veterinary Research*, **47** (12), 2666–2668.
- DiBartola, S.P., Tarr, M.J. & Benson, M.D. (1986b) Tissue distribution of amyloid deposits in Abyssinian cats with familial amyloidosis. *Journal of Comparative Pathology*, **96** (4), 387–398.
- DiBartola, S.P., Tarr, M.J., Parker, A.T. *et al.* (1989) Clinicopathologic findings in dogs with renal amyloidosis: 59 cases (1976–1986). *Journal of the American Veterinary Medical Association*, **195** (3), 358–364.
- DiBartola, S.P., Tarr, M.J., Webb, D.M. & Giger, U. (1990) Familial renal amyloidosis in Chinese Shar Pei dogs. *Journal of the American Veterinary Medical Association*, **197** (4), 483–487.
- Di Bella, A., Maurella, C., Cauvin, A. *et al.* (2013) Proteinuria in canine patients with lymphoma. *Journal of Small Animal Practice*, **54** (1), 28–32.
- Dickie, A.M., Schwarz, T. & Sullivan, M. (2002) Temporomandibular joint morphology in Cavalier King Charles Spaniels. *Veterinary Radiology and Ultrasound*, **43**, 260–266.
- Diesel, G., Pfeiffer, D., Crispin, S. & Brodbelt, D. (2010) Risk factors for tail injuries in dogs in Great Britain. *Veterinary Record*, **166**, 812–817.
- Dillberger, J.E. & McAtee, S.A. (2017) Osteosarcoma inheritance in two families of Scottish Deerhounds. *Canine Genetics and Epidemiology*, **4** (1), 3.
- Dinev, I. (2002) Incidence of canine neoplasms: a retrospective histopathological study. II. Tumours of the skin and associated structures. *Bulgarian Journal of Veterinary Medicine*, **5** (4), 269–278.
- Diters, R.W. & Walsh, K.M. (1984) Feline basal cell tumors: a review of 124 cases. *Veterinary Pathology*, **21** (1), 51–56.
- Docampo, M.J., Zanna, G., Fondevila, D. *et al.* (2011) Increased HAS2-driven hyaluronic acid synthesis in Shar-pei dogs with hereditary cutaneous hyaluronosis (mucinosis). *Veterinary Dermatology*, **22** (6), 535–545.
- Dodgson, S.E., Day, R. & Fyfe, J.C. (2012) Congenital hypothyroidism with goiter in Tenterfield Terriers. *Journal of Veterinary Internal Medicine*, **26** (6), 1350–1357.
- Dodman, N. H., Knowles, K. E., Shuster, L. *et al.* (1996) Behavioral changes associated with suspected complex partial seizures in Bull Terriers. *Journal of the American Veterinary Medical Association*, **208**, 688–691.

- Doerr, K.A., Outerbridge, C.A., White, S.D. *et al.* (2013) Calcinosis cutis in dogs: histopathological and clinical analysis of 46 cases. *Veterinary Dermatology*, **24** (3), 355–361.
- Dohoo, I., Martin, W. & Stryhn, H. (2009). *Veterinary Epidemiologic Research*. VER Inc., Charlottetown, Canada.
- Dole, R.S. & Spurgeon, T.L. (1998) Frequency of supernumerary teeth in a dolichocephalic canine breed, the Greyhound. *American Journal of Veterinary Research*, **59**, 16–17.
- Donaldson, D., Sansom, J., Scase, T. *et al.* (2006) Canine limbal melanoma: 30 cases (1992–2004) Part 1. Signalment, clinical and histological features and pedigree analysis. *Veterinary Ophthalmology*, **9** (2), 115–119.
- Donzel, E., Arti, L. & Chahory, S. (2016) Epidemiology and clinical presentation of canine cataracts in France: a retrospective study of 404 cases. *Veterinary Ophthalmology*, **20** (2), 131–139.
- Dorn, C.R. & Priester, W.A. (1976) Epidemiologic analysis of oral and pharyngeal cancer in dogs, cats, horses and cattle. *Journal of the American Veterinary Medical Association*, **169** (11), 1202–1206.
- Dostal, J., Hrdlicova, A. & Horak, P. (2011) Progressive rod-cone degeneration (PRCD) in selected dog breeds and variability in its phenotypic expression. *Veterinární Medicína*, **56** (5), 243–247.
- Dougherty, S.A. & Center, S.A. (1990) Heritable polyarthritis in adolescent akitas. *Journal of Veterinary Internal Medicine*, **4** (2), 112.
- Dowling, P. (2006) Pharmacogenetics: It's not just about ivermectin in collies. *Canadian Veterinary Journal*, **47**, 1165–1168.
- Downs, L.M., Bell, J.S., Freeman, C. *et al.* (2013) Late-onset progressive retinal atrophy in Gordon and Irish Setters is associated with a frameshift mutation in *C2orf71*. *Animal Genetics*, **44** (2), 169–177.
- Downs, L.M., Hitti, R., Pregnotato, S. & Mellersh, C. (2014a) Genetic screening for PRA-associated mutations in multiple dog breeds shows that PRA is heterogeneous within and between breeds. *Veterinary Ophthalmology*, **17** (2), 126–130.
- Downs, L.M., Wallin-Håkansson, B., Bergström, T. & Mellersh, C.S. (2014b) A novel mutation in *TTC8* is associated with progressive retinal atrophy in the Golden Retriever. *Canine Genetics and Epidemiology*, **1**, 4.
- Dreyfus, J., Schobert, C.S. & Dubielzig, R.R. (2011) Superficial corneal squamous cell carcinoma occurring in dogs with chronic keratitis. *Veterinary Ophthalmology*, **14** (3), 161–168.
- Driehuys, S., Van Winkle, T.J., Sammarco, C.D. & Drobatz, K.J. (1998) Myocardial infarction in dogs and cats: 37 cases (1985–1994). *Journal of the American Veterinary Medical Association*, **213** (10), 1444–1448.
- Driver, C.J., Chandler, K., Walmsley, G., Shihab, N. & Volk, H.A. (2013) The association between Chiari-like malformation, ventriculomegaly and seizures in Cavalier King Charles Spaniels. *Veterinary Journal*, **195**, 235–237.
- Drobatz, K.J. & Casey, K.K. (2000) Eclampsia in dogs: 31 cases (1995–1998). *Journal of the American Veterinary Medical Association*, **217** (2), 216–219.
- Drögemüller, C., Becker, D., Kessler, B. *et al.* (2008a) A deletion in the N-Myc downstream regulated gene 1 (*NDRG1*) gene in Greyhounds with polyneuropathy. *PLoS One*, **5** (6), e11258.
- Drögemüller, C., Karlsson, E.K., Hytönen, M.K. *et al.* (2008b) A mutation in hairless dogs implicates *FOXI3* in ectodermal development. *Science*, **321** (5895), 1462.
- Drögemüller, M., Jagannatharaj, V., Becker, D. *et al.* (2014) A mutation in the FAM83G gene in dogs with hereditary footpad hyperkeratosis (HFH). *PLoS One*, **10** (5), e1004370.
- Drost, W.T., Guillermo Couto, C., Fischetti, A.J. *et al.* (2006) Comparison of glomerular filtration rate between Greyhounds and non-Greyhound dogs. *Journal of Veterinary Internal Medicine*, **20** (3), 544–546.
- Duclos, D. (2013) Canine pododermatitis. *Veterinary Clinics of North America: Small Animal Practice*, **43** (1), 57–87.
- Duffy, D.L., Hsu, Y. & Serpell, J.A. (2008) Breed differences in canine aggression. *Applied Animal Behaviour Science*, **114** (3–4), 441–460.
- Dukes-McEwan, J., Borgarelli, M., Tidholm, A., Vollmar, A.C. & Häggström, J. (2003) Proposed guidelines for the diagnosis of canine idiopathic dilated cardiomyopathy. *Journal of Veterinary Cardiology*, **5** (2), 7–19.

- Duncan, I.D. & Griffiths, I.R. (1981) Canine giant axonal neuropathy; some aspects of its clinical, pathological and comparative features. *Journal of Small Animal Practice*, **22** (8), 491–501.
- Duval, J.M., Budberg, S.C., Flo, G.L. *et al.* (1999) Breed, sex, and body weight as risk factors for rupture of the cranial cruciate ligament in young dogs. *Journal of the American Veterinary Medical Association*, **215**, 811–814.
- Dyson, D., Maxie, M. & Schnurr, D. (1998) Morbidity and mortality associated with anesthetic management in small animal veterinary practice in Ontario. *Journal of the American Animal Hospital Association*, **34** (4), 325–335.
- Eagleson, J.S., Diaz, J., Platt, S.R. *et al.* (2009) Cervical vertebral malformation-malarticulation syndrome in the Bernese Mountain Dog: clinical and magnetic resonance imaging features. *Journal of Small Animal Practice*, **50** (4), 186–193.
- Eaton, K.A., Biller, D.S., DiBartola, S.P. *et al.* (1997) Autosomal dominant polycystic kidney disease in Persian and Persian-cross cats. *Veterinary Pathology*, **34** (2), 117–126.
- Eberle, N., Babo, V., von, Nolte, I., Baumgärtner, W. & Betz, D. (2012) Splenic masses in dogs. Part 1: Epidemiologic, clinical characteristics as well as histopathologic diagnosis in 249 cases (2000–2011). *Tierärztliche Praxis, Ausgabe K, Kleintiere/Heimtiere*, **40** (4), 250–260.
- Eckardt, J., Kluth, S., Dierks, C., Philipp, U. & Distl, O. (2013) Population screening for the mutation associated with osteogenesis imperfecta in Dachshunds. *Veterinary Record*, **172**, 364–364.
- Edelmann, M.L., Miyadera, K., Iwabe, S. & Komáromy, A.M. (2013) Investigating the inheritance of prolapsed nictitating membrane glands in a large canine pedigree. *Veterinary Ophthalmology*, **16** (6), 416–422.
- Edney, A. & Smith, P. (1986) Study of obesity in dogs visiting veterinary practices in the United Kingdom. *Veterinary Record*, **118** (14), 391–396.
- Edwards, D.S., Henley, W.E., Harding, E.F. *et al.* (2003) Breed incidence of lymphoma in a UK population of insured dogs. *Veterinary and Comparative Oncology*, **1** (4), 200–206.
- Egenvall, A., Bonnett, B.N. & Haggstrom, J. (2006) Heart disease as a cause of death in insured Swedish dogs younger than 10 years of age. *Journal of Veterinary Internal Medicine*, **20** (4), 894–903.
- Egenvall, A., Bonnett, B.N., Öhagen, P. *et al.* (2005) Incidence of and survival after mammary tumors in a population of over 80,000 insured female dogs in Sweden from 1995 to 2002. *Preventive Veterinary Medicine*, **69** (1–2), 109–127.
- Egenvall, A., Bonnett, B.N., Olson, P. & Hedhammar, Å. (1998) Validation of computerized Swedish dog and cat insurance data against veterinary practice records. *Preventive Veterinary Medicine*, **36** (1), 51–65.
- Egenvall, A., Hagman, R., Bonnett, B.N. *et al.* (2001) Breed risk of pyometra in insured dogs in Sweden. *Journal of Veterinary Internal Medicine*, **15** (6), 530–538.
- Egenvall, A., Nødtvedt, A. & von Euler, H. (2007) Bone tumors in a population of 400 000 insured Swedish dogs up to 10 y of age: incidence and survival. *Canadian Journal of Veterinary Research*, **71** (4), 292–299.
- Ekenstedt, K.J. & Oberbauer, A.M. (2013) Inherited epilepsy in dogs. *Topics in Companion Animal Medicine*, **28** (2), 51–58.
- Ekenstedt, K.J., Becker, D., Minor, K.M. *et al.* (2014) An ARHGEF10 deletion is highly associated with a juvenile-onset inherited polyneuropathy in Leonberger and Saint Bernard dogs. *PLoS Genetics*, **10** (10), e1004635.
- Ekenstedt, K.J., Patterson, E.E., Minor, K.M. *et al.* (2011) Candidate genes for idiopathic epilepsy in four dog breeds. *BMC Genetics*, **12**, 38.
- Ekensten, B. & Torrång, I. (1995) Heritability of the depth of the opening of the ciliary cleft in Samoyeds. *American Journal of Veterinary Research*, **56** (9), 1138–1143.
- Ekstrand, C. & Linde-Forsburg, C. (1994) Dystocia in the cat: a retrospective study of 155 cases. *Journal of Small Animal Practice*, **35** (9), 459–464.
- Ellinwood, N.M., Wang, P., Skeen, T. *et al.* (2003) A model of mucopolysaccharidosis IIIB (Sanfilippo syndrome type IIIB): N-acetyl- α -D-glucosaminidase deficiency in Schipperke dogs. *Journal of Inherited Metabolic Disease*, **26** (5), 489–504.
- Ellison, G. & Halling, K.B. (2004) Atypical pectus excavatum in two Welsh Terrier littermates. *Journal of Small Animal Practice*, **45** (6), 311–314.

- Elvers, I., Turner-Maier, J., Swofford, R. *et al.* (2015) Exome sequencing of lymphomas from three dog breeds reveals somatic mutation patterns reflecting genetic background. *Genome Research*, **25** (11), 1634–1645.
- Emmerson, T. (2014) Brachycephalic obstructive airway syndrome: a growing problem. *Journal of Small Animal Practice*, **55** (11), 543–544.
- Eneroth, A., Linde-Forsberg, C., Uhlhorn, M. *et al.* (1999) Radiographic pelvimetry for assessment of dystocia in bitches: a clinical study in two terrier breeds. *Journal of Small Animal Practice*, **40** (6), 257–264.
- Engvall, E. & Bushnell, N. (1990) Patellar luxation in Abyssinian cats. *Feline Practice*, **18** (4), 20–22.
- Eom, K., Moon, K., Seong, Y. *et al.* (2008) Ultrasonographic evaluation of tracheal collapse in dogs. *Journal of Veterinary Science*, **9** (4), 401–405.
- Er, J. & Sutton, R. (1989) A survey of skin neoplasms in dogs from the Brisbane region. *Australian Veterinary Journal*, **66** (7), 225–227.
- Ergul Ekiz, E., Arslan, M., Ozcan, M. *et al.* (2011) Frequency of dog erythrocyte antigen 1.1 in 4 breeds native to different areas in Turkey. *Veterinary Clinical Pathology*, **40** (4), 518–523.
- Eriksen, K. & Grøndalen, J. (1984) Familial renal disease in Soft-coated Wheaten Terriers. *Journal of Small Animal Practice*, **25** (8), 489–500.
- Ernst, T., Kessler, M., Lautscham, E., Willimzig, L. & Neiger, R. (2016) Multicentric lymphoma in 411 dogs: an epidemiological study. *Tierärztliche Praxis Ausgabe K: Kleintiere / Heimtiere*, **44**, 245–251.
- Evans, J. M., Noorai, R. E., Tsai, K. L. *et al.* (2017) Beyond the MHC: a canine model of dermatomyositis shows a complex pattern of genetic risk involving novel loci, *PLoS Genetics*, **13** (2), e1006604.
- Evans, J., Katz, M. L., Levesque, D. *et al.* (2005) A variant form of neuronal ceroid lipofuscinosis in American Bulldogs. *Journal of Veterinary Internal Medicine*, **19**, 44–51.
- Evans, J., Levesque, D. & Shelton, G.D. (2004) Canine inflammatory myopathies: a clinicopathologic review of 200 cases *Journal of Veterinary Internal Medicine*, **18** (5), 679–691.
- Evans, K.M. & Adams, V.J. (2010a) Proportion of litters of purebred dogs born by caesarean section. *Journal of Small Animal Practice*, **51** (2), 113–118.
- Evans, K.M. & Adams, V.J. (2010b) Mortality and morbidity due to gastric dilatation-volvulus syndrome in pedigree dogs in the UK. *Journal of Small Animal Practice*, **51** (7), 376–381.
- Evans, S.M. & Biery, D.N. (1980) Congenital peritoneopericardial diaphragmatic hernia in the dog and cat: a literature review and 17 additional case histories. *Veterinary Radiology*, **21** (3), 108–116.
- Fabrizio, F., Baumwart, R., Iazbik, M.C. *et al.* (2006) Left basilar systolic murmur in retired Racing Greyhounds. *Journal of Veterinary Internal Medicine*, **20**, 78–82.
- Fabrizio, F., Calam, A.E., Dobson, J.M. *et al.* (2014) Feline mediastinal lymphoma: a retrospective study of signalment, retroviral status, response to chemotherapy and prognostic indicators. *Journal of Feline Medicine and Surgery*, **16** (8), 637–644.
- Fall, T., Hamlin, H.H., Hedhammar, Å., Kämpe, O. & Engvall, A. (2007) Diabetes mellitus in a population of 180,000 insured dogs: incidence, survival, and breed distribution. *Journal of Veterinary Internal Medicine*, **21** (6), 1209–1216.
- Fall, T., Hedhammar, Å., Wallberg, A., Fall, N. *et al.* (2010) Diabetes mellitus in Elkhounds is associated with diestrus and pregnancy. *Journal of Veterinary Internal Medicine*, **24** (6), 1322–1328.
- Famula, T., Belanger, J. & Oberbauer, A. (2003) Heritability and complex segregation analysis of hypoadrenocorticism in the Standard Poodle. *Journal of Small Animal Practice*, **44** (1), 8–12.
- Famula, T.R., Siemens, L.M., Davidson, A.P. *et al.* (2002) Evaluation of the genetic basis of tricuspid valve dysplasia in Labrador Retrievers. *American Journal of Veterinary Research*, **63** (6), 816–820.
- Farias, F.H.G., Zeng, R., Johnson, G.S. *et al.* (2011) A truncating mutation in ATP13A2 is responsible for adult-onset neuronal ceroid lipofuscinosis in Tibetan Terriers. *Neurobiology of Disease*, **42** (3), 468–474.
- Farrow, B.R.H. & Malik, R. (1981) Hereditary myotonia in the Chow Chow. *Journal of Small Animal Practice*, **22**, 451–465.
- Fasanella, F.J., Shively, J.M., Wardlaw, J.L. & Givaruangsawat, S. (2010) Brachycephalic airway obstructive syndrome in dogs: 90 cases (1991–2008). *Journal of the American Veterinary Medical Association*, **237**, 1048–1051.

- Favrot, C., Steffan, J., Seewald, W. & Picco, F. (2010) A prospective study on the clinical features of chronic canine atopic dermatitis and its diagnosis. *Veterinary Dermatology*, **21** (1), 23–31.
- Featherstone, H.J., Renwick, P., Heinrich, C. L. *et al.* (2009) Efficacy of lamellar resection, cryotherapy, and adjunctive grafting for the treatment of canine limbal melanoma. *Veterinary Ophthalmology*, **12** (suppl. 1), 65–72.
- Featherstone, H.J. & Sansom, J. (2004) Feline corneal sequestra: a review of 64 cases (80 eyes) from 1993 to 2000. *Veterinary Ophthalmology*, **7** (4), 213–227.
- Feliu-Pascual, A.L., Shelton, G.D., Targett, M.P. *et al.* (2006) Inherited myopathy of Great Danes. *Journal of Small Animal Practice*, **47**, 249–254.
- Fels, L. & Distl, O. (2014) Identification and validation of quantitative trait loci (QTL) for canine hip dysplasia (CHD) in German Shepherd Dogs. *PLoS One*, **9** (5), e96618.
- Fenn, J., Drees, R., Volk, H.A. & De Decker, S. (2016) Comparison of clinical signs and outcomes between dogs with presumptive ischemic myelopathy and dogs with acute noncompressive nucleus pulposus extrusion. *Journal of the American Veterinary Medical Association*, **249**, 767–775.
- Ferguson, E.A., Cerundolo, R., Lloyd, D.H., Cappello, R. & Rest, J. (2000) Dermatomyositis in five Shetland Sheepdogs in the United Kingdom. *Veterinary Record*, **146** (8), 214–217.
- Ferm, K., Björnerfeldt, S., Karlsson, Å. *et al.* (2009) Prevalence of diagnostic characteristics indicating canine autoimmune lymphocytic thyroiditis in Giant Schnauzer and Hovawart dogs. *Journal of Small Animal Practice*, **50** (4), 176–179.
- Ferreira, R.R., Gopegui, R.R. & Matos, A.J.F. (2011) Frequency of dog erythrocyte antigen 1.1 expression in dogs from Portugal. *Veterinary Clinical Pathology*, **40** (2), 198–201.
- Fialová, I., Paninárová, M., Nečas, A., Stehlík, L. & Proks, P. (2014) Prevalence of lumbosacral transitional vertebrae in dogs in the Czech Republic. *Acta Veterinaria Brno*, **83** (4), 399–403.
- Fiani, N., Verstraete, F.J.M., Kass, P.H. *et al.* (2011) Clinicopathologic characterization of odontogenic tumors and focal fibrous hyperplasia in dogs: 152 cases (1995–2005). *Journal of the American Veterinary Medical Association*, **238** (4), 495–500.
- Fietan, H., Biourge, V.C., Watson, A.L. *et al.* (2014) Nutritional management of inherited copper-associated hepatitis in the Labrador Retriever. *Veterinary Journal*, **199** (3), 429–433.
- Fietan, H., Biourge, V.C., Watson, A.L. *et al.* (2015) Dietary management of Labrador Retrievers with subclinical hepatic copper accumulation. *Journal of Veterinary Internal Medicine*, **29** (3), 822–827.
- Fife, M., Blocker, T., Fife, T. *et al.* (2011) Canine conjunctival mast cell tumors: a retrospective study. *Veterinary Ophthalmology*, **14** (3), 153–160.
- Finco, D.R., Duncan, J.D., Crowell, W.A. *et al.* (1977) Familial renal disease in Norwegian Elkhound dogs: morphologic examinations. *American Journal of Veterinary Research*, **38** (7), 941–947.
- Finster, S.T., DeFrancesco, T.C., Atkins, C.E. *et al.* (2008) Supraventricular tachycardia in dogs: 65 cases (1990–2007) *Journal of Veterinary Emergency Critical Care*, **18** (5), 503–510.
- Firdova, Z., Turnova, E., Bielikova, M., Turna, J. & Dudas, A. (2016) The prevalence of ABCB1:c.227_230delATAG mutation in affected dog breeds from European countries. *Research in Veterinary Science*, **106**, 89–92.
- Fitzpatrick, N., Smith, T.J., Evans, R.B. & Yeadon, R. (2009) Radiographic and arthroscopic findings in the elbow joints of 263 dogs with medial coronoid disease. *Veterinary Surgery*, **38** (2), 213–223.
- Flagstad, A., Trojaborg, W. & Gammeltoft, S. (1989) Congenital myasthenic syndrome in the dog breed Gammel Dansk Hønsehund. An animal model for research in neuromuscular disorders. *Scandinavian Journal of Laboratory Animal Science*, **16**, 89–102.
- Fleming, J.M., Creevy, K.E. & Promislow, D.E.L. (2011) Mortality in North American dogs from 1984 to 2004: an investigation into age-, size-, and breed-related causes of death. *Journal of Veterinary Internal Medicine*, **25** (2), 187–198.
- Fletcher, S.M. & Pinkerton, P.H. (1972) An inherited anaemia associated with hereditary chondrodysplasia in the Alaskan Malamute. *Canadian Veterinary Journal*, **13**, 270–271.
- Fletcher, J.L., Williamson, P., Horan, D. & Taylor, R.M. (2010) Clinical signs and neuropathologic abnormalities in working Australian Kelpies with globoid cell leukodystrophy (Krabbe disease). *Journal of the American Veterinary Medical Association*, **237**, 682–688.

- Fluehmann, G., Doherr, M.G. & Jaggy, A. (2006) Canine neurological diseases in a referral hospital population between 1989 and 2000 in Switzerland. *Journal of Small Animal Practice*, **47**, 582–587.
- Foale, R.D., Herrtage, M.E. & Day, M.J. (2003) Retrospective study of 25 young Weimaraners with low serum immunoglobulin concentrations and inflammatory disease. *Veterinary Record*, **153** (18), 553–558.
- Foley, J.E. & Pedersen, N.C. (1996) The inheritance of susceptibility to feline infectious peritonitis in purebred catteries. *Feline Practice*, **24** (1), 14–22.
- Fonda, D., Gualtieri, M. & Scanziani, E. (1989) Gastric carcinoma in the dog: a clinicopathological study of 11 cases. *Journal of Small Animal Practice*, **30** (6), 353–360.
- Fonfara, S., de la Heras Alegret, L., German, A.J. *et al.* (2011) Underlying diseases in dogs referred to a veterinary teaching hospital because of dyspnea: 229 cases (2003–2007). *Journal of the American Veterinary Medical Association*, **239** (9), 1219–1224.
- Forcada, Y., Guitian, J. & Gibson, G. (2007) Frequencies of feline blood types at a referral hospital in the south east of England. *Journal of Small Animal Practice*, **48**, 570–573.
- Ford, R.B. (1993) Idiopathic hyperchylomicronaemia in Miniature Schnauzers. *Journal of Small Animal Practice*, **34** (10), 488–492.
- Forman, O.P., De Risio, L., Matiassek, K., Platt, S. & Mellersh, C. (2015) Spinocerebellar ataxia in the Italian Spinone dog is associated with an intronic GAA repeat expansion in ITPR1. *Mammalian Genome*, **26** (1), 108–117.
- Forman, O.P., De Risio, L. & Mellersh, C.S. (2013) Missense mutation in CAPN1 is associated with spinocerebellar ataxia in the Parson Russell Terrier dog breed. *PLoS One*, **8** (5), e64627.
- Forrester, S.D., Boothe, D. M. & Troy, G.C. (1989) Current concepts in the management of canine epilepsy. *Compendium on Continuing Education for the Practicing Veterinarian*, **11** (7), 811–820.
- Forsberg, M. (1986) Breed differences in the predisposition for diabetes mellitus and Cushing's syndrome in the dog. *Svensk Veterinärtidning*, **38**, 177–180.
- Forsee, K.M., Davis, G.J., Mouat, E.E., Salmeri, K.R. & Bastian, R.P. (2013) Evaluation of the prevalence of urinary incontinence in spayed female dogs: 566 cases (2003–2008). *Journal of the American Veterinary Medical Association*, **242** (7), 959–962.
- Fossum, T.W., Birchard, S.J. & Jacobs, R.M. (1986) Chylothorax in 34 dogs. *Journal of the American Veterinary Medical Association*, **188**, 1315–1318.
- Foster, J.D., Sample, S., Köhler, R. *et al.* (2014) Serum biomarkers of clinical and cytologic response in dogs with idiopathic immune-mediated polyarthropathy. *Journal of Veterinary Internal Medicine*, **28** (3), 905–911.
- Foster, S.J., Curtis, R. & Barnett, K.C. (1986) Primary lens luxation in the Border Collie. *Journal of Small Animal Practice*, **27** (1), 1–6.
- Foureman, P., Whiteley, M. & Giger, U. (2002) Canine leukocyte adhesion deficiency: presence of the Cys36Ser β -2 integrin mutation in an affected US Irish Setter cross-breed dog and in US Irish Red and White Setters. *Journal of Veterinary Internal Medicine*, **16** (5), 518–523.
- Fournel, C., Chabanne, L., Caux, C. *et al.* (1992) Canine systemic lupus erythematosus. I. A study of 75 cases. *Lupus*, **1**, 133–139.
- Fracassi, F., Pietra, M., Boari, A., Aste, G., Giunti, M. & Famigli-Bergamini, P. (2004) Breed distribution of canine diabetes mellitus in Italy. *Veterinary Research Communications*, **28** (1), 339–342.
- Fracassi, F., Zagnoli, L., Rosenberg, D. *et al.* (2014) Spontaneous acromegaly: a retrospective case control study in German shepherd dogs. *Veterinary Journal*, **202** (1), 69–95.
- Frank, L.A., Hnilica, K.A. & Oliver, J.W. (2004) Adrenal steroid hormone concentrations in dogs with hair cycle arrest (Alopecia X) before and during treatment with melatonin and mitotane. *Veterinary Dermatology*, **15** (5), 278–284.
- Frank, L.A., Hnilica, K.A., Rohrbach, B.W. & Oliver, J.W. (2003) Retrospective evaluation of sex hormones and steroid hormone intermediates in dogs with alopecia. *Veterinary Dermatology*, **14** (2), 91–97.
- Franklin, R.J., Jeffery, N.D. & Ramsey, I.K. (1995) Neuroaxonal dystrophy in a litter of Papillon pups. *Journal of Small Animal Practice*, **36** (10), 441–444.
- Franklin, R.J., Ramsey, I.K. & McKerrell, R.E. (1997) An inherited neurological disorder of the St Bernard dog characterised by unusual cerebellar cortical dysplasia. *Veterinary Record*, **140** (25), 656–657.

- Frazer, M. M., Schick, A. E., Lewis, T. P. & Jazic, E. (2011) Sebaceous adenitis in Havanese dogs: a retrospective study of the clinical presentation and incidence. *Veterinary Dermatology*, **22** (3), 267–274.
- Freeman, A.C., Platt, S.R., Kent, M. *et al.* (2014) Chiari-like malformation and syringomyelia in American Brussels Griffon dogs. *Journal of Veterinary Internal Medicine*, **28** (5), 1551–1559.
- Freeman, W.E. (2005) Idiosyncrasies in Greyhounds that can affect their medical care. *Veterinary Medicine*, **100** (8), 592–600.
- French, A.T., Ogden, R., Eland, C. *et al.* (2012) Genome-wide analysis of mitral valve disease in Cavalier King Charles Spaniels. *Veterinary Journal*, **193**, 283–286.
- Frgelecová, L., Škorič, M., Fictum, P. *et al.* (2013) Canine gastrointestinal tract tumours: a retrospective study of 74 cases. *Acta Veterinaria Brno*, **82** (4), 387–392.
- Fricker, G.V., Smith, K., & Gould, D.J. (2016) Survey of the incidence of pectinate ligament dysplasia and glaucoma in the UK Leonberger population. *Veterinary Ophthalmology*, **19** (5), 379–385.
- Fritsch, R. & Ost, P. (1983) Hereditary tail anomalies in the Dachshund. *Berliner und Münchener Tierärztliche Wochenschrift*, **96**, 444–450.
- Fritsche, R., Dolf, G., Schelling, C. *et al.* (2014) Inheritance of ectopic ureters in Entlebucher Mountain Dogs. *Journal of Animal Breeding and Genetics*, **131** (2), 146–152.
- Fritz, T.E., Zeman, R.C. and Zelle, M.R. (1970) Pathology and familial incidence of thyroiditis in a closed Beagle colony. *Experimental and Molecular Pathology*, **12**, 14–30.
- Fujise, H., Higa, K. & Nakayama, T. (1997) Incidence of dogs possessing red blood cells with high K in Japan and East Asia. *Journal of Veterinary Medical Science*, **59** (6), 495–497.
- Fujitu, K. & Sakai, T. (1999) Factors affecting the occurrence of feline gingivostomatitis. *Journal of the Japan Veterinary Medical Association*, **52** (3), 175–179.
- Fulmer, A.K. & Mauldin, G.E. (2007) Canine histiocytic neoplasia: an overview. *Canadian Veterinary Journal*, **48** (10), 1041–1043.
- Furrow, E., Minor, K.M., Taylor S.M. *et al.* (2013) Relationship between dynamin 1 mutation status and characteristics of recurrent episodes of exercise-induced collapse in Labrador Retrievers. *Journal of the American Veterinary Medical Association*, **242** (6), 786–791.
- Fyfe, J.C., Giger, U., Hall, C.A. *et al.* (1991) Inherited selective intestinal cobalamin malabsorption and cobalamin deficiency in dogs. *Pediatric Research*, **29** (1), 24–31.
- Fyfe, J.C., Kampschmidt, K., Dang, V. *et al.* (2003) Congenital hypothyroidism with goitre in Toy Fox Terriers. *Journal of Veterinary Internal Medicine*, **17** (1), 50–57.
- Fyfe, J.C., Kurzhals, R.L., Hawkins, M.G. *et al.* (2007) A complex rearrangement in *GBE1* causes both perinatal hypoglycemic collapse and late-juvenile-onset neuromuscular degeneration in glycogen storage disease type IV of Norwegian Forest Cats. *Molecular Genetics and Metabolism*, **90** (4), 383–392.
- Gabriel, A., Poncelet, L., van Ham, L. *et al.* (2006) Laryngeal paralysis-polyneuropathy complex in young related Pyrenean Mountain Dogs. *Journal of Small Animal Practice*, **47** (3), 144–149.
- Galis, F., Van Der Sluijs, I., Van Dooren, T.J.M., Metz, J.A.J. & Nussbaumer, M. (2007) Do large dogs die young? *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, **308B** (2), 119–126.
- Gandolfi, B., Grahn, R.A., Creighton, E.K. *et al.* (2015) COLQ variant associated with Devon Rex and Sphynx feline hereditary myopathy. *Animal Genetics*, **46** (6), 711–715.
- Gandolfi, B., Liu, H., Griffioen, L., & Pedersen, N.C. (2013) Simple recessive mutation in ENAM is associated with amelogenesis imperfecta in Italian Greyhounds. *Animal Genetics*, **44** (5), 568–578.
- Garcia-Sancho, M., Villaescusa, A., Sainz, A. *et al.* (2013) Importance of double endoscopic exploration (respiratory and digestive) in French Bulldogs. *Archivos de Medicina Veterinaria*, **45** (3), 283–289.
- Garden, O.A., Pidduck, H., Lakhani, K.H. *et al.* (2000) Inheritance of gluten-sensitive enteropathy in Irish Setters. *American Journal of Veterinary Research*, **61** (4), 462–468.
- Garosi, L.S., Rossmeisl, J.H., de Lahunta, A. *et al.* (2005) Primary orthostatic tremor in Great Danes. *Journal of Veterinary Internal Medicine*, **19**, 606–609.
- Gaudet, A.D. (1985) Retrospective study of 128 cases of canine dystocia. *Journal of the American Animal Hospital Association*, **21**, 813–818.
- Gaynor, A.R., Shofer, F.S. & Washabau, R.J. (1997) Risk factors for acquired megaesophagus in dogs. *Journal of the American Veterinary Medical Association*, **211** (11), 1406–1412.

- Geisen, V., Stengel, C. & Hartmann, K. (2008) Epidemiologic situation of leptospirosis in dogs in the southern states of Germany. *Tierärztliche Praxis, Ausgabe K, Kleintiere/Heimtiere*, **36**, 329–336.
- Gelatt, K.N. & MacKay, E.O. (2004a) Prevalence of the breed-related glaucomas in pure-bred dogs in North America. *Veterinary Ophthalmology*, **7** (2), 97–111.
- Gelatt, K.N. & MacKay, E.O. (2004b) Secondary glaucomas in the dog in North America. *Veterinary Ophthalmology*, **7** (4), 245–259.
- Gelatt, K.N. & MacKay, E.O. (2005) Prevalence of primary breed-related cataracts in the dog in North America. *Veterinary Ophthalmology*, **8** (2), 101–111.
- Gelatt, K.N., Powell, N.G. & Huston, K. (1981) Inheritance of microphthalmia with coloboma in the Australian shepherd dog. *American Journal of Veterinary Research*, **42** (10), 1686–1690.
- Genevois, J.-P., Remy, D., Viguier, E. *et al.* (2008) Prevalence of hip dysplasia according to official radiographic screening, among 31 breeds of dogs in France. *Veterinary and Comparative Orthopaedics and Traumatology*, **21**, 21–24.
- Gentilini, F. & Turba, M.E. (2013) Two novel real-time PCR methods for genotyping the von Willebrand disease type I mutation in Doberman Pinscher dogs. *Veterinary Journal* **197** (2), 457–460.
- Gentilini, F., Rovesti, G.L. & Turba, M.E. (2009) Real-time detection of the mutation responsible for progressive rod-cone degeneration in Labrador Retriever dogs using locked nucleic acid TaqMan probes. *Journal of Veterinary Diagnostic Investigation* **21** (5) 689–692.
- Gerber, B., Eichenberger, S., Wittenbrink, M.M. & Reusch, C. E. (2007) Increased prevalence of *Borrelia burgdorferi* infections in Bernese Mountain Dogs: a possible breed predisposition. *BMC Veterinary Research*, **3** (1), 15.
- Gerber, K., Harvey, J.W., D'Agorne, S., Wood, J. & Giger, U. (2009) Hemolysis, myopathy, and cardiac disease associated with hereditary phosphofructokinase deficiency in two Whippets. *Veterinary Clinical Pathology*, **38** (1), 46–51.
- German, A.J., Hall, E.J., Kelly, D.F. *et al.* (2000a) An immunohistochemical study of histiocytic ulcerative colitis in Boxer dogs. *Journal of Comparative Pathology*, **122** (2–3), 163–175.
- German, A.J., Helps, C.R., Hall, E.J. *et al.* (2000b) Cytokine mRNA expression in mucosal biopsies from German Shepherd Dogs with small intestinal enteropathies. *Digestive Diseases and Sciences*, **45** (1), 7–17.
- Geyer, J., Döring, B., Godoy, J.R. *et al.* (2005) Frequency of the nt230 (del4) MDR1 mutation in Collies and related dog breeds in Germany. *Journal of Veterinary Pharmacology and Therapeutics*, **28** (6), 545–551.
- Geyer, J., Klintzsch, S., Meerkamp, K. *et al.* (2007) Detection of the nt230 (del4) MDR1 mutation in White Swiss Shepherd dogs: case reports of doramectin toxicosis, breed predisposition, and microsatellite analysis. *Journal of Veterinary Pharmacology and Therapeutics*, **30** (5), 482–485.
- Gharahkhani, P., O'Leary, C., Duffy, D., Bernays, M. & Kyaw-Tanner, M. (2012) Primary lens luxation in Australian Tenterfield and Miniature Bull Terriers is due to an old ADAMTS17 mutation and is an additive trait. *Open Genomics Journal*, **5**, 7–13.
- Gianella, P., Pfammatter, N.S. & Burgener, I.A. (2009) Oesophageal and gastric endoscopic foreign body removal: complications and follow-up of 102 dogs. *Journal of Small Animal Practice*, **50** (12), 649–654.
- Gibbons, S.E., Macias, C., Tonzing, G.L. *et al.* (2006) Patellar luxation in 70 large breed dogs. *Journal of Small Animal Practice*, **47** (1), 3–9.
- Gibson, A., Dean, R., Yates, D. *et al.* (2013) A retrospective study of pyometra at five RSPCA hospitals in the UK: 1728 cases from 2006 to 2011. *Veterinary Record*, **173** (16), 396.
- Giesel, O., Kusch, S. & Klein, H. (1986) Incidence, primary location and metastases of malignant haemangioendothelioma in dogs. Analysis of post mortem data for 1970–1985. *Tierärztliche Praxis*, **14**, 389–395.
- Giger, U., Bucheler, J. & Patterson, D.F. (1991a) Frequency and inheritance of A and B blood types in feline breeds of the United States. *Journal of Heredity*, **82**, 15–20.
- Giger, U., Griot-Wenk, M., Bucheler, J. *et al.* (1991b) Geographical variation of the feline blood type frequencies in the United States. *Feline Practice*, **19** (6), 21–27.
- Giger, U., Reilly, M.P., Asakura, T., Baldwin, C.J. & Harvey, J.W. (1986) Autosomal recessive inherited phosphofructokinase deficiency in English Springer Spaniel dogs. *Animal Genetics*, **17** (1), 15–23.

- Gilger, B.C., Hamilton, H.L., Wilkie, D.A. *et al.* (1995) Traumatic ocular proptoses in dogs and cats: 84 cases (1980–1993). *Journal of the American Veterinary Medical Association*, **206** (8), 1186–1190.
- Gill, J. L., Tsai, K. L., Krey, C. *et al.* (2012) A canine BCAN microdeletion associated with episodic falling syndrome. *Neurobiology of Disease*, **45**, 130–136.
- Gilliam, D., O'Brien, D.P., Coates, J.R. *et al.* (2014) A homozygous KCNJ10 mutation in Jack Russell Terriers and related breeds with spinocerebellar ataxia with myokymia, seizures, or both. *Journal of Veterinary Internal Medicine*, **28** (3), 871–877.
- Ginja, M.M.D., Silvestre, A.M., Colaço, J. *et al.* (2009) Hip dysplasia in Estrela Mountain Dogs: prevalence and genetic trends 1991–2005. *Veterinary Journal*, **182**, (2) 275–282.
- Ginja, M.M.D., Silvestre, A.M., Gonzalo-Orden, J.M. & Ferreira, A.J.A. (2010) Diagnosis, genetic control and preventive management of canine hip dysplasia: A review. *Veterinary Journal*, **184** (3), 269–276.
- Ginn, J.A., Kumar, M.S.A., McKiernan, B.C. & Powers, B.E. (2008) Nasopharyngeal turbinates in brachycephalic dogs and cats. *Journal of the American Animal Hospital Association*, **44** (5), 243–249.
- Giori, L., Gironi, S., Scarpa, P. *et al.* (2011) Grey eosinophils in sighthounds: frequency in 3 breeds and comparison of eosinophil counts determined manually and with 2 hematology analyzers. *Veterinary Clinical Pathology*, **40** (4), 475–483.
- Giuliano, E.A., Chappell, R., Fischer, B. *et al.* (1999) A matched observational study of canine survival with primary intraocular melanocytic neoplasia. *Veterinary Ophthalmology*, **2** (3), 185–190.
- Glickman, L.T., Domanski, L.M., Patronek, G.J. & Visintainer, F. (1985) Breed-related risk factors for canine parvovirus enteritis. *Journal of the American Veterinary Medical Association*, **187** (6), 589–594.
- Glickman, L.T., Glickman, N.W., Schellenberg, D.B., Raghavan, M. & Lee, T.L. (2000a) Incidence of and breed-related risk factors for gastric dilatation-volvulus in dogs. *Journal of the American Veterinary Medical Association*, **216** (1), 40–45.
- Glickman, L.T., Glickman, N.W., Schellenberg, D.B., Raghavan, M. & Lee, T.L. (2000b) Non-dietary risk factors for gastric dilatation-volvulus in large and giant breed dogs. *Journal of the American Veterinary Medical Association*, **217** (10), 1492–1499.
- Glickman, L.T., Glickman, N.W., Pérez, C.M. *et al.* (1994) Analysis of risk factors for gastric dilatation and dilatation-volvulus in dogs. *Journal of the American Veterinary Medical Association*, **204** (9), 1465–1471.
- Gnudi, G., Volta, A., Morgan, J.P. *et al.* (2009) Radiographic features of pelvis and hip joint development of English Bulldogs. *Veterinary and Comparative Orthopaedics and Traumatology*, **23**, 19–27.
- Gobar, G.M. (1998) Program for surveillance of causes of death of dogs, using the Internet to survey small animal veterinarians. *Journal of the American Veterinary Medical Association*, **213** (2), 251–256.
- Gödde, T. & Steffen, F. (2007) Surgical treatment of lumbosacral foraminal stenosis using a lateral approach in twenty dogs with degenerative lumbosacral stenosis. *Veterinary Surgery*, **36** (7), 705–713.
- Godiksen, M.T.N., Granström, S., Koch, J. *et al.* (2011) Hypertrophic cardiomyopathy in young Maine Coon cats caused by the p.A31P cMyBP-C mutation--the clinical significance of having the mutation. *Acta Veterinaria Scandinavica*, **53**, 7.
- Goebel, H.H., Bilzer, T., Dahme, E. *et al.* (1988) Morphological studies in canine (Dalmatian) neuronal ceroid-lipofuscinosis. *American Journal of Medical Genetics*, **5**, 127–139.
- Goldschmidt, M.H. & Mcmanus, P. (2000). *Dermatopathology. Skin Tumors of the Dog and Cat*. University of Pennsylvania School of Veterinary Medicine. <http://cal.vet.upenn.edu/projects/derm> (accessed October 2017).
- Goldstein, O., Guyon, R., Kukekova, A. *et al.* (2010) COL9A2 and COL9A3 mutations in canine autosomal recessive osseous dysplasia. *Mammalian Genome*, **21** (7), 398–408.
- Goldstein, O., Mezey, J.G., Schweitzer, P.A. *et al.* (2013) IQCB1 and PDE6B mutations cause similar early onset retinal degenerations in two closely related terrier dog breeds. *Investigative Ophthalmology and Visual Science*, **54** (10), 7005–7019.
- Goldstein, O., Zangerl, B., Pearce-Kelling, S. *et al.* (2006) Linkage disequilibrium mapping in domestic dog breeds narrows the progressive rod–cone degeneration interval and identifies ancestral disease-transmitting chromosome. *Genomics*, **88** (5), 541–550.

- Goldstein, R.E., Atwater, D.Z., Cazolli, D.M. *et al.* (2007) Inheritance, mode of inheritance, and candidate genes for primary hyperparathyroidism in Keeshonden. *Journal of Veterinary Internal Medicine*, **21** (1), 199–203.
- Gomez, S.M., Morris, D.O., Rosenbaum, M.R. *et al.* (2004). Outcome and complications associated with treatment of pemphigus foliaceus in dogs: 43 cases (1994–2000). *Journal of the American Veterinary Medical Association*, **224**, 1312–1316.
- Good, K.L., Komáromy, A.M., Kass, P.H. *et al.* (2016) Novel retinopathy in related Gordon Setters: a clinical, behavioral, electrophysiological, and genetic investigation. *Veterinary Ophthalmology*, **19** (5), 398–408.
- Goodfellow, M., Papasouliotis, K., Cue, S., Crawford, E. & Hall, E. (2008) Effect of storage on microcytosis observed in dogs with portosystemic vascular anomalies. *Research in Veterinary Science*, **84** (3), 490–493.
- Goodrich, K.R., Kyles, A.E., Kass, P.H. & Campbell, F. (2007) Retrospective comparison of surgical ligation and transarterial catheter occlusion for treatment of patent ductus arteriosus in two hundred and four dogs (1993–2003). *Veterinary Surgery*, **36** (1), 43–49.
- Gookin, J.L., Bunch, S.E., Rush, L.J. (1998) Evaluation of microcytosis in 18 Shibas. *Journal of the American Veterinary Medical Association*, **212** (8), 1258–1259.
- Goto, A., Arata, S., Kiyokawa, Y., Takeuchi, Y. & Mori, Y. (2012) Risk factors for canine tail chasing behaviour in Japan. *Veterinary Journal*, **192** (3), 445–448.
- Gough, A. & Thomas, A. (2004). *Breed Predispositions to Disease in Dogs and Cats*. Blackwell, Oxford.
- Gould, D., Pettitt, L., McLaughlin, B. *et al.* (2011) ADAMTS17 mutation associated with primary lens luxation is widespread among breeds. *Veterinary Ophthalmology*, **14** (6), 378–384.
- Gould, D.J., Petersen-Jones, S.M., Lin, C.T. & Sargan, D.R. (1997) Cloning of canine rom-1 and its investigation as a candidate gene for generalized progressive retinal atrophies in dogs. *Animal Genetics*, **28** (6), 391–396.
- Goulle, F. (2010) Sudden acquired retinal degeneration in the dog: report of seventeen cases and literature review. *Revue de Médecine Vétérinaire*, **161**, 79–83.
- Gow, A.G., Marques, A.I., Yool, D.A. *et al.* (2012) Dogs with congenital porto-systemic shunting (cPSS) and hepatic encephalopathy have higher serum concentrations of C-reactive protein than asymptomatic dogs with cPSS. *Metabolic Brain Disease*, **27** (2), 227–229.
- Graham, K.L., White, J.D. & Billson, F.M. (2016) Feline corneal sequestra: outcome of corneoconjunctival transposition in 97 cats (109 eyes). *Journal of Feline Medicine and Surgery*, **19** (6), 710–716.
- Graham, P.A., Refsal, K.R. & Nachreiner, R.F. (2007) Etiopathologic findings of canine hypothyroidism. *Veterinary Clinics of North America: Small Animal Practice*, **37** (4), 617–631.
- Grahn, B.H. & Cullen, C.L. (2001) Retinopathy of Great Pyrenees dogs: fluorescein angiography, light microscopy and transmitting and scanning electron microscopy. *Veterinary Ophthalmology*, **4** (3), 191–199.
- Grahn, B.H., Philibert, H., Cullen, C.L. *et al.* (1998) Multifocal retinopathy of Great Pyrenees dogs. *Veterinary Ophthalmology*, **1** (4), 211–221.
- Grahn, B.H., Sandmeyer, L.L. & Breaux, C. (2008) Retinopathy of Coton de Tulear dogs: clinical manifestations, electroretinographic, ultrasonographic, fluorescein and indocyanine green angiographic, and optical coherence tomographic findings. *Veterinary Ophthalmology*, **11**, 242–249.
- Grahn, B.H., Sandmeyer, L.S. & Breaux, C.B. (2006) Multifocal bullous retinopathy of Coton de Tulear dogs. *Canadian Veterinary Journal*, **47** (9), 929–930.
- Grahn, B.H., Storey, E.S. & McMillan, C. (2004) Inherited retinal dysplasia and persistent hyperplastic primary vitreous in Miniature Schnauzer dogs. *Veterinary Ophthalmology*, **7** (3), 151–158.
- Grahn, R.A., Grahn, J.C., Penedo, M.C., Helps, C.R. & Lyons, L.A. (2012) Erythrocyte pyruvate kinase deficiency mutation identified in multiple breeds of domestic cats. *BMC Veterinary Research*, **8**, 207.
- Grall, A., Guaguère, E., Planchais, S. *et al.* (2012) PNPLA1 mutations cause autosomal recessive congenital ichthyosis in golden retriever dogs and humans. *Nature Genetics*, **44** (2), 140–147.
- Gramer, I., Leidolf, R., Döring, B. *et al.* (2011) Breed distribution of the nt230 (del4) MDR1 mutation in dogs. *Veterinary Journal*, **189** (1), 67–71.

- Granger, N. (2011) Canine inherited motor and sensory neuropathies: an updated classification in 22 breeds and comparison to Charcot-Marie-Tooth disease. *Veterinary Journal*, **188** (3), 274–285.
- Granger, N., Smith, P.M. & Jeffery, N.D. (2010) Clinical findings and treatment of non-infectious meningoencephalomyelitis in dogs: a systematic review of 457 published cases from 1962 to 2008. *Veterinary Journal*, **184** (3), 290–297.
- Granström, S., Nyberg Godiksen, M.T., Christiansen, M. *et al.* (2011) Prevalence of hypertrophic cardiomyopathy in a cohort of British Shorthair cats in Denmark. *Journal of Veterinary Internal Medicine*, **25** (4), 866–871.
- Gray, J.M., Kirberger, R.M. & Spotswood, T.C. (2003) Cervical spondylomyelopathy (wobbler syndrome) in the Boerboel. *Journal of the South African Veterinary Association*, **74**, 104–110.
- Greco, D.S., Feldman, E.C., Peterson, M.E. *et al.* (1991) Congenital hypothyroid dwarfism in a family of Giant Schnauzers. *Journal of Veterinary Internal Medicine*, **5** (2), 57–65.
- Green, J.L., Cimino Brown, D. & Agnello, K.A. (2012) Preoperative thoracic radiographic findings in dogs presenting for gastric dilatation-volvulus (2000–2010): 101 cases. *Journal of Veterinary Emergency and Critical Care*, **22** (5), 595–600.
- Greenfield, C.L., Messick, J.B., Solter, P.F. *et al.* (1999) Leukopenia in 6 healthy Belgian Tervuren. *Journal of the American Veterinary Medical Association*, **215** (8), 1121–1122.
- Greenwell, C.M. & Brain, P.H. (2014) Aspiration pneumonia in the Irish Wolfhound: a possible breed predisposition. *Journal of Small Animal Practice*, **55** (10), 515–520.
- Greer, K., Schatzberg, S., Porter, B. *et al.* (2009) Heritability and transmission analysis of necrotizing meningoencephalitis in the Pug. *Research in Veterinary Science*, **86** (3), 438–442.
- Greer, K.A., Canterberry, S.C. & Murphy, K.E. (2007) Statistical analysis regarding the effects of height and weight on life span of the domestic dog. *Research in Veterinary Science*, **82** (2), 208–214.
- Greer, K.A., Wong, A.K., Liu, H. *et al.* (2010) Necrotizing meningoencephalitis of Pug Dogs associates with dog leukocyte antigen class II and resembles acute variant forms of multiple sclerosis. *Tissue Antigens*, **76** (2), 110–118.
- Gregory, B.L., Shelton, G.D., Bali, D.S. *et al.* (2007) Glycogen storage disease type IIIa in Curly-Coated Retrievers. *Journal of Veterinary Internal Medicine*, **21**, 40–46.
- Grieco, V., Riccardi, E., Greppi, G.F. *et al.* (2008) Canine testicular tumours: a study on 232 dogs. *Journal of Comparative Pathology*, **138** (2–3), 86–89.
- Griffiths, I.R., McCulloch, M.C. & Abrahams, S. (1985) Progressive axonopathy: an inherited neuropathy of Boxer dogs. II. The nature and distribution of the pathological changes. *Neuropathology and Applied Neurobiology*, **11**, 431–446.
- Griffiths, L.G., Tiruneh, R., Sullivan, M. & Reid, S.W.J. (2000) Oropharyngeal penetrating injuries in 50 dogs: A retrospective study. *Veterinary Surgery*, **29** (5), 383–388.
- Grindem, C.B., Breitschwerdt, E.B., Corbett, W.T. & Jans, H.E. (1991) Epidemiologic survey of thrombocytopenia in dogs: A report on 987 cases. *Veterinary Clinical Pathology*, **20** (2), 38–43.
- Grüntzig, K., Graf, R., Boo, G. *et al.* (2016) Swiss Canine Cancer Registry 1955–2008: Occurrence of the most common tumour diagnoses and influence of age, breed, body size, sex and neutering status on tumour development. *Journal of Comparative Pathology*, **155**, 156–170.
- Grützner, N., Bishop, M.A., Suchodolski, J.S. & Steiner, J.M. (2010) Association study of cobalamin deficiency in the Chinese Shar Pei. *Journal of Heredity*, **101** (2), 211–217.
- Grützner, N., Cranford, S.M., Norby, B. *et al.* (2012) Evaluation of serum cobalamin concentrations in dogs of 164 dog breeds (2006–2010). *Journal of Veterinary Diagnostic Investigation*, **24** (6), 1105–1114.
- Grützner, N., Heilmann, R.M., Bridges, C.S., Suchodolski, J.S. & Steiner, J.M. (2013) Serum concentrations of canine alpha1-proteinase inhibitor in cobalamin-deficient Yorkshire Terrier dogs. *Journal of Veterinary Diagnostic Investigation*, **25** (3), 376–385.
- Guandalini, A., Di Girolamo, N., Santillo, D. *et al.* (2016) Epidemiology of ocular disorders presumed to be inherited in three large Italian dog breeds in Italy. *Veterinary Ophthalmology*, **20** (5), 420–426.
- Gubbels, E.J., Scholten, J., Janss, L. & Rothuizen, J. (2009) Relationship of cryptorchidism with sex ratios and litter sizes in 12 dog breeds. *Animal Reproduction Science*, **113** (1–4), 187–195.
- Guilliard, M. (2007) Central tarsal bone fracture in the Border Collie. *Journal of Small Animal Practice*, **48**, 414–417.

- Guilliard, M.J., Segboer, I. & Shearer, D.H. (2010) Corns in dogs: signalment, possible aetiology and response to surgical treatment. *Journal of Small Animal Practice*, **51** (3), 162–168.
- Guillot, J., Bensignor, E., Jankowski, F. *et al.* (2003) Comparative efficacies of oral ketoconazole and terbinafine for reducing *Malassezia* population sizes on the skin of Basset Hounds. *Veterinary Dermatology*, **14**, 153–157.
- Guiot, L.P., Lansdowne, J.L., Rouppert, P. & Stanley, B.J. (2008) Hiatal hernia in the dog: A clinical report of four Chinese Shar Peis. *Journal of the American Animal Hospital Association*, **44** (6), 335–341.
- Gulløv, C.H., Toft, N., Baadsager, M.M.N. & Berendt, M. (2011) Epilepsy in the Petit Basset Griffon Vendeen: Prevalence, Semiology, and Clinical Phenotype. *Journal of Veterinary Internal Medicine*, **25** (6), 1372–1378.
- Gunby, J.M., Hardie, R.J. & Bjorling, D.E. (2004) Investigation of the potential heritability of persistent right aortic arch in Greyhounds. *Journal of the American Veterinary Medical Association*, **224** (7), 1120–1122.
- Gunn-Moore, D.A. & Crispin, S.M. (1998) Unusual ocular condition in young Burmese cats. *Veterinary Record*, **142**, 376.
- Gunn-Moore, D.A. & Thrusfield, M.V. (1995) Feline dystocia: prevalence and association with cranial conformation and breed. *Veterinary Record*, **136** (14), 350–353.
- Gunn-Moore, D.A. (2003) Feline lower urinary tract disease. *Journal of Feline Medicine and Surgery*, **5** (2), 133–138.
- Gunn-Moore, D.A., Dodkin, S.J. & Sparkes, A.H. (2002) An unexpected high prevalence of azotemia in Birman cats. *Journal of Feline Medicine and Surgery*, **4** (3), 165–166.
- Gunn-Moore, D.A., Farland, S.E., Bewer, J.I. *et al.* (2011) Mycobacterial disease in cats in Great Britain: I. Culture results, geographical distribution and clinical presentation of 339 cases. *Journal of Feline Medicine and Surgery*, **13** (12), 934–944.
- Gunn-Moore, D.A., McCann, T.M., Reed, N. *et al.* (2007) Prevalence of *Tritrichomonas foetus* infection in cats with diarrhoea in the UK. *Journal of Feline Medicine and Surgery*, **9** (3), 214–218.
- Gunn-Moore, D.A., Simpson, K.E. & Day, M.J. (2009) Blood types in Bengal cats in the UK *Journal of Feline Medicine and Surgery*, **11** (10), 826–828.
- Guptill, L., Glickman, L. & Glickman, N. (2003) Time trends and risk factors for diabetes mellitus in dogs: analysis of veterinary medical data base records (1970–1999). *Veterinary Journal*, **165** (3), 240–247.
- Guthrie, J.W., Keeley, B.J., Maddock, E., Bright, S.R. & May, C. (2012) Effect of signalment on the presentation of canine patients suffering from cranial cruciate ligament disease. *Journal of Small Animal Practice*, **53** (5), 273–277.
- Guy, N.C., Luescher, U.A., Dohoo, S.E. *et al.* (2001) Demographic and aggressive characteristics of dogs in a general veterinary caseload. *Applied Animal Behaviour Science*, **74** (1), 15–28.
- Guziewicz, K.E., Zangerl, B., Lindauer, S.J. *et al.* (2007) Bestrophin gene mutations cause canine multifocal retinopathy: A novel animal model for best disease. *Investigative Ophthalmology and Visual Science*, **48** (5), 1959–1967.
- Hadži-Milić, M., Đorđević, J. & Krstić, N. (2013) Cyclosporine A (CsA) treatment of chronic autoimmune-mediated keratoconjunctivitis sicca in dogs. *Acta veterinaria*, **63** (5–6), 677–685.
- Hagman, R., Ström Holst, B., Möller, L. *et al.* (2014) Incidence of pyometra in Swedish insured cats. *Theriogenology*, **82** (1), 114–120.
- Hague, D.W., Humphries, H.D., Mitchell, M.A. *et al.* (2015) Risk factors and outcomes in cats with acquired myasthenia gravis (2001–2012). *Journal of Veterinary Internal Medicine*, **29** (5), 1307–1312.
- Haines, D.M., Lording, P.M. & Penhale, W.J. (1984) Survey of thyroglobulin autoantibodies in dogs. *American Journal of Veterinary Research*, **45** (8), 1493–1497.
- Hakozaki, T., Iwata, M., Kanno, N. *et al.* (2015) Cervical intervertebral disk herniation in chondrodystrophoid and nonchondrodystrophoid small-breed dogs: 187 cases (1993–2013). *Journal of the American Veterinary Medical Association*, **247**, 1408–1411.
- Hall, S.J.G. & Wallace, M.E. (1996) Canine epilepsy: a genetic counselling programme for Keeshonds. *Veterinary Record*, **138** (15), 358–360.

- Hamamoto, Y., Hasegawa, D., Mizoguchi, S. *et al.* (2016) Retrospective epidemiological study of canine epilepsy in Japan using the International Veterinary Epilepsy Task Force classification 2015 (2003–2013): etiological distribution, risk factors, survival time, and lifespan. *BMC Veterinary Research*, **12** (1), 248.
- Hamelin, A., Begon, D., Conchou, F. *et al.* (2017) Clinical characterisation of polydactyly in Maine Coon cats. *Journal of Feline Medicine and Surgery*, **19** (4), 382–393.
- Hammer, A., Getzy, D., Upton, M. *et al.* (2001) Salivary gland neoplasia in the dog and cat: survival times and prognostic factors. *Journal of the American Animal Hospital Association*, **37** (5), 478–482.
- Hamor, R.E. (2000) Evaluation of results for Schirmer tear tests conducted with and without application of a topical anesthetic in clinically normal dogs of 5 breeds. *American Journal of Veterinary Research*, **61** (11), 1422.
- Hamp, S.E., Hamp, M., Olsson, S.E., Lindberg, R. & Schauman, P. (1997) Radiography of spontaneous periodontitis in dogs. *Journal of Periodontal Research*, **32** (7), 589–597.
- Hampson, E.C.G.M., Smith, R.I.E. & Bernays, M.E. (2002) Primary glaucoma in Burmese cats. *Australian Veterinary Journal*, **80** (11), 672–680.
- Hanna, F.Y. (2001) Lumbosacral osteochondrosis: radiological features and surgical management in 34 dogs. *Journal of Small Animal Practice*, **42**, 272–278.
- Hansen, N., Foster, S.F., Burrows, A.K. *et al.* (2015) Cutaneous asthenia (Ehlers-Danlos-like syndrome) of Burmese cats. *Journal of Feline Medicine and Surgery*, **17** (11), 954–963.
- Hanson, J.M., Tengvall, K., Bonnett, B.N. & Hedhammar, Å. (2016) Naturally occurring adrenocortical insufficiency: an epidemiological study based on a swedish-insured dog population of 525,028 dogs. *Journal of Veterinary Internal Medicine*, **30** (1), 76–84.
- Hanson, S.M., Smith, M.O., Walker, T.L. & Shelton, G.D. (1998) Juvenile-onset distal myopathy in Rottweiler dogs. *Journal of Veterinary Internal Medicine*, **12** (2), 103–108.
- Hanssen, I. (1991) Hip dysplasia in dogs in relation to their month of birth. *Veterinary Record*, **128** (18), 425–426.
- Hanssen, I., Falck, G., Grammeltved, A.T. *et al.* (1998) Hypochondroplastic dwarfism in the Irish Setter. *Journal of Small Animal Practice*, **39** (1), 10–14.
- Hansson-Hamlin, H. & Lilliehöök, L. (2009) A possible systemic rheumatic disorder in the Nova Scotia Duck Tolling Retriever. *Acta Veterinaria Scandinavica*, **51**, 16.
- Hansson-Hamlin, H. & Lilliehöök, L. (2013) Steroid-responsive meningitis-arteritis in Nova Scotia Duck Tolling Retrievers. *Veterinary Record*, **173** (21), 527.
- Harari, J., Patterson, J.S. & Rosenthal, R.C. (1986) Clinical and pathologic features of thyroid tumours in 26 dogs. *Journal of the American Veterinary Medical Association*, **188** (10), 1160–1164.
- Harasen, G. (2009) Sesamoid disease. *Canadian Veterinary Journal*, **50** (10), 1095.
- Hargis, A.M., Brignac, M.M., Al-Bagdadi, F.A.K., Muggli, F. & Mundell, A. (1991) Black hair follicular dysplasia in Black and White Saluki Dogs: Differentiation from color mutant alopecia in the Doberman Pinscher by microscopic examination of hairs. *Veterinary Dermatology*, **2** (2), 69–83.
- Hargis, A.M., Ihrke, P.J., Spangler, W.L. & Stannard, A.A. (1992) A retrospective clinicopathologic study of 212 dogs with cutaneous hemangiomas and hemangiosarcomas. *Veterinary Pathology*, **29** (4), 316–328.
- Hargis, A.M., Myers, S., Gortel, K. *et al.* (2013) Proliferative, lymphocytic, infundibular mural folliculitis and dermatitis with prominent follicular apoptosis and parakeratotic casts in four Labrador Retrievers: preliminary description and response to therapy. *Veterinary Dermatology*, **24** (3), 346–e77.
- Harnevik, L., Hoppe, A. & Söderkvist, P. (2006) SLC7A9 cDNA cloning and nutritional analysis of SLC3A1 and SLC7A9 in canine cystinuria. *Mammalian Genome*, **17** (7), 769–776.
- Harrison, R.L., Clark, L. & Corletto, F. (2012) Comparison of mean heart rate in anaesthetized dachshunds and other breeds of dog undergoing spinal magnetic resonance imaging. *Veterinary Anaesthesia and Analgesia*, **39**, 230–235.
- Hart, B.L., Hart, L.A., Thigpen, A.P. & Willits, N.H. (2016) Neutering of German Shepherd Dogs: associated joint disorders, cancers and urinary incontinence. *Veterinary Medicine and Science*, **2** (3), 191–199.

- Hartley, C., Donaldson, D., Smith, K.C. *et al.* (2012) Congenital keratoconjunctivitis sicca and ichthyosiform dermatosis in 25 Cavalier King Charles Spaniel dogs. Part I: clinical signs, histopathology, and inheritance. *Veterinary ophthalmology*, **15** (5), 315–326.
- Hartley, W.J., Barker, J.S.F., Wanner, R.A. & Farrow, B.R.H. (1978) Inherited cerebellar degeneration in the Rough Coated Collie. *Australian Veterinary Practitioner*, **8**, 79–85.
- Hartley, W.J. & Palmer, A.C. (1973) Ataxia in Jack Russell Terriers. *Acta Neuropathologica*, **26** (1), 71–74.
- Harvey, R.G. & Maxwell, P.J. (1996) The mineral composition of nails in normal dogs and comparison with shed nails in canine idiopathic onychomadesis. *Veterinary Dermatology*, **7** (1), 29–33.
- Haupt, K.H., Prieur, D.J., Hargis, A.M. *et al.* (1985) Familial canine dermatomyositis: clinicopathologic, immunologic and serologic studies. *American Journal of Veterinary Research*, **46**, 1870–1875.
- Hawkins, E.C., Basseches, J., Berry, C.R., Stebbins, M.E. & Ferris, K.K. (2003) Demographic, clinical, and radiographic features of bronchiectasis in dogs: 316 cases (1988–2000). *Journal of the American Veterinary Medical Association*, **223** (11), 1628–1635.
- Haworth, K., Putt, W., Cattanaach, B. *et al.* (2001) Canine homolog of the T-box transcription factor T; failure of the protein to bind to its DNA target leads to a short-tail phenotype. *Mammalian Genome*, **12** (3), 212–218.
- Hawthorne, J., Wallace, L., Fenner, W. & Waters, D. (2001) Fibrocartilaginous embolic myelopathy in Miniature Schnauzers. *Journal of the American Animal Hospital Association*, **37** (4), 374–383.
- Hayes, G. (2009) Gastrointestinal foreign bodies in dogs and cats: a retrospective study of 208 cases. *Journal of Small Animal Practice*, **50** (11), 576–583.
- Hayes, H.M. (1974a) Congenital umbilical and inguinal hernias in cattle, horses, swine, dogs, and cats: risk by breed and sex among hospital patients. *American Journal of Veterinary Research*, **35**, 839–842.
- Hayes, H.M. (1974b) Ectopic ureter in dogs: epidemiologic features. *Teratology*, **10** (2), 129–132.
- Hayes, H.M. (1984) Breed associations of canine ectopic ureter: a study of 217 female cases. *Journal of Small Animal Practice*, **25** (8), 501–504.
- Hayes H.M., Milne, K.L. & Mandell, C.P. (1981) Epidemiological features of feline mammary carcinoma. *Veterinary Record*, **108** (22), 476–479.
- Hayes, H.M. & Pendergrass, T.W. (1976) Canine testicular tumours and epidemiologic features of 410 dogs. *International Journal of Cancer*, **18** (4), 482–487.
- Hayes, H.M. & Sass, B. (1988) Chemoreceptor neoplasia: a study of the epidemiological features of 357 canine cases. *Journal of Veterinary Medicine Series A*, **35**, 401–408.
- Hayes, H.M., Selby, L.A., Wilson, G.P. & Hohn, R.B. (1979) Epidemiologic observations of canine elbow disease (emphasis on dysplasia). *Journal of the American Animal Hospital Association*, **14**, 449–453.
- Hayes, H.M. & Wilson, G.P. (1977) Hormone-dependent neoplasms of the canine perianal gland. *Cancer Research*, **37** (7 pt 1), 2068–2071.
- Hayes, H.M. & Wilson, G.P. (1986) Hospital incidence of hypospadias in dogs in North America. *Veterinary Record*, **118** (22), 605–607.
- Hayes, H.M., Wilson, G.P. & Fraumeni, H.F. (1982) Carcinoma of the nasal cavity and paranasal sinuses in dogs: descriptive epidemiology. *Cornell Veterinarian*, **72** (2), 168–179.
- Hayes, H.M., Wilson, G.P., Pendergrass, T.W. & Cox, V.S. (1985) Canine cryptorchism and subsequent testicular neoplasia: case-control study with epidemiologic update. *Teratology*, **32** (1), 51.
- Hayes, H.M. & Young, J.L. (1978) Epidemiologic features of canine ovarian neoplasms, *Gynecologic Oncology*, **6**, 348–353.
- Haywood, S., Rutgers, H.C. & Christian, M.K. (1988) Hepatitis and copper accumulation in Skye Terriers. *Veterinary Pathology*, **25** (6), 408–414.
- He, Q., Madsen, M., Kilkenney, A. *et al.* (2005) Amnionless function is required for cubilin brush-border expression and intrinsic factor-cobalamin (vitamin B12) absorption in vivo, *Blood. American Society of Hematology*, **106**, 1447–1453.
- Headley, S.A. & Graça, D.L. (2000) Canine distemper: epidemiological findings of 250 cases. *Brazilian Journal of Veterinary Research and Animal Science*, **37** (2), 136–140.
- Hearon, K., Berg, J.M., Bonczynski, J.J. *et al.* (2014) Upper thoracic disc disease (T1–T9) in large-breed dogs. *Journal of the American Animal Hospital Association*, **50** (2), 105–111.

- Heckler, M.C.T., Sakate, M., Amorim, R.L. *et al.* (2011) Lissencephaly in dogs and cats: a review. *Clinica Veterinaria*, **16** (90), 76–80.
- Heejin Oui, Juyeon Oh, Seoyeon Keh *et al.* (2015) Measurements of the pulmonary vasculature on thoracic radiographs in healthy dogs compared to dogs with mitral regurgitation. *Veterinary Radiology and Ultrasound*, **56** (3), 251–256.
- Hegstad-Davies, R.L., Torres, S.M.F., Sharkey, L.C. *et al.* (2015) Breed-specific reference intervals for assessing thyroid function in seven dog breeds. *Journal of Veterinary Diagnostic Investigation*, **27**, 716–727.
- Heidner, G.L., Kornegay, J.N., Page, R.L., Dodge, R.K. & Thrall, D.E. (1991) Analysis of survival in a retrospective study of 86 dogs with brain tumors. *Journal of Veterinary Internal Medicine*, **5** (4), 219–226.
- Heiene, R., Indrebø, A., Kvart, C., Skaalnes, H.M. & Ulstad, A.K. (2000) Prevalence of murmurs consistent with aortic stenosis among boxer dogs in Norway and Sweden. *Veterinary Record*, **147**, 152–156.
- Heilmann, R.M., Grützner, N., Iazbik, M.C. *et al.* (2017) Hyperhomocysteinemia in Greyhounds and its association with hypofolatemia and other clinicopathologic variables. *Journal of Veterinary Internal Medicine* **31** (1), 109–116.
- Heinrich, C.L., Lakhani, K.H., Featherstone, H.J. *et al.* (2006) Cataract in the UK Leonberger population. *Veterinary Ophthalmology*, **9** (5), 350–356.
- Heitmann, M., Hamann, H., Brahm, R. *et al.* (2005) Analysis of prevalence of presumed inherited eye disease in Entlebucher Mountain Dogs. *Veterinary Ophthalmology*, **8** (3), 145–151.
- Heller, A.R., van der Woerd, A., Gaarder, J.E. *et al.* (2017) Sudden acquired retinal degeneration in dogs: breed distribution of 495 canines. *Veterinary Ophthalmology*, **20** (2), 103–106.
- Heller, D.A., Stebbins, M.E., Reynolds, T.L. *et al.* (2005) A retrospective study of 87 cases of canine soft tissue sarcomas, 1986–2001. *International Journal of Applied Research in Veterinary Medicine*, **3** (2), 81–87.
- Hendrix, D.V., Nasisse, M.P., Cowen, P. & Davidson, M.G. (1993) Clinical signs, concurrent diseases and risk factors associated with retinal detachment in dogs. *Progress in Veterinary and Comparative Ophthalmology*, **3**, 87–91.
- Henke, D., Böttcher, P., Doherr, M.G., Oechtering, G. & Flegel, T. (2008) Computer-assisted magnetic resonance imaging brain morphometry in American Staffordshire Terriers with cerebellar cortical degeneration. *Journal of Veterinary Internal Medicine*, **22** (4), 969–975.
- Henrich, E., Hildebrandt, N., Schneider, C., Hassdenteufel, E. & Schneider, M. (2011) Transvenous coil embolization of patent ductus arteriosus in small (≤ 3.0 kg) dogs. *Journal of Veterinary Internal Medicine*, **25**, 65–70.
- Henry, C.J., William, G.B., Whitley, E.M. *et al.* (2005) Canine digital tumors: a veterinary cooperative oncology group retrospective study of 64 dogs. *Journal of Veterinary Internal Medicine*, **19** (5), 720–724.
- Herrera, H.D. (2007) Severe, unilateral, unresponsive keratoconjunctivitis sicca in 16 juvenile Yorkshire Terriers. *Veterinary Ophthalmology*, **10** (5), 285–288.
- Herrtage, E. and Houlton, J. E. (1979) Collapsing Clumber Spaniels. *Veterinary Record*, **105**, 334.
- Hertel, E., Bergström, T., Kell, Ü. *et al.* (2010) Retinal degeneration in nine Swedish Jämthund dogs. *Veterinary Ophthalmology*, **13** (2), 110–116.
- Heske, L., Körberg, I.B., Nødtvedt, A. & Jäderlund, K.H. (2015) Clinical characteristics of epilepsy of unknown origin in the Rottweiler breed. *Acta Veterinaria Scandinavica*, **57** (1), 75.
- Heske, L., Nødtvedt, A., Jäderlund, K.H., Berendt, M. & Egenvall, A. (2014) A cohort study of epilepsy among 665,000 insured dogs: Incidence, mortality and survival after diagnosis. *Veterinary Journal*, **202** (3), 471–476.
- Hess, R.S., Kass, P.H., Shofer, F.S., Van Winkle, T.J. & Washabau, R.J. (1999) Evaluation of risk factors for fatal acute pancreatitis in dogs. *Journal of the American Veterinary Medical Association*, **214** (1), 46–51.
- Hess, R.S., Kass, P.H. & Van Winkle, T.J. (2003) Association between diabetes mellitus, hypothyroidism or hyperadrenocorticism, and atherosclerosis in dogs. *Journal of Veterinary Internal Medicine*, **17** (4), 489–494.

- Hess, R.S., Kass, P.H. & Ward, C.R. (2000) Breed distribution of dogs with diabetes mellitus admitted to a tertiary care facility. *Journal of the American Veterinary Medical Association*, **216** (9), 1414–1417.
- Hesse, A. (1990) Canine urolithiasis: epidemiology and analysis of urinary calculi. *Journal of Small Animal Practice*, **31** (12), 599–604.
- Heyman, S.J., Diefenderfer, D.L., Goldschmidt, M.H. *et al.* (1992) Canine axial skeletal osteosarcoma a retrospective study of 116 cases (1986 to 1989). *Veterinary Surgery*, **21** (4), 304–310.
- Higgins, R.J., Dickinson, P.J., Kube, S.A. *et al.* (2008) Necrotising meningoencephalitis in five Chihuahua dogs. *Veterinary Pathology*, **45**, 336–346.
- Higgins, R.J., LeCouteur, R.A., Kornegay, J.N. *et al.* (1998) Late-onset progressive spinocerebellar degeneration in Brittany Spaniel dogs. *Acta Neuropathologica*, **96**, 97–101.
- Hill, R.C. & Van Winkle, T.J. (1993) Acute necrotizing pancreatitis and acute suppurative pancreatitis in the cat. A retrospective study of 40 cases (1976–1989). *Journal of Veterinary Internal Medicine*, **7** (1), 25–33.
- Hillbertz, N.H.C.S. (2005) Inheritance of dermoid sinus in the Rhodesian Ridgeback. *Journal of Small Animal Practice*, **46** (2), 71–74.
- Hinton, L.E., McLoughlin, M.A., Johnson, S.E. & Weisbrode, S.E. (2002) Spontaneous gastroduodenal perforation in 16 dogs and seven cats (1982–1999). *Journal of the American Animal Hospital Association*, **38** (2), 176–187.
- Hirsch, V.M. & Cunningham, T.A. (1984) Hereditary anomaly of neutrophil granulation in Birman cats. *American Journal of Veterinary Research*, **45**, 2170–2174.
- Hirth, R.S., Greenstein, E.T. & Peer, R.L. (1974) Anterior capsular opacities (spurious cataracts) in Beagle Dogs. *Veterinary Pathology*, **11**, 181–194.
- Ho, L.K., Troy, G.C. & Waldron, D.R. (2011) Clinical outcomes of surgically managed ectopic ureters in 33 dogs. *Journal of the American Animal Hospital Association*, **47** (3), 196–202.
- Hoffmann, G., van den Ingh, T.S., Bode, P. *et al.* (2006) Copper-associated chronic hepatitis in Labrador Retrievers. *Journal of Veterinary Internal Medicine*, **20** (4), 856–861.
- Hogg, G. G., Horton, B. J. & Brown, H. (1978) Inherited pyruvate kinase deficiency and normal haematologic values in Australian Basenji dogs. *Australian Veterinary Journal*, **54**, 367–370.
- Holder, A.L., Kennedy, L.J., Ollier, W.E.R. & Catchpole, B. (2015) Breed differences in development of anti-insulin antibodies in diabetic dogs and investigation of the role of dog leukocyte antigen (DLA) genes. *Veterinary Immunology and Immunopathology*, **167**, 130–138.
- Holder, A.L., Price, J.A., Adams, J.P. *et al.* (2014) A retrospective study of the prevalence of the canine degenerative myelopathy associated superoxide dismutase 1 mutation (SOD1:c.118G>A) in a referral population of German Shepherd Dogs from the UK. *Canine Genetics and Epidemiology*, **1**, 10.
- Holle, D.M., Stankovics, M.E., Sarna, C.S. *et al.* (1999) The geographic form of retinal dysplasia in dogs is not always a congenital abnormality. *Veterinary Ophthalmology*, **2** (1), 61–66.
- Höllmer, M., Willesen, J.L., Jensen, A.T. *et al.* (2008) Aortic stenosis in the Dogue de Bordeaux. *Journal of Small Animal Practice*, **49**, 432–437.
- Holly, V.L., Sandmeyer, L.S., Bauer, B.S., Verges, L., Grahn, B.H. (2016) Golden Retriever cystic uveal disease: a longitudinal study of iridociliary cysts, pigmentary uveitis, and pigmentary/cystic glaucoma over a decade in western Canada. *Veterinary Ophthalmology*, **19** (3), 237–244.
- Holm, B.R. (2003) Efficacy of milbemycin oxime in the treatment of canine generalized demodicosis: a retrospective study of 99 dogs (1995–2000). *Veterinary Dermatology*, **14** (4), 189–195.
- Holm, B.R., Rest, J.R. & Seewald, W. (2004) A prospective study of the clinical findings, treatment and histopathology of 44 cases of pyotraumatic dermatitis. *Veterinary Dermatology*, **15** (6), 369–376.
- Holm, K.S., Morris, D.O., Gomez, S.M. *et al.* (1999) Eosinophilic dermatitis with edema in nine dogs, compared with eosinophilic cellulitis in humans. *Journal of the American Veterinary Medical Association*, **215**, 649–652.
- Holmes, M.A. (2007) Evaluation of the evidence. *Veterinary Clinics of North America: Small Animal Practice*, **37** (3), 447–462.
- Holmes, M.A. & Ramey, D.W. (2007) An introduction to evidence-based veterinary medicine. *Veterinary Clinics of North America: Equine Practice*, **23** (2), 191–200.
- Holst, B.S., Axné, E., Öhlund, M. *et al.* (2017) Dystocia in the cat evaluated using an insurance database. *Journal of Feline Medicine and Surgery*, **19** (1), 42–47.

- Holt, P. & Moore, A.H. (1995) Canine ureteral ectopia: an analysis of 175 cases and comparison of surgical treatments. *Veterinary Record*, **136** (14), 345–349.
- Holt, P.E. & Thrusfield, M.V. (1993) Association in bitches between breed, size, neutering and docking, and acquired urinary incontinence due to incompetence of the urethral sphincter mechanism. *Veterinary Record*, **133** (8), 177–180.
- Holt, P.E. (1985) Urinary incontinence in the bitch due to sphincter mechanism incompetence: prevalence in referred dogs and retrospective analysis of sixty cases. *Journal of Small Animal Practice*, **26** (4), 181–190.
- Holt, P.E., Hotston Moore, A. & Thrusfield, M.V. (2000) Breed predisposition to ureteral ectopia in bitches in the UK. *Veterinary Record*, **146** (19), 561.
- Hood, J.C., Dowling, J., Bertram, J.F. *et al.* (2002a) Correlation of histopathological features and renal impairment in autosomal dominant Alport syndrome in Bull Terriers. *Nephrology, Dialysis, Transplantation*, **17** (11), 1897–1908.
- Hood, J.C., Huxtable, C., Naito, I. *et al.* (2002b) A novel model of autosomal dominant Alport syndrome in Dalmatian dogs. *Nephrology, Dialysis, Transplantation*, **17** (12), 2094–2098.
- Höpfner, R., Glaus, T., Gardelle, O. *et al.* (2010) Prävalenz von Herzgeräuschen, Aorten- und Pulmonalstenosen bei Zuchtuntersuchungen von Boxern in der Schweiz. *Schweizer Archiv für Tierheilkunde*, **152**, 319–324.
- Hoppe, A., Swenson, L., Jönsson, L. & Hedhammar, A. (1990) Progressive nephropathy due to renal dysplasia in Shih Tzu dogs in Sweden: a clinical pathological and genetic study. *Journal of Small Animal Practice*, **31** (2), 83–91.
- Hoppe, F. & Svalastoga, E. (1980) Temporomandibular dysplasia in American Cocker Spaniels. *Journal of Small Animal Practice*, **21**, 675–678.
- Hornok, S., Edelhofer, R. & Farkas, R. (2006) Seroprevalence of canine babesiosis in Hungary suggesting breed predisposition. *Parasitology Research*, **99** (6), 638–642.
- Hosgood, G., Hedlund, C., Pechman, R. & Dean, P. (1995) Perineal herniorrhaphy: perioperative data from 100 dogs. *Journal of the American Animal Hospital Association*, **31**, 331–342.
- Houston, D.M. & Moore, A.E. (2009) Canine and feline urolithiasis: examination of over 50 000 urolith submissions to the Canadian veterinary urolith centre from 1998 to 2008. *Canadian Veterinary Journal*, **50** (12), 1263–1268.
- Houston, D.M., Moore, A.E., Favrin, M.G. & Hoff, B. (2004) Canine urolithiasis: a look at over 16 000 urolith submissions to the Canadian Veterinary Urolith Centre from February 1998 to April 2003. *Canadian Veterinary Journal*, **45** (3), 225–230.
- Houston, D.M., Ribble, C.S. & Head, L.L. (1996) Risk factors associated with parvovirus enteritis in dogs: 283 cases (1982–1991). *Journal of the American Veterinary Medical Association*, **208** (4), 542–546.
- Houston, D.M., Vanstone, N.P., Moore, A.E.P. *et al.* (2016) Evaluation of 21 426 feline bladder urolith submissions to the Canadian Veterinary Urolith Centre (1998–2014). *Canadian Veterinary Journal*, **57** (2), 196–201.
- Hughes, A.M., Jokinen, P., Bannasch, D.L. *et al.* (2010) Association of a dog leukocyte antigen class II haplotype with hypoadrenocorticism in Nova Scotia Duck Tolling Retrievers. *Tissue Antigens*, **75** (6), 684–690.
- Hughes, A.M., Nelson, R.W., Famula, T.R. *et al.* (2007) Clinical features and heritability of hypoadrenocorticism in Nova Scotia Duck Tolling Retrievers: 25 cases (1994–2006). *Journal of the American Veterinary Medical Association*, **231** (3), 407–412.
- Hulsmeyer, V., Zimmermann, R., Brauer, C., Sauter-Louis, C. & Fischer, A. (2010) Epilepsy in Border Collies: clinical manifestation, outcome, and mode of inheritance. *Journal of Veterinary Internal Medicine*, **24**, 171–178.
- Hultgren, B.D., Stevens, J.B. & Hardy, R.M. (1986) Inherited, chronic, progressive hepatic degeneration in Bedlington terriers with increased liver copper concentrations: clinical and pathologic observations and comparison with other copper-associated liver diseases. *American Journal of Veterinary Research*, **47**, 365–377.
- Hungs, M., Fan, J., Lin, L. (2001) Identification and functional analysis of mutations in the hypocretin (orexin) genes of narcoleptic canines. *Genome Research*, **11** (4), 531–539.

- Hunt, G.B. (2004) Effect of breed on anatomy of portosystemic shunts resulting from congenital diseases in dogs and cats: a review of 242 cases. *Australian Veterinary Journal*, **82** (12), 746–749.
- Huxtable, C.R. & Davis, P.E. (1976) The pathology of polyarthritis in young Greyhounds. *Journal of Comparative Pathology*, **86**, 11–21.
- Hyttönen, M.K., Grall, A., Hédan, B. *et al.* (2009) Ancestral T-Box mutation is present in many, but not all, short-tailed dog breeds. *Journal of Heredity*, **100** (2), 236–240.
- Iazbik, M.C. & Couto, C.G. (2005) Morphologic characterization of specific granules in Greyhound eosinophils. *Veterinary Clinical Pathology*, **34** (2), 140–143.
- Iazbik, M.C., O'Donnell, M., Marin, L. *et al.* (2010) Prevalence of dog erythrocyte antigens in retired racing Greyhounds. *Veterinary Clinical Pathology*, **39** (4), 433–435.
- Igarashi, H., Ohno, K., Fujiwara-Igarashi, A. *et al.* (2015) Inflammatory colorectal polyps in miniature Dachshunds frequently develop ventrally in the colorectal mucosa. *Veterinary Journal*, **203**, 256–258.
- Igna, C., Dascălu, R., Bumb, D., Sicoe, B. & Schüsler, L. (2016) The incidence of panosteitis in dogs admitted in surgery clinic of the Faculty of Veterinary Medicine Timisoara: Retrospective study (2000–2015). *Terra Nova*, **2**, 1.29.
- Ihrke, P.J., Stannard, A.A., Ardans, A.A. & Griffin, C.E. (1985) Pemphigus foliaceus in dogs: a review of 37 cases. *Journal of the American Veterinary Medical Association*, **186** (1), 59–66.
- Imre, M., Farkas, R., Ilie, M.S., Imre, K. & Dărăbuș, G. (2013) Survey of babesiosis in symptomatic dogs from Romania: Occurrence of *Babesia gibsoni* associated with breed. *Ticks and Tick-borne Diseases*, **4**, 500–502.
- Inal Gultekin, G., Raj, K., Foureman, P. *et al.* (2012a) Erythrocytic pyruvate kinase mutations causing hemolytic anemia, osteosclerosis, and secondary hemochromatosis in dogs. *Journal of Veterinary Internal Medicine*, **26**, 935–944.
- Inal Gultekin, G., Raj, K., Lehman, S., Hillström, A. & Giger, U. (2012b) Missense mutation in PFKM associated with muscle-type phosphofructokinase deficiency in the Wachtelhund dog. *Molecular and Cellular Probes*, **26** (6), 243–247.
- Indrebø, A., Langeland, M., Juul, H.M. *et al.* (2008) A study of inherited short tail and taillessness in Pembroke Welsh Corgi. *Journal of Small Animal Practice*, **49** (5), 220–224.
- Inoue, M., Hasegawa, A., Hosoi, Y. & Sugiura, K. (2016) Association between breed, gender and age in relation to cardiovascular disorders in insured dogs in Japan. *Journal of Veterinary Medical Science*, **78**, 347–350.
- Israel, S.K., Levine, J.M., Kerwin, S.C., Levine, G.J. & Fosgate, G.T. (2009) The relative sensitivity of computed tomography and myelography for identification of thoracolumbar intervertebral disk herniations in dogs. *Veterinary Radiology and Ultrasound*, **50** (3), 247–252.
- Israeli, I., Steiner, J., Segev, G., Kass, P.H. *et al.* (2012) Serum pepsinogen-A, canine pancreatic lipase immunoreactivity, and C-reactive protein as prognostic markers in dogs with gastric dilatation-volvulus. *Journal of Veterinary Internal Medicine*, **26** (4), 920–928.
- It, V., Barrientos, L., López Gappa, J. *et al.* (2010) Association of canine juvenile generalized demodicosis with the dog leukocyte antigen system. *Tissue Antigens*, **76** (1), 67–70.
- Itoh, H., Hara, Y., Yoshimi, N. *et al.* (2008) A retrospective study of intervertebral disc herniation in dogs in Japan: 297 cases. *Journal of Veterinary Medical Science*, **70** (7), 701–706.
- Itoh, T., Uchida, K., Ishikawa, K. *et al.* (2005) Clinicopathological survey of 101 canine mammary gland tumors: differences between small-breed dogs and others. *Journal of Veterinary Medical Science*, **67** (3), 345–347.
- Jackson, H.A. (2004) Eleven cases of vesicular cutaneous lupus erythematosus in Shetland Sheepdogs and Rough Collies: clinical management and prognosis. *Veterinary Dermatology*, **15** (1), 37–41.
- Jackson, H.A. & Olivry, T. (2001) Ulcerative dermatosis of the Shetland Sheepdog and Rough Collie may represent a novel vesicular variant of cutaneous lupus erythematosus. *Veterinary Dermatology*, **12**, 19–27.
- Jackson, H.A., Olivry, T., Berget, F. *et al.* (2004) Immunopathology of vesicular cutaneous lupus erythematosus in the Rough Collie and Shetland sheepdog: a canine homologue of subacute cutaneous lupus erythematosus in humans. *Veterinary Dermatology*, **15** (4), 230–239.
- Jacobs, G., Mahaffey, M. & Rawlings, C.A. (1990) Valvular pulmonic stenosis in four Boykin Spaniels. *Journal of the American Animal Hospital Association*, **26**, 247–252.

- Jaderlund, K.H., Orvind, E., Johnsson, E. *et al.* (2007) A neurologic syndrome in Golden Retrievers presenting as a sensory ataxic neuropathy. *Journal of Veterinary Internal Medicine*, **21**, 1307–1315.
- Jae, H. K., Kyung, I. K., Hyun, J. S. *et al.* (2005) Color-dilution alopecia in dogs. *Journal of Veterinary Science*, **6** (3), 259–261.
- Jaeger, G.H., Marcellin-Little, D.J. & Ferretti, A. (2007) Morphology and correction of distal tibial valgus deformities. *Journal of Small Animal Practice*, **48** (12), 678–682.
- Jaeger, J.Q., Mattoon, J.S., Bateman, S.W. & Morandi, F. (2003) Combined use of ultrasonography and contrast enhanced computed tomography to evaluate acute necrotizing pancreatitis in two dogs. *Veterinary Radiology and Ultrasound*, **44** (1), 72–79.
- Jaeger, K., Linek, M., Power, H.T. *et al.* (2010) Breed and site predispositions of dogs with atopic dermatitis: a comparison of five locations in three continents. *Veterinary Dermatology*, **21** (1), 119–123.
- Jagannathan, V., Bannoehr, J., Plattet, P. *et al.* (2013) A mutation in the SUV39H2 gene in Labrador Retrievers with hereditary nasal parakeratosis (HNPK) provides insights into the epigenetics of keratinocyte differentiation. *PloS Genetics*, **9** (10), e1003848.
- Jaggy, A. & Vandevelde, M. (1988) Multisystem neuronal degeneration in Cocker Spaniels. *Journal of Veterinary Internal Medicine*, **2**, 117–120.
- Jaggy, A., Faissler, D., Gaillard, C. *et al.* (1998) Genetic aspects of idiopathic epilepsy in Labrador Retrievers. *Journal of Small Animal Practice*, **39** (6), 275–280.
- Jaggy, A., Gaillard, C., Lang, J. & Vandevelde, M. (1988) Hereditary cervical spondylopathy (wobbler syndrome) in the Borzoi dog. *Journal of the American Animal Hospital Association*, **24**, 453–460.
- Jaggy, A., Land, J. & Schawalder, P. (1987) Cauda equina syndrome in the dog. *Schweizer Archiv für Tierheilkunde*, **129** (4), 171–192.
- Jagielski, D., Lechowski, R., Hoffmann-Jagielska, M. & Winiarczyk, S. (2002) A retrospective study of the incidence and prognostic factors of multicentric lymphoma in dogs (1998–2000). *Journal of Veterinary Medicine. Series A*, **49** (8), 419–424.
- James, R.W. (1991) Persistent pupillary membrane in Basenji dogs. *Veterinary Record*, **128**, 287–288.
- Jamne, Ø., Nafstad, P., Skancke, E. & Grøndalen, J. (1998) Primary ciliary dyskinesia. A case of respiratory problems in Bichon Frise dogs. *Norsk Veterinærtidsskrift*, **110**, 199–202.
- Jansen, B., Valli, V.E., Thorner, P., Bauman, R. & Lumsden, J.H. (1987) Samoyed hereditary glomerulopathy: serial, clinical and laboratory (urine, serum biochemistry and hematology) studies. *Canadian Journal of Veterinary Research*, **51** (3), 387–393.
- Jansen, J.H. & Arnesen, K. (1990) Oxalate nephropathy in a Tibetan Spaniel litter. A probable case of primary hyperoxaluria. *Journal of Comparative Pathology*, **103** (1), 79–84.
- Janssens, S.D., Haagsman, A.N. & Haar, G. (2017) Middle ear polyps: results of traction avulsion after a lateral approach to the ear canal in 62 cats (2004–2014). *Journal of Feline Medicine and Surgery*, **19** (8), 803–808.
- Jardim, L., Andrade-Neto, J. & Alessi, A. (1999) Neuronal vacuolation and spongiform lesions in young Rottweiler dogs. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, **51** (5), 449–452.
- Järvinen, A.K., Saario, E., Andresen, E. *et al.* (1995) Lung injury leading to respiratory distress syndrome in young Dalmatian dogs. *Journal of Veterinary Internal Medicine*, **9**, 162–168.
- Jeanson-Leh, L., Lameth, J., Krimi, S. *et al.* (2014) Serum profiling identifies novel muscle 16miRNA and cardiomyopathy-related miRNA biomarkers in Golden Retriever muscular dystrophy dogs and Duchenne muscular dystrophy patients. *American Journal of Pathology* **184** (11) 2885–2898.
- Jefferies, R., Ryan, U.M., Jardine, J. *et al.* (2007) Blood, Bull Terriers and babesiosis: further evidence for direct transmission of *Babesia gibsoni* in dogs. *Australian Veterinary Journal*, **85** (11), 459–463.
- Jeffery, N. (2015) Liberating the (data) population from subjugation to the 5% (P-value). *Journal of Small Animal Practice*, **56** (8), 483–484.
- Jenkins, W.L., van Dyk, E. & McDonald, C.B. (1976) Myasthenia gravis in a Fox Terrier litter. *Journal of the South Africa Veterinary Association*, **47** (1), 59–62.
- Jepsen-Grant, K., Pollard, R. & Johnson, L. (2013) Vertebral heart scores in eight dog breeds. *Veterinary Radiology and Ultrasound*, **54** (1), 3–8.
- Jian-guo, L., Fang, W. & Dapeng, Z. (2009) Characteristic and structure of pulmonary artery blood vessel of Tibetan Mastiff living at different altitudes. *Journal of Anhui Agricultural Sciences*, **1**, 064.

- Jitpean, S., Hagman, R., Ström Holst, B. *et al.* (2012) Breed variations in the incidence of pyometra and mammary tumours in Swedish dogs. *Reproduction in Domestic Animals*, **47** (suppl. 6), 347–350.
- Jochman-Edwards, C.M., Tilley, L.P., Lichtenberger, M., Smith, F.W.K. & Kirby, R. (2002) Electrocardiographic findings in Miniature Schnauzers with syncope. *Journal of Veterinary Emergency and Critical Care*, **12** (4), 253–259.
- John, J., Wu, M.-F., Maidment, N.T. *et al.* (2004) Developmental changes in CSF hypocretin-1 (orexin-A) levels in normal and genetically narcoleptic Doberman Pinschers. *Journal of Physiology*, **560** (2), 587–592.
- Johnsen, D.A.J., Maggs, D.J. & Kass, P.H. (2006) Evaluation of risk factors for development of secondary glaucoma in dogs: 156 cases (1999–2004). *Journal of the American Veterinary Medical Association*, **229** (8), 1270–1274.
- Johnson, B.W. (1991) Congenitally abnormal visual pathways of Siamese cats. *Compendium on Continuing Education for the Practicing Veterinarian*, **13** (3), 374–378.
- Johnson, G.F., Sternlieb, I., Twedt, D.C. *et al.* (1980) Inheritance of copper toxicosis in Bedlington Terriers. *American Journal of Veterinary Research*, **41** (11), 1865–1866.
- Johnson, G.R. & Feeney, D.A. (1984) Recurring lung lobe torsion in three Afghan Hounds. *Journal of the American Veterinary Medical Association*, **184**, 842–845.
- Johnson, L.R. & Fales, W.H. (2001) Clinical and microbiologic findings in dogs with bronchoscopically diagnosed tracheal collapse: 37 cases (1990–1995). *Journal of the American Veterinary Medical Association*, **219** (9), 1247–1250.
- Johnson, L.R., Mayhew, P.D., Steffey, M.A. *et al.* (2013) Upper airway obstruction in Norwich Terriers: 16 cases. *Journal of Veterinary Internal Medicine*, **27** (6), 1409–1415.
- Johnson, M.S., Martin, M., Edwards, D., French, A. & Henley, W. (2004) Pulmonic stenosis in dogs: balloon dilation improves clinical outcome. *Journal of Veterinary Internal Medicine*, **18** (5), 656–662.
- Johnson, M.S. & Martin, M.W.S. (2007) Successful medical treatment of 15 dogs with pyothorax. *Journal of Small Animal Practice*, **48** (1), 12–16.
- Johnson, P., De Risio, L., Sparkes, A. (2012) Clinical, morphologic, and morphometric features of cranial thoracic spinal stenosis in large and giant breed dogs. *Veterinary Radiology and Ultrasound*, **53** (5), 524–534.
- Johnson, R.P., Neer, T.M., Partington, B.P. *et al.* (2001) Familial cerebellar ataxia with hydrocephalus in Bull Mastiffs. *Veterinary Radiology and Ultrasound*, **42**, 246–249.
- Jokinen, P., Rusanen, E.M., Kennedy, L.J. *et al.* (2011) MHC class II risk haplotype associated with canine chronic superficial keratitis in German Shepherd Dogs. *Veterinary Immunology and Immunopathology*, **140**, 37–41.
- Jolly, R.D., Hopwood, J.J., Marshall, N.R. *et al.* (2012) Mucopolysaccharidosis type VI in a Miniature Poodle-type dog caused by a deletion in the arylsulphatase B gene. *New Zealand Veterinary Journal*, **60** (3), 183–188.
- Jones, B. & Alley, M. (1985) Primary idiopathic hypoparathyroidism in St. Bernard dogs. *New Zealand Veterinary Journal*, **33** (6), 94–97.
- Jones, B.R., Anderson, L.J., Barnes, G.R.G., Johnstone, A.C. & Juby, W.D. (1977), Myotonia in related Chow Chow dogs. *New Zealand Veterinary Journal*, **25**, 217–220.
- Jones, B.R., Omodo-Eluk, A.J., Larkin, H., Rogers, K.D. & Sperrin, M. (2001) Canine uroliths: analysis of uroliths from dogs in Ireland. *Irish Veterinary Journal*, **54**, 629–632.
- Jones, B.R., Kirkman, J.H., Hogan, J. & Holmes, S. (1998) Analysis of uroliths from cats and dogs in New Zealand, 1993–96. *New Zealand Veterinary Journal*, **46** (6), 233–236.
- Juraschko, K., Meyer-Lindenberg, A., Nolte, I. *et al.* (2003) Analysis of systematic effects on congenital sensorineural deafness in German Dalmatian dogs. *Veterinary Journal*, **166** (2), 164–169.
- Jurina, K. & Grevel, V. (2004) Spinal arachnoid pseudocysts in 10 Rottweilers. *Journal of Small Animal Practice*, **45** (1), 9–15.
- Kaae, J.A., Callan, M.B. & Brooks, M. (2007) Hereditary factor VII deficiency in the Alaskan Klee Kai. *Journal of Veterinary Internal Medicine*, **21**, 976–981.
- Kahn, S.A., Mullin, C.M., de Lorimier, L-P. *et al.* (2013) Doxorubicin and deracoxib adjuvant therapy for canine splenic hemangiosarcoma: A pilot study. *Canadian Veterinary Journal*, **54** (3), 237–242.

- Kaiser, E., Krauser, K. & Schwartz-Porsche, D. (1991) Lafora disease (progressive myoclonic epilepsy) in the Bassett Hound – possibility of early diagnosis using muscle biopsy? *Tierärztliche Praxis*, **19**, 290–295.
- Kalaher, K.M., Anderson, W.I. & Scott, D.W. (1990) Neoplasms of the apocrine sweat glands in 44 dogs and 10 cats. *Veterinary Record*, **127** (16), 400–403.
- Kaldrymidou, H., Leontides, L., Koutinas, A.F., Saridomichelakis, M.N. & Karayannopoulou, M. (2002) Prevalence, distribution and factors associated with the presence and the potential for malignancy of cutaneous neoplasms in 174 dogs admitted to a clinic in Northern Greece. *Journal of Veterinary Medicine. Series A*, **49** (2), 87–91.
- Karbe, E. & Schiefer, B. (1967) Familial amaurotic idiocy in male German Shorthair Pointers. *Pathologia Veterinaria*, **4** (3), 223–232.
- Karlsson, E.K., Sigurdsson, S., Ivansson, E. *et al.* (2013) Genome-wide analyses implicate 33 loci in heritable dog osteosarcoma, including regulatory variants near CDKN2A/B. *Genome Biology*, **14** (12), R132.
- Karmi, N., Brown, E.A., Hughes, S.S. *et al.* (2010a) Estimated frequency of the canine hyperuricosuria mutation in different dog breeds. *Journal of Veterinary Internal Medicine*, **24** (6), 1337–1342.
- Karmi, N., Safra, N., Young, A. & Bannasch, D.L. (2010b) Validation of a urine test and characterization of the putative genetic mutation for hyperuricosuria in Bulldogs and Black Russian Terriers. *American Journal of Veterinary Research*, **71** (8), 909–914.
- Kaswan, R.L. & Salisbury, M.A. (1990) A new perspective on canine keratoconjunctivitis sicca. Treatment with ophthalmic cyclosporine. *Veterinary Clinics of North America: Small Animal Practice*, **20** (3), 583–613.
- Kathmann, I., Cizinauskas, S., Doherr, M. *et al.* (2006) Daily controlled physiotherapy increases survival time in dogs with suspected degenerative myelopathy. *Journal of Veterinary Internal Medicine*, **20** (4), 927–932.
- Kathmann, I., Jaggy, A., Busato, A. *et al.* (1999) Clinical and genetic investigations of idiopathic epilepsy in the Bernese Mountain Dog. *Journal of Small Animal Practice*, **40**, 319–325.
- Kathrani, A., Werling, D. & Allenspach, K. (2011) Canine breeds at high risk of developing inflammatory bowel disease in the south-eastern UK. *Veterinary Record*, **169** (24), 635.
- Kato, K., Sasaki, N., Matsunaga, S., Nishimura, R. & Ogawa, H. (2006) Incidence of canine glaucoma with goniodysplasia in Japan: a retrospective study. *Journal of Veterinary Medical Science*, **68** (8), 853–858.
- Katz, M.L., Khan, S., Awano, T. (2005) A mutation in the CLN8 gene in English Setter dogs with neuronal ceroid-lipofuscinosis. *Biochemical and Biophysical Research Communications*, **327** (2), 541–547.
- Kawabe, M., Mori, T., Ito, Y. *et al.* (2015) Outcomes of dogs undergoing radiotherapy for treatment of oral malignant melanoma: 111 cases (2006–2012). *Journal of the American Veterinary Medical Association*, **247** (10), 1146–1153.
- Kearsley-Fleet, L., O'Neill, D.G., Volk, H.A., Church, D.B. & Brodbelt, D.C. (2013) Prevalence and risk factors for canine epilepsy of unknown origin in the UK. *Veterinary Record*, **172** (13), 338.
- Keller, E.T., MacEwen, E.G., Rosenthal, R.C., Helfand, S.C. & Fox, L.E. (1993) Evaluation of prognostic factors and sequential combination chemotherapy with doxorubicin for canine lymphoma. *Journal of Veterinary Internal Medicine*, **7** (5), 289–295.
- Kelley, J., Sharkey, L.C., Christopherson, P.W. & Rendahl, A. (2014) Platelet count and plateletcrit in Cavalier King Charles Spaniels and Greyhounds using the Advia 120 and 2120. *Veterinary Clinical Pathology*, **43**, 43–49.
- Kemp, C., Thiele, H., Dankof, A. *et al.* (2009) Cleft lip and/or palate with monogenic autosomal recessive transmission in Pyrenees Shepherd Dogs. *Cleft Palate-Craniofacial Journal*, **46** (1), 81–88.
- Kennedy, L.J., Huson, H.J., Leonard, J. *et al.* (2006a) Association of hypothyroid disease in Doberman Pinscher dogs with a rare major histocompatibility complex DLA class II haplotype. *Tissue Antigens*, **67** (1), 53–56.
- Kennedy, L.J., Lunt, M., Barnes, A. *et al.* (2007) Factors influencing the antibody response of dogs vaccinated against rabies. *Vaccine*, **25** (51), 8500–8507.

- Kennedy, L.J., O'Neill, T., House, A. *et al.* (2008) Risk of anal furunculosis in German Shepherd dogs is associated with the major histocompatibility complex. *Tissue Antigens*, **71** (1), 51–56.
- Kennedy, L.J., Quarmby, S., Happ, G.M. *et al.* (2006b) Association of canine hypothyroidism with a common major histocompatibility complex DLA class II allele. *Tissue Antigens*, **68** (1), 82–86.
- Kennel Club (2016). *Dog Health Group Annual Report 2015*. <http://www.thekennelclub.org.uk/for-vets-and-researchers/dog-health-group-annual-report> (accessed October 2017).
- Kent, A.C.C., Constantino-Casas, F., Rusbridge, C. *et al.* (2016) Prevalence of pancreatic, hepatic and renal microscopic lesions in post-mortem samples from Cavalier King Charles Spaniels. *Journal of Small Animal Practice*, **57**, 188–193.
- Kent, M., Glass, E.N. & Haley, A.C. (2014) Ischemic stroke in Greyhounds: 21 cases (2007–2013). *Journal of the American Veterinary Medical Association*, **245** (1), 113–117.
- Kerl, M.E. (2003) Update on canine and feline fungal diseases. *Veterinary Clinics of North America: Small Animal Practice*, **33** (4), 721–747.
- Kerlin, R.L. & Van Winkle, T.J. (1995) Renal dysplasia in Golden Retrievers. *Veterinary Pathology*, **32** (3), 327–329.
- Kern, T. & Riis, R. (1981) Optic nerve hypoplasia in three Miniature Poodles. *Journal of the American Veterinary Medical Association*, **178** (1), 49.
- Kessler, M., Bomhard, D. von, Matis, U., (1997) Canine and feline mast cell tumours: epidemiology, diagnosis and therapy. *Kleintierpraxis*, **42**, 361–374.
- Ketteritzsch, K., Hamann, H., Brahm, R. *et al.* (2004) Genetic analysis of presumed inherited eye diseases in Tibetan Terriers. *Veterinary Journal*, **168** (2), 151–159.
- Khanna, C., Lund, E.M., Raffae, M. & Armstrong, P. J. (1998) Hypomagnesemia in 188 dogs: a hospital population-based prevalence study. *Journal of Veterinary Internal Medicine*, **12**, 304–309.
- KiChang, L., KyungJin, S., HoJung, C. *et al.* (2004) Retrospective study of tracheal collapse: 99 cases diagnosed by radiography in dogs (2002–2003). *Journal of Veterinary Clinics*, **21**, 276–279.
- Kidd, L.B., Salavaggione, O.E., Szumlanski, C.L. *et al.* (2004) Thiopurine methyltransferase activity in red blood cells of dogs. *Journal of Veterinary Internal Medicine*, **18** (2), 214–218.
- Kienle, R.D., Thomas, W.P. & Pion, P.D. (1994) The natural clinical history of canine congenital subaortic stenosis. *Journal of Veterinary Internal Medicine*, **8** (6), 423–431.
- Kijas, J.M., Juneja, R.K., Gäfvert, S. & Anderson, L. (2000) Detection of the causal mutation for canine leukocyte adhesion deficiency (CLAD) using pyrosequencing. *Animal Genetics*, **31** (5), 326–328.s.
- Kijas, J.W., Miller, B.J., Pearce-Kelling, S.E. *et al.* (2003) Canine models of ocular disease: outcross breedings define a dominant disorder present in the English mastiff and bullmastiff breeds. *Journal of Heredity*, **94** (1), 27–30.
- Killingsworth, C.R., Walshaw, R., Dunstan, R.W. *et al.* (1988) Bacterial population and histologic changes in dogs with perianal fistula. *American Journal of Veterinary Research*, **49** (10), 1736–1741.
- Kim, C.-G., Lee, S.-Y., Kim, J.-W. & Park, H.-M. (2013) Assessment of dental abnormalities by full-mouth radiography in small breed dogs. *Journal of the American Animal Hospital Association*, **49** (1), 23–30.
- Kim, H.J., Chang, H.-S., Choi, C.-B. *et al.* (2005) Infiltrative lipoma in cervical bones in a dog. *Journal of Veterinary Medical Science*, **67** (10), 1043–1046.
- Kim, J., Won, H. & Jeong, S. (2009) A retrospective study of ulcerative keratitis in 32 dogs. *International Journal of Applied Research in Veterinary Medicine*, **7** (1/2), 27–31.
- Kim, S.E., Liptak, J.M., Gall, T.T. *et al.* (2007) Epirubicin in the adjuvant treatment of splenic hemangiosarcoma in dogs: 59 cases (1997–2004). *Journal of the American Veterinary Medical Association*, **23** (10), 1550–1557.
- Kimmel, S.E., Waddell, L.S. & Michel, K.E. (2000) Hypomagnesemia and hypocalcemia associated with protein-losing enteropathy in Yorkshire Terriers: five cases (1992–1998). *Journal of the American Veterinary Medical Association*, **217** (5), 703–706.
- Kimmel, S.E., Ward, C.R., Henthorn, P.S. & Hess, R.S. (2002) Familial insulin-dependent diabetes mellitus in Samoyed dogs. *Journal of the American Animal Hospital Association*, **38** (3), 235–238.
- Kimura, T. & Doi, K. (1996) Spontaneous comedones on the skin of hairless descendants of Mexican Hairless Dogs. *Experimental Animals*, **45**, 377–384.

- Kimura, T., Ohshima, S. & Doi, K. (1993) The inheritance and breeding results of hairless descendants of Mexican Hairless Dogs. *Laboratory Animals*, **27**, 55–58.
- Kinzel, S., Hein, S., Buecker, A., Krombach, G.A. & Kuepper, W. (2003) Diagnosis and treatment of arthrosis of cervical articular facet joints in Scottish Deerhounds: 9 cases (1998–2002). *Journal of the American Veterinary Medical Association*, **223** (9), 1311–1315.
- Kirberger, R.M. & Stander, N. (2007) Incidence of canine elbow dysplasia in South Africa. *Journal of the South African Veterinary Association*, **78** (2), 59–62.
- Kirberger, R.M., Steenkamp, G., Spotswood, T.C. *et al.* (2006) Stenotic nasopharyngeal dysgenesis in the dachshund: seven cases (2002–2004). *Journal of the American Animal Hospital Association*, **42**, 290–297.
- Kirsch, J.A., Hauptman, J.G. & Walshaw, R. (2002) A urethropexy technique for surgical treatment of urethral prolapse in the male dog. *Journal of the American Animal Hospital Association*, **38** (4), 381–384.
- Kitaa, J.M.A., Mulei, C.M., Mande, J.D. *et al.* (2014) A retrospective study of canine ehrlichiosis in Kenya. *International Journal of Veterinary Science*, **3** (3), 122–124.
- Kittleson, M.D., Keene, B., Pion, P.D. & Loyer, C.G. (1997) Results of the multicenter spaniel trial (MUST): taurine- and carnitine-responsive dilated cardiomyopathy in American cocker spaniels with decreased plasma taurine concentration. *Journal of Veterinary Internal Medicine*, **11**, 204–211.
- Kjær, D. S., Haaland, M., Bjerkås, E. (2010) Multifocal retinal degeneration in the Border Collie. *Norsk Veterinærtidsskrift*, **122**, 295–300.
- Klein, H.E., Krohne, S.G., Moore, G.E. & Stiles, J. (2011) Postoperative complications and visual outcomes of phacoemulsification in 103 dogs (179 eyes): 2006–2008. *Veterinary Ophthalmology*, **14**, 114–120.
- Kloene, J., Sewell, A.C., Hamann, H., Distl, O. & Tipold, A. (2008) Clinical investigations of seizures in Border Terriers. *Kleintierpraxis*, **53**, 5–12.
- Klopfenstein, M., Howard, J., Rossetti, M. & Geissbühler, U. (2016) Life expectancy and causes of death in Bernese mountain dogs in Switzerland. *BMC Veterinary Research*, **12**, 153.
- Kluger, E.K., Hardman, C., Govendir, M. *et al.* (2009) Triglyceride response following an oral fat tolerance test in Burmese cats, other pedigree cats and domestic crossbred cats. *Journal of Feline Medicine and Surgery*, **11** (2), 82–90.
- Kluth, S., Eckardt, J. & Distl, O. (2014) Selection response to DNA testing for canine ceroid lipofuscinosis in Tibetan Terriers. *Veterinary Journal*, **201** (3), 433–434.
- Knottenbelt, C.M., Addie, D.D., Day, M.J. *et al.* (1999) Determination of the prevalence of feline blood types in the UK. *Journal of Small Animal Practice*, **40** (3), 115–118.
- Knowler, C., Giger, U., Dodds, W.J. *et al.* (1994) Factor XI deficiency in Kerry Blue Terriers. *Journal of the American Veterinary Medical Association*, **205** (11), 1557–1561.
- Koch, S.A. (1972) Cataracts in interrelated Old English Sheepdogs. *Journal of the American Veterinary Medical Association*, **160** (3), 299–301.
- Kogan, D.A., Johnson, L.R., Jandrey, K.E. & Pollard, R.E. (2008) Clinical, clinicopathologic, and radiographic findings in dogs with aspiration pneumonia: 88 cases (2004–2006). *Journal of the American Veterinary Medical Association*, **233** (11), 1742–1747.
- Kohn, B., Goldschmidt, M.H., Hohenhaus, A.E. *et al.* (2000) Anemia, splenomegaly, and increased osmotic fragility of erythrocytes in Abyssinian and Somali cats. *Journal of the American Veterinary Medical Association*, **217** (10), 1483–1491.
- Kohyama, M., Tada, N., Mitsui, H. *et al.* (2016) Real-time PCR genotyping assay for canine progressive rod-cone degeneration and mutant allele frequency in Toy Poodles, Chihuahuas and Miniature Dachshunds in Japan. *Journal of Veterinary Medical Science*, **78**, 481–484.
- Kolbjørnsen, O., Press, C.M. & Landsverk, T. (1994) Gastropathies in the Lundehund. I. Gastritis and gastric neoplasia associated with intestinal lymphangiectasia. *Acta Pathologica, Microbiologica et Immunologica Scandinavica*, **102** (9), 647–661.
- Kolicheski, A., Johnson, G.S., O'Brien, D.P. *et al.* (2016) Australian Cattle Dogs with neuronal ceroid lipofuscinosis are homozygous for a CLN5 nonsense mutation previously identified in Border Collies. *Journal of Veterinary Internal Medicine*, **30**, 1149–1158.

- Kornegay, J.N., Goodwin, M.A. & Spyridakis, L.K. (1987) Hypomyelination in Weimaraner dogs. *Acta Neuropathologica*, **72** (4), 394–401.
- Kortz, G.D., Meier, W.A., Higgins, R.J. *et al.* (1997) Neuronal vacuolation and spinocerebellar degeneration in young Rottweiler dogs. *Veterinary Pathology*, **34** (4), 296–302.
- Koskinen, L.L.E., Seppälä, E.H., Belanger, J.M. *et al.* (2015) Identification of a common risk haplotype for canine idiopathic epilepsy in the ADAM23 gene. *BMC Genomics*, **16** (1), 465.
- Kraijer-Huwer, I.M.G., Gubbels, E.J., Scholten, J. *et al.* (2008) Characterization and prevalence of cataracts in Labrador Retrievers in the Netherlands. *American Journal of Veterinary Research*, **69** (10), 1336–1340.
- Kramer, J.W., Davis, W.C. & Prieur, D.J. (1977) The Chediak–Higashi syndrome of cats. *Laboratory Investigation*, **36** (5), 554–562.
- Kramer, J.W., Klassen, J.H., Baskin, D.G. *et al.* (1988) Inheritance of diabetes mellitus in Keshond dogs. *American Journal of Veterinary Research*, **4** (3), 428–431.
- Kramer, J.W., Nottingham, S., Robinette, J. *et al.* (1980) Inherited, early onset, insulin-requiring diabetes mellitus of Keeshond dogs. *Diabetes*, **29** (7), 558–565.
- Kramer, J.W., Schiffer, S.P., Sande, R.D., Rantanen, N.W. & Whitener, E.K. (1982) Characterization of heritable thoracic hemivertebra of the German Shorthaired Pointer. *Journal of the American Veterinary Medical Association*, **181** (8), 814–815.
- Kramer, J.W., Venta, P.J., Klein, S.R. *et al.* (2004) A von Willebrand's factor genomic nucleotide variant and polymerase chain reaction diagnostic test associated with inheritable type-2 von Willebrand's disease in a line of german shorthaired pointer dogs. *Veterinary Pathology*, **41** (3), 221–228.
- Kranenburg, H.C., Voorhout, G., Grinwis, G.C., Hazewinkel, H.A. & Meij, B.P. (2011) Diffuse idiopathic skeletal hyperostosis (DISH) and spondylosis deformans in purebred dogs: a retrospective radiographic study. *Veterinary Journal*, **190** (2), e84–e90.
- Kraus, C., Pavard, S. & Promislow, D.E.L. (2013) The size-life span trade-off decomposed: why large dogs die young. *American Naturalist*, **181** (4), 492–505.
- Krawiec, D.R. & Heflin, D. (1992) Study of prostatic disease in dogs: 177 cases (1981–1986). *Journal of the American Veterinary Medical Association*, **200** (8), 1119–1122.
- Krebs, I.A., Lindsley, S., Shaver, S. & MacPhail, C. (2014) Short- and long-term outcome of dogs following surgical correction of a persistent right aortic arch. *Journal of the American Animal Hospital Association*, **50** (3), 181–186.
- Krecny, M., Tichy, A., Rushton, J. & Nell, B. (2015) A retrospective survey of ocular abnormalities in pugs: 130 cases. *Journal of Small Animal Practice*, **56** (2), 96–102.
- Kreutzer, R., Leeb, T., Müller, G. *et al.* (2005) A duplication in the canine β -galactosidase gene *GLB1* causes exon skipping and GM₁-gangliosidosis in Alaskan huskies. *Genetics*, **170** (4), 1857–1861.
- Kristiansen, E., Revold, T., Lingaas, F. *et al.* (2017) Cataracts in the Norwegian Buhund: current prevalence and characteristics. *Veterinary Ophthalmology*, **20** (5), 460–467.
- Krontveit, R.I. & Moe, L. (2013) A prospective study of risk factors for and long-term effects of radiological canine hip dysplasia in four breeds. *Norsk Veterinærtidsskrift*, **125** (5), 259–268.
- Krontveit, R.I., Nødtvedt, A., Sævik, B.K. *et al.* (2010) A prospective study on canine hip dysplasia and growth in a cohort of four large breeds in Norway (1998–2001). *Preventative Veterinary Medicine*, **97** (3–4), 252–263.
- Kropatsch, R., Akkad, D.A., Frank, M. *et al.* (2016) A large deletion in RPGR causes XLPRA in Weimaraner dogs. *Canine Genetics and Epidemiology*, **3** (1), 7.
- Kropatsch, R., Petrasch-Prwez, E., Seelow, D. *et al.* (2010) Generalized progressive retinal atrophy in the Irish Glen of Imaal Terrier is associated with a deletion in the ADAM9 gene. *Molecular and Cellular Probes*, **24** (6), 357–363.
- Krotscheck, V., Adin, C.A., Hunt, G.B. *et al.* (2007) Epidemiologic factors associated with the anatomic location of intrahepatic portosystemic shunts in dogs. *Veterinary Surgery*, **36** (1), 31–36.
- Kruth, S.A., Feldman, E.C. & Kennedy, P.C. (1982) Insulin-secreting islet tumours: establishing a diagnosis and the clinical course for 25 days. *Journal of the American Veterinary Medical Association*, **181** (1), 54–58.

- Ksiazek, M., Gruszczynska, J. & Grzegorzolka, B. (2014) Problems in cooperation with breeders and dog owners on the example of entropion frequency estimation in polish hounds. *Acta Scientiarum Polonorum. Zootechnica*, **13** (4).
- Kubai, M.A., Bentley, E., Miller, P.E., Mutti, D.O. & Murphy, C.J. (2008) Refractive states of eyes and association between ametropia and breed in dogs. *American Journal of Veterinary Research*, **69** (7), 946–951.
- Kučera, J. (2007) Urate urolithiasis in dogs: a retrospective study. *Veterinářství*, **57**, 142–148.
- Kuehner, K.A., Marks, S.L., Kass, P.H. *et al.* (2011) *Tritrichomonas foetus* infection in purebred cats in Germany: prevalence of clinical signs and the role of co-infection with other enteroparasites. *Journal of Feline Medicine and Surgery*, **13** (4), 251–258.
- Kuhl, K.A., Shofer, F.S. & Goldschmidt, M.H. (1994) Comparative histopathology of pemphigus foliaceus and superficial folliculitis in the dog. *Veterinary Pathology*, **31** (1), 19–27.
- Kuiper, J.D., van der Gaag, I. & Mouwen, J.M.V.M. (1982) Polyodontia and abnormal forms of teeth in dogs. *Tijdschrift voor Diergeneeskunde*, **107**, 451–457.
- Kull, P.A., Hess, R.S., Craig, L.E., Saunders, H.M. & Washabau, R.J. (2001) Clinical, clinicopathologic, radiographic, and ultrasonographic characteristics of intestinal lymphangiectasia in dogs: 17 cases (1996–1998). *Journal of the American Veterinary Medical Association*, **219** (2), 197–202.
- Kumar, K.S., Kumar, V.V.V.A., Nagaraj, P. *et al.* (2011) Idiopathic pericardial effusion in dogs: a three-year study. *VetScan*, **6** (1), 78.
- Kunkle, G.A., White, S.D., Calderwood-Mays, M. *et al.* (1993) Focal metatarsal fistulas in 5 dogs. *Journal of the American Veterinary Medical Association*, **202**, 756–757.
- Kuster, N., Hässig, M. & Spiess, B. (2011) [Hereditary eye disease in the Entlebucher Mountain Dog in Switzerland: a retrospective study from 1999 to 2009]. *Schweizer Archiv für Tierheilkunde*, **153** (6), 269–275.
- Kutsunai, M., Kanemoto, H., Fukushima, K. *et al.* (2014) The association between gall bladder mucocoeles and hyperlipidaemia in dogs: a retrospective case control study. *Veterinary Journal*, **199** (1), 76–79.
- Kuwamura, M., Adachi, T., Yamate, J. *et al.* (2002) Necrotising encephalitis in the Yorkshire terrier: a case report and literature review. *Journal of Small Animal Practice*, **43** (10), 459–463.
- Kuznetsova, E., Bettenay, S., Nikolaeva, L., Majzoub, M. & Mueller, R. (2012) Influence of systemic antibiotics on the treatment of dogs with generalized demodicosis. *Veterinary Parasitology*, **188** (1–2), 148–155.
- Kyöstilä, K., Cizinauskas, S., Seppala, E.H. *et al.* (2012) A SEL1L mutation links a canine progressive early-onset cerebellar ataxia to the endoplasmic reticulum-associated protein degradation (ERAD) machinery. *PLoS Genetics*, **8** (6), e1002759.
- Kyöstilä, K., Lappalainen, A.K. & Lohi, H. (2013) Canine chondrodysplasia caused by a truncating mutation in collagen-binding integrin alpha subunit 10. *PLOS One*, **8**, 9 (2013): e75621.
- Labelle, A.L., Dresser, C.B., Hamor, R.E., Allender, M.C. & Disney, J.L. (2013) Characteristics of, prevalence of, and risk factors for corneal pigmentation (pigmentary keratopathy) in Pugs. *Journal of the American Veterinary Medical Association*, **243** (5), 667–674.
- Lachaume, P., Hitte, C., Jouquand, S. *et al.* (1998) Identification and analysis of the dog keratin 9 (KRT9) gene. *Animal Genetics*, **29**, 173–177.
- Laffort-Dassot, C., Beco, L. & Carloti, D.N. (2002) Follicular dysplasia in five Weimaraners. *Veterinary Dermatology*, **13** (5), 253–260.
- LaFond, E., Breur, G.J. & Austin, C.C. (2002) Breed susceptibility for developmental orthopedic diseases in dogs. *Journal of the American Veterinary Medical Association*, **38** (5), 467–477.
- Lamb, C.R., Wikeley, H., Boswood, A. *et al.* (2001) Use of breed-specific ranges for the vertebral heart scale as an aid to the radiographic diagnosis of cardiac disease in dogs. *Veterinary Record*, **148** (23), 707–711.
- Landsverk, T. & Gamlem, H. (1984) Intestinal lymphangiectasia in the Lundehund. Scanning electron microscopy of intestinal mucosa. *Acta Pathologica, Microbiologica et Immunologica Scandinavica*, **92** (5), 353–362.

- Langenbach, A., Anderson, M., Dambach, D., Sorenmo, K. & Shofer, F. (1998) Extraskelatal osteosarcomas in dogs: a retrospective study of 169 cases (1986–1996). *Journal of the American Animal Hospital Association*, **34** (2), 113–120.
- Lannek, E.B. & Miller, P.E. (2001) Development of glaucoma after phacoemulsification for removal of cataracts in dogs: 22 cases (1987–1997). *Journal of the American Veterinary Medical Association*, **218**, 70–76.
- Lappalainen, A.K., Hyvärinen, T., Junnila, J. & Laitinen-Vapaavuori, O. (2016) Radiographic evaluation of elbow incongruity in Skye terriers. *Journal of Small Animal Practice*, **57** (2), 96–99.
- Lara-Garcia, A., Couto, C.G. & Iazbik, M.C. (2008) Postoperative bleeding in retired Racing Greyhounds. *Journal of Veterinary Internal Medicine*, **22**, 525–533.
- Laratta, L.J., Riis, R.C., Kern, T.J. *et al.* (1985) Multiple congenital ocular defects in the Akita dog. *Cornell Veterinarian*, **75** (3), 381–392.
- Larsson, M. (1986) The breed, sex and age distribution of dogs with primary hypothyroidism. *Svensk Veterinärtidning*, **38**, 181–183.
- Lavoué, R., Trumel, C., Smets, P.M.Y. *et al.* (2015) Characterization of proteinuria in Dogue de Bordeaux Dogs, a breed predisposed to a familial glomerulonephropathy: a retrospective study. *PLoS One*, **10** (7), e0133311.
- Lavoué, R., van der Lugt, J.J., Day, M.J. *et al.* (2010) Progressive juvenile glomerulonephropathy in 16 related French Mastiff (Bordeaux) dogs. *Journal of Veterinary Internal Medicine*, **24** (2), 314–322.
- Lavrijsen, I.C.M., Heuven, H.C.M., Voorhout, G. (2012) Phenotypic and genetic evaluation of elbow dysplasia in Dutch Labrador Retrievers, Golden Retrievers, and Bernese Mountain dogs. *Veterinary Journal*, **193** (2), 486–492.
- Lavrijsen, I.C.M., Leegwater, P.A.J. *et al.* (2014) Genome wide analysis indicates genes for basement membrane and cartilage matrix proteins as candidates for hip dysplasia in Labrador Retrievers. *PLoS One*, **9** (1), e87735.
- Lawler, D.F., Keltner, D.G., Hoffman, W.E. *et al.* (1996) Benign familial hyperphosphatasemia in Siberian huskies. *American Journal of Veterinary Research*, **57** (5), 612–617.
- Lawrence, Y.A., Ruaux, C.G., Nemanic, S. & Milovancev, M. (2015) Characterization, treatment, and outcome of bacterial cholecystitis and bactibilia in dogs. *Journal of the American Veterinary Medical Association*, **246**, 982–989.
- Laws, E.J., Harcourt-Brown, T.R., Granger, N. & Rose, J.H. (2017) An exploratory study into factors influencing development of acute canine polyradiculoneuritis in the UK. *Journal of Small Animal Practice*, **58** (8), 437–443.
- Lawson, D.D. (1973) Canine distichiasis. *Journal of Small Animal Practice*, **14** (8), 469–478.
- Lazarus, J.A., Pickett, J.P. & Champagne, E.S. (1998) Primary lens luxation in the Chinese Shar Pei: clinical and hereditary characteristics. *Veterinary Ophthalmology*, **1** (2–3), 101–107.
- Lederer, R., Bennett, D. & Parkin, T. (2014) Survey of tail injuries sustained by working gundogs and terriers in Scotland. *Veterinary Record*, **174**, 451–451.
- Lee, A., Kathrani, A., Priestnall, S.L. *et al.* (2015) Lack of correlation between mucosal immunoglobulin A-positive plasma cell numbers and TLR5 genotypes in German shepherd Dogs with idiopathic chronic enteropathy. *Journal of Comparative Pathology*, **2–3**, 201–205.
- Lee, F.F., Bradley, C.W., Cain, C.L. *et al.* (2016) Localized parakeratotic hyperkeratosis in sixteen Boston Terrier dogs. *Veterinary Dermatology*, **27**, 384–e96.
- Lee, S., Lee, S., Moon, H. *et al.* (2007) Isolation, characterization and genetic analysis of canine GATA4 gene in a family of Doberman Pinschers with an atrial septal defect. *Journal of Genetics*, **86**, 241–247.
- Lees, G.E., Hilman, R.G., Kashtan, C.E. *et al.* (1998) A model of autosomal recessive Alport syndrome in English Cocker Spaniel dogs. *Kidney International*, **54** (3), 706–719.
- Lefebvre, F. (2013) Epidemiology of canine parvovirus infection in the USA: an update. *Veterinary Focus*, **23** (2), 37–38.
- Legge, C.H., López, A., Hanna, P. *et al.* (2013) Histological characterization of dilated cardiomyopathy in the juvenile toy Manchester Terrier. *Veterinary Pathology*, **50** (6), 1043–1052.
- Leidinger, E., Freeman, K., Kirtz, G., Hooijberg, E. & Sick, K. (2014) Breed related odds ratio and anatomic distribution of canine mast cell tumours in Austria. *Tierärztliche Praxis Kleintiere*, **42** (6), 367–373.

- Leipold, H.W., Huston, K., Blauch, B. *et al.* (1974) Congenital defects on the caudal vertebral column and spinal cord in Manx cats. *Journal of the American Veterinary Medical Association*, **164** (5), 520–523.
- Leisewitz, A.L., Spenser, J.A., Jacobsen, L.S. & Schroeder, H. (1997) Suspected primary immunodeficiency syndrome in three related Irish Wolfhounds. *Journal of Small Animal Practice*, **38** (5), 209–212.
- Lekcharoensuk, C., Lulich, J.P., Osborne, C.A. *et al.* (2000a) Patient and environmental factors associated with calcium oxalate urolithiasis in dogs. *Journal of the American Veterinary Medical Association*, **217** (4), 515–519.
- Lekcharoensuk, C., Lulich, J.P., Osborne, C.A. *et al.* (2000b) Association between patient-related factors and risk of calcium oxalate and magnesium ammonium phosphate urolithiasis in cats. *Journal of the American Veterinary Medical Association*, **217** (4), 520–525.
- Lekcharoensuk, C., Osborne, C.A. & Lulich, J.P. (2001) Epidemiologic study of risk factors for lower urinary tract diseases in cats. *Journal of the American Veterinary Medical Association*, **218** (9), 1429–1435.
- Lem, K.Y., Fosgate, G.T., Norby, B. & Steiner, J.M. (2008) Associations between dietary factors and pancreatitis in dogs. *Journal of the American Veterinary Medical Association*, **233** (9), 1425–1431.
- Lenz, J.A., Furrow, E., Craig, L.E. & Cannon, C.M. (2017) Histiocytic sarcoma in 14 Miniature Schnauzers: a new breed predisposition? *Journal of Small Animal Practice*, **58** (8), 461–467.
- Leppänen, M., Martenson, J. & Mäki, K. (2001) Results of ophthalmologic screening examinations of German Pinschers in Finland: a retrospective study. *Veterinary Ophthalmology*, **4** (3), 165–169.
- Leppänen, M. & Saloniemi, H. (1998) Screening and controlling canine inherited ocular diseases in Finland: epidemiological, economical and health promotional aspect. *Veterinary Ophthalmology*, **1** (4), 203–210.
- Leppänen, M. & Saloniemi, H. (1999) Controlling canine hip dysplasia in Finland. *Preventative Veterinary Medicine*, **42** (2), 121–131.
- Lessa, T.B., de Abreu, D.K., Rodrigues, M.N. *et al.* (2014) Morphological and ultrastructural evaluation of the Golden Retriever muscular dystrophy trachea, lungs, and diaphragm muscle. *Microscopy Research and Technique*, **77** (11), 857–861.
- Levine, G.J., Levine, J.M., Walker, M.A. *et al.* (2006) Evaluation of the association between spondylosis deformans and clinical signs of intervertebral disk disease in dogs: 172 cases (1999–2000). *Journal of the American Veterinary Medical Association*, **228**, 96–100.
- Levine, J.M., Fosgate, G.T., Porter, B., Schatzberg, S.J. & Greer, K. (2008) Epidemiology of necrotizing meningoencephalitis in Pug dogs. *Journal of Veterinary Internal Medicine*, **22** (4), 961–968.
- Levy, I., Hall, C., Trentacosta, N., Percival, M. (2009) A preliminary retrospective survey of injuries occurring in dogs participating in canine agility. *Veterinary and Comparative Orthopaedics and Traumatology*, **22**, 321–324.
- Lewis, D.D., Shelton, G.D., Piras, A. *et al.* (1997) Gracilis or semitendinosus myopathy in 18 dogs. *Journal of the American Animal Hospital Association*, **33**, 177–188.
- Lewis, D.G. (1989) Cervical spondylomyelopathy ('wobbler' syndrome) in the dog: a study based on 224 cases. *Journal of Small Animal Practice*, **30** (12), 657–665.
- Lewis, D.G., Kelly, D.F. & Sansom, J. (1986) Congenital microphthalmia and other developmental ocular anomalies in the Dobermann. *Journal of Small Animal Practice*, **27** (9), 559–566.
- Lewis, L.D. & Morris, M.L. (1984) Canine urolithiasis: diagnosis and treatment. *Modern Veterinary Practice*, **65** (5), 375–378.
- Lewis, T., Rusbridge, C., Knowler, P., Blott, S. and Woolliams, J. A. (2010) Heritability of syringomyelia in Cavalier King Charles Spaniels. *Veterinary Journal*, **183**, 345–347.
- Lewis, T., Swift, S., Woolliams, J. A. and Blott, S. (2011a) Heritability of premature mitral valve disease in Cavalier King Charles Spaniels. *Veterinary Journal*, **188**, 73–76.
- Lewis, T.W., Ilksa, J.J., Blott, S.C. & Woolliams, J.A. (2011b) Genetic evaluation of elbow scores and the relationship with hip scores in UK Labrador Retrievers. *Veterinary Journal*, **189** (2), 227–233.
- Li, A., Bennett, D., Gibbs, C. *et al.* (2000) Radial carpal bone fractures in 15 dogs. *Journal of Small Animal Practice*, **41**, 74–79.
- Li, F.-Y., Cuddon, P.A., Song, J. *et al.* (2006) Canine spongiform leukoencephalomyelopathy is associated with a missense mutation in cytochrome b. *Neurobiology of Disease*, **21** (1), 35–42.

- Li, Y., Wu, D.-D., Boyko, A.R. *et al.* (2014) Population variation revealed high-altitude adaptation of Tibetan Mastiffs. *Molecular Biology and Evolution*, **31** (5), 1200–1205.
- Liao, A.T., Chu, P.-Y., Yeh, L.-S. *et al.* (2009) A 12-year retrospective study of canine testicular tumors. *Journal of Veterinary Medical Science*, **71** (7), 919–923.
- Licht, B.G., Licht, M.H., Harper, K.M. *et al.* (2002) Clinical presentations of naturally occurring canine seizures: similarities to human seizures. *Epilepsy and Behaviour*, **3** (5), 460–470.
- Lightfoot, R.M., Cabral, L., Gooch, L. *et al.* (1996) Retinal pigment epithelial dystrophy in Briard dogs. *Research in Veterinary Science*, **60** (1), 17–23.
- Lilliehöök, I., Gunnarsson, L., Zakrisson, G. & Tvedten, H. (2000) Diseases associated with pronounced eosinophilia: a study of 105 dogs in Sweden. *Journal of Small Animal Practice*, **41** (6), 248–253.
- Lind, D.L., Fidel, J.L., Gay, J.M. & Mealey, K.L. (2013) Evaluation of vincristine-associated myelosuppression in Border Collies. *American Journal of Veterinary Research*, **74**, 257–261.
- Linde Forsberg, C. & Persson, G. (2007) A survey of dystocia in the Boxer breed, Acta Veterinaria Scandinavica. *BioMed Central*, **49**, 8.
- Linek, M., Boss, C., Haemmerling, R. *et al.* (2005) Effects of cyclosporine A on clinical and histologic abnormalities in dogs with sebaceous adenitis. *Journal of the American Veterinary Medical Association*, **226** (1), 59–64.
- Ling, G.V., Franti, C.E., Johnson, D.L. & Ruby, A.L. (1998a) Urolithiasis in dogs. IV: Survey of interrelations among breed, mineral composition, and anatomic location of calculi, and presence of urinary tract infection. *American Journal of Veterinary Research*, **59**, 650–660.
- Ling, G.V., Franti, C.E., Ruby, A.L. & Johnson, D.L. (1998b) Urolithiasis in dogs. II: Breed prevalence, and interrelations of breed, sex, age, and mineral composition. *American Journal of Veterinary Research*, **59**, 630–642.
- Ling, G.V., Ruby, A.L., Johnson, D.L., Thurmond, M. & Franti, C.E. (1998c) Renal calculi in dogs and cats: prevalence, mineral type, breed, age, and gender interrelationships (1981–93). *Journal of Veterinary Internal Medicine*, **12** (1), 11–21.
- Ling, G.V., Stabenfeldt, G.H., Comer, K.M. *et al.* (1979) Canine hyperadrenocorticism: pretreatment clinical and laboratory evaluation of 117 cases. *Journal of the American Veterinary Medical Association*, **174** (2), 159–163.
- Ling, G.V., Thurmond, M.C., Choi, Y.K. *et al.* (2003) Changes in proportion of canine urinary calculi composed of calcium oxalate or struvite in specimens analyzed from 1981 through 2001. *Journal of Veterinary Internal Medicine*, **17** (6), 817–823.
- Lingaas, F., Aarskaug, T., Sletten, M. *et al.* (1998) Genetic markers linked to neuronal ceroid lipofuscinosis in English Setter dogs. *Animal Genetics*, **29** (5), 371–376.
- Linton, M., Nimmo, J.S., Norris, J.M. *et al.* (2015) Feline gastrointestinal eosinophilic sclerosing fibroplasia: 13 cases and review of an emerging clinical entity. *Journal of Feline Medicine and Surgery*, **17** (5), 392–404.
- Lippmann, T., Jonkisz, A., Dobosz, T. *et al.* (2007) Haplotype-defined linkage region for gPRA in Schapendoes dogs. *Molecular Vision*, **13**, 174–180.
- Lipsitz, D., Berry, J.L. & Shelton, G.D. (1999) Inherited predisposition to myasthenia gravis in Newfoundlands. *Journal of the American Veterinary Medical Association*, **215**, 956–958.
- Littlewood, J.D. (1989) Inherited bleeding disorders of dogs and cats. *Journal of Small Animal Practice*, **30** (3), 140–143.
- Littman, M.P. (2013) Lyme nephritis. *Journal of Veterinary Emergency and Critical Care*, **23** (2), 163–173.
- Littman, M.P., Dambach, D.M., Vaden, S.L. & Giger, U. (2000) Familial protein-losing enteropathy and protein-losing nephropathy in Soft Coated Wheaten Terriers: 222 cases (1983–1997). *Journal of Veterinary Internal Medicine*, **14** (1), 68–80.
- Littman, M.P., Wiley, C.A., Raducha, M.G. & Henthorn, P.S. (2013) Glomerulopathy and mutations in NPHS1 and KIRREL2 in Soft-Coated Wheaten Terrier dogs. *Mammalian Genome*, **24** (3), 119–126.
- Liu, N.-C., Sargan, D.R., Adams, V.J. *et al.* (2015) Characterisation of brachycephalic obstructive airway syndrome in French Bulldogs using whole body barometric plethysmography. *PLoS One*, **10** (6), e0130741.

- Liu, S.K. & Tilley, L.P. (1975) Malformation of the canine mitral valve complex. *Journal of the American Veterinary Medical Association*, **167** (6), 465–471.
- Liu, S.K., Tilley, L.P., Tappe, J.P. & Fox, P.R. (1986) Clinical and pathologic findings in dogs with atherosclerosis: 21 cases (1970–1983). *Journal of the American Veterinary Medical Association*, **189** (2), 227–232.
- Lium, B. & Moe, L. (1985) Hereditary multifocal renal cystadenocarcinomas and nodular dermatofibrosis in the German Shepherd Dog: macroscopic and histopathologic changes. *Veterinary Pathology*, **22** (5), 447–455.
- Lobetti, R. (2000) Common variable immunodeficiency in Miniature Dachshunds affected with *Pneumocystis carinii* pneumonia. *Journal of Veterinary Diagnostic Investigation*, **12**, 39–45.
- Lobetti, R.G., Leisewitz, A. & Spencer, J.A. (1996) *Pneumocystis carinii* in the Miniature Dachsund: case report and literature review. *Journal of Small Animal Practice*, **37** (6), 280–285.
- Loder, R.T. & Todhunter, R.J. (2017) Demographics of hip dysplasia in the Maine Coon cat. *Journal of Feline Medicine and Surgery*. doi: 10.1177/1098612X17705554 [Epub ahead of print].
- Long, S.E. & Crispin, S.M. (1999) Inheritance of multifocal retinal dysplasia in the Golden Retriever in the UK. *Veterinary Record*, **145** (24), 702–704.
- Lorinson, D. & Bright, R.M. (1998) Long-term outcome of medical and surgical treatment of hiatal hernias in dogs and cats: 27 cases (1978–1996). *Journal of the American Veterinary Medical Association*, **213**, 381–384.
- Lortz, J., Favrot, C., Mecklenburg, L. *et al.* (2010) A multicentre placebo-controlled clinical trial on the efficacy of oral ciclosporin A in the treatment of canine idiopathic sebaceous adenitis in comparison with conventional topical treatment. *Veterinary Dermatology*, **21** (6), 593–601.
- Lothrup, C.D. (1988) Pathophysiology of canine growth-hormone responsive alopecia. *Compendium on Continuing Education for the Practicing Veterinarian*, **10** (12), 1346–1349.
- Louly, C.C.B., Soares, S.F., da Nóbrega Silveira, D., Guimarães, M.S. & Borges, L.M.F. (2010) Differences in the behavior of *Rhipicephalus sanguineus* tested against resistant and susceptible dogs. *Experimental and Applied Acarology*, **51** (4), 353–362.
- Louly, C.C.B., Soares, S.F., Silveira, D.N. *et al.* (2009) Differences in the susceptibility of two breeds of dogs, English Cocker Spaniel and Beagle, to *Rhipicephalus sanguineus* (Acari: Ixodidae). *International Journal of Acarology*, **35**, 25–32.
- Loures, F.H. & Conceição, L. G. (2009) Nevi and cutaneous hamartomas in dogs: retrospective clinical and epidemiologic study of 81 cases, *Ciência Rural*, **39**, 2527–2532.
- Louwerens, M., London, C.A., Pedersen, N.C. *et al.* (2005) Feline lymphoma in the post-feline leukaemia virus era. *Journal of Veterinary Internal Medicine*, **19** (3), 329–335.
- Low, W.W., Uhl, J.M., Kass, P.H., Ruby, A.L. & Westropp, J.L. (2010) Evaluation of trends in urolith composition and characteristics of dogs with urolithiasis: 25,499 cases (1985–2006). *Journal of the American Veterinary Medical Association*, **236** (2), 193–200.
- Lowe, J.K., Kukekova, A.V., Kirkness, E.F. *et al.* (2003) Linkage mapping of the primary disease locus for collie eye anomaly. *Genomics*, **82** (1), 86–95.
- Lowrie, M. & Garosi, L. (2016) Natural history of canine paroxysmal movement disorders in Labrador retrievers and Jack Russell terriers. *Veterinary Journal*, **213**, 33–37.
- Lowrie, M., Bessant, C., Harvey, R.J. *et al.* (2015a) Audiogenic reflex seizures in cats. *Journal of Feline Medicine and Surgery*, **18** (4), 328–336.
- Lowrie, M., Penderis, J., McLaughlin, M., Eckersall, P.D. & Anderson, T.J. (2009) Steroid responsive meningitis-arteritis: A prospective study of potential disease markers, prednisolone treatment, and long-term outcome in 20 dogs (2006–2008). *Journal of Veterinary Internal Medicine*, **23** (4), 862–870.
- Lowrie, M., Smith, P. M., Keuster, T. de & Garosi, L. (2015b) Trance-like syndrome in Bull Terriers. *Veterinary Record*, **177**, 223.
- Luckschander, N., Allenspach, K., Hall, J. *et al.* (2006) Perinuclear antineutrophilic cytoplasmic antibody and response to treatment in diarrheic dogs with food responsive disease or inflammatory bowel disease. *Journal of Veterinary Internal Medicine*, **20** (2), 221–227.
- Lulich, J.P., Osborne, C.A., Thumchai, R. *et al.* (1999) Epidemiology of canine calcium oxalate uroliths. Identifying risk factors. *Veterinary Clinics of North America: Small Animal Practice*, **29** (1), 113–122.

- Lund, E. (2012) Epidemiology of periodontal disease in older cats. *Veterinary Focus*, **22** (3), 23–24.
- Lund, E.M., Armstrong, P.J., Kirk, C.A. & Klausner, J.S. (2006) Prevalence and risk factors for obesity in adult dogs from private US veterinary practices. *International Journal of Applied Research in Veterinary Medicine*, **4** (2), 177–186.
- Lund, J.D., Agger, J.F. & Vestergaard, K.S. (1996) Reported behaviour problems in pet dogs in Denmark: age distribution and influence of breed and gender. *Preventive Veterinary Medicine*, **28** (1), 33–48.
- Lurie, D.M., Lucrov, M.D., Griffey, S.M. *et al.* (2004) T-cell-derived malignant lymphoma in the Boxer breed. *Veterinary and Comparative Oncology*, **2** (3), 171–175.
- Lutz, S., Sewell, A.C., Bigler, B. *et al.* (2012) Serum cobalamin, urine methylmalonic acid, and plasma total homocysteine concentrations in Border Collies and dogs of other breeds. *American Journal of Veterinary Research*, **73**, 1194–1199.
- Lutz, S., Sewell, A.C., Reusch, C.E. and Kook, P.H. (2013) Clinical and laboratory findings in Border Collies with presumed hereditary juvenile cobalamin deficiency. *Journal of the American Animal Hospital Association*, **49**, 197–203.
- Lyons, L.A., Erdman, C.A., Grahn, R.A. *et al.* (2016a) Aristaless-like Homeobox protein 1 (ALX1) variant associated with craniofacial structure and frontonasal dysplasia in Burmese cats. *Developmental Biology*, **409** (2), 451–458.
- Lyons, L.A., Grahn, R.A., Genova, F. *et al.* (2016b) Mucopolysaccharidosis VI in cats: clarification regarding genetic testing. *BMC Veterinary Research*, **12** (1), 136.
- MacDonald, K.A., Cagney, O. & Magne, M.L. (2009) Echocardiographic and clinicopathologic characterization of pericardial effusion in dogs: 107 cases (1985–2006). *Journal of the American Veterinary Medical Association*, **235** (12), 1456–1461.
- Macedo, A.S., Casagrande, A.J., Friesen, R. & Minto, B.W. (2015) Craniomandibular osteopathy in a West Highland White Terrier. *Acta Scientiae Veterinariae*, **43** (2), 1–4.
- Mackenzie, G., Barnhart, M., Kennedy, S., DeHoff, W. & Schertel, E. (2010) A retrospective study of factors influencing survival following surgery for gastric dilatation-volvulus syndrome in 306 dogs. *Journal of the American Animal Hospital Association*, **46** (2), 97–102.
- MacLachlan, N.J., Breitschwerdt, E.B., Chambers, J.M., Argenzio, R.A. & De Buysscher, E.V. (1988) Gastroenteritis of Basenji Dogs. *Veterinary Pathology*, **25**, 36–41.
- MacMillan, A.D. & Lipton, D.E. (1978) Heritability of multifocal retinal dysplasia in American Cocker Spaniels. *Journal of the American Veterinary Medical Association*, **172** (5), 568–572.
- MacMillan, A.D., Waring, G.O., Spangler, W.L. & Roth, A.M. (1979) Crystalline corneal opacities in the Siberian Husky. *Journal of the American Veterinary Medical Association*, **175** (8), 829–832.
- Macready, D.M., Johnson, L.R. & Pollard, R.E. (2007) Fluoroscopic and radiographic evaluation of tracheal collapse in dogs: 62 cases (2001–2006). *Journal of the American Veterinary Medical Association*, **230** (12), 1870–1876.
- MacVean, D.W., Monlux, A.W., Anderson, P.S., Silberg, S.L. & Roszel, J.F. (1978) Frequency of canine and feline tumors in a defined population. *Veterinary Pathology*, **15** (6), 700–715.
- Maddox, T.W., May, C., Keeley, B.J. & McConnell, J.F. (2013) Comparison between shoulder computed tomography and clinical findings in 89 dogs presented for thoracic limb lameness. *Veterinary Radiology and Ultrasound*, **54**, 358–364.
- Maddux, J.M., Edwards, D.F., Barnhill, M.A. & Sanders, W.L. (1991) Neutrophil function in dogs with congenital ciliary dyskinesia. *Veterinary Pathology*, **28**, 347–353.
- Maggio-Price, L., Emerson, C.L., Hinds, T.R., Vincenzi, F.F., Hammond, W.R. (1988) Hereditary non-spherocytic hemolytic anemia in Beagles. *American Journal of Veterinary Research*, **49**, 1020–1025.
- Mahaffey, M.B., Yarbrough, K.M. & Munnell, J.F. (1978) Focal loss of pigment in the Belgian Tervuren Dog. *Journal of the American Veterinary Medical Association*, **173**, 390–396.
- Mahony, O.M., Knowles, K.E., Braund, K.G., Averill, D.R. & Frimberger, A.E. (1998) Laryngeal paralysis-polyneuropathy complex in young Rottweilers. *Journal of Veterinary Internal Medicine*, **12** (5), 330–337.
- Mai, W., Seiler, G.S., Lindl-Bylicki, B.J. & Zwingenberger, A.L. (2015) CT AND MRI features of carotid body paragangliomas in 16 dogs. *Veterinary Radiology and Ultrasound*, **56**, 374–383.

- Malik, R. & Church, D.B. (1988) Congenital mitral insufficiency in Bull Terriers. *Journal of Small Animal Practice*, **29**, 549–557.
- Malik, R., Dill-Macky, E., Martin, P. *et al.* (1995) Cryptococcosis in dogs: a retrospective study of 20 consecutive cases. *Journal of Medical Mycology*, **33** (5), 291–297.
- Malik, R., Foster, S.F., Martin, P. *et al.* (2002) Acute febrile neutrophilic vasculitis of the skin of young Shar-Pei dogs. *Australian Veterinary Journal*, **80** (4), 200–206.
- Malik, R., Griffin, D.L., White, J.D. *et al.* (2005) The prevalence of feline A/B blood types in the Sydney region. *Australian Veterinary Journal*, **83**, 38–44.
- Malik, R., Musca, N.G., Gunew, M.N. *et al.* (2015) Periodic hypokalaemic polymyopathy in Burmese and closely related cats: a review including the latest genetic data. *Journal of Feline Medicine and Surgery*, **17** (5), 417–426.
- Malm, S., Fikse, W.F., Danell, B. & Strandberg, E. (2008) Genetic variation and genetic trends in hip and elbow dysplasia in Swedish Rottweiler and Bernese Mountain Dog. *Journal of Animal Breeding and Genetics*, **125** (6), 403–412.
- Manchester, A.C., Hill, S., Sabatino, B. *et al.* (2013) Association between granulomatous colitis in French Bulldogs and invasive *Escherichia coli* and response to fluoroquinolone antimicrobials. *Journal of Veterinary Internal Medicine*, **27** (1), 56–61.
- Mandigers, P.F., Van Nes, J.J., Knol, B.W. *et al.* (1993) Hereditary necrotizing myelopathy in Kooiker dogs. *Research in Veterinary Science*, **54**, 118–123.
- Mandigers, P.J.J., van den Ingh, T.S., Bode, P. *et al.* (2004) Association between liver copper concentration and subclinical hepatitis in Doberman Pinschers. *Journal of Veterinary Internal Medicine*, **18** (5), 647–650.
- Mankowska, M., Krzeminska, P., Graczyk, M. *et al.* (2017) Confirmation that a deletion in the POMC gene is associated with body weight of Labrador Retriever dogs. *Research in Veterinary Science*, **112**, 116–118.
- Mao, J., Xia, Z., Chen, J. & Yu, J. (2013) Prevalence and risk factors for canine obesity surveyed in veterinary practices in Beijing, China. *Preventive Veterinary Medicine*, **112** (3–4), 438–442.
- Marcellin-Little, D.J., De Young, D.J., Ferris, K.K. *et al.* (1994) Incomplete ossification of the humeral condyle in Spaniels. *Veterinary Surgery*, **23**, 475–487.
- March, P.A., Coates, J.R., Abyad, R.J. *et al.* (2009) Degenerative myelopathy in 18 Pembroke Welsh Corgi dogs. *Veterinary Pathology Online*, **46** (2), 241–250.
- Marchant, T.W., Johnson, E.J., McTeir, L. *et al.* (2017) Canine brachycephaly is associated with a retrotransposon-mediated missplicing of SMOC2. *Current Biology*, **27** (11), 1573–1584.e6.
- Marin, L.M., Brown, J., McBrien, C. *et al.* (2007) Vertebral heart size in retired racing Greyhounds. *Veterinary Radiology and Ultrasound*, **48** (4), 332–334.
- Marmor, M., Willeberg, P., Glickman, L.T. *et al.* (1982) Epizootiologic patterns of diabetes mellitus in dogs. *American Journal of Veterinary Research*, **43** (3), 465–470.
- Marolf, A., Blaik, M. & Specht, A. (2007) A retrospective study of the relationship between tracheal collapse and bronchiectasis in dogs. *Veterinary Radiology and Ultrasound*, **48** (3), 199–203.
- Martin, C. & Leach, R. (1970) Everted membrana nictitans in German Shorthaired Pointers. *Journal of the American Veterinary Medical Association*, **157**, 1229–1232.
- Martin, M.W., Stafford Johnson, M.J. & Celona, B. (2009) Canine dilated cardiomyopathy: a retrospective study of signalment, presentation and clinical findings in 369 cases. *Journal of Small Animal Practice*, **50** (1), 23–29.
- Martin, M.W., Stafford Johnson, M.J., Strehlau, G. & King, J.N. (2010) Canine dilated cardiomyopathy: a retrospective study of prognostic findings in 367 clinical cases. *Journal of Small Animal Practice*, **51** (8), 428–436.
- Martin, P.T., Shelton, G.D., Dickinson, P.J. *et al.* (2008) Muscular dystrophy associated with alpha-dystroglycan deficiency in Sphynx and Devon Rex cats. *Neuromuscular disorders*, **18** (12), 942–952.
- Martinez, J.T., Rogers, L.K., Kellogg, C. *et al.* (2016) Plasma vasoprotective eicosanoid concentrations in healthy Greyhounds and non-Greyhound dogs. *Journal of Veterinary Internal Medicine*, **30** (2), 583–590.

- Martinez, S., Fajardo, R., Valdés, J. *et al.* (2007) Histopathologic study of long-bone growth plates confirms the Basset Hound as an osteochondrodysplastic breed. *Canadian Journal of Veterinary Research*, **71**, 66–69.
- Martinez, S., Valdes, J. & Alonso, R.A. (2000) Achondroplastic dog breeds have no mutations in the transmembrane domain of the FGFR-3 gene. *Canadian Journal of Veterinary Research*, **64**, 243–245.
- Martinez-Ruzafa, I., Kruger, J.M., Miller, R. *et al.* (2012) Clinical features and risk factors for development of urinary tract infections in cats. *Journal of Feline Medicine and Surgery*, **14** (10), 729–740.
- Martinoli, S., Neilssen, P. & White, R.A.S. (2014) The outcome of combined urethropexy and colposuspension for management of bitches with urinary incontinence associated with urethral sphincter mechanism incompetence. *Veterinary Surgery*, **43** (1), 52–57.
- Martin-Vaquero, P. & da Costa, R.C. (2014) Evaluation of traditional and novel radiographic vertebral ratios in Great Danes with versus without cervical spondylomyelopathy. *Veterinary Radiology and Ultrasound*, **55** (5), 488–495.
- Martins-Bessa, A., Cardoso, L., Costa, T. *et al.* (2015) Reproductive emergencies in the bitch: a retrospective study. *Journal of the Hellenic Veterinary Medical Society*, **66**, 231–240.
- März, I., Wilkie, L.J., Harrington, N. *et al.* (2015) Familial cardiomyopathy in Norwegian Forest cats. *Journal of Feline Medicine and Surgery*, **17** (8), 681–691.
- Mason, I.S. & Jones, J. (1989) Juvenile cellulitis in Gordon Setters. *Veterinary Record*, **124** (24), 642.
- Mason, T.A. (1976) Persistent pupillary membrane in the Basenji. *Australian Veterinary Journal*, **52** (8), 343–344.
- Massey, J., Rothwell, S., Rusbridge, C. *et al.* (2013) Association of an MHC class II haplotype with increased risk of polymyositis in Hungarian Vizsla dogs. *PLoS One*, **8** (2), e56490.
- Massey, J., Short, A.D., Catchpole, B. (2014) Genetics of canine anal furunculosis in the German Shepherd Dog. *Immunogenetics*, **66** (5), 311–324.
- Masuda, K., Hashizume, C., Kikusui, T., Takeuchi, Y. & Mori, Y. (2004) Breed differences in genotype and allele frequency of catechol o-methyltransferase gene polymorphic regions in dogs. *Journal of Veterinary Medical Science*, **66** (2), 183–187.
- Matic, S.E. (1988) Congenital heart disease in the dog. *Journal of Small Animal Practice*, **29**, 743–759.
- Mattin, M.J., Boswood, A., Church, D.B. *et al.* (2015a) Degenerative mitral valve disease: Survival of dogs attending primary-care practice in England. *Preventive Veterinary Medicine*, **122**, 436–442.
- Mattin, M.J., Boswood, A., Church, D.B. *et al.* (2015b) Prevalence of and risk factors for degenerative mitral valve disease in dogs attending primary-care veterinary practices in England. *Journal of Veterinary Internal Medicine*, **29** (3), 847–854.
- Mattin, M., O'Neill, D., Church, D. *et al.* (2014) An epidemiological study of diabetes mellitus in dogs attending first opinion practice in the UK. *Veterinary Record*, **174** (14), 349.
- Mauldin, E.A., Credille, K.M., Dunstan, R.W. *et al.* (2008) The clinical and morphological features of nonepidermolytic ichthyosis in the golden retriever. *Veterinary Pathology*, **45**, 174–180.
- Mauldin, E.A., Palmeiro, B.S., Goldschmidt, M.H. *et al.* (2006) Comparison of clinical history and dermatologic findings in 29 dogs with severe eosinophilic dermatitis: a retrospective analysis. *Veterinary Dermatology*, **17** (5), 338–347.
- Mauldin, E.A., Scott, D.W., Miller, W.H. & Smith, C.A. (1997) *Malassezia* dermatitis in the dog: a retrospective histopathological and immunopathological study of 86 cases (1990–95). *Veterinary Dermatology*, **8** (3), 191–202.
- Mauldin, E.A., Wang, P., Evans, E. *et al.* (2015) Autosomal recessive congenital ichthyosis in American Bulldogs is associated with NIPAL4 (ICHTHYIN) deficiency. *Veterinary Pathology*, **52**, 654–662.
- Mauler, D.A., De Decker, S., De Risio, L. *et al.* (2014) Signalment, clinical presentation, and diagnostic findings in 122 dogs with spinal arachnoid diverticula. *Journal of Veterinary Internal Medicine*, **28** (1), 175–181.
- Maurer, M., Mary, J., Guillaud, L. *et al.* (2012) Centronuclear myopathy in Labrador Retrievers: a recent founder mutation in the PTPLA gene has rapidly disseminated worldwide. *PLoS One*, **7** (10), e46408.
- Mausberg, E.M., Leeb, T., Dolf, G. *et al.* (2007) Evaluation of the CTS2L gene as a candidate gene for alopecia X in Pomeranians and Keeshonden. *Animal Biotechnology*, **18** (4), 291–296.

- Mauser, A.E., Whitlark, J., Whitney, K.M. *et al.* (1996) A deletion mutation causes hemophilia B in Lhasa Apso dogs. *Blood*, **88**, 3451–3455.
- Mayhew, P.D., Bush, W.W. & Glass, E.N. (2002) Trigeminal neuropathy in dogs: a retrospective study of 29 cases (1991–2000). *Journal of the American Animal Hospital Association*, **38** (3), 262–270.
- Mayhew, P.D., McLearn, R.C., Ziemer, S.L. *et al.* (2004) Risk factors for recurrence of clinical signs associated with thoracolumbar intervertebral disk herniation in dogs: 229 cases (1994–2000). *Journal of the American Veterinary Medical Association*, **225**, 1231–1236.
- Mayousse, V., Desquilbet, L., Jeandel, A. & Blot, S. (2017) Prevalence of neurological disorders in French Bulldog: a retrospective study of 343 cases (2002–2016). *BMC Veterinary Research*, **13** (1), 212.
- Mazaki-Tovi, M., Baneth, G., Aroch, I. *et al.* (2002) Canine spirocercosis: clinical, diagnostic, pathologic, and epidemiologic characteristics. *Veterinary Parasitology*, **107** (3), 235–250.
- Mazrier, H., Vogelnest, L.J., Thomson, P.C. *et al.* (2016) Canine atopic dermatitis: breed risk in Australia and evidence for a susceptible clade. *Veterinary Dermatology*, **27** (3), 167–e42.
- Mazzucchelli, S., Vaillant, M.D., Wéverberg, F. *et al.* (2012) Retrospective study of 155 cases of prolapse of the nictitating membrane gland in dogs. *Veterinary Record*, **170** (17), 443.
- McAlees, T.J. (2010) Immune-mediated haemolytic anaemia in 110 dogs in Victoria, Australia. *Australian Veterinary Journal*, **88** (1–2), 25–28.
- McCue, J., Langston, C., Palma, D. *et al.* (2009) Urate urolithiasis. *Compendium on Continuing Education for the Practicing Veterinarian*, **31** (10), 468–475.
- McCutcheon, K.A. & Fleming, S.J. (2001/2002) Grief resulting from euthanasia and natural death of companion animals. *Journal of Death and Dying*, **44** (2), 169–188.
- McEwan, N.A., Huang, H.P. & Mellor, D.J. (2003) Immunoglobulin levels in Bull Terriers suffering from lethal acrodermatitis. *Veterinary Immunology and Immunopathology*, **96**, 235–238.
- McEwan, N.A., McNeil, P.E., Thompson, H. & McCandush, I.A.P. (2000) Diagnostic features, confirmation and disease progression in 28 cases of lethal acrodermatitis of Bull Terriers. *Journal of Small Animal Practice*, **41**, 501–507.
- McGraw, R.A. & Carmichael, K.P. (2006) Molecular basis of globoid cell leukodystrophy in Irish Setters. *Veterinary Journal*, **171**, 370–372.
- McGrotty, Y.L., Ramsey, I.K. & Knottenbelt, C.M. (2003) Diagnosis and management of hepatic copper accumulation in a Skye Terrier. *Journal of Small Animal Practice*, **44** (2), 85–89.
- McGwin, G. (2010) Causation in epidemiology. *American Journal of Ophthalmology*, **150** (5), 599–601.
- McKay, L.W., Seguin, M.A., Ritchey, J.W. & Levy, J.K. (2004) Juvenile nephropathy in two related Pembroke Welsh Corgi puppies. *Journal of Small Animal Practice*, **45** (11), 568–571.
- McKeever, P.J., Torres, S.M.F. & O'Brien, T.D. (1992) Spiculosis. *Journal of the American Animal Hospital Association*, **28** (3), 257–262.
- McLellan, G.J., Aquino, S., Sigle, K. *et al.* (2004) Congenital glaucoma and microphakia in Siamese cats. *Veterinary Ophthalmology*, **7** (6), 428.
- McLellan, G.J., Elks, R., Lybaert, P. *et al.* (2002) Vitamin E deficiency in dogs with retinal pigment epithelial dystrophy. *Veterinary Record*, **151** (22), 663–667.
- McLellan, G.J. & Teixeira, L.B.C. (2015) Feline glaucoma. *Veterinary Clinics of North America: Small Animal Practice*, **45** (6), 1307–1333.
- McNiel, E.A., Prink, A.L. & O'Brien, T.D. (2006) Evaluation of risk and clinical outcome of mast cell tumours in Pug dogs. *Veterinary and Comparative Oncology*, **4** (1), 2–8.
- Mealey, K.L. & Meurs, K.M. (2008) Breed distribution of the ABCB1–1Δ (multidrug sensitivity) polymorphism among dogs undergoing ABCB1 genotyping. *Journal of the American Veterinary Medical Association*, **233** (6), 921–924.
- Mealey, K.L., Minch, J.D., White, S.N., Snekvik, K.R. & Mattoon, J.S. (2010) An insertion mutation in ABCB4 is associated with gallbladder mucocele formation in dogs. *Comparative Hepatology*, **9** (1), 6.
- Mealey, K.L., Munyard, K.A. & Bentien, S.A. (2005) Frequency of the mutant *MDR1* allele associated with multidrug sensitivity in a sample of herding breed dogs living in Australia. *Veterinary Parasitology*, **131**, 193–196.

- Meek, K., Klenker, L., Dallas, C. (2001) SCID in Jack Russell Terriers: a new animal model of DNA-PKcs deficiency. *Journal of Immunology*, **167** (4), 2142–2150.
- Melián, C. & Peterson, M.E. (1996) Diagnosis and treatment of naturally occurring hypoadrenocorticism in 42 dogs. *Journal of Small Animal Practice*, **37** (6), 268–275.
- Mellanby, R.J. & Herrtage, M.E. (2005) Long-term survival of 23 dogs with pericardial effusions. *Veterinary Record*, **156** (18), 568–571.
- Mellema, M. S. & Hoareau, G. L. (2014) Hypomagnesemia in brachycephalic dogs. *Journal of Veterinary Internal Medicine*, **28** (5), 1418–1423.
- Mellersh, C.S., Graves, K.T., McLoughlin, B. *et al.* (2007) Mutation in HSF4 associated with early but not late-onset hereditary cataract in the Boston Terrier. *Journal of Heredity*, **98** (5), 531–533.
- Mellersh, C.S., Pettitt, L., Forman, O.P., Vaudin, M. & Barnett, K.C. (2006) Identification of mutations in HSF4 in dogs of three different breeds with hereditary cataracts. *Veterinary Ophthalmology*, **9** (5), 369–378.
- Melville, K., Smith, K.C. & Dobromyiskij, M.J. (2015) Feline cutaneous mast cell tumours: a UK-based study comparing signalment and histological features with long-term outcomes. *Journal of Feline Medicine and Surgery*, **17** (6), 486–493.
- Melville, S.A., Wilson, C.L., Chiang, C.S. *et al.* (2005) A mutation in canine CLN5 causes neuronal ceroid lipofuscinosis in Border collie dogs. *Genomics*, **86** (3), 287–294.
- Menaut, P., Bélanger, M.C., Beauchamp, G., Ponzio, N.M. & Moise, N.S. (2005) Atrial fibrillation in dogs with and without structural or functional cardiac disease: a retrospective study of 109 cases. *Journal of Veterinary Cardiology*, **7** (2), 75–83.
- Menotti-Raymond, M., David, V.A., Pflueger, S. *et al.* (2010) Widespread retinal degenerative disease mutation (rdAc) discovered among a large number of popular cat breeds. *Veterinary Journal*, **186**, 32–38.
- Menzel, J. & Distl, O. (2011) Unusual vascular ring anomaly associated with a persistent right aortic arch and an aberrant left subclavian artery in German Pinschers. *Veterinary Journal*, **187** (3), 352–355.
- Meola, S.D. (2013) Brachycephalic airway syndrome. *Topics in Companion Animal Medicine*, **28** (3), 91–96.
- Merlini, N.B., Guberman, Ú.C., Gandolfi, M.G. *et al.* (2014) Retrospective study of 71 cases of prolapsed third eyelid gland (2009–2013). *Arquivos de Ciências Veterinárias e Zoologia da UNIPAR*, **17**, 175–178.
- Merola, V., Khan, S. & Gwaitney-Brant, S. (2009) Ivermectin toxicosis in dogs: a retrospective study. *Journal of the American Animal Hospital Association*, **45** (3), 106–111.
- Merveille, A.C., Battaille, G., Billem, F. *et al.* (2014) Clinical findings and prevalence of the mutation associated with primary ciliary dyskinesia in Old English Sheepdogs. *Journal of Veterinary Internal Medicine*, **28** (3), 771–778.
- Mesa-Sánchez, I., Zaldivar-López, S., Couto, C.G. *et al.* (2012) Haematological, blood gas and acid-base values in the Galgo Español (Spanish greyhound). *Journal of Small Animal Practice*, **53** (7), 398–403.
- Metzger, J., Wöhlke, A., Mischke, R. *et al.* (2015) A novel SLC27A4 splice acceptor site mutation in Great Danes with ichthyosis. *PLoS One*, **10** (10), e0141514.
- Meurs, K.M., Stern, J.A., Sisson, D.D. *et al.* (2013) Association of dilated cardiomyopathy with the striatin mutation genotype in Boxer Dogs. *Journal of Veterinary Internal Medicine*, **27**, 1437–1440.
- Meurs, K.M. (2010) Genetics of cardiac disease in the small animal patient. *Veterinary Clinics of North America: Small Animal Practice*, **40** (4), 701–715.
- Meurs, K.M., Miller, M.W. & Wright, N.A. (2001) Clinical features of dilated cardiomyopathy in Great Danes and results of a pedigree analysis: 17 cases (1990–2000). *Journal of the American Veterinary Medical Association*, **218**, 729–732.
- Meurs, K.M., Norgard, M.M., Kuan, M., Haggstrom, J. & Kittleson, M. (2009) Analysis of 8 sarcomeric candidate genes for feline hypertrophic cardiomyopathy mutations in cats with hypertrophic cardiomyopathy. *Journal of Veterinary Internal Medicine*, **23** (4), 840–843.
- Meyer-Lindenberg, A., Langhann, A., Fehr, M. & Nolte, I. (2002) Prevalence of fragmented medial coronoid process of the ulna in lame adult dogs. *Veterinary Record*, **151** (8), 230–234.
- Meyers-Wallen, V.N. & Patterson, D.F. (1988) XX sex reversal in the American cocker spaniel dog: phenotypic expression and inheritance. *Human Genetics*, **80** (1), 23–30.

- Meyers-Wallen, V.N., Bowman, L., Acland, G.M. *et al.* (1995) Sry-Negative XX sex reversal in the German Shorthaired Pointer dog. *Journal of Heredity*, **86** (5), 369–374.
- Mhlanga-Mutangadura, T., Johnson, G.S., Schnabel, R.D. *et al.* (2016) A mutation in the Warburg syndrome gene, RAB3GAP1, causes a similar syndrome with polyneuropathy and neuronal vacuolation in Black Russian Terrier dogs. *Neurobiology of Disease*, **86**, 75–85.
- Michalska, Z. & Michalski, Z. (1997) Prevalence of neoplasms in animals in Wrocław and Lower Silesia in 1976–1995. *Medycyna Weterynaryjna*, **53** (5), 263–267.
- Michell, A.R. (1999) Longevity of British breeds of dog and its relationships with sex, size, cardiovascular variables and disease. *Veterinary Record*, **145** (22), 625–629.
- Michelsen, J. (2013) Canine elbow dysplasia: Aetiopathogenesis and current treatment recommendations. *Veterinary Journal*, **196** (1), 12–19.
- Micun, J., Sobczak-Filipiak, M., Winnicka, A. *et al.* (2009) Thrombocytopenia as a characteristic trait in the Polish Ogar dog. *Polish Journal of Veterinary Sciences*, **12** (4), 523.
- Millán, Y., Mascort, J., Blanco, A., Costa, C. *et al.* (2010) Hypomyelination in three Weimaraner dogs. *Journal of Small Animal Practice*, **51** (11), 594–598.
- Miller, D.M. (1995) The occurrence of mast cell tumors in young Shar-Peis. *Journal of Veterinary Diagnostic Investigation*, **7** (3), 360–363.
- Miller, L.M., Hegreberg, G.A., Prieur, D.J. *et al.* (1984) Inheritance of congenital myasthenia gravis in Smooth Fox Terrier dogs. *Journal of Heredity*, **75**, 163–166.
- Miller, L.M., Lennon, V.A., Lambert, E.H. *et al.* (1983) Congenital myasthenia gravis in 13 Smooth Fox Terriers. *Journal of the American Veterinary Medical Association*, **182** (7), 694–697.
- Miller, M.A. & Dunstan, R.W. (1993) Seasonal flank alopecia in boxers and Airedale Terriers: 24 cases (1985–1992). *Journal of the American Veterinary Medical Association*, **203**, 1567.
- Miller, M.A., Nelson, S.L. & Turk, J.R. *et al.* (1991) Cutaneous neoplasia in 340 cats. *Veterinary Pathology*, **28** (5), 389–395.
- Miller, P.E., Miller, L.M. & Schoster, J.V. (1990) Feline blastomycosis: a report of three cases and literature review (1961–1988). *Journal of the American Animal Hospital Association*, **26** (4), 417–424.
- Miller, S.A., Hohenhaus, A.E. & Hale, A.S. (2004) Case-control study of blood type, breed, sex, and bacteremia in dogs with immune-mediated hemolytic anemia. *Journal of American Veterinary Medical Association*, **224**, 232–235.
- Miller, W.H. (1990a) Colour dilution alopecia in Doberman Pinschers with blue or fawn coat colours: a study on the incidence and histopathology of this disorder. *Veterinary Dermatology*, **1**, 113.
- Miller, W.H. (1990b) Follicular dysplasia in adult black and red Doberman pinschers. *Veterinary Dermatology*, **1**, 181.
- Miller, W.H. & Scott, D.W. (1995) Follicular dysplasia of the Portuguese Water Dog. *Veterinary Dermatology*, **6** (2), 67–74.
- Miller, W.H., Wellington, J.R. & Scott, D.W. (1992) Dermatologic disorders of Chinese Shar Peis: 58 cases (1981–1989). *Journal of the American Veterinary Medical Association*, **200** (7), 986–990.
- Millichamp, N.J., Curtis, R. & Barnett, K.C. (1988) Progressive retinal atrophy in Tibetan Terriers. *Journal of the American Veterinary Medical Association*, **192** (6), 769–776.
- Millis, D.L., Hauptman, J.G. & Johnson, C.A. (1992) Cryptorchidism and monorchism in cats: 25 cases (1980–1989). *Journal of the American Veterinary Medical Association*, **200** (8), 1128–1130.
- Mills, J.N., Day, M.J. & Shaw, S.E. *et al.* (1985) Autoimmune haemolytic anaemia in dogs. *Australian Veterinary Journal*, **62** (4), 121–123.
- Milne, K.L. & Hayes, H.M. (1981) Epidemiologic features of canine hypothyroidism. *Cornell veterinarian*, **71** (1), 3–14.
- Minkus, G., Breuer, W., Wanke, R. *et al.* (1994) Familial nephropathy in Bernese Mountain dogs. *Veterinary Pathology*, **31** (4), 421–428.
- Minor, K.M., Patterson, E.E., Keating, M.K. *et al.* (2011) Presence and impact of the exercise-induced collapse associated DNM1 mutation in Labrador Retrievers and other breeds. *Veterinary Journal*, **189** (2), 214–219.
- Miranda, S., Roura, X., Picado, A., Ferrer, L. & Ramis, A. (2008) Characterization of sex, age, and breed for a population of canine leishmaniosis diseased dogs. *Research in Veterinary Science*, **85** (1), 35–38.

- Mircean, V., Mircean, M., Gavrea, R. & Cozma, V. (2008) Epidemiological aspects of otitis externa in dogs. *Lucrari Stiintifice*, **41**, 427–436.
- Mircean, V., Titilincu, A., Mircean, M. & Magdaş, C. (2010) *Malassezia pachydermatis* infection in dogs: a retrospective of its epidemiology, clinical and cytological results in Cluj, Romania. *Scientia Parasitologica*, **11** (3), 153–163.
- Mischke, R., Kühnlein, P., Kehl, A. *et al.* (2011) G244E in the canine factor IX gene leads to severe haemophilia B in Rhodesian Ridgebacks. *Veterinary Journal* **187** (1) 113–118.
- Misdrop, W. & Hart, A.A. (1979) Some prognostic and epidemiologic factors in canine osteosarcoma. *Journal of the National Cancer Institute*, **62** (3), 537–545.
- Mitchell, A.L. & Pearce, S.H.S. (2012) Autoimmune Addison disease: pathophysiology and genetic complexity. *Nature Reviews Endocrinology*, **8** (5), 306–316.
- Miyama, T., Sakata, Y., Shimada, Y. *et al.* (2005) Epidemiological survey of Babesia gibsoni infection in dogs in eastern Japan. *Journal of Veterinary Medical Science*, **67** (5), 467–471.
- Mizukami, K., Chang, H.-S., Ota, M. *et al.* (2012) Collie eye anomaly in Hokkaido dogs: case study. *Veterinary Ophthalmology*, **15** (2), 128–132.
- Mizukami, K., Chang, H.-S., Yabuki, A. *et al.* (2011) Novel rapid genotyping assays for neuronal ceroid lipofuscinosis in Border Collie dogs and high frequency of the mutant allele in Japan. *Journal of Veterinary Diagnostic Investigation*, **23**, 1131–1139.
- Mizukami, K., Yabuki, A., Kawamichi, T. *et al.* (2013) Real-time PCR genotyping assay for canine trapped neutrophil syndrome and high frequency of the mutant allele in Border Collies. *Veterinary Journal*, **195**, 260–261.
- Mochizuki, H., Motsinger-Reif, A., Bettini, C., Moroff, S. & Breen, M. (2016) Association of breed and histopathological grade in canine mast cell tumours. *Veterinary and Comparative Oncology*, **15** (3), 829–839.
- Moe, L. (2001) Population-based incidence of mammary tumours in some dog breeds. *Journal of Reproduction and Fertility, Supplement*, **57**, 439–443.
- Moe, L., Gamlem, H., Dahl, K. & Glatte, E. (2008) Canine neoplasia: population-based incidence of vascular tumours. *APMIS. Supplementum*, **(125)**, 63–68.
- Moesser, C.F. & Wade, C.M. (2017) Relationship between transitional lumbosacral vertebrae and eight lumbar vertebrae in a breeding colony of Labrador Retrievers and Labrador Crosses. *Australian Veterinary Journal*, **95** (1), 33–36.
- Moise, N.S., Wiedenkiller, D., Yeager, A.E. *et al.* (1989) Clinical, radiographic, and bronchial cytologic features of cats with bronchial disease: 65 cases (1980–1986). *Journal of the American Veterinary Medical Association*, **194** (10), 1467–1473.
- Moneva-Jordan, A., Corcoran, B.M., French, A. *et al.* (2001) Sick sinus syndrome in nine West Highland White Terriers. *Veterinary Record*, **148** (5), 142–147.
- Monobe, M.M., Araújo, J.P., Lunsford, K.V., Silva, R. & Bulla, C. (2015) Frequency of the MDR1 mutant allele associated with multidrug sensitivity in dogs from Brazil. *Veterinary Medicine: Research and Reports*, **6**, 111–117.
- Monteiro, R., Adams, V., Keys, D. & Platt, S. R. (2012) Canine idiopathic epilepsy: prevalence, risk factors and outcome associated with cluster seizures and status epilepticus. *Journal of Small Animal Practice*, **53**, 526–530.
- Montgomery, D.L. (1982) Hereditary striatonigral and cerebello-olivary degeneration of the Kerry Blue Terrier. *Dissertation Abstracts International B*, **42** (7), 2732.
- Montgomery, K.W., van der Woerd, A. & Cottrill, N.B. (2008) Acute blindness in dogs: sudden acquired retinal degeneration syndrome versus neurological disease (140 cases, 2000–2006). *Veterinary Ophthalmology*, **11** (5), 314–320.
- Montgomery, R.D., Milton, J.L., Hathcock, J.T. *et al.* (1994) Osteochondritis dissecans of the canine tarsal joint. *Compendium on Continuing Education for the Practicing Veterinarian*, **16** (7), 835–845.
- Montgomery, R.D., Milton, J.L., Henderson, R.A. *et al.* (1989) Osteochondritis dissecans of the canine stifle. *Compendium on Continuing Education for the Practicing Veterinarian*, **11** (10), 1199–1205.
- Montoya, J.A., Morales, M., Ferrer, O., Molina, J.M. & Corbera, J.A. (1998) The prevalence of *Dirofilaria immitis* in Gran Canaria, Canary Islands, Spain (1994–1996). *Veterinary Parasitology*, **75** (2–3), 221–226.

- Montoya, J.A., Morales, M., Juste, M.C., Bañares, A., Simon, F. & Genchi, C. (2006) Seroprevalence of canine heartworm disease (*Dirofilaria immitis*) on Tenerife Island: an epidemiological update. *Parasitology Research*, **100** (1), 103–105.
- Moody, J.A., Famula, T.R. & Sampson, R.C. (2005) Identification of microsatellite markers linked to progressive retinal atrophy in American Eskimo Dogs. *American Journal of Veterinary Research*, **66** (11), 1900–1902.
- Moon, P.F., Erb, H.N., Ludders, J.W., Gleed, R.D. & Pascoe, P.J. (1998) Perioperative management and mortality rates of dogs undergoing cesarean section in the United States and Canada. *Journal of the American Veterinary Medical Association*, **213**, 365–369.
- Moon-Fanelli, A.A. & Dodman, N.H. (1998) Description and development of compulsive tail chasing in terriers and response to clomipramine treatment. *Journal of the American Veterinary Medical Association*, **212**, 1252–1257.
- Moon-Fanelli, A.A., Dodman, N.H. & Cottam, N. (2007) Blanket and flank sucking in Doberman Pinschers. *Journal of the American Veterinary Medical Association*, **23**, 907–912.
- Moore, F.M. & Thornton, G.W. (1983) Telangiectasia of Pembroke Welsh Corgi dogs. *Veterinary Pathology*, **20** (2), 203–208.
- Moore, G.E., Guptill, L.F., Ward, M.P., Glickman, N.W., Faunt, K.K., Lewis, H.B. & Glickman, L.T. (2005) Adverse events diagnosed within three days of vaccine administration in dogs. *Journal of the American Veterinary Medical Association*, **227** (7), 1102–1108.
- Moore, P.F. (1984) Systemic histiocytosis of Bernese Mountain Dogs. *Veterinary Pathology*, **21** (6), 554–563.
- Moore, P.F. & Rosin, A. (1986) Malignant histiocytosis of Bernese Mountain Dogs. *Veterinary Pathology*, **23** (1), 1–10.
- Morales, M., Ynaraja, E. & Montoya, J.A. (2001) Dilated cardiomyopathy in Presa Canario Dogs: ECG findings. *Journal of Veterinary Medicine. Series A*, **48** (10), 577–580.
- Morgan, J.P., Med, V., Ackerman, N., Bailey, C.S. & Pool, R.R. (1980) Vertebral tumors in the dog: A clinical radiologic, and pathologic study of 61 primary and secondary lesions. *Veterinary Radiology*, **21** (5), 197–212.
- Morgan, J.P., Wind, A. & Davidson, A.P. (1999) Bone dysplasias in the Labrador Retriever: a radiographic study. *Journal of the American Animal Hospital Association*, **35** (4), 332–340.
- Morgan, R.V. (1994) Feline corneal sequestration: a retrospective study of 42 cases (1987–1991). *Journal of the American Animal Hospital Association*, **30** (1), 24–28.
- Morgan, R.V., Duddy, J.M. & McClurg, K. (1993) Prolapse of the gland of the third eyelid in dogs: a retrospective study of 89 cases (1980–1990). *Journal of the American Animal Hospital Association*, **29** (1), 56–60.
- Mori, N., Lee, P., Muranaka, S. *et al.* (2010) Predisposition for primary hyperlipidemia in Miniature Schnauzers and Shetland Sheepdogs as compared to other canine breeds. *Research in Veterinary Science*, **88** (3), 394–399.
- Moriello, K.A. (2004) A new look at ulcerative dermatosis of Shetland Sheepdogs and Rough Collies. *Veterinary Medicine*, **99**, 1003–1004.
- Morita, T., Shimada, A., Takeuchi, T. *et al.* (2002) Cliniconeuropathologic findings of familial frontal lobe epilepsy in Shetland Sheepdogs. *Canadian Journal of Veterinary Research = Revue Canadienne de Recherche Veterinaire*, **66** (1), 35–41.
- Morris, J.S., McInnes, E.F., Bostock, D.E. *et al.* (2002) Immunohistochemical and histopathologic features of 14 malignant fibrous histiocytomas from Flat-Coated Retrievers. *Veterinary Pathology*, **39** (4), 473–479.
- Morrison, W.B., Wilsman, N.J., Fox, L.E. & Farnum, C.E. (1987) Primary ciliary dyskinesia in the dog. *Journal of Veterinary Internal Medicine*, **1** (2), 67–74.
- Morrow, M., Ottobre, J., Ottobre, A. *et al.* (2015) Breed-dependent differences in the onset of fear-related avoidance behavior in puppies. *Journal of Veterinary Behavior: Clinical Applications and Research*, **10**, 286–294.
- Mortari, A.C., Rahal, S.C., Vulcano, L.C., da Silva, V.C. & Volpi, R.S. (2009) Use of radiographic measurements in the evaluation of dogs with medial patellar luxation. *Canadian Veterinary Journal*, **50** (10), 1064–1068.

- Mortier, F., Strohmeier, K., Hartmann, K. *et al.* (2015) Acute haemorrhagic diarrhoea syndrome in dogs: 108 cases. *Veterinary Record*, **176** (24), 627.
- Morton, L.D., Sanecki, R.K., Gordon, D.E. *et al.* (1990) Juvenile renal disease in Miniature Schnauzer dogs. *Veterinary Pathology*, **27** (6), 455–458.
- Mosher, D.S., Quignon, P., Bustamante, C.D. *et al.* (2007) A mutation in the myostatin gene increases muscle mass and enhances racing performance in heterozygote dogs. *PLoS Genetics*, **3** (5), e79.
- Mueller, R.S., Krebs, I., Power, H.T. & Fieseler, K.V. (2006) Pemphigus foliaceus in 91 dogs. *Journal of the American Animal Hospital Association*, **42** (3), 189–196.
- Mueller, R.S., Meyer, D., Bensignor, E. & Sauter-Louis, C. (2009) Treatment of canine generalized demodicosis with a 'spot-on' formulation containing 10% moxidectin and 2.5% imidacloprid (Advocate®, Bayer Healthcare). *Veterinary Dermatology*, **20** (5–6), 441–446.
- Muldoon, L.L., Neuwelt, E.A., Pagel, M.A. *et al.* (1994) Characterization of the molecular defect in a feline model for type II GM2-gangliosidosis (Sandhoff disease). *American Journal of Pathology*, **144** (5), 1109–1118.
- Multari, D., Perazzi, A., Contiero, B., De Mattia, G. & Iacopetti, I. (2016) Pocket technique or pocket technique combined with modified orbital rim anchorage for the replacement of a prolapsed gland of the third eyelid in dogs: 353 dogs. *Veterinary Ophthalmology*, **19**, 214–219.
- Mulvihill, J.J., Mulvihill, C.G. & Priester, W.A. (1980) Cleft palate in domestic animals: Epidemiologic features. *Teratology*, **21** (1), 109–112.
- Muñana, K.R., Nettifee-Osborne, J.A., Bergman, R.L. & Mealey, K.L. (2012) Association between ABCB1 genotype and seizure outcome in collies with epilepsy. *Journal of Veterinary Internal Medicine*, **26**, 1358–1364.
- Munday, J.S., Dyer, C.B., Hartman, A.C. *et al.* (2006) A possible predisposition to dilated cardiomyopathy in Huntaway dogs. *New Zealand Veterinary Journal*, **54**, 231–234.
- Munjar, T.A., Austin, C.C. & Breur, G.J. (1998) Comparison of risk factors for hypertrophic osteodystrophy, craniomandibular osteopathy and canine distemper virus infection. *Veterinary and Comparative Orthopaedics and Traumatology*, **11**, 37–43.
- Munro, H.M.C. & Thrusfield, M.V. (2001) Battered pets': non-accidental physical injuries found in dogs and cats. *Journal of Small Animal Practice*, **42** (6), 279–290.
- Munyard, K.A., Sherry, C.R. & Sherry, L. (2007) A retrospective evaluation of congenital ocular defects in Australian Shepherd dogs. *Veterinary Ophthalmology* **10** (1), 19–22.
- Murphy, C.J., Marfurt, C.F., McDermott, A. *et al.* (2001) Spontaneous chronic corneal epithelial defects (SCCED) in dogs: clinical features, innervation, and effect of topical SP, with or without IGF-1. *Investigative Ophthalmology and Visual Science*, **42** (10), 2252–2261.
- Murphy, C.J., Zadnik, K. & Mannis, M.J. (1992) Myopia and refractive error in dogs. *Investigative Ophthalmology and Visual Science*, **33** (8), 2459–2463.
- Murphy, K.A. & Brisson, B.A. (2006) Evaluation of lung lobe torsion in Pugs: 7 cases (1991–2004). *Journal of the American Veterinary Medical Association*, **228** (1), 86–90.
- Murphy, S., Sparkes, A.H., Blunden, A.S. *et al.* (2006) Effects of stage and number of tumours on prognosis of dogs with cutaneous mast cell tumours. *Veterinary Record*, **158** (9), 287–291.
- Muste, A., Papuc, I., Beteg, F., Lăcătuș, R. & Donisa, A. (2010) Observations about osteosarcoma incidence in dogs. *Cluj Veterinary Journal*, **17** (1), 51–54.
- Načeradská, M. (2009) [Fanconi syndrome in Basenji in the Czech Republic and Germany.] *Veterinářství*, **59**, 7–11.
- Nachreiner, R.F., Refsal, K.R., Graham, P.A. & Bowman, M.M. (2002) Prevalence of serum thyroid hormone autoantibodies in dogs with clinical signs of hypothyroidism. *Journal of the American Veterinary Medical Association*, **220** (4), 466–471.
- Nakamoto, Y., Ozawa, T., Katakabe, K. *et al.* (2009) Fibrocartilaginous embolism of the spinal cord diagnosed by characteristic clinical findings and magnetic resonance imaging in 26 dogs. *Journal of Veterinary Medical Science*, **71** (2), 171–176.
- Narama, I., Kobayashi, Y., Yamagami, T., Ozaki, K. & Ueda, Y. (2005) Pigmented cutaneous papillomatosis (pigmented epidermal nevus) in three Pug dogs; histopathology, electron microscopy and analysis of viral DNA by the Polymerase Chain Reaction. *Journal of Comparative Pathology*, **132** (2), 132–138.

- Naranjo, C., Dubielzig, R.R. & Friedrichs, K.R. (2007) Canine ocular histiocytic sarcoma. *Veterinary Ophthalmology* **10** (3), 632–645.
- Narfström, K. (1985) Progressive retinal atrophy in the Abyssinian cat. Clinical characteristics. *Investigative Ophthalmology and Visual Science*, **26** (2), 193–200.
- Narfström, K. (1999a) Retinal dystrophy or 'congenital stationary night blindness' in the Briard. *Veterinary Ophthalmology*, **2** (1), 75–76.
- Narfström, K. (1999b) Herediatry and congenital ocular diseases in the cat. *Journal of Feline Medicine and Surgery*, **1**, 135–141.
- Narfström, K. & Ekesten, B. (1998) Electroretinographic evaluation of Papillons with and without hereditary retinal degeneration. *American Journal of Veterinary Research*, **59** (2), 221–226.
- Narfström, K. & Wrigstad, A. (1999) Clinical, electrophysiological and morphological changes in a case of hereditary retinal degeneration in the Papillon dog. *Veterinary Ophthalmology*, **2** (1), 67–74.
- Narfström, K., Wrigstad, A., Ekesten, B. & Berg, A.L. (2007) Neuronal ceroid lipofuscinosis: clinical and morphologic findings in nine affected Polish Owczarek Nizinny (PON) dogs. *Veterinary Ophthalmology*, **10** (2), 111–120.
- Narojek, T., Fiszdon, K. & Hanysz, E. (2008) Canine elbow dysplasia in different breeds. *Bulletin of the Veterinary Institute in Pulawy*, **52**, 169–173.
- Nash, A.S. (1989) Familial renal disease in dogs. *Journal of Small Animal Practice*, **30**, 178–183.
- Nayak, D.C., Tripathy, S.B. & Dey, P.C. (1997) Prevalence of canine demodicosis in Orissa (India). *Veterinary Parasitology*, **73** (3/4), 347–352.
- Neath, P. J., Brockman, D. J. and King, L. G. (2000) Lung lobe torsion in dogs: 22 cases (1981–1999). *Journal of the American Veterinary Medical Association*, **217**, 1041–1044.
- Nečas, A. (1999) Clinical aspects of surgical treatment of thoracolumbar disc disease in dogs: a retrospective study of 300 cases. *Acta Veterinaria Brno*, **68** (2), 121–130.
- Nečas, A., Dvořák, M. & Zatloukal, J. (1999) Incidence of osteochondrosis in dogs and its late diagnosis. *Acta Veterinaria Brno*, **68** (2), 131–139.
- Nečas, A., Zatloukal, J., Kecova, H. & Dvořák, M. (2000) Predisposition of dog breeds to rupture of the cranial cruciate ligament. *Acta Veterinaria Brno*, **69** (4), 305–310.
- Neff, M. W., Robertson, K. R., Wong, A. K. *et al.* (2004) Breed distribution and history of canine *mdr1-1Δ*, a pharmacogenetic mutation that marks the emergence of breeds from the collie lineage. *Proceedings of the National Academy of Sciences of the United States of America*, **101** (32), 11725–11730.
- Neff, M.W. & Rine, J. (2006) A fetching model organism. *Cell*, **124** (2), 229–231.
- Nemzek, J.A., Agrodnia, M.D. & Hauptman, J.G. (2007) Breed-specific pro-inflammatory cytokine production as a predisposing factor for susceptibility to sepsis in the dog. *Journal of Veterinary Emergency and Critical Care*, **17** (4), 368–372.
- Nesbitt, G.H., Izzo, J., Peterson, L. *et al.* (1980) Canine hypothyroidism: a retrospective study of 108 cases. *Journal of the American Veterinary Medical Association*, **177** (11), 1117–1122.
- Nestor, D.D., Holan, K.M., Johnson, C.A., Schall, W. & Kaneene, J.B. (2006) Serum alkaline phosphatase activity in Scottish Terriers versus dogs of other breeds. *Journal of the American Veterinary Medical Association*, **228** (2), 222–224.
- Ng, C.Y. & Mills, J.N. (1985) Clinical and haematological features of haemangiosarcomas in dogs. *Australian Veterinary Journal*, **62** (1), 1–4.
- Nganvongpanit, K. & Yano, T. (2011) Prevalence of and risk factors of patellar luxation in dogs in Chiang Mai, Thailand, during the years 2006–2011. *Thai Journal of Veterinary Medicine*, **41** (4), 449.
- Nganvongpanit, K. & Yano, T. (2013) Prevalence of swimming puppy syndrome in 2,443 puppies during the years 2006–2012 in Thailand. *Veterinary Medicine International*, 617803.
- Nibe, K., Kita, C., Morozumi, M. *et al.* (2007) Clinicopathological features of canine neuroaxonal dystrophy and cerebellar cortical abiotrophy in Papillon and Papillon-related dogs. *Journal of Veterinary Medical Science*, **69** (10), 1047–1052.
- Nibe, K., Nakayama, H. & Uchida, K. (2009) Immunohistochemical features of dystrophic axons in Papillon dogs with neuroaxonal dystrophy. *Veterinary Pathology Online*, **46** (3), 474–483.
- Nicolet, D.C. (2006) Study on the frequency of occurrence of keratoconjunctivitis sicca in predisposed dog breeds. *Dansk Veterinærtidsskrift*, **89**, 30–34.

- Nicorescu, V. & Crivineanu, V. (2007) Researches regarding the diagnosis of atopic dermatitis in dog. *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cui-Napoca Veterinary Medicine*, **64** (1/2), 595.
- Nielen, A.L.J., Janss, L.L.G. & Knol, B.W. (2001) Heritability estimations for diseases, coat colour, body weight and height in a birth cohort of Boxers. *American Journal of Veterinary Research*, **62**, 1198–1206.
- Nielsen, L., Andreasen, S.N., Andersen, S.D. & Kristensen, A.T. (2010) Malignant histiocytosis and other causes of death in Bernese Mountain Dogs in Denmark. *Veterinary record*, **166**, 199–202.
- Nielsen, L.N., Wiinberg, B., Kjølgaard-Hansen, M. & Kristensen, A. T. (2011) The presence of antiphospholipid antibodies in healthy Bernese Mountain Dogs. *Journal of Veterinary Internal Medicine*, **25**, 1258–1263.
- Niewold, T.A., van der Linde-Sipman, J.S., Murphy, C. *et al.* (1999) Familial amyloidosis in cats: Siamese and Abyssinian AA proteins differ in primary sequence and pattern of deposition. *Amyloid*, **6** (3), 205–209.
- Niskanen, M. & Thrusfield, M.V. (1998) Associations between age, parity, hormonal therapy and breed, and pyometra in Finnish dogs. *Veterinary Record*, **143** (18), 493–498.
- Nivy, R., Lyons, L.A., Aroch, I. *et al.* (2015) Polycystic kidney disease in four British shorthair cats with successful treatment of bacterial cyst infection. *Journal of Small Animal Practice* **56** (9) 585–589.
- Njikam Nsangou, I., Huault, M., Pirson, V. & Detilleux, J. (2009) The influence of phylogenetic origin on the occurrence of brachycephalic airway obstruction syndrome in a large retrospective study. *International Journal of Applied Research in Veterinary Medicine*, **7** (3).
- Noden, D.M. & Evans, H.E. (1986) Inherited homeotic midfacial malformations in Burmese cats. *Journal of Craniofacial Genetics and Developmental Biology Supplement*, **2**, 249–266.
- Nødtvedt, A., Egenvall, A., Bergvall, K. & Hedhammar, A. (2006) Incidence of and risk factors for atopic dermatitis in a Swedish population of insured dogs. *Veterinary Record*, **159** (8), 241–246.
- Nødtvedt, A., Gamlem, H., Gunnes, G. *et al.* (2011) Breed differences in the proportional morbidity of testicular tumours and distribution of histopathologic types in a population-based canine cancer registry. *Veterinary and Comparative Oncology*, **9** (1), 45–54.
- Noli, C., Colombo, S., Abramo, F. *et al.* (2004) Papular eosinophilic/mastocytic dermatitis (feline urticaria pigmentosa) in Devon Rex cats: a distinct disease entity or a histopathological reaction pattern? *Veterinary Dermatology* **15**, 253–259.
- Noonan, C.H. & Kay, J.M. (1990) Prevalence and geographic distribution of Fanconi syndrome in Basenjis in the United States. *Journal of the American Veterinary Medical Association*, **197** (3), 345–349.
- Norris, A.J., Naydan, D.K. & Wilson, D.W. (2005a) Interstitial lung disease in West Highland White Terriers. *Veterinary Pathology Online*, **42** (1), 35–41.
- Norris, A.M., Laing, E.J., Valli, V.E.O. *et al.* (1992) Canine bladder and urethral tumors: A retrospective study of 115 cases (1980–1985). *Journal of Veterinary Internal Medicine*, **6** (3), 145–153.
- Norris, C.R., Williams, B.J., Ling, G.V. *et al.* (2000) Recurrent and persistent urinary tract infections in dogs: 383 cases (1969–1995). *Journal of the American Animal Hospital Association*, **36** (6), 484–492.
- Norris, J.M., Boswars, K.L., White, J.D. *et al.* (2005b) Clinicopathological findings associated with feline infectious peritonitis in Sydney, Australia: 42 cases (1990–2002). *Australian Veterinary Journal*, **83** (11), 666–673.
- Noszczyk-Nowak, A., Michałek, M., Kałuża, E., Cepiel, A. & Paślawska, U. (2017) Prevalence of arrhythmias in dogs examined between 2008 and 2014. *Journal of Veterinary Research*, **61** (1), 103.
- Notari, L. & Goodwin, D. (2007) A survey of behavioural characteristics of pure-bred dogs in Italy. *Applied Animal Behaviour Science*, **103** (1–2), 118–130.
- Nuttall, T.J., German, A.J., Holden, S.L. *et al.* (2008) Successful resolution of dermatophyte mycetoma following terbinafine treatment in two cats. *Veterinary Dermatology*, **19** (6), 405–410.
- Oberbauer, A.M., Bell, J.S., Belanger, J.M. & Famula, T.R. (2006) Genetic evaluation of Addison's disease in the Portuguese Water Dog. *BMC Veterinary Research*, **2**, 15.
- Oberbauer, A.M., Bineman, K.S., Belanger, J.M. *et al.* (2002) Inheritance of hypoadrenocorticism in Bearded Collies. *American Journal of Veterinary Research*, **63** (5), 643–647.

- Oberbauer, A.M., Hollingsworth, S.R., Belanger, J.M. *et al.* (2008) Inheritance of cataracts and primary lens luxation in Jack Russell Terriers. *American Journal of Veterinary Research*, **69** (2), 222–227.
- O'Brien, C.R., Krockenberger, M.B., Wigney, D.I., Martin, P. & Malik, R. (2004). Retrospective study of feline and canine cryptococcosis in Australia from 1981 to 2001: 195 cases. *Medical Mycology*, **42** (5), 449–460.
- O'Brien, D.P., Johnson, G.S., Schnabel, R.D. *et al.* (2005) Genetic mapping of canine multiple system degeneration and ectodermal dysplasia loci. *Journal of Heredity*, **96**, 727–734.
- O'Brien, M.G., Berg, J. & Engler, S.J. (1992) Treatment by digital amputation of subungual squamous cell carcinoma in dogs: 21 cases (1987–1988). *Journal of the American Veterinary Medical Association*, **201** (5), 759–761.
- O'Brien, T.D., Osborne, C.A., Yano, B.L. *et al.* (1982) Clinicopathologic manifestations of progressive renal disease in Lhasa Apso and Shih Tzu dogs. *Journal of the American Veterinary Medical Association*, **180** (6), 658–664.
- Ochoa, R., Breitschwerdt, E.B. & Lincoln, K.L. (1984) Immunoproliferative small intestinal disease in Basenji dogs: morphologic observations. *American Journal of Veterinary Research*, **45** (3), 482–490.
- Ofri, R., Lambrou, G.N., Allgoewer, I. *et al.* (2009) Clinical evaluation of pimecrolimus eye drops for treatment of canine keratoconjunctivitis sicca: a comparison with cyclosporine A. *Veterinary Journal*, **179** (1), 70–77.
- Ogata, N., Gillis, T., Liu, X. *et al.* (2013) Brain structural abnormalities in Doberman Pinschers with canine compulsive disorder. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, **45**, 1–6.
- O'Grady, M.R., Minors, S.L. & O'Sullivan, M.L. (2008) Effect of pimobendan on case fatality rate in Doberman Pinschers with congestive heart failure caused by dilated cardiomyopathy. *Journal of Veterinary Internal Medicine*, **22** (4), 897–904.
- Ohad, D.G., Avrahami, A., Waner, T. *et al.* (2013) The occurrence and suspected mode of inheritance of congenital subaortic stenosis and tricuspid valve dysplasia in Dogue de Bordeaux dogs. *Veterinary Journal*, **197** (2), 351–357.
- Oheida, A.H., Philip, C.J., Yen, H.-H. *et al.* (2016) Observations of sacrocaudal fusion in Greyhounds and other dogs. *Veterinary Comparative Orthopaedics and Traumatology*, **29** (1), 61–67.
- Ohlerth, S., Senn, S., Geissbühler, U., Kircher, P. & Flückiger, M. (2016) Prevalence of humeral head osteochondrosis in the Greater Swiss Mountain dog and the Border Collie in Switzerland. *Schweizer Archiv für Tierheilkunde*, **158** (11), 749–754.
- Öhlund, M., Fall, T., Ström Hölst, B. *et al.* (2015) Incidence of diabetes mellitus in insured swedish cats in relation to age, breed and sex. *Journal of Veterinary Internal Medicine*, **29** (5), 1342–1347.
- Ohmi, A., Tsukamoto, A., Ohno, K. *et al.* (2012) A retrospective study of inflammatory colorectal polyps in Miniature Dachshunds. *Journal of Veterinary Medical Science*, **74**, 59–64.
- Okafor, C.C., Lefebvre, S.L., Pearl, D.L. *et al.* (2014) Risk factors associated with calcium oxalate urolithiasis in dogs evaluated at general care veterinary hospitals in the United States. *Preventive Veterinary Medicine*, **115** (3–4), 217–228.
- Olby, N., Blot, S., Thibaud, J.-L. *et al.* (2004) Cerebellar cortical degeneration in adult American Staffordshire Terriers. *Journal of Veterinary Internal Medicine*, **18** (2), 201–208.
- O'Leary, C.A. & Atwell, R.B. (2003) Control of familial and renal cardiac diseases in English Bull Terriers: how to repair a damaged breed? *Australian Veterinary Practitioner*, **33** (2), 75–80.
- O'Leary, C.A., Duffy, D.L., Gething, M.A. *et al.* (2013) Investigation of diabetes mellitus in Burmese cats as an inherited trait: a preliminary study. *New Zealand Veterinary Journal*, **61** (6), 354–358.
- O'Leary, C.A., Ghoddusi, M. & Huxtable, C.R. (2002) Renal pathology of polycystic kidney disease and concurrent hereditary nephritis in Bull Terriers. *Australian Veterinary Journal*, **80** (6), 353–361.
- O'Leary, C.A., Mackay, B.M., Malik, R. *et al.* (1999) Polycystic kidney disease in Bull Terriers: an autosomal dominant inherited disorder. *Australian Veterinary Journal*, **77** (6), 361–366.
- O'Leary, C.A., Mackay, B.M., Taplin, R.H. *et al.* (2005) Auscultation and echocardiographic findings in Bull Terriers with and without polycystic kidney disease. *Australian Veterinary Journal*, **83** (5), 270–275.
- O'Leary, C.A., Parslow, A., Malik, R. *et al.* (2014) The inheritance of extra-hepatic portosystemic shunts and elevated bile acid concentrations in Maltese dogs. *Journal of Small Animal Practice*, **55** (1), 14–21.

- O'Leary, C.A. & Wilkie, I. (2009) Cardiac valvular and vascular disease in Bull Terriers. *Veterinary Pathology*, **46**, 1149–1155.
- Olesen, H.P., Jensen, O.A. & Norn, M.S. (1974) Congenital hereditary cataract in Cocker Spaniels. *Journal of Small Animal Practice*, **15** (12), 741–750.
- Oliveira, P., Domenech, O., Silva, J. *et al.* (2011) Retrospective review of congenital heart disease in 976 dogs. *Journal of Veterinary Internal Medicine*, **25** (3), 477–483.
- Oliver, J.A., Forman, O. P., Pettitt, L. & Mellersh, C. S. (2015) Two independent mutations in ADAMTS17 are associated with primary open angle glaucoma in the Basset Hound and Basset Fauve de Bretagne breeds of dog. *PLoS One*, **10**, e0140436.
- Oliver, J.A., Ekiri, A. & Mellersh, C.S. (2016a) Prevalence of pectinate ligament dysplasia and associations with age, sex and intraocular pressure in the Basset hound, Flatcoated retriever and Dandie Dinmont terrier. *Canine Genetics and Epidemiology*, **3**, 1.
- Oliver, J.A., Ekiri, A. & Mellersh, C.S. (2016b) Prevalence and progression of pectinate ligament dysplasia in the Welsh Springer Spaniel. *Journal of Small Animal Practice*, **57** (8), 416–421.
- Olivera, D.K., Riis, R.C., Dutton, A.G. *et al.* (1991) Feline lens displacement: a retrospective analysis of 345 cases. *Progress in Veterinary and Comparative Ophthalmology*, **1**, 239–244.
- Olsen, L.H., Fredholm, M. & Pedersen, H.D. (1999) Epidemiology and inheritance of mitral valve prolapse in Dachshunds. *Journal of Veterinary Internal Medicine*, **13**, 448–456.
- Olsson, M., Frankowiack, M., Tengvall, K. *et al.* (2014) The dog as a genetic model for immunoglobulin A (IgA) deficiency: Identification of several breeds with low serum IgA concentrations. *Veterinary Immunology and Immunopathology*, **160** (4), 255–259.
- Olsson, M., Meadows, J.R.S., Truvé, K. *et al.* (2011) A novel unstable duplication upstream of HAS2 predisposes to a breed-defining skin phenotype and a periodic fever syndrome in Chinese Shar-Pei dogs. *PLoS Genetics*, **7** (3), e1001332.
- Olsson, M., Tintle, L., Kierczak, M. *et al.* (2013) Thorough investigation of a Canine Autoinflammatory Disease (AID) confirms one main risk locus and suggests a modifier locus for amyloidosis. *PLoS One*, **8** (10), e75242.
- O'Marra, S.K., Delaforcade, A.M. and Shaw, S.P. (2011) Treatment and predictors of outcome in dogs with immune-mediated thrombocytopenia. *Journal of the American Veterinary Medical Association*, **238**, 346–352.
- Ondreka, N., Amort, K.H., Stock, K.F. *et al.* (2012) Skeletal morphology and morphometry of the lumbosacral junction in German Shepherd Dogs and an evaluation of the possible genetic basis for radiographic findings. *Veterinary Journal*, **196**, 64–70.
- O'Neill, D.G., Church, D.B., McGreevy, P.D., Thomson, P.C. & Brodbelt, D.C. (2013a) Longevity and mortality of owned dogs in England. *Veterinary Journal*, **198** (3), 638–643.
- O'Neill, D.G., Church, D.B., McGreevy, P.D., Thomson, P.C. & Brodbelt, D.C. (2014a) Approaches to canine health surveillance. *Canine Genetics and Epidemiology*, **1** (1), 2.
- O'Neill, D.G., Church, D.B., McGreevy, P.D., Thomson, P.C. & Brodbelt, D.C. (2014b) Prevalence of disorders recorded in dogs attending primary-care veterinary practices in England. *PLoS One*, **9** (3), 1–16.
- O'Neill, D.G., Church, D.B., McGreevy, P.D., Thomson, P.C. & Brodbelt, D.C. (2015a) Longevity and mortality of cats attending primary care veterinary practices in England. *Journal of Feline Medicine and Surgery*, **17** (2), 125–133.
- O'Neill, D.G., Coulson, N.R., Church, D.B. & Brodbelt, D.C. (2017a). Demography and disorders of German Shepherd Dogs under primary veterinary care in the UK. *Canine Genetics and Epidemiology*, **4** (1), 7.
- O'Neill, D.G., Darwent, E.C., Church, D.B. & Brodbelt, D.C. (2016a) Demography and health of Pugs under primary veterinary care in England. *Canine Genetics and Epidemiology*, **3** (1), 1–12.
- O'Neill, D.G., Elliott, J., Church, D. B. *et al.* (2013b) Chronic kidney disease in dogs in UK veterinary practices: prevalence, risk factors, and survival. *Journal of Veterinary Internal Medicine*, **27**, 814–821.
- O'Neill, D.G., Gostelow, R., Orme, C. *et al.* (2016b) Epidemiology of diabetes mellitus among 193,435 cats attending primary care practices in England. *Journal of Veterinary Internal Medicine*, **30** (4), 964–972.

- O'Neill, D.G., Jackson, C., Guy, J. H. *et al.* (2015b) Epidemiological associations between brachycephaly and upper respiratory tract disorders in dogs attending veterinary practices in England. *Canine Genetics and Epidemiology*, **2**, 10.
- O'Neill, D.G., Lee, M.M., Brodbelt, D.C., Church, D.B. & Sanchez, R.F. (2017b) Corneal ulcerative disease in dogs under primary veterinary care in England: epidemiology and clinical management. *Canine Genetics and Epidemiology*, **4** (1), 5.
- O'Neill, D.G., Meeson, R.L., Sheridan, A., Church, D.B. & Brodbelt, D.C. (2016c) The epidemiology of patellar luxation in dogs attending primary-care veterinary practices in England. *Canine Genetics and Epidemiology*, **3** (1), 1–12.
- O'Neill, D.G., O'Sullivan, A.M., Manson, E.A. *et al.* (2017c) Canine dystocia in 50 UK first-opinion emergency-care veterinary practices: prevalence and risk factors. *Veterinary Record*, **181** (4), 88.
- O'Neill, D.G., Scudder, C., Faire, J. M. *et al.* (2016d) Epidemiology of hyperadrenocorticism among 210,824 dogs attending primary-care veterinary practices in the UK from 2009 to 2014. *Journal of Small Animal Practice*, **57** (7), 365–373.
- Onions, D.E. (1984) A prospective survey of familial canine lymphosarcoma. *Journal of the National Cancer Institute*, **72** (4), 909–912.
- Orthopedic Foundation for Animals (2015). *Patellar Luxation Statistics*. http://www.offa.org/stats_pl.html (accessed October 2017).
- Orthopedic Foundation for Animals (2016). [Disease Information]. <http://www.offa.org> (accessed October 2017).
- Osborne, C.A., Hammer, R.F. & Klausner, J.S. (1981) Canine silica urolithiasis. *Journal of the American Veterinary Medical Association*, **178** (8), 809–813.
- Osborne, C.A., Lulich, J.P., Polzin, D.J. *et al.* (1999) Analysis of 77,000 canine uroliths. Perspectives from the Minnesota Urolith Center. *Veterinary Clinics of North America: Small Animal Practice*, **29** (1), 17–38.
- Oshima, Y., Bjerkås, E. & Peiffer, R.L. (2004) Ocular histopathologic observations in Norwegian Elkhounds with primary open-angle, close-cleft glaucoma. *Veterinary Ophthalmology*, **7** (3), 185–188.
- Oswald, G.P. & Orton, E.C. (1993) Patent ductus arteriosus and pulmonary hypertension in related Pembroke Welsh Corgis. *Journal of the American Veterinary Medical Association*, **202** (5), 761–764.
- O'Toole, D., Young, S., Severin, G.A. & Neumann, S. (1983) Retinal dysplasia of English Springer Spaniel dogs: light microscopy of the postnatal lesions. *Veterinary Pathology*, **20** (3), 298–311.
- O'Toole, D.O. & Roberts, S. (1984) Generalized progressive retinal atrophy in two Akita dogs. *Veterinary Pathology*, **21** (5), 457–462.
- Outerbridge, C.A., Marks, Stanley L. & Rogers, Quinton R. (2002) Plasma amino acid concentrations in 36 dogs with histologically confirmed superficial necrolytic dermatitis. *Veterinary Dermatology*, **13** (4), 177–186.
- Øvrebø Bohnhorst, J., Hanssen, I. & Moen, T. (2001) Antinuclear antibodies (ANA) in Gordon setters with symmetrical lupoid onychodystrophy and black hair follicular dysplasia. *Acta Veterinaria Scandinavica*, **42**, 323–329.
- Owen, M.C., Lamb, C.R., Lu, D. & Targett, M.P. (2004) Material in the middle ear of dogs having magnetic resonance imaging for investigation of neurologic signs. *Veterinary Radiology and Ultrasound*, **45**, 149–155.
- Owen, T.J., Bruvette, D.S. & Lavton, C.E. (1996) Chemodectoma in dogs. *Compendium on Continuing Education for the Practicing Veterinarian*, **18** (3), 253–265.
- Packer, R.M.A., Hendricks, A., Tivers, M.S. & Burn, C.C. (2015) Impact of facial conformation on canine health: Brachycephalic Obstructive Airway Syndrome. *PLoS One*, **10** (10), e0137496.
- Packer, R.M.A., Hendricks, A., Volk, H. A., Shihab, N. K. & Burn, C. C. (2013) How long and low can you go? Effect of conformation on the risk of thoracolumbar intervertebral disc extrusion in domestic dogs, *PLoS One*, **8** (7), e69650.
- Packer, R.M.A., Seath, I. J., O'Neill, D. G., De Decker, S. & Volk, H. A. (2016a) DachsLife 2015: an investigation of lifestyle associations with the risk of intervertebral disc disease in Dachshunds. *Canine Genetics and Epidemiology*, **3**, 8.

- Packer, R.M.A., Shihab, N.K., Torres, B.B.J. *et al.* (2016b) Risk factors for cluster seizures in canine idiopathic epilepsy. *Research in Veterinary Science*, **105**, 136–138.
- Padgett, G.A., Madewell, B.R., Keller, E.T. *et al.* (1995) Inheritance of histiocytosis in Bernese Mountain Dogs. *Journal of Small Animal Practice*, **36** (3), 93–98.
- Page, A., Edwards, G. & Atwell, R.B. (1993) Echocardiographic values in the Greyhound. *Australian Veterinary Journal*, **70**, 361–364.
- Pagé, N., Paradis, M., Lapointe, J.M. *et al.* (2003) Hereditary nasal parakeratosis in Labrador Retrievers. *Veterinary Dermatology*, **14**, 103–110.
- Pagès, J. P. (1988) Amylose rénale chez le chien: analyse de 35 observations. *Pratique Médicale et Chirurgicale de l'Animal de Compagnie*, **23**, 311–318.
- Palmer, A.C., Payne, J.E. & Lallace, M.E. (1973) Hereditary quadriplegia and amblyopia in the Irish Setter. *Journal of Small Animal Practice*, **14** (6), 343–352.
- Paltrinieri, S., Ibba, F. & Rossi, G. (2014) Haematological and biochemical reference intervals of four feline breeds. *Journal of Feline Medicine and Surgery*, **16** (2), 125–136.
- Panakova, L., Koch, H., Kolb, S. & Mueller, R.S. (2008) Thyroid testing in Sloughis. *Journal of Veterinary Internal Medicine*, **22** (5), 1144–1148.
- Panciera, D.L. (1994) Hypothyroidism in dogs: 66 cases (1987–1992). *Journal of the American Veterinary Medical Association*, **204** (5), 761–767.
- Panich, R., Scott, D.W. & Miller, W.H. (1991) Canine cutaneous sterile pyogranuloma/granuloma syndrome: a retrospective analysis of 29 cases (1976–1988). *Journal of the American Animal Hospital Association*, **27**, 519–528.
- Pápa, K., Máthé, A., Abonyi-Tóth, Z. (2011) Occurrence, clinical features and outcome of canine pancreatitis (80 cases). *Acta Veterinaria Hungarica*, **59** (1), 37–52.
- Papaoiannou, N.G. & Dubielzig, R. R. (2013) Histopathological and immunohistochemical features of vitreoretinopathy in Shih Tzu dogs. *Journal of Comparative Pathology*, **148** (2–3), 230–235.
- Pape, L.A., Price, J.M. & Rippe, J.M. (1986) Hemodynamics and left ventricular function: a comparison between adult Racing Greyhounds and Greyhounds completely untrained from birth. *Basic Research in Cardiology*, **81** (4), 417–424.
- Paradis, M., De Jaham, C., Page, N. *et al.* (2005) Acral mutilation and analgesia in 13 French Spaniels. *Veterinary Dermatology*, **16**, 87–93.
- Paradis, M., Scott, D.W. & Breton, L. (1989) Squamous cell carcinoma of the nail bed in three related Giant Schnauzers. *Veterinary Record*, **125** (12), 322–324.
- Park, S.A., Yi, N.Y., Jeong, M.B. *et al.* (2009) Clinical manifestations of cataracts in small breed dogs. *Veterinary Ophthalmology*, **12** (4), 205–210.
- Parker, H.G., Chase, K., Cadieu, E., Lark, K.G. & Ostrander, E.A. (2010) An insertion in the *rspo2* gene correlates with improper coat in the Portuguese Water Dog. *Journal of Heredity*, **101** (5), 612–617.
- Parker, H.G., Dreger, D.L., Rimbault, M. *et al.* (2017) Genomic analyses reveal the influence of geographic origin, migration, and hybridization on modern dog breed development. *Cell Reports*, **19** (4), 697–708.
- Parker, H.G. & Kilroy-Glynn, P. (2012) Myxomatous mitral valve disease in dogs: does size matter? *Journal of Veterinary Cardiology*, **14** (1), 19–29.
- Parker, H.G., Kukekova, A.V., Akey, D.T. *et al.* (2007) Breed relationships facilitate fine-mapping studies: a 7.8-kb deletion cosegregates with Collie eye anomaly across multiple dog breeds. *Genome Research*, **17** (11), 1562–1571.
- Parker, H.G., VonHoldt, B.M., Quignon, P. *et al.* (2009) An expressed *fgf4* retrogene is associated with breed-defining chondrodysplasia in domestic dogs. *Science*, **325** (5943), 995–998.
- Parker, J.E., Knowler, S.P., Rusbridge, C., Noorman, E. & Jeffery, N.D. (2011) Prevalence of asymptomatic syringomyelia in Cavalier King Charles Spaniels. *Veterinary Record*, **168** (25), 667.
- Parker, W.M. & Foster, R.A. (1996) Cutaneous vasculitis in five Jack Russell Terriers. *Veterinary Dermatology*, **7**, 109–115.
- Parry, B.W., Howard, M.A., Mansell, P.D. *et al.* (1988) Haemophilia A in German Shepherd Dogs. *Australian Veterinary Journal*, **65**, 276–279.

- Parshall, C.J., Wyman, M., Nitroy, S., Acland, G.M. & Aguirre, G.D. (1991) Photoreceptor dysplasia: an inherited progressive retinal atrophy of Miniature Schnauzer dogs. *Progress in Veterinary and Comparative Ophthalmology*, **1** (3), 187.
- Pasquini, A., Luchetti, E. & Cardini, G. (2008) Plasma lipoprotein concentrations in the dog: the effects of gender, age, breed and diet. *Journal of Animal Physiology and Animal Nutrition*, **92** (6), 718–722.
- Paster, E.R. (2005) Estimates of prevalence of hip dysplasia in Golden Retrievers and Rottweilers and the influence of bias on published prevalence figures. *Journal of the American Veterinary Medical Association*, **226** (3), 387–392.
- Pastor, M., Chalvet-Monfray, K., Marchal, T. *et al.* (2009) Genetic and environmental risk indicators in canine non-hodgkin's lymphomas: Breed associations and geographic distribution of 608 cases diagnosed throughout France over 1 year. *Journal of Veterinary Internal Medicine*, **23** (2), 301–310.
- Paterson, S. (1995) Sterile idiopathic pedal panniculitis in the German Shepherd Dog: clinical presentation and response to treatment of four cases. *Journal of Small Animal Practice*, **36** (11), 498–501.
- Pathak, E.J. (2004) Type 3 von Willebrand's disease in a Shetland Sheepdog. *Canadian Veterinary Journal*, **45** (8), 685–687.
- Patnaik, A.K., Ehler, W.J. & MacEwen, E.G. (1984a) Canine cutaneous mast cell tumor: Morphologic grading and survival time in 83 dogs. *Veterinary Pathology*, **21** (5), 469–474.
- Patnaik, A.K., Lieberman, P.H., Erlandson, R.A. & Liu, S.K. (1984b) Canine sinonasal skeletal neoplasms: chondrosarcomas and osteosarcomas. *Veterinary Pathology*, **21** (5), 475–482.
- Patronek, G.J., Waters, D.J. & Glickman, L.T. (1997) Comparative longevity of pet dogs and humans: implications for gerontology research. *Journal of Gerontology: Biological Sciences*, **52** (3), B171–B178.
- Patterson, D.F. (1989) Hereditary congenital heart defects in dogs. *Journal of Small Animal Practice*, **30** (3), 153–165.
- Patterson, E.E., Armstrong, P.J., O'Brien, D.P. *et al.* (2005) Clinical description and mode of inheritance of idiopathic epilepsy in English Springer Spaniels. *Journal of the American Veterinary Medical Association*, **226** (1), 54–58.
- Patterson, E.E., Mickelson, J.R., Da, Y. *et al.* (2003) Clinical characteristics and inheritance of idiopathic epilepsy in Vizslas. *Journal of Veterinary Internal Medicine*, **17**, 319–325.
- Patterson, W.R., Estry, D.W., Schwartz, K.A. *et al.* (1989) Absent platelet aggregation with normal fibrinogen binding in Basset Hound hereditary thrombopathy. *Thrombosis and Haemostasis*, **62**, 1011–1015.
- Paulsen, M. E., Lavach, J. D., Snyder, S. P., Severin, G. A. & Eichenbaum, J. D. (1987) Nodular granulomatous episclerokeratitis in dogs: 19 cases (1973–1985). *Journal of the American Veterinary Medical Association*, **190**, 1581–1587.
- Payne, J., Luis Fuentes, V., Boswood, A. *et al.* (2010) Population characteristics and survival in 127 referred cats with hypertrophic cardiomyopathy (1997 to 2005). *Journal of Small Animal Practice*, **51** (10), 540–547.
- Pearson, H., Gaskell, C.J., Gibbs, C. & Waterman, A. (1974) Pyloric and oesophageal dysfunction in the cat. *Journal of Small Animal Practice*, **15**, 487–501.
- Pedersen, H.D., Häggström, J., Olsen, L.H. *et al.* (2002) Idiopathic asymptomatic thrombocytopenia in Cavalier King Charles Spaniels is an autosomal recessive trait. *Journal of Veterinary Internal Medicine*, **16**, 169–173.
- Pedersen, K. & Scott, D.W. (1991) Idiopathic pyogranulomatous inflammation and leukocytoclastic vasculitis of the nasal planum, nostrils and nasal mucosa in Scottish Terriers in Denmark. *Veterinary Dermatology*, **2** (2), 85–89.
- Pedersen, N.C., Liu, H., Greenfield, D.L. *et al.* (2012a) Multiple autoimmune diseases syndrome in Italian Greyhounds: preliminary studies of genome-wide diversity and possible associations within the dog leukocyte antigen (DLA) complex. *Veterinary Immunology and Immunopathology*, **145** (1–2), 264–276.
- Pedersen, N., Liu, H., McLaughlin, B. & Sacks, B. (2012b) Genetic characterization of healthy and sebaceous adenitis affected Standard Poodles from the United States and the United Kingdom. *Tissue Antigens*, **80** (1), 46–57.

- Pedersen, N.C., Liu, H., Leonard, A. *et al.* (2015) A search for genetic diversity among Italian Greyhounds from Continental Europe and the USA and the effect of inbreeding on susceptibility to autoimmune disease. *Canine Genetics and Epidemiology* **2**, 17.
- Peeters, M.E., Haagen, A.J.V., Goedegebuure, S.A. & Wolvekamp, W.T.C. (1991) Dysphagia in Bouviers associated with muscular dystrophy; evaluation of 24 cases. *Veterinary Quarterly*, **13**, 65–73.
- Peeters, M.E. & Ubbink, G.J. (1994) Dysphagia-associated muscular dystrophy: a familial trait in the Bouvier des Flandres. *Veterinary Record*, **134**, 444–446.
- Pekow, C.A., Hinds, T.R., Maggio-Price, L. *et al.* (1992) Osmotic stress in red blood cells from Beagles with hemolytic anemia. *American Journal of Veterinary Research*, **53**, 1457–1461.
- Pelander, L., Ljungvall, I., Egenvall, A. *et al.* (2015) Incidence of and mortality from kidney disease in over 600,000 insured Swedish dogs. *Veterinary Record*, **176** (25), 656.
- Pellegrini, B., Acland, G.M., & Ray, J. (2002) Cloning and characterization of opticin cDNA: evaluation as a candidate for canine oculo-skeletal dysplasia. *Gene*, **282** (1–2), 121–131.
- Pellegrino, F.C., Pacheco, E.L. & Vazzoler, M.L. (2009) Hearing disorders in dogs: literature review and clinical study in Dogo Argentino. *Anales de Veterinaria de Murcia*, **25**, 5–20.
- Pemberton, T.J., Choi, S., Mayer, J.A. *et al.* (2014) A mutation in the canine gene encoding folliculin-interacting protein 2 (FNIP2) associated with a unique disruption in spinal cord myelination. *Glia*, **62** (1), 39–51.
- Pendergrass, T.W. & Hayes, H.M. (1975) Cryptorchism and related defects in dogs: epidemiologic comparison with man. *Teratology*, **12** (1), 51–55.
- Penderis, J., Calvin, J., Abramson, C. *et al.* (2007) L-2-hydroxyglutaric aciduria: characterisation of the molecular defect in a spontaneous canine model. *Journal of Medical Genetics*, **44** (5), 334–340.
- Peng, J.Y., Stephenson, N., Foley, J.E. *et al.* (2015) Incidence rates and risk factors for owner-reported adverse events following vaccination of dogs that did or did not receive a *Leptospira* vaccine. *Journal of the American Veterinary Medical Association*, **247** (10), 1139–1145.
- Pereira, J.D.S., Giannico, A.T., Rincón, E.M. *et al.* (2014) Pigmentary uveitis associated with iridociliary cysts in American Bulldogs. *Archives of Veterinary Science*, **19**, 15–21.
- Perez-Guisado, J. & Munoz-Serrano, A. (2009) Factors linked to territorial aggression in dogs. *Journal of Animal and Veterinary Advances*, **8** (7), 1412–1418.
- Perryman, L.E. (2004) Molecular pathology of severe combined immunodeficiency in mice, horses, and dogs. *Veterinary Pathology*, **41**, 95–100.
- Pesteanu-Somogyi, L.D., Radzai, C. & Pressler, B.M. (2006) Prevalence of feline infectious peritonitis in specific cat breeds. *Journal of Feline Medicine and Surgery*, **8** (1), 1–5.
- Petazzoni, M., Nicetto, T., Vezzoni, A., Piras, A. & Palmer, R. (2012) Treatment of pes varus using locking plate fixation in seven Dachshund dogs. *Veterinary and Comparative Orthopaedics and Traumatology*, **25**, 231–238.
- Peters, J.A. (1969) Canine mastocytoma: excess risk as related to ancestry. *Journal of the National Cancer Institute*, **42** (3), 435–443.
- Petersen, T., Proschowsky, H.F., Hardon, T., Rasch, S.N. & Fredholm, M. (2015) Prevalence and heritability of distichiasis in the English Cocker Spaniel. *Canine Genetics and Epidemiology*, **2**, 11.
- Petersen-Jones, S.M. & Entz, D.D. (2002) An improved DNA-based test for detection of the codon 616 mutation in the alpha cyclic GMP phosphodiesterase gene that causes progressive retinal atrophy in the Cardigan Welsh Corgi. *Veterinary Ophthalmology*, **5** (2), 103–106.
- Petersen-Jones, S.M., Entz, D.D. & Sargan, D.R. (1999) cGMP phosphodiesterase- α mutation causes progressive retinal atrophy in the Cardigan Welsh Corgi Dog. *Investigative Ophthalmology and Visual Science*, **40** (8), 1637–1644.
- Petersen-Jones, S.M., Forcier, J. & Mentzer, A.L. (2007) Ocular melanosis in the Cairn Terrier: clinical description and investigation of mode of inheritance. *Veterinary Ophthalmology*, **10** (1), 63–69.
- Peterson, M.E., Kintzer, P.P. & Kass, P.H. (1996) Pretreatment clinical and laboratory findings in dogs with hypoadrenocorticism: 225 cases (1979–1993). *Journal of the American Veterinary Medical Association*, **208** (1), 85–91.
- Patrick, S.W. (1996) The incidence of eye disease in dogs in a veterinary academic hospital: 1772 cases. *Journal of the South African Veterinary Association*, **67** (3), 108–110.

- Pettigrew, R., Fyee, J.C., Gregory, B.L. *et al.* (2007) CNS hypomyelination in Rat Terrier Dogs with congenital goiter and a mutation in the thyroid peroxidase gene. *Veterinary Pathology*, **44** (1), 50–56.
- Pfahler, S. & Distl, O. (2015) Effective population size, extended linkage disequilibrium and signatures of selection in the rare dog breed Lundehund. *PLoS One*, **10** (4), e0122680.
- Pfahler, S., Menzel, J., Brahm, R. *et al.* (2015) Prevalence and formation of primary cataracts and persistent hyperplastic tunica vasculosa lentis in the German Pinscher population in Germany. *Veterinary Ophthalmology*, **18** (2), 135–140.
- Pfeiffer, D.U. (2010) *Veterinary Epidemiology: an Introduction*. Wiley-Blackwell, Oxford.
- Pfeiffer, I. & Brenig, B. (2005) Frequency of the canine leucocyte adhesion deficiency (CLAD) mutation among Irish Red Setters in Germany. *Journal of Animal Breeding and Genetics*, **122** (2), 140–142.
- Philipp, U., Hamann, H., Mecklenberg, L. *et al.* (2005) Polymorphisms within the canine MLPH gene are associated with dilute coat colour in dogs. *BMC Genetics*, **6**, 34.
- Philipp, U., Vollmar, A., Häggström, J., Thomas, A. & Distl, O. (2012) Multiple loci are associated with dilated cardiomyopathy in Irish Wolfhounds. *PLoS One*, **7** (6), e36691.
- Phillips, D.E. & Harkin, K.R. (2003) Hypothyroidism and myocardial failure in two Great Danes. *Journal of the American Animal Hospital Association*, **39** (2), 133–1337.
- Phillips, J.C. (2007) Heritability and segregation analysis of osteosarcoma in the Scottish Deerhound. *Genomics*, **90** (3), 354.
- Phillips, J.C., Lembcke, L. & Chamberlin, T. (2010) A novel locus for canine osteosarcoma (OSA1) maps to CFA34, the canine orthologue of human 3q26. *Genomics*, **96** (4), 220–227.
- Picavet, P., Detilleux, J., Verschuren, S. *et al.* (2007) Analysis of 4495 canine and feline uroliths in the Benelux. A retrospective study: 1994–2004. *Journal of Animal Physiology and Animal Nutrition*, **91** (5–6), 247–251.
- Picco, F., Zini, E., Nett, C. *et al.* (2008) A prospective study on canine atopic dermatitis and food-induced allergic dermatitis in Switzerland. *Veterinary Dermatology*, **19** (3), 150–155.
- Picut, C.A. & Lewis, R.M. (1987a) Comparative pathology of canine hereditary nephropathies: an interpretive review. *Veterinary Research Communications*, **11** (6), 561–587.
- Picut, C.A. & Lewis, R.M. (1987b) Microscopic features of canine renal dysplasia. *Veterinary Pathology*, **24** (2), 156–163.
- Pidduck, H. & Webbon, P.M. (1978) The genetic control of Perthes' disease in Toy Poodles: a working hypothesis. *Journal of Small Animal Practice*, **19** (1–12), 729–733.
- Piek, C.J., Hazewinkel, H.A.W., Wolvekamp, W.T.C., Nap, R.C. & Mey, B.P. (1996) Long term follow-up of avascular necrosis of the femoral head in the dog. *Journal of Small Animal Practice*, **37** (1), 12–18.
- Piirsalu, K., McLean, R., Zuber, R. *et al.* (1994) Role of I-123 serum amyloid protein in the detection of familial amyloidosis in Oriental cats. *Journal of Small Animal Practice*, **35** (11), 581–586.
- Pinter, L., Jurak, Ž., Ukalović, M. & Sušić, V. (1999) Epidemiological and clinical features of dermatophytoses in dogs and cats in Croatia between 1990 and 1998. *Veterinarski Arhiv*, **69**, 261–270.
- Pipan, M., Brown, D., Battaglia, C. *et al.* (2012) An internet-based survey of risk factors for surgical gastric dilatation-volvulus in dogs. *Journal of the American Veterinary Medical Association*, **240** (12), 1456–1462.
- Pirie, C.G., Knollinger, A.M., Thomas, C.B. *et al.* (2006) Canine conjunctival hemangioma and hemangiosarcoma: a retrospective evaluation of 108 cases (1989–2004). *Veterinary Ophthalmology*, **9** (4), 215–226.
- Plant, J.D., Lund, E.M. & Yang, M. (2011) A case-control study of the risk factors for canine juvenile-onset generalized demodicosis in the USA. *Veterinary Dermatology*, **22** (1), 95–99.
- Platt, S., Freeman, J., di Stefani, A., Wieczorek, L. & Henley, W. (2006) Prevalence of unilateral and bilateral deafness in Border Collies and association with phenotype. *Journal of Veterinary Internal Medicine*, **20**, 1355–1362.
- Platt, S.R., Chambers, J.N. & Cross, A. (2004) A modified ventral fixation for surgical management of atlantoaxial subluxation in 19 dogs. *Veterinary Surgery*, **33** (4), 349–354.
- Plutchik, R. (1971) Individual and breed differences in approach and withdrawal in dogs. *Behaviour*, **40**, 302–311.

- Počta, S. & Svoboda, M. (2007) Incidence of canine hypersensitivity in the region of North Eastern Bohemia. *Acta Veterinaria Brno*, **76** (3), 451–459.
- Podberscek, A. and Blackshaw, J. (1991) Dog attacks on postal delivery officers in Queensland. *Australian Veterinary Journal*, **68**, 215–216.
- Poldervaart, J.H., Favier, R.P., Penning, L.C., van den Ingh, T.S. & Rothuizen, J. (2009) Primary hepatitis in dogs: a retrospective review (2002–2006). *Journal of Veterinary Internal Medicine*, **23** (1), 72–80.
- Polisca, A., Troisi, A., Fontaine, E. *et al.* (2016) A retrospective study of canine prostatic diseases from 2002 to 2009 at the Alfort Veterinary College in France. *Theriogenology*, **85** (5), 835–840.
- Pollard, R.E., Koehne, A.L., Peterson, C.B. *et al.* (2015) Japanese Bobtail: vertebral morphology and genetic characterization of an established cat breed. *Journal of Feline Medicine and Surgery*, **17** (8), 719–726.
- Polton, G.A., Mowat, V., Lee, H.C., McKee, K.A. & Scase, T.J. (2006) Breed, gender and neutering status of British dogs with anal sac gland carcinoma. *Veterinary and Comparative Oncology*, **4** (3), 125–131.
- Polnet, C.M., Dupre, G.P., Freiche, V.G. *et al.* (2005) Prevalence of gastrointestinal tract lesions in 73 brachycephalic dogs with upper respiratory syndrome. *Journal of Small Animal Practice*, **46**, 273–279.
- Poole, C. (2001) Low P-values or narrow confidence intervals: which are more durable? *Epidemiology (Cambridge, Mass.)*, **12** (3), 291–294.
- Post, K., Van Haaften, B. & Okkens, A.C. (1991) Vaginal hyperplasia in the bitch: literature review and commentary. *Canadian Veterinary Journal*, **32** (1), 35–37.
- Potanas, C.P., Padgett, S. & Gamblin, R.M. (2015) Surgical excision of anal sac apocrine gland adenocarcinomas with and without adjunctive chemotherapy in dogs: 42 cases (2005–2011). *Journal of the American Veterinary Medical Association*, **246** (8), 877–884.
- Power, H. & Prélaud, P. (2011) The role of breed in canine skin disease. *Veterinary Focus*, **21** (3), 10–17.
- Power, J.W. (1976) Avulsion of the tibial tuberosity in the Greyhound. *Australian Veterinary Journal*, **52** (11), 491–495.
- Prélaud, P. & Guaguere, E. (1998) Diagnostic de la dermatite par allergie aux piqures de puces. *Pratique Médicale et Chirurgicale de l'Animal de Compagnie*, **33** (suppl), 373.
- Prémont, J.E., Monclin, S., Farnir, F. *et al.* (2012) Perilimbic pocket technique for surgical repositioning of prolapsed nictitans gland in dogs. *Veterinary Record*, **171** (10), 247.
- Priester, W.A. (1976) Canine intervertebral disc disease: occurrence by age, breed, and sex among 8,117 cases. *Theriogenology*, **6** (2/3), 293–303.
- Proschowsky, H.F., Rugbjerg, H. & Ersbøll, A.K. (2003a) Morbidity of purebred dogs in Denmark. *Preventive Veterinary Medicine*, **58** (1–2), 53–62.
- Proschowsky, H.F., Rugbjerg, H. & Ersbøll, A.K. (2003b) Mortality of purebred and mixed-breed dogs in Denmark. *Preventive Veterinary Medicine*, **58** (1–2), 63–74.
- Proverbio, D., Spada, E., Perego, R. *et al.* (2013) Assessment of blood types of Ragdoll cats for transfusion purposes. *Veterinary Clinical Pathology*, **42** (2), 157–162.
- Prymak, C., McKee, L.J., Goldschmidt, M.H. *et al.* (1988) Epidemiologic, clinical, pathologic and prognostic characteristics of splenic hemangiosarcoma and splenic haematoma: 217 cases (1985). *Journal of the American Veterinary Medical Association*, **193** (6), 706–712.
- Puerto, D.A., Brockman, D.J., Lindquist, C. & Drobatz, K. (2002) Surgical and nonsurgical management of and selected risk factors for spontaneous pneumothorax in dogs: 64 cases (1986–1999). *Journal of the American Veterinary Medical Association*, **220** (11), 1670–1674.
- Pugh, C.A., Bronsvoort, B.M. de C., Handel, I.G. (2016) Cumulative incidence and risk factors for limber tail in the Dogslife Labrador Retriever cohort. *Veterinary Record*, **179** (11), 275.
- Pujar, S., Kothapalli, K.S., Kirkness, E. *et al.* (2005) Exclusion of Lhx9 as a candidate gene for Sry-negative XX sex reversal in the American Cocker Spaniel model. *Journal of Heredity*, **96** (4), 452–454.
- Pujar, S. & Meyers-Wallen, V.N. (2009) A molecular diagnostic test for persistent müllerian duct syndrome in Miniature Schnauzer dogs. *Sexual Development*, **3** (6), 326–328.
- Qvigstad, G., Kolbjørnsen, O., Skancke, E. *et al.* (2008) Gastric neuroendocrine carcinoma associated with atrophic gastritis in the Norwegian Lundehund. *Journal of Comparative Pathology*, **139** (4), 194–201.

- Rachid, M.A., Demaula, C.D., Scott, D.W. *et al.* (2003) Concurrent follicular dysplasia and interface dermatitis in Boxer dogs. *Veterinary Dermatology*, **14**, 159–166.
- Radasch, R.M., Lewis, D.F., McDonald, D.E. *et al.* (2008) Pes varus correction in Dachshunds using a hybrid external fixator. *Veterinary Surgery*, **37**, 71–81.
- Raffan, E., Dennis, R.J., O'Donovan, C.J. *et al.* (2016) A deletion in the canine pomc gene is associated with weight and appetite in obesity-prone labrador retriever dogs. *Cell Metabolism*, **23** (5), 893–900.
- Raffan, P.J. (1993) A new surgical technique for repair of perineal hernias in the dog. *Journal of Small Animal Practice*, **34** (1), 13–19.
- Rah, H., Maggs, D.J., Blankenship, T.N. *et al.* (2005) Early-onset, autosomal recessive, progressive retinal atrophy in Persian cats. *Investigative Ophthalmology and Visual Science*, **46** (5), 1742–1747.
- Ralphs, S.C. & Beale, B. (2000) Canine idiopathic erosive polyarthritis. *Compendium on Continuing Education for the Practicing Veterinarian*, **22** (7), 671–677.
- Ramírez, G.A., Altamira, J. & Vilafranca, M. (2015) Cartilaginous tumors of the larynx and trachea in the dog. *Veterinary Pathology*, **52**, 1019–1026.
- Ramos, R.V., Monteiro-Steagall, B.P. & Steagall, P.V.M. (2014) Management and complications of anaesthesia during balloon valvuloplasty for pulmonic stenosis in dogs: 39 cases (2000 to 2012). *Journal of Small Animal Practice*, **55**, 207–212.
- Ramos-Vara, J.A., Beissenherz, M.E., Miller, M.A. *et al.* (2000) Retrospective study of 338 canine oral melanomas with clinical, histologic and immunohistochemical review of 129 cases. *Veterinary Pathology*, **37** (6), 597–608.
- Ranen, E., Dank, G., Lavy, E. *et al.* (2008) Oesophageal sarcomas in dogs: histological and clinical evaluation. *Veterinary Journal*, **178** (1), 78–84.
- Ravens, P.A., Xu, B.J., & Vogelnest, L.J. (2014) Feline atopic dermatitis: a retrospective study of 45 cases (2001–2012). *Veterinary Dermatology*, **25** (2), 95–e28.
- Raw, M.-E. & Gaskell, C.J. (1985) A review of one hundred cases of presumed canine epilepsy. *Journal of Small Animal Practice*, **26** (11), 645–652.
- Raymond, S.L., Jones, D.W., Brooks, M.B. & Dodds, W.J. (1990) Clinical and laboratory features of a severe form of von Willebrand disease in Shetland Sheepdogs. *Journal of the American Veterinary Medical Association*, **197** (10), 1342–1346.
- Raymond-Letron, I., Bourges-Abella, N., Rousseau, T. *et al.* (2012) Histopathologic features of canine distichiasis. *Veterinary Ophthalmology*, **15** (2), 92–97.
- RCVS Knowledge (2017). *RCVS Knowledge: Veterinary Knowledge Online*. <http://knowledge.rcvs.org.uk> (accessed October 2017).
- Read, R.A. & Broun, H.C. (2007) Entropion correction in dogs and cats using a combination Hotz–Celsus and lateral eyelid wedge resection: results in 311 eyes. *Veterinary Ophthalmology*, **10** (1), 6–11.
- Read, R.A., Black, A.P., Armstrong, S.J., MacPherson, G.C. & Peek, J. (1992) Incidence and clinical significance of sesamoid disease in Rottweilers. *Veterinary Record*, **130** (24), 533–535.
- Read, R.A., Wood J.L.N. & Lakhani, K.H. (1998) Pectinate ligament dysplasia (PLD) and glaucoma in Flat Coated Retrievers. I. Objectives, technique and results of a PLD survey. *Veterinary Ophthalmology*, **1** (2–3), 85–90.
- Reeves, M.P. (2004) A retrospective report of 90 dogs with suspected cane toad (*Bufo marinus*) toxicity. *Australian Veterinary Journal*, **82** (10), 608–611.
- Reichler, I.M., Eckrich Specker, C., Hubler, M. *et al.* (2012) Ectopic ureters in dogs: clinical features, surgical techniques and outcome. *Veterinary Surgery*, **41** (4), 515–522.
- Reichler, I.M., Hauser, B., Schiller, I. *et al.* (2001) Sebaceous adenitis in the Akita: clinical observations, histopathology and heredity. *Veterinary Dermatology*, **12** (5), 243–253.
- Reimer, M.E., Troy, G.C. & Warnick, L.D. (1999) Immune-mediated hemolytic anemia: 70 cases (1988–1996). *Journal of the American Animal Hospital Association*, **35** (5), 384–391.
- Reimer, S.B., Kyles, A.E., Filipowicz, D.E. *et al.* (2004) Long-term outcome of cats treated conservatively or surgically for peritoneopericardial diaphragmatic hernia: 66 cases (1987–2002). *Journal of the American Veterinary Medical Association*, **224**, 728–732.
- Reineke, E.L., Burkett, D.E. & Drobatz, K.J. (2008) Left atrial rupture in dogs: 14 cases (1990–2005). *Journal of Veterinary Emergency and Critical Care*, **18** (2), 158–164.

- Reisner, I.R., Houpt, K.A. & Shofer, F.S. (2005) National survey of owner-directed aggression in English Springer Spaniels. *Journal of the American Veterinary Medical Association*, **227** (10), 1594–1603.
- Rest, J.R., Forrester, D. & Hopkins, J.N. (1996) Familial vasculopathy of German Shepherd Dogs. *Veterinary Record*, **138** (6), 144.
- Rettenmaier, J.L., Keller, G.G., Lattimer, J.C., Corley, E.A. & Ellersieck, M.R. (2002) Prevalence of canine hip dysplasia in a veterinary teaching hospital population. *Veterinary Radiology and Ultrasound*, **43** (4), 313–318.
- Reusch, C., Hoerauf, A., Lechner, J. *et al.* (1994) A new familial glomerulopathy in Bernese Mountain dogs. *Veterinary Record*, **134** (16), 411–415.
- Reusch, C.E. & Feldman E.C. (1991) Canine hyperadrenocorticism due to adrenocortical neoplasia. Pretreatment evaluation of 41 dogs. *Journal of Veterinary Internal Medicine*, **5** (1), 3–10.
- Rhodes, T.H., Vite, C.H., Giger, U., Patterson, D.F., Fahlke, C. & George, A.L. (1999) A missense mutation in canine CLC-1 causes recessive myotonia congenita in the dog 1. *FEBS Letters*, **456** (1), 54–58.
- Ricciarelli, A. (2016) Evaluation of behavioral characteristics and diseases incidence of the English Bulldog. *Dog Behavior*, **2**, 13–20.
- Richter, K.P., Kallet, A.J., Feldman, E.C. *et al.* (1990) Primary hyperparathyroidism in seven cats. *Journal of Veterinary Internal Medicine*, **4** (2), 115.
- Richtsmeier, J.T., Sack, G.H., Grausz, H.M. *et al.* (1994) Cleft palate with autosomal recessive transmission in Brittany Spaniels. *Cleft Palate Cranio-facial Journal*, **31** (5), 364–371.
- Riecks, T.W., Birchard, S.J. & Stephens, J.A. (2007) Surgical correction of brachycephalic syndrome in dogs: 62 cases (1991–2004). *Journal of the American Veterinary Medical Association*, **230** (9), 1324–1328.
- Riehl, J., Okura, M., Mignot, E. *et al.* (2000) Inheritance of von Willebrand's disease in a colony of Doberman Pinschers. *American Journal of Veterinary Research*, **61** (2), 115–120.
- Riis, R.C., Cummings, J.F., Loew, E.R. & de Lahunta, A. (1992) Tibetan Terrier model of canine ceroid lipofuscinosis. *American Journal of Medical Genetics*, **42** (4), 615–621.
- Riond, B., Schuler, E., Rogg, E., Hofmann-Lehmann, R. & Lutz, H. (2011) Prevalence of dog erythrocyte antigen 1.1 in dogs in Switzerland evaluated with the gel column technique. *Schweizer Archiv für Tierheilkunde*, **153**, 369–374.
- Risotto, K., Villamil, J.A., Selting, K.A. *et al.* (2011) Recent trends in feline intestinal neoplasia: an epidemiologic study of 1,129 cases in the Veterinary Medical Database from 1964 to 2004. *Journal of the American Animal Hospital Association*, **47** (1), 28–36.
- Rivas, A.L., Tintle, L., Argentieri, D. *et al.* (1995) A primary immunodeficiency syndrome in Shar-Pei dogs. *Clinical Immunology and Immunopathology*, **74** (3), 243–251.
- Rivas, A.L., Tintle, L., Meyers-Wallen, V. *et al.* (1993) Inheritance of renal amyloidosis in Chinese Shar-pei dogs. *Journal of Heredity*, **84** (6), 438–442.
- Roach, W. & Hecht, S. (2007) What is your diagnosis? Gastroesophageal intussusception. *Journal of the American Veterinary Medical Association*, **231** (3), 381–382.
- Robat, C.S., Cesario, L., Gaeta, R. *et al.* (2013) Clinical features, treatment options, and outcome in dogs with thymoma: 116 cases (1999–2010). *Journal of the American Veterinary Medical Association*, **243** (10), 1448–1454.
- Robbins, G.R. (1965) Unilateral renal agenesis in the beagle. *Veterinary Record*, **77** (46), 1345–1347.
- Roberts, S.M., Severin, G.A. & Lavach, J.D. (1986) Prevalence and treatment of palpebral neoplasms in the dog: 200 cases (1975–1983). *Journal of the American Veterinary Medical Association*, **189** (10), 1355–1359.
- Robertson, J.J. (1984) Perineal hernia repair in dogs. *Modern Veterinary Practice*, **65**, 365–368.
- Robinson, E.P., Sams, R.A. & Muir, W.W. (1986) Barbuturate anaesthesia in Greyhound and mixed-breed dogs: comparative cardiopulmonary effects, anesthetic effects, and recovery rates. *American Journal of Veterinary Research*, **47**, 2105–2112.
- Robinson, R. (1985) Chinese Crested Dog. *Journal of Heredity*, **76**, 217–218.
- Robinson, R. (1992) Legg–Calvé–Perthes disease in dogs: Genetic aetiology. *Journal of Small Animal Practice*, **33** (6), 275–276.

- Rodarte-Almeida, A.C.V., Petersen-Jones, S., Langohr, I.M. *et al.* (2016) Retinal dysplasia in American Pit Bull Terriers: phenotypic characterization and breeding study. *Veterinary Ophthalmology*, **19**, 11–21.
- Roe, K., Pratt, A., Lulich, J., Osborne, C. & Syme, H.M. (2012) Analysis of 14,008 uroliths from dogs in the UK over a 10-year period. *Journal of Small Animal Practice*, **53** (11), 634–640.
- Roedler, F.S., Pohl, S. & Oechtering, G.U. (2013) How does severe brachycephaly affect dog's lives? Results of a structured preoperative owner questionnaire. *Veterinary Journal*, **198** (3), 606–610.
- Rogers, K.D., Jones, B., Roberts, L. *et al.* (2011) Composition of uroliths in small domestic animals in the United Kingdom. *Veterinary Journal*, **188** (2), 228–230.
- Rohdin, C., Gilliam, D., O'Leary, C.A. *et al.* (2015) A KCNJ10 mutation previously identified in the Russell group of terriers also occurs in Smooth-Haired Fox Terriers with hereditary ataxia and in related breeds. *Acta Veterinaria Scandinavica*, **57** (1), 26.
- Rolim, V.M., Casagrande, R.A., Watanabe, T.T. *et al.* (2012) Melanoma amelanótico em cães: estudo retrospectivo de 35 casos (2004–2010) e caracterização imuno-histoquímica. *Pesquisa Veterinária Brasileira*, **32**, 340–346.
- Roll, A. & Unshelm, J. (1997) Aggressive conflicts amongst dogs and factors affecting them. *Applied Animal Behaviour Science*, **52** (3), 229–242.
- Romagnoli, S.E. (1991) Canine cryptorchidism. *Veterinary Clinics of North America: Small Animal Practice*, **21** (3), 533–544.
- Romão, F.G., Leitão, L.M.M., Machado, L.H. de A. *et al.* (2011) Hypoadrenocorticism in dogs: review. *Clinica Veterinaria*, **16**, 86–92.
- Roperto, F., Cerundolo, R., Restucci, B., Vincensi, M.R. *et al.* (1995) Colour dilution alopecia (CDA) in ten Yorkshire Terriers. *Veterinary Dermatology*, **6** (4), 171–178.
- Ropstad, E.O., Narfström, K., Lingaas, F. *et al.* (2008) Functional and structural changes in the retina of wire-haired dachshunds with early-onset cone-rod dystrophy. *Investigative Ophthalmology and Visual Science*, **49** (3), 1106–1115.
- Roque, J.B., O'Leary, C.A., Kyaw-Tanner, M. *et al.* (2011) High allergen-specific serum immunoglobulin E levels in nonatopic West Highland White Terriers. *Veterinary Dermatology*, **22** (3), 257–266.
- Rørvik, A.M. (1993) Risk factors for humeral condylar fractures in the dog: a retrospective study. *Journal of Small Animal Practice*, **34** (6), 277–282.
- Rose, J.H., Kwiatkowska, M., Henderson, E.R. *et al.* (2014) The impact of demographic, social, and environmental factors on the development of steroid-responsive meningitis-arthritis (SRMA) in the United Kingdom. *Journal of Veterinary Internal Medicine*, **28** (4), 1199–1202.
- Rosenberger, J.A., Pablo, N.V. & Crawford, P.C. (2007) Prevalence of and intrinsic risk factors for appendicular osteosarcoma in dogs: 179 cases (1996–2005). *Journal of the American Veterinary Medical Association*, **231** (7), 1076–1080.
- Rosner, A.L. (2012) Evidence-based medicine: revisiting the pyramid of priorities. *Journal of Bodywork and Movement Therapies*, **16** (1), 42–49.
- Rostaher, A., Hofer-Inteeworn, N., Kummerle-Fraune, C., Fischer, N.M. & Favrot C. (2017) Triggers, risk factors and clinico-pathological features of urticaria in dogs: a prospective observational study of 24 cases. *Veterinary Dermatology*, **28** (1), 38–e9.
- Roth, A.M., Ekins, M.B., Waring, G.O. *et al.* (1981) Oval corneal opacities in Beagles. III Histochemical demonstration of stromal lipids without hyperlipidaemia. *Investigative Ophthalmology and Visual Science*, **21** (1 pt 1), 95–106.
- Rothstein, E., Scott, D.W., Miller, W.H. & Bagladi, M.S. (1998) A retrospective study of dysplastic hair follicles and abnormal melanization in dogs with follicular dysplasia syndromes or endocrine skin disease. *Veterinary Dermatology*, **9** (4), 235–241.
- Rothwell, T.L., Howlett, C.R., Middleton, D.A. *et al.* (1987) Skin neoplasms of dogs in Sydney. *Australian Veterinary Journal*, **64** (6), 161–164.
- Roudebush, P., Maslin, W.R., & Cooper, R.C. (1988) Canine tumoral calcinosis. *Compendium on Continuing Education for the Practicing Veterinarian*, **10** (10), 1162–1164.
- Ru, G., Terracini, B. & Glickman, L.T. (1998) Host related risk factors for canine osteosarcoma. *Veterinary Journal*, **156** (1), 31–39.

- Rudd, R., Whitehair, J. & Margolis, J. (1990) Results of management of osteochondritis dissecans of the humeral head in dogs: 44 cases (1982 to 1987). *Journal of the American Animal Hospital Association*, **26** (2), 173–178.
- Rudling, E.H., Schlamowitz, S., Pipper, C.B. *et al.* (2016) The prevalence of the electrocardiographic J wave in the Petit Basset Griffon Vendéen compared to 10 different dog breeds. *Journal of Veterinary Cardiology*, **18** (1), 26–33.
- Rudmann, D.G., Coolman, B.R., Perez, C.M. & Glickman, L.T. (1992) Evaluation of risk factors for blastomycosis in dogs: 857 cases (1980–1990). *Journal of the American Veterinary Medical Association*, **201** (11), 1754–1759.
- Rugbjerg, H., Proschowsky, H.F., Ersbøll, A.K. & Lund, J.D. (2003) Risk factors associated with interdog aggression and shooting phobias among purebred dogs in Denmark. *Preventive Veterinary Medicine*, **58**, 85–100.
- Rusbridge, C. (2005) Neurological diseases of the Cavalier King Charles Spaniel. *Journal of Small Animal Practice*, **46**, 265–272.
- Rusbridge, C. & Knowler, S. P. (2004) Inheritance of occipital bone hypoplasia (Chiari type I malformation) in Cavalier King Charles Spaniels. *Journal of Veterinary Internal Medicine*, **18**, 673–678.
- Rusbridge, C., Heath, S., Gunn-Moore, D.A. *et al.* (2010) Feline orofacial pain syndrome (FOPS): a retrospective study of 113 cases. *Journal of Feline Medicine and Surgery*, **12** (6), 498–508.
- Rusbridge, C., Knowler, S.P., Pieterse, L. *et al.* (2009) Chiari-like malformation in the Griffon Bruxellois. *Journal of Small Animal Practice*, **50**, 386–393.
- Russell, N.J., Bond, K.A., Robertson, I.D., Parry, B.W. & Irwin, P.J. (2006) Primary hypoparathyroidism in dogs: a retrospective study of 17 cases. *Australian Veterinary Journal*, **84** (8), 285–290.
- Ryan, C.T., Glass, E.N., Seiler, G., Zwingenberger, A.L. & Mai, W. (2014) Magnetic resonance imaging findings associated with lateral cerebral ventriculomegaly in English Bulldogs. *Veterinary Radiology and Ultrasound*, **55**, 292–299.
- Ryan, R., Gutierrez-Quintana, R., ter Haar, G. & De Decker, S. (2017) Prevalence of thoracic vertebral malformations in French Bulldogs, Pugs and English Bulldogs with and without associated neurological deficits. *Veterinary Journal*, **221**, 25–29.
- Sackett, D.L., Rosenberg, W.M.C., Gray, J.A.M., Haynes, R.B. & Richardson, W.S. (1996) Evidence based medicine: what it is and what it isn't. *British Medical Journal*, **312** (7023), 71–72.
- Sacks, J.J., Sinclair, L., Gilchrist, J., Golab, G.C. & Lockwood, R. (2000) Breeds of dogs involved in fatal human attacks in the United States between 1979 and 1998. *Journal of the American Veterinary Medical Association*, **217** (6), 836–840.
- Safra, N., Hitchens, P.L., Maverakis, E. *et al.* (2016) Serum levels of innate immunity cytokines are elevated in dogs with metaphyseal osteopathy (hypertrophic osteodystrophy) during active disease and remission. *Veterinary Immunology and Immunopathology*, **179** (0), 32–35.
- Safra, N., Johnson, E.G., Lit, L. *et al.* (2013) Clinical manifestations, response to treatment, and clinical outcome for Weimaraners with hypertrophic osteodystrophy: 53 cases (2009–2011). *Journal of the American Veterinary Medical Association*, **242** (9), 1260–1266.
- Safra, N., Pedersen, N.C., Wolf, Z. *et al.* (2011) Expanded dog leukocyte antigen (DLA) single nucleotide polymorphism (SNP) genotyping reveals spurious class II associations. *Veterinary Journal*, **189** (2), 220–226.
- Saito, M., Sharp, N.J.H., Kortz, G.D. *et al.* (2002) Magnetic resonance imaging features of lissencephaly in 2 Lhasa Apsos. *Veterinary Radiology and Ultrasound*, **43**, 331–337.
- Sakai, T., Harashima, T. & Yamamura, H. (1994) Two cases of hereditary quadriplegia and amblyopia in a litter of Irish Setters. *Journal of Small Animal Practice*, **35**, 221–223.
- Salas, Y., Márquez, A., Diaz, D. & Romero, L. (2015) Epidemiological study of mammary tumors in female dogs diagnosed during the period 2002–2012: a growing animal health problem. *PLoS One*, **10**, e0127381.
- Sale, C.S.H. & Williams, J.M. (2006) Results of transthoracic esophagotomy retrieval of esophageal foreign body obstructions in dogs: 14 cases (2000–2004). *Journal of the American Animal Hospital Association*, **42** (6), 450–456.

- Salmon Hillbertz, N.H.C., Isaksson, M., Karlsson, E.K. *et al.* (2007) Duplication of FGF3, FGF4, FGF19 and ORAOV1 causes hair ridge and predisposition to dermoid sinus in Ridgeback dogs. *Nature Genetics*, **39** (11), 1318–1320.
- Salvin, H.E., McGreevy, P.D., Sachdev, P.S. & Valenzuela, M.J. (2012) The effect of breed on age-related changes in behavior and disease prevalence in cognitively normal older community dogs, *Canis lupus familiaris*. *Journal of Veterinary Behavior: Clinical Applications and Research*, **7** (2), 61–69.
- Sanchez, R.F., Innocent, G., Mould, J. & Billson, F.M. (2007) Canine keratoconjunctivitis sicca: disease trends in a review of 229 cases. *Journal of Small Animal Practice*, **48** (4), 211–217.
- Sánchez-Molano, E., Woolliams, J.A., Blott, S.C. & Wiener, P. (2014) Assessing the impact of genomic selection against hip dysplasia in the Labrador Retriever dog. *Journal of Animal Breeding and Genetics*, **131** (2), 134–145.
- Sanchis-Mora, S., Pelligand, L., Thomas, C.L. *et al.* (2016) Dogs attending primary-care practice in England with clinical signs suggestive of Chiari-like malformation/syringomyelia. *Veterinary Record*, **179**, 436.
- Sande, R.D., Alexander, J.E., Spencer, G.R., Padgett, G.A. & Davis, W.C. (1982) Dwarfism in Alaskan Malamutes: a disease resembling metaphyseal dysplasia in human beings. *American Journal of Pathology*, **106**, 224–236.
- Sanders, D.N., Zeng, R., Wenger, D.A. *et al.* (2013) GM2 gangliosidosis associated with a HEXA missense mutation in Japanese Chin dogs: a potential model for Tay Sachs disease. *Molecular Genetics and Metabolism*, **108** (1), 70–75.
- Sanders, S.G., Bagley, R.S., Silver, G.M., Moore, M. & Tucker, R.L. (2004) Outcomes and complications associated with ventral screws, pins, and polymethyl methacrylate for atlantoaxial instability in 12 dogs. *Journal of the American Animal Hospital Association*, **40** (3), 204–210.
- Sansom, J. & Barnett, K.C. (1985) Keratoconjunctivitis sicca in the dog: a review of two hundred cases. *Journal of Small Animal Practice*, **26** (3), 121–131.
- Santilli, R.A., Bontempi, L.V. & Perego, M. (2011) Ventricular tachycardia in English Bulldogs with localised right ventricular outflow tract enlargement. *Journal of Small Animal Practice*, **52**, 574–580.
- Santilli, R.A., Critelli, M. & Toaldo, M.B. (2010) ECG of the month. Accessory atrioventricular pathway-mediated tachycardia. *Journal of the American Veterinary Medical Association*, **237** (10), 1142–1144.
- Santilli, R.A., Ramera, L., Toaldo, M.B. & Perego, M. (2014) Arrhythmogenic right ventricular cardiomyopathy in English Bulldog: an emerging disease. *Veterinaria (Cremona)*, **28**, 7–15.
- Sapienza, J.S., Simó, F.J. & Prades-Sapienza, A. (2000) Golden Retriever uveitis: 75 cases (1994–1999). *Veterinary Ophthalmology*, **3** (4), 241–246.
- Sapierzyński, R. (2007) Neoplasms of the urinary and genital systems in dogs and cats. Part III. Ovarian, uterine and vaginal tumors. *Zycie Weterynaryjne* **82**, 35–38.
- Sapierzyński, R. (2009) Otitis externa in dogs. *Medycyna Weterynaryjna*, **65**, 552–556.
- Sapierzyński, R., Malicka, E., Bielecki, W. *et al.* (2007a) Tumours of the urogenital system in dogs and cats. Retrospective review of 138 cases. *Polish Journal of Veterinary Sciences*, **10** (2), 97–103.
- Sapierzyński, R., Malicka, E., Bielecki, W. *et al.* (2007b) Nowotwory jamy ustnej u psów i kotów, retrospektywny przegląd 143 przypadków. *Medycyna Weterynaryjna*, **63**, 1196–1199.
- Sargan, D.R., Withers, D., Pettitt, L. *et al.* (2007) Mapping the mutation causing lens luxation in several terrier breeds. *Journal of Heredity*, **98** (5), 534–538.
- Saridomichelakis, M.N., Farmaki, R., Leontides, L.S. & Koutinas, A.F. (2007) Aetiology of canine otitis externa: a retrospective study of 100 cases. *Veterinary Dermatology*, **18**, 341–347.
- Saridomichelakis, M.N., Koutinas, A.F., Gioulekas, D. & Leontidis, L. (1999) Canine atopic dermatitis in Greece: clinical observations and the prevalence of positive intradermal test reactions in 91 spontaneous cases. *Veterinary Immunology and Immunopathology*, **69** (1), 61–73.
- Saroğlu, M., Devecioğlu, Y. & Altunatmaz, K. (2005) Fundoscopic normal variations of the retina in Turkish Sheepdogs and multifocal retinal dysplasia: A comparative study in Akbash and Kangal breeds. *Turkish Journal of Veterinary and Animal Sciences*, **29** (2), 551–556.

- Sartor, A.J., Bentley, A.M. & Brown, D.C. (2013) Association between previous splenectomy and gastric dilatation-volvulus in dogs: 453 cases (2004–2009). *Journal of the American Veterinary Medical Association*, **242** (10), 1381–1384.
- Sato, K., Agoh, H., Kaneshige, T., Hikasa, Y. & Kagota, K. (2000) Hypercholesterolemia in Shetland Sheepdogs. *Journal of Veterinary Medical Science*, **62** (12), 1297–1301.
- Saunders, G.K., Wood, P.A., Myers, R.K., Shell, L.G. & Carithers, R. (1988) GM1 gangliosidosis in Portuguese Water Dogs: Pathologic and biochemical findings. *Veterinary Pathology*, **25** (4), 265–269.
- Scansen, B.A., Simpson, E.M., López-Alvarez, J. *et al.* (2015) Pulmonary artery dissection in eight dogs with patent ductus arteriosus. *Journal of Veterinary Cardiology*, **17** (2), 107–119.
- Scanziani, E., Giusti, A.M., Gualtieri, M. & Fonda, D. (1991) Gastric carcinoma in the Belgian Shepherd Dog. *Journal of Small Animal Practice*, **32**, 465–469.
- Schaaf, O.R., Eaton-Wells, R. & Mitchell, R.S. (2009) Biceps brachii and brachialis tendon of insertion injuries in eleven Racing Greyhounds. *Veterinary Surgery*, **38** (7), 825–833.
- Schafer, G., Nolte, I., Reinhard, F. & Rudolph, R. (1982) Congenital patellar luxation in the dog. 1. Experimental studies. 2. Clinical features, aetiology and results of therapy. *Kleintierpraxis*, **27**, 121–130.
- Schalamon, J., Ainoedhofer, H., Singer, G. *et al.* (2006) Analysis of dog bites in children who are younger than 17 years. *Pediatrics*, **117** (3), e374–e379.
- Schantz, P.M., Steurer F.J., Duprey Z.H. *et al.* (2005) Autochthonous visceral leishmaniasis in dogs in North America. *Journal of the American Veterinary Medical Association*, **226** (8), 1316–1322.
- Schatzberg, S.J., Haley, N.J., Barr, S.C., de Lahunta, A. & Sharp, N.J.H. (2005) Polymerase chain reaction screening for DNA viruses in paraffin-embedded brains from dogs with necrotizing meningoencephalitis, necrotizing leukoencephalitis, and granulomatous meningoencephalitis. *Journal of Veterinary Internal Medicine*, **19** (4), 553–559.
- Schellenberg, D., Yi, Q., Glickman, N.W. *et al.* (1998) Influence of thoracic conformation and genetics on the risk of gastric dilatation-volvulus in Irish Setters. *Journal of the American Animal Hospital Association*, **34** (1), 64–73.
- Schermerhorn, T., Center, S.A., Dykes, N.L. *et al.* (1996) Characterization of hepatoportal microvascular dysplasia in a kindred of Cairn Terriers. *Journal of Veterinary Internal Medicine*, **10**, 219–230.
- Schick, R.O. & Fadok, V.A. (1986) Responses of atopic dogs to regional allergens: 268 cases (1981–1984). *Journal of the American Veterinary Medical Association*, **189** (11), 1493–1496.
- Schlensker, E. & Distl, O. (2012) Appearance and heredity of hemivertebra in dogs. *Praktische Tierarzt*, **93**, 14–23.
- Schlensker, E. & Distl, O. (2013) Prevalence, grading and genetics of hemivertebrae in dogs. *European Journal of Companion Animal Practice*, **23** (3), 119–123.
- Schlensker, E. & Distl, O. (2016) Heritability of hemivertebrae in the French Bulldog using an animal threshold model. *Veterinary Journal*, **207**, 188–189.
- Schlueter, C., Budras, K.D., Ludewig, E. *et al.* (2009) Brachycephalic feline noses: CT and anatomical study of the relationship between head conformation and the nasolacrimal drainage system. *Journal of Feline Medicine and Surgery*, **11** (11), 891–900.
- Schmutz, S.M. (2014) An analysis of the inheritance pattern of an adult-onset hearing loss in Border Collie dogs. *Canine Genetics and Epidemiology*, **1**, 6.
- Schmutz, S.M., Moker, J.S., Clark, E.G. & Shewfelt, R. (1998) Black hair follicular dysplasia, an autosomal recessive condition in dogs. *Canadian Veterinary Journal*, **39** (10), 644–646.
- Schneider, M., Hildebrandt, N., Schweigl, T. *et al.* (2001) Transvenous embolization of small patent ductus arteriosus with single detachable coils in dogs. *Journal of Veterinary Internal Medicine*, **15** (3), 222–228.
- Schneider, S.M., Cianciolo, R.E., Nabity, M.B. *et al.* (2013) Prevalence of immune-complex glomerulonephritides in dogs biopsied for suspected glomerular disease: 501 cases (2007–2012). *Journal of Veterinary Internal Medicine*, **27**, S67–S75.
- Schober, K.E. & Baade, H. (2006) Doppler echocardiographic prediction of pulmonary hypertension in West Highland White Terriers with chronic pulmonary disease. *Journal of Veterinary Internal Medicine*, **20** (4), 912–920.

- Schoning, P.R. & Cowan, L.A. (2000) Bald thigh syndrome of Greyhound dogs: gross and microscopic findings. *Veterinary Dermatology*, **11** (1), 49–51.
- Schrauwen, I., Barber, R.M., Schatzberg, S.J. *et al.* (2014) Identification of novel genetic risk loci in Maltese dogs with necrotizing meningoencephalitis and evidence of a shared genetic risk across toy dog breeds. *PLoS ONE*, **9** (11), e112755.
- Schrempp, D.R., Childress, M.O., Stewart, J.C. *et al.* (2013) Metronomic administration of chlorambucil for treatment of dogs with urinary bladder transitional cell carcinoma. *Journal of the American Veterinary Medical Association*, **242** (11), 1534–1538.
- Schrope, D.P. & Kelch, W.J. (2006) Signalment, clinical signs, and prognostic indicators associated with high-grade second- or third-degree atrioventricular block in dogs: 124 cases (January 1, 1997–December 31, 1997). *Journal of the American Veterinary Medical Association*, **228** (11), 1710–1717.
- Schultheiss, P.C. (2004) A retrospective study of visceral and nonvisceral hemangiosarcoma and hemangiomas in domestic animals. *Journal of Veterinary Diagnostic Investigation*, **16** (6), 522–526.
- Schultheiss, P.C. (2006) Histologic features and clinical outcomes of melanomas of lip, haired skin and nail bed locations of dogs. *Journal of Veterinary Diagnostic Investigation*, **18** (4), 422–425.
- Schultz, R.M., Johnson, E.G., Wisner, E.R. *et al.* (2008) Clinicopathologic and diagnostic imaging characteristics of systemic aspergillosis in 30 dogs. *Journal of Veterinary Internal Medicine*, **22** (4), 851–859.
- Schultz, R.M., Puchalski, S.M., Kent, M. & Moore, P.F. (2007) Skeletal lesions of histiocytic sarcoma in nineteen dogs. *Veterinary Radiology and Ultrasound*, **48** (6), 539–543.
- Schulze, C., Meyer, H.P., Blok, A.L. *et al.* (1998) Renal dysplasia in three young adult Dutch Kooiker dogs. *Veterinary Quarterly*, **20** (4), 146–148.
- Schwarz, P.D., Withrow, S.J., Curtis, C.R. *et al.* (1991) Mandibular resection as a treatment for oral cancer in 81 dogs. *Journal of the American Animal Hospital Association*, **27** (6), 601–610.
- Schwarz, T., Sullivan, M., Störk, C.K. *et al.* (2002) Aortic and cardiac mineralization in the dog. *Veterinary Radiology and Ultrasound*, **43** (5), 419–427.
- Scott, D.W. (1981) Observations on canine atopy. *Journal of the American Animal Hospital Association*, **17** (1), 91–100.
- Scott, D.W. (1986) Vitamin A-responsive dermatosis in the Cocker Spaniel. *Journal of the American Animal Hospital Association*, **22**, 125.
- Scott, D.W. & Anderson, W.I. (1990) Canine sebaceous gland tumors: a retrospective analysis of 172 cases. *Canine Practice*, **15** (1), 19–27.
- Scott, D.W. & Anderson, W.I. (1991) Canine hair follicle neoplasms: a retrospective analysis of 80 cases (1986–1987). *Veterinary Dermatology*, **2** (3–4), 143–150.
- Scott, D.W. & Buerger, R.G. (1988) Idiopathic calcinosis circumscripta in the dog: a retrospective analysis of 130 cases. *Journal of the American Animal Hospital Association*, **24**, 651.
- Scott, D.W. & Miller, W.H. (1996) Primary seborrhoea in English springer spaniels: A retrospective study of 14 cases. *Journal of Small Animal Practice*, **37** (4), 173–178.
- Scott, D.W. & Miller, W.H. (1999) Erythema multiforme in dogs and cats: literature review and case material from the Cornell University College of Veterinary Medicine (1988–96). *Veterinary Dermatology*, **10** (4), 297–309.
- Scott, D.W. & Miller, W.H. (2012a) Idiopathic nasodigital hyperkeratosis in dogs: a retrospective analysis of 35 cases (1988–1998). *Japanese Journal of Veterinary Dermatology*, **18**, 169–170.
- Scott, D.W. & Miller, W.H. (2012b) Localized demodicosis in dogs: a retrospective study of 46 cases (1988–1998). *Japanese Journal of Veterinary Dermatology*, **18**, 87–91.
- Scott, D.W. & Miller, W.H. (2013) Cutaneous food allergy in cats: a retrospective study of 48 cases (1988–2003). *Japanese Journal of Veterinary Dermatology*, **19** (4), 203–210.
- Scott, D.W. & Paradis, M. (1990) A survey of canine and feline skin disorders seen in a university practice: small Animal Clinic, University of Montreal, Saint-Hyacinthe, Quebec (1987–1988). *Canadian Veterinary Journal*, **31**, 830–835.
- Scott, D.W., Rouselle, S., & Miller, W.H. (1995) Symmetrical lupoid onychodystrophy in dogs: a retrospective analysis of 18 cases (1989–93). *Journal of the American Animal Hospital Association*, **31** (3), 194–200.

- Scott, D.W. & Walton, D. K. (1986) Hyposomatotropism in the mature dog: discussion of 22 cases. *Journal of the American Animal Hospital Association*, **22**, 467–473.
- Scott, E.M., Esson, D.W., Fritz, K.J. & Dubielzig, R.R. (2013) Major breed distribution of canine patients enucleated or eviscerated due to glaucoma following routine cataract surgery as well as common histopathologic findings within enucleated globes. *Veterinary Ophthalmology*, **16**, 64–72.
- Scott-Moncrieff, J.C., Snyder, P.W., Glickman, L.T. *et al.* (1992) Systemic necrotizing vasculitis in nine young Beagles. *Journal of the American Veterinary Medical Association*, **201**, 1553–1558.
- Seavers, A., Snow, D., Mason, K. & Malik, R. (2008) Evaluation of the thyroid status of Basenji dogs in Australia. *Australian Veterinary Journal*, **86**, 429–434.
- Seddon, J.M., Hampson, E.C.G.M., Smith, R.I.E. & Hughes, I.P. (2006) Genetic heterogeneity of day blindness in Alaskan Malamutes. *Animal Genetics*, **37**, 407–410.
- Seeliger, F., Leeb, T., Peters, M. *et al.* (2003) Osteogenesis imperfect in 2 litters of Dachshunds. *Veterinary Pathology*, **40**, 530–539.
- Segalini, V., Hericher, T., Grellet, A. *et al.* (2009) Thyroid function and infertility in the dog: a survey in five breeds. *Reproduction in Domestic Animals*, **44** (2), 211–213.
- Segev, G., Cowgill, L.D., Jessen, S. *et al.* (2012) Renal amyloidosis in dogs: a retrospective study of 91 cases with comparison of the disease between Shar-Pei and non-Shar-Pei dogs. *Journal of Veterinary Internal Medicine*, **26** (2), 259–268.
- Seiler, G.S., Rhodes, J., Cianciolo, R. & Casal, M.L. (2010) Ultrasonographic findings in Cairn Terriers with preclinical renal dysplasia. *Veterinary Radiology and Ultrasound*, **51**, 453–457.
- Seiler, R.J. (1979) Colorectal polyps of the dog: a clinicopathologic study of 17 cases. *Journal of the American Veterinary Medical Association*, **174**, 72–75.
- Seim-Wikse, T., Jörundsson, E., Nødtvedt, A. *et al.* (2013) Breed predisposition to canine gastric carcinoma: a study based on the Norwegian canine cancer register. *Acta Veterinaria Scandinavica*, **55** (1), 25.
- Selby, L.A., Becker, S.V. & Hayes, H.W. (1981) Epidemiologic risk factors associated with canine systemic mycoses. *American Journal of Epidemiology*, **113**, 133–139.
- Selby, L.A., Hayes, H.M. & Becker, S.V. (1979) Epizootiologic features of canine hydrocephalus. *American Journal of Veterinary Research*, **40** (3), 411–413.
- Selmic, L.E., Nelson, D.A., Saunders, A.B., Hobson, H.P. & Saunders, W.B. (2013) An intrapericardial technique for PDA ligation: surgical description and clinical outcome in 35 dogs. *Journal of the American Animal Hospital Association*, **49** (1), 31–40.
- Seppälä, E.H., Reuser, A.J.J. & Lohi, H. (2013) A nonsense mutation in the acid α -glucosidase gene causes Pompe disease in Finnish and Swedish Lapphunds. *PLoS One*, **8** (2), e56825.
- Serfass, P., Chetboul, V., Sampedrano, C.C. *et al.* (2006) Retrospective study of 942 small-sized dogs: Prevalence of left apical systolic heart murmur and left-sided heart failure, critical effects of breed and sex. *Journal of Veterinary Cardiology*, **8** (1), 11–18.
- Serres, F., Chetboul, V., Tissier, R. *et al.* (2007) Chordae tendineae rupture in dogs with degenerative mitral valve disease: prevalence, survival, and prognostic factors (114 Cases, 2001–2006). *Journal of Veterinary Internal Medicine*, **21** (2), 258–264.
- Serres, F., Chetboul, V., Tissier, R. *et al.* (2008) Comparison of 3 ultrasound methods for quantifying left ventricular systolic function: correlation with disease severity and prognostic value in dogs with mitral valve disease. *Journal of Veterinary Internal Medicine*, **22** (3), 566–577.
- Seton, R. & Seton, J. (1990) Canine parvovirus in Rottweiler pups. *Australian Veterinary Practitioner*, **20** (4).
- Sevelius, E., Andersson, M. & Jönsson, L. (1994) Hepatic accumulation of alpha-1-antitrypsin in chronic liver disease in the dog. *Journal of Comparative Pathology*, **111** (4), 401–412.
- Sharman, J.R. & Wilton, A.N. (2007) Elimination of neutrophil elastase and the genes for (corrected) adaptor protein complex 3 subunits (corrected) as the cause of trapped neutrophil syndrome in Border Collies. *Animal Genetics*, **38** (2), 188–189.
- Shaw, S.C., Wood, J.L., Freeman J. *et al.* (2004) Estimation of heritability of atopic dermatitis in Labrador and Golden Retrievers. *American Journal of Veterinary Research*, **65** (7), 1014–1020.

- Shearin, A.L. & Ostrander, E.A. (2010) Canine morphology: hunting for genes and tracking mutations. *PLoS Biology*, **8** (3), e1000310.
- Shearman, J.R., Cook, R.W., McCowan, C. *et al.* (2011) Mapping cerebellar abiotrophy in Australian Kelpies. *Animal Genetics*, **42** (6), 675–678.
- Sheerer, K.N., Couto, C.G., Marin, L.M. *et al.* (2013) Haematological and biochemical values in North American Scottish Deerhounds. *Journal of Small Animal Practice*, **54** (7), 354–360.
- Shell, L.G., Berezowski, J., Rishniw, M. *et al.* (2015) Clinical and breed characteristics of idiopathic head tremor syndrome in 291 dogs: a retrospective study. *Veterinary Medicine International*, **2015**, 165463.
- Shell, L.G., Potthoff, A.I., Katherman, A. *et al.* (1989) Neuronal-visceral GM1 gangliosidosis in Portuguese Water Dogs. *Journal of Veterinary Internal Medicine*, **3** (1), 1–7.
- Shelton, G.D. (1999) Neuromuscular disorders affecting young dogs and cats. *Veterinary Neurology and Neurosurgery Journal*, **1**, 1.
- Shelton, G.D., Johnson, G.C., O'Brien, D.P. *et al.* (2012) Degenerative myelopathy associated with a missense mutation in the superoxide dismutase 1 (SOD1) gene progresses to peripheral neuropathy in Pembroke Welsh Corgis and Boxers. *Journal of the Neurological Sciences*, **318** (1–2), 55–64.
- Shelton, G.D., Podel, L.M., Poncelet, L. *et al.* (2003) Inherited polyneuropathy in Leonberger dogs: a mixed or intermediate form of Charcot-Marie-Tooth disease? *Muscle Nerve*, **27**, 471–477.
- Shelton, G.D., Schule, A. & Kass, P.H. (1997) Risk factors for acquired myasthenia gravis in dogs: 1,154 cases (1991–1995). *Journal of the American Veterinary Medical Association*, **211** (11), 1428–1431.
- Shiel, R.E., Brennan, S.F. & Omodo-Eluk, A.J. (2007a) Thyroid hormone concentrations in young, healthy, pretraining Greyhounds. *Veterinary Record*, **161**, 616–619.
- Shiel, R.E., Brennan, S.F. & O'Rourke, L.G. (2007b) Hematologic values in young pretraining healthy Greyhounds. *Veterinary Clinical Pathology*, **36**, 274–277.
- Shiel, R.E., Kennedy, L.J., Nolan, C.M. *et al.* (2014) Major histocompatibility complex class II alleles and haplotypes associated with non-suppurative meningoencephalitis in Greyhounds. *Tissue Antigens*, **84** (3), 271–276.
- Shiel, R.E., Mooney, C.T., Brennan, S.F. *et al.* (2010a) Clinical and clinicopathological features of non-suppurative meningoencephalitis in young Greyhounds in Ireland. *Veterinary Record*, **167** (9), 333–337.
- Shiel, R.E., Sist, M., Nachreiner, R.F., Ehrlich, C.P. & Mooney, C.T. (2010b) Assessment of criteria used by veterinary practitioners to diagnose hypothyroidism in sighthounds and investigation of serum thyroid hormone concentrations in healthy Salukis. *Journal of the American Veterinary Medical Association*, **236** (3), 302–308.
- Shih, J.L., Keating, J.H., Freeman, L.M. *et al.* (2007) Chronic hepatitis in Labrador Retrievers: clinical presentation and prognostic factors. *Journal of Veterinary Internal Medicine*, **21** (1), 33–39.
- Shilby, S., Karl, S., Hittmair, M. *et al.* (2014) Acute gastroesophageal intussusception in a juvenile Australian Shepherd Dog: endoscopic treatment and long-term follow-up. *BMC Veterinary Research*, **10**, 109.
- Shiokawa, Y., Mori, T., Hoshino, Y. *et al.* (2013) Breed predilection of canine histiocytic sarcoma at Gifu University. *Journal of the Japan Veterinary Medical Association*, **66** (5), 321–324.
- Shoop, S., Marlow, S., Church, D. *et al.* (2015) Prevalence and risk factors for mast cell tumours in dogs in England. *Canine Genetics and Epidemiology*, **2** (1), 1.
- Short, A.D., Boag, A., Catchpole, B. *et al.* (2013) A candidate gene analysis of canine hypoadrenocorticism in 3 dog breeds. *Journal of Heredity*, **104** (6), 807–820.
- Short, A.D., Dunne, A., Lohi, H. *et al.* (2011) Characteristics of epileptic episodes in UK dog breeds: an epidemiological approach. *Veterinary Record*, **169** (2), 48.
- Short, A.D., Mellersh, C.S., Platt, H. *et al.* (2010) Exonic mutations in the L2HGDH gene in Staffordshire Bull Terriers. *Veterinary Record*, **167** (12), 455–457.
- Sidjanin, D.J., Lowe, J.K., McElwee, J.L. *et al.* (2002) Canine CNGB3 mutations establish cone degeneration as orthologous to the human achromatopsia locus ACHM3. *Human Molecular Genetics*, **11** (16), 1823–1833.

- Silva-Hidalgo, G., Juárez-Barranco, F., López-Valenzuela, M. & Dávila-Paredes, M. (2015) Carcinoma de células escamosas en caninos de Culiacán, Sinaloa, México: estudio retrospectivo (2006–2014). *Revista Científica*, **25** (4), 304–310.
- Silvestre-Ferreira, A.C., Pastor, J., Almeida, O. *et al.* (2004) Frequencies of feline blood types in northern Portugal. *Veterinary Clinical Pathology*, **33**, 240–243.
- Simmerson, S.M., Armstrong, P.J., Wünschmann, A. *et al.* (2014) Clinical features, intestinal histopathology, and outcome in protein-losing enteropathy in Yorkshire Terrier dogs. *Journal of Veterinary Internal Medicine*, **28** (2), 331–337.
- Sisó, S., Botteron, C., Muhie, A. *et al.* (2005) A novel leucodystrophy in a dog. *Journal of Comparative Pathology*, **132** (2–3), 232–236.
- Skelly, B.J. & Franklin, R.J.M. (2002) Recognition and diagnosis of lysosomal storage diseases in the cat and dog. *Journal of Veterinary Internal Medicine*, **16** (2), 133–141.
- Skelly, B.J., Sargan, D.R., Herrtage, M.E. & Winchester, B.G. (1996) The molecular defect underlying canine fucosidosis. *Journal of Medical Genetics*, **33** (4), 284–288.
- Skelly, B.J., Wallace, M., Rajpurohit, Y.R., Wang, P. & Giger, U. (1999) Identification of a 6 base pair insertion in West Highland White Terriers with erythrocyte pyruvate kinase deficiency. *American Journal of Veterinary Research*, **60** (9), 1169–1172.
- Skelly, C.M., McAllister, H. & Donnelly, W.J. (1997) Avulsion of the tibial tuberosity in a litter of Greyhound puppies. *Journal of Small Animal Practice*, **38** (10), 445–449.
- Skibild, E., Dahlgaard, K., Rajpurohit, Y., Smith, B.F. & Giger, U. (2001) Haemolytic anaemia and exercise intolerance due to phosphofructokinase deficiency in related Springer Spaniels. *Journal of Small Animal Practice*, **42** (6), 298–300.
- Slappendel, R.J., Beijer, E.G. & Van Leeuwen, M. (1998) Type III von Willebrand's disease in Dutch Kookier dogs. *Veterinary Quarterly*, **20**, 93–97.
- Slappendel, R.J., van der Gaag, I., van Nes, J.J. *et al.* (1991) Familial stomatocytosis-hypertrophic gastritis (FSHG), a newly recognised disease in the dog (Drentse patrijshond). *Veterinary Quarterly*, **13** (1), 30–40.
- Slappendel, R.J., van Zwieten, R., van Leeuwen, M. *et al.* (2005) Hereditary spectrin deficiency in Golden Retriever Dogs. *Journal of Veterinary Internal Medicine*, **19**, 187–192.
- Slater, M. R. & Erb, H. N. (1986) Effects of risk factors and prophylactic treatment on primary glaucoma in the dog. *Journal of the American Veterinary Medical Association*, **188**, 1028–1030.
- Slatter, D.H., Lavach, J.D., Severin, G.A. & Young, S. (1977) Überreiter's syndrome (chronic superficial keratitis) in dogs in the Rocky Mountain area: a study of 463 cases. *Journal of Small Animal Practice*, **18** (12), 757–772.
- Slocombe, R.F., Mitten, R. & Mason, T.A. (1989) Leucoencephalomyelopathy in Australian Rottweiler dogs. *Australian Veterinary Journal*, **66** (5), 147–150.
- Smith, B.F., Stedman, H., Rajpurohit, Y. *et al.* (1996a) Molecular basis of canine muscle type phosphofructokinase deficiency. *Journal of Biological Chemistry*, **271** (33), 20070–20074.
- Smith, F.O. (2006) Canine pyometra. *Theriogenology*, **66** (3), 610–612.
- Smith, M.O., Wenger, D.A., Hill, S.L. & Matthews, J. (1996b) Fucosidosis in a family of American-bred English Springer Spaniels. *Journal of the American Veterinary Medical Association*, **209** (12), 2088–2090.
- Smith, S.A., Tobias, A.H., Jacob, K.A. *et al.* (2003) Arterial thromboembolism in cats: acute crisis in 127 cases (1992–2001) and long-term management with low-dose aspirin in 24 cases. *Journal of Veterinary Internal Medicine*, **17** (1), 73–83.
- Snead, E.C.R., Taylor, S.M., van der Kooij, M. *et al.* (2015) Clinical phenotype of X-linked myotubular myopathy in Labrador Retriever puppies. *Journal of Veterinary Internal Medicine*, **29** (1), 254–260.
- Snelling, S.R. & Edwards, G.A. (2003) A retrospective study of unilateral arytenoid lateralisation in the treatment of laryngeal paralysis in 100 dogs (1992–2000). *Australian Veterinary Journal*, **81** (8), 464–468.
- Snyder, J.M., Shofer, F.S., Van Winkle, T.J. *et al.* (2006) Canine intracranial primary neoplasia: 173 cases (1986–2003). *Journal of Veterinary Internal Medicine*, **20**, 669–675.

- Solanki, J., Hasnani, J., Patel, D., Patel, P. & Raval, S. (2007) Canine demodicosis in Anand. *Journal of Veterinary Parasitology*, **21** (1), 79–80.
- Solanti, S., Laitinen, O. & Atroshi, F. (2002) Hereditary and clinical characteristics of lateral luxation of the superficial digital flexor tendon in Shetland Sheepdogs. *Veterinary Therapeutics*, **3** (1), 97.
- Somma, A.T., Moreno, J.C.D., Sato, M.T. *et al.* (2017) Characterization of a novel form of progressive retinal atrophy in Whippet dogs: a clinical, electroretinographic, and breeding study. *Veterinary Ophthalmology*, **20** (5), 450–459.
- Sommerlad, S.F., Morton, J.M., Haile-Mariam, M. *et al.* (2012) Prevalence of congenital hereditary sensorineural deafness in Australian Cattle Dogs and associations with coat characteristics and sex. *BMC Veterinary Research*, **8**, 202.
- Song, R.B., Glass, E.N. & Kent, M. (2016) Spina bifida, meningocele, and meningocele. *Veterinary Clinics of North America: Small Animal Practice*, **46** (2), 327–345.
- Song, R.B., Vite, C.H., Bradley, C.W. & Cross, J.R. (2013) Postmortem evaluation of 435 cases of intracranial neoplasia in dogs and relationship of neoplasm with breed, age, and body weight. *Journal of Veterinary Internal Medicine*, **27** (5), 1143–1152.
- Soontornvipart, K., Wangdee, C., Kalpravidh, M. *et al.* (2013) Incidence and genetic aspects of patellar luxation in Pomeranian dogs in Thailand. *Veterinary Journal*, **196** (1), 122–125.
- Soute, B.A., Ulrich, M.M., Watson, A.D. *et al.* (1992) Congenital deficiency of all vitamin K-dependent blood coagulation factors due to a defective vitamin K-dependent carboxylase in Devon Rex cats. *Thrombosis and Haemostasis*, **68** (5), 521–525.
- Spada, E., Antognoni, M.T., Proverbio, D. *et al.* (2015) Haematological and biochemical reference intervals in adult Maine Coon cat blood donors. *Journal of Feline Medicine and Surgery*, **17** (12), 1020–1027.
- Spada, E., Miglio, A., Proverbio, D. *et al.* (2014) Signalment and blood types in cats being evaluated as blood donors at two Italian university blood banks. *Veterinary Medicine International*, **2014** (0): 704836.
- Spadari, A., Spinella, G., Morini, M. *et al.* (2008) Sartorius muscle contracture in a German Shepherd Dog. *Veterinary Surgery*, **37** (2), 149–152.
- Sparkes, A.H., Gruffydd-Jones, T.J., Shaw, S.E., Wright, A.I. & Stokes, C.R. (1993) Epidemiological and diagnostic features of canine and feline dermatophytosis in the United Kingdom from 1956 to 1991. *Veterinary Record*, **133** (3), 57–61.
- Sparkes, A.H., Rogers, K., Henley, W.E. *et al.* (2006) A questionnaire-based study of gestation, parturition and neonatal mortality in pedigree breeding cats in the UK. *Journal of Feline Medicine and Surgery*, **8** (3), 145–157.
- Spaull, R.F.V. & Friend, E.J. (2014) Laryngeal collapse as a primary disease in English Bull Terriers. *Veterinary Record*, **175**, 510.
- Speeti, M., Stahls, A., Meri, S. *et al.* (2003) Upregulation of major histocompatibility complex class II antigens in hepatocytes in Doberman hepatitis. *Veterinary Immunology and Immunopathology*, **96** (1–2), 1–12.
- Speiss, B.M. (1994) Inherited eye disease in the Entlebucher Mountain dog. *Schweizer Archiv für Tierheilkunde*, **136** (3), 105–110.
- Speiss, B.M., Bolliger, J.O., Guscetti, F. *et al.* (1998) Multiple ciliary body cysts and secondary glaucoma in the Great Dane: a report of nine cases. *Veterinary Ophthalmology*, **1** (1), 41–45.
- Sponenberg, D.P. & de Lahunta, A. (1981) Hereditary hypertrophic neuropathy in Tibetan Mastiff dogs. *Journal of Heredity*, **72** (4), 287–287.
- Sponenberg, D.P. & Graf-Webster, E. (1986) Hereditary meningoencephalocoele in Burmese cats. *Journal of Heredity*, **77** (1), 60.
- Spurling, N.W., Burton, L.K., Peacock, R. & Pilling, T. (1972) Hereditary factor-VII deficiency in the Beagle. *British Journal of Haematology*, **23**, 59–67.
- Spurling, N.W., Peacock, R. & Pilling, T. (1974) The clinical aspects of canine factor-VII deficiency including some case histories. *Journal of Small Animal Practice*, **15**, 229–239.
- Srivastava, A.K., Singh, B., Srivastava, A.K., Sinha, N. & Sharma, A.K. (2009) Canine neoplasms: a study on occurrence and distribution pattern. *Indian Journal of Veterinary Surgery*, **30** (1), 45–46.

- Stafford Johnson, M., Martin, M., Binns, S. *et al.* (2004) A retrospective study of clinical findings, treatment and outcome in 143 dogs with pericardial effusion. *Journal of Small Animal Practice*, **45** (11), 546–552.
- Stahl, C., Wacker, C., Weber, U. *et al.* (2010) MRI features of gastrocnemius musculotendinopathy in herding dogs. *Veterinary Radiology and Ultrasound*, **51**, 380–385.
- Stannard, A.A. & Pulley, L.T. (1975) Intracutaneous cornifying epithelioma (keratoacanthoma) in the dog: a retrospective study of 25 cases. *Journal of the American Veterinary Medical Association*, **167** (5), 355–358.
- Starr, A.N., Famula, T.R., Markward, N.J. *et al.* (2007) Hereditary evaluation of multiple developmental abnormalities in the Havanese dog breed. *Journal of Heredity*, **98**, 510–517.
- Stefanello, D., Avallone, G., Ferrari, R. *et al.* (2011) Canine cutaneous perivascular wall tumors at first presentation: clinical behavior and prognostic factors in 55 cases. *Journal of Veterinary Internal Medicine*, **25** (6), 1398–1405.
- Steinberg, K.S., Van Winkle, T., Bell, J.S. *et al.* (2000) Cerebellar degeneration in Old English Sheepdogs. *Journal of the American Veterinary Medical Association*, **217**, 1162–1165.
- Steiss, J.E. (2002) Muscle disorders and rehabilitation in canines. *Veterinary Clinics of North America: Small Animal Practice*, **32** (1), 267–285.
- Stenner, V. J., MacKay, B., King, T. *et al.* (2007) Protothecosis in 17 Australian dogs and a review of the canine literature. *Medical Mycology*, **45**, 249–266.
- Stephenson, H.M., Fonfara, S., Lopez-Alvarez, J. *et al.* (2012) Screening for dilated cardiomyopathy in Great Danes in the United Kingdom. *Journal of Veterinary Internal Medicine*, **26** (5), 1140–1147.
- Stern, J.A., White, S.N., Lehmkuhi, L.B. *et al.* (2014) A single codon insertion in PICALM is associated with development of familial subvalvular aortic stenosis in Newfoundland dogs. *Human Genetics*, **133** (9), 1139–1148.
- Stern-Bertholtz, W., Sjostrom, L. & Hakanson, N.W. (2003) Primary secretory otitis media in the Cavalier King Charles Spaniel: a review of 61 cases. *Journal of Small Animal Practice*, **44**, 253–256.
- Stevenson, A.E. & Markwell, P.J. (2001) Comparison of urine composition of healthy Labrador Retrievers and Miniature Schnauzers. *American Journal of Veterinary Research*, **62** (11), 1782–1786.
- Stinckens, A., Georges, M. & Buys, N. (2011) Mutations in the myostatin gene leading to hypermuscularity in mammals: indications for a similar mechanism in fish? *Animal Genetics*, **42** (3), 229–234.
- Stock, K.F., Klein, S., Tellhelm, B. *et al.* (2011) Genetic analyses of elbow and hip dysplasia in the German Shepherd Dog. *Journal of Animal Breeding and Genetics*, **128** (3), 219–229.
- Stokol, T., Parry, B.W. & Mansell, P.D. (1995) von Willebrand's disease in Scottish Terriers in Australia. *Australian Veterinary Journal*, **72** (11), 404–407.
- Storengen, L.M. & Lingaas, F. (2015) Noise sensitivity in 17 dog breeds: Prevalence, breed risk and correlation with fear in other situations. *Applied Animal Behaviour Sciences*, **171** (0), 152–160.
- Storey, E.S., Grahn, B.H. & Alcorn, J. (2005) Multifocal chorioretinal lesions in Borzoi dogs. *Veterinary Ophthalmology*, **8** (5), 337–347.
- Störk, C.K., Petite, A.F., Norrie, R.A., Polton, G.A. & Rayward, R.M. (2009) Variation in position of the medial fabella in West Highland White Terriers and other dogs. *Journal of Small Animal Practice*, **50** (5), 236–240.
- Strafuss, A.C. (1976) Sebaceous gland carcinoma in dogs. *Journal of the American Veterinary Medical Association*, **169**, 325–326.
- Strain, G.M. (2004) Deafness prevalence and pigmentation and gender associations in dog breeds at risk. *Veterinary Journal*, **167**, 23–32.
- Strain, G.M., Clark, L.A., Wahl, J.M. *et al.* (2009) Prevalence of deafness in dogs heterozygous or homozygous for the merle allele. *Journal of Veterinary Internal Medicine*, **23** (2), 282–286.
- Strom, A.R., Hässig, M., Iburg, T.M. & Spiess, B.M. (2011a) Epidemiology of canine glaucoma presented to University of Zurich from 1995 to 2009. Part 1: Congenital and primary glaucoma (4 and 123 cases). *Veterinary Ophthalmology*, **14** (2), 121–126.
- Strom, A.R., Hässig, M., Iburg, T.M. & Spiess, B.M. (2011b) Epidemiology of canine glaucoma presented to University of Zurich from 1995 to 2009. Part 2: Secondary glaucoma (217 cases). *Veterinary Ophthalmology*, **14** (2), 127–132.

- Stull, J.W., Evason, M., Carr, A.P. & Waldner, C. (2008) Canine immune-mediated polyarthritis: clinical and laboratory findings in 83 cases in western Canada (1991–2001). *Canadian Veterinary Journal*, **49** (12), 1195–1203.
- Sturaro, E., Menegazzo, L., Piccinini, P. *et al.* (2006) Prevalence and genetic parameters for hip dysplasia in Italian population of purebred dogs. *Italian Journal of Animal Science*, **5**, 107–116.
- Sturges, B.K., Dickinson, P.J., Bollen, A.W. *et al.* (2008) Magnetic resonance imaging and histological classification of intracranial meningiomas in 112 dogs. *Journal of Veterinary Internal Medicine*, **22**, 586–595.
- Sturgess, C.P., Waters, L., Gruffydd-Jones, T.J. *et al.* (1997) Investigation of the association between whole blood and tissue taurine levels and the development of thoracic deformities in neonatal Burmese kittens. *Veterinary Record*, **141**, 566–570.
- Sueiro, F.A.R., Alessi, A.C. & Vassallo, J. (2004) Canine lymphomas: a morphological and immunohistochemical study of 55 cases, with observations on p53 immunoreexpression. *Journal of Comparative Pathology*, **131** (2–3), 207–213.
- Sullivan, M., Cox, F., Pead, M.J. *et al.* (1987a) Thyroid tumours in the dog. *Journal of Small Animal Practice*, **28** (6), 505–512.
- Sullivan, M., Lee, R., Fisher, E.W., Nash, A.S. & McCandlish, I.A. (1987b) A study of 31 cases of gastric carcinoma in dogs. *Veterinary Record*, **120** (4), 79–83.
- Surman, S., Couto, C.G., DiBartola, S.P. *et al.* (2014) Arterial blood pressure, proteinuria, and renal histopathology in clinically healthy retired Racing Greyhounds. *Journal of Veterinary Internal Medicine*, **26** (6), 1320–1329.
- Suwankong, N., Meij, B.P., Voorhout, G. *et al.* (2008) Review and retrospective analysis of degenerative lumbosacral stenosis in 156 dogs treated by dorsal laminectomy. *Veterinary and Comparative Orthopaedics and Traumatology*, **21**, 284–293.
- Suzuki, M., Uchida, K., Morozumi, M. *et al.* (2003) A comparative pathological study on canine necrotizing meningoencephalitis and granulomatous meningoencephalomyelitis. *Journal of Veterinary Medical Science*, **65** (11), 1233–1239.
- Švara, T., Gombač, M., Pogorevc, E. *et al.* (2014) A retrospective study of canine testicular tumours in Slovenia. *Slovenian Veterinary Research*, **24**, 81–88.
- Svensson, M., Olsén, L., Winkler, P.A. *et al.* (2016) Progressive retinal atrophy in the Polski Owczarek Nizinny dog: a clinical and genetic study. *Veterinary Ophthalmology*, **19** (3), 195–205.
- Syrjä, P., Heikkilä, H.P., Lilja-Maula, L. *et al.* (2013) The histopathology of idiopathic pulmonary fibrosis in West Highland White Terriers shares features of both non-specific interstitial pneumonia and usual interstitial pneumonia in man. *Journal of Comparative Pathology*, **149** (2–3), 303–313.
- Syrjä, P., Saari, S., Rajamäki, M. *et al.* (2009) Pulmonary histopathology in Dalmatians with familial acute respiratory distress syndrome (ARDS). *Journal of Comparative Pathology*, **141** (4), 254–259.
- Szczepanik, M., Wilkolek, P., Adamek, L. *et al.* (2012) Sebaceous adenitis in dogs. *Zapalenie gruczołow łojowych u psów*, **87** (4), 296–299.
- Taboada, J., Harvey, J.W., Levy, M.G. (1992) Seroprevalence of babesiosis in Greyhounds in Florida. *Journal of the American Veterinary Medical Association*, **200** (1), 47–50.
- Tafti, A.K., Hanna, P. & Bourque, A.C. (2005) Calcinosis circumscripta in the dog: A retrospective pathological study. *Journal of Veterinary Medicine. Series A*, **52** (1), 13–17.
- Takada, K., Kitamura, H., Takiguchi, M., Saito, M. & Hashimoto, A. (2002) Cloning of canine 21-hydroxylase gene and its polymorphic analysis as a candidate gene for congenital adrenal hyperplasia-like syndrome in Pomeranians. *Research in Veterinary Science*, **73** (2), 159–163.
- Takahashi, M., Tomiyasu, H., Hotta, E. *et al.* (2014) Clinical characteristics and prognostic factors in dogs with histiocytic sarcomas in Japan. *Journal of Veterinary Medical Science*, **76** (5), 661–666.
- Takanosu, M., Takanosu, T., Suzuki, H. *et al.* (2008) Incomplete dominant osteochondrodysplasia in heterozygous Scottish Fold cats. *Journal of Small Animal Practice*, **49**, 197–199.
- Takeuchi, Y. & Mori, Y. (2006) A comparison of the behavioral profiles of purebred dogs in Japan to profiles of those in the United States and the United Kingdom. *Journal of Veterinary Medical Science*, **68** (8), 789–796.

- Takeuchi, Y., Kaneko, F., Hashizume, C. *et al.* (2009) Association analysis between canine behavioural traits and genetic polymorphisms in the Shiba Inu breed. *Animal Genetics*, **40** (5), 616–622.
- Tamura, S., Tamura, Y., Uchida, K. *et al.* (2010) GM2 Gangliosidosis Variant 0 (Sandhoff-Like Disease) in a family of Toy Poodles. *Journal of Veterinary Internal Medicine*, **24** (5), 1013–1019.
- Tanaka, H., Nakayama, M. & Takase, K. (2003) Histiocytic ulcerative colitis in a French Bulldog. *Journal of Veterinary Medical Science*, **65** (3), 431–433.
- Tappin, S., Goodfellow, M., Peters, I. *et al.* (2012) Frequency of the mutant MDR1 allele in dogs in the UK. *Veterinary Record*, **171** (3), 72.
- Tarpataki, N., Pápa, K., Reiczigel, J., Vajdovich, P. & Vörös, K. (2006) Prevalence and features of canine atopic dermatitis in Hungary. *Acta Veterinaria Hungarica*, **54** (3), 353–366.
- Tauro, A., Addicott, D., Foale, R.D. *et al.* (2015) Clinical features of idiopathic inflammatory polymyopathy in the Hungarian Vizsla. *BMC Veterinary Research*, **11** (1), 97.
- Tavares, S.B., Magalhães, A., de Sousa, L. (2015) Labrador retrievers are more attracted to water than to social stimuli: A pilot study. *Journal of Veterinary Behaviour*, **10** (6), 465–470.
- Taylor, S., Minor, K., Shmon, C.L. *et al.* (2016) Border Collie collapse: owner survey results and veterinary description of videotaped episodes. *Journal of the American Animal Hospital Association*, **52**, 364–370.
- Taylor, S., Shmon, C., Su, L. *et al.* (2016) Evaluation of dogs with Border Collie collapse, including response to two standardized strenuous exercise protocols. *Journal of the American Animal Hospital Association*, **52**, 281–290.
- Taylor-Brown, F.E., Meeson, R.L., Brodbelt, D.C. *et al.* (2015) Epidemiology of cranial cruciate ligament disease diagnosis in dogs attending primary-care veterinary practices in England. *Veterinary Surgery*, **44** (6), 777–783.
- Tcherneva, T. & Giger, U. (2007) Molecular base of coagulation factor XI deficiency in Kerry Blue Terrier. *Bulgarian Journal of Veterinary Medicine*, **10** (4), 247–255.
- Teikari, S. (1996) A survey on the frequency and nature of seizures in Finnish Belgian Shepherd dogs. *Suomen Eläinlääkärilehti*, **102**, 697–699.
- Temwichitr, J., Leegwater, P.A.J. & Hazewinkel, H.A.W. (2010) Fragmented coronoid process in the dog: a heritable disease. *Veterinary Journal*, **185**, 123–129.
- Tengvall, K., Kierczak, M., Bergvall, K. *et al.* (2013) Genome-wide analysis in German Shepherd Dogs reveals association of a locus on CFA 27 with atopic dermatitis. *PLOS Genetics*, **9** (5), e1003475.
- Teske, E., Naan, E.C., van Dijk, E.M., Van Garderen, E. & Schalken, J.A. (2002) Canine prostate carcinoma: epidemiological evidence of an increased risk in castrated dogs. *Molecular and Cellular Endocrinology*, **197** (1–2), 251–255.
- Teske, E., Van Heerde, P., Rutteman, G.R. *et al.* (1994) Prognostic factors for treatment of malignant lymphoma in dogs. *Journal of the American Veterinary Medical Association*, **205** (12), 1722–1728.
- Tetas Pont, R., Matas Riera, M., Newton, R. & Donaldson, D. (2015) Corneal and anterior segment foreign body trauma in dogs: a review of 218 cases. *Veterinary Ophthalmology*, **19** (5), 386–397.
- Tevell, E.H., Bergvall, K. & Egenvall, A. (2008) Sebaceous adenitis in Swedish dogs, a retrospective study of 104 cases. *Acta Veterinaria Scandinavica*, **50**, 11.
- Thøfner, M.S., Stougaard, C.L., Westrup, U. *et al.* (2015) Prevalence and heritability of symptomatic syringomyelia in Cavalier King Charles Spaniels and long-term outcome in symptomatic and asymptomatic littermates. *Journal of Veterinary Internal Medicine*, **29**, 243–250.
- Thomas, R., Borst, L. & Rotroff, D. (2014) Genomic profiling reveals extensive heterogeneity in somatic DNA copy number aberrations of canine hemangiosarcoma. *Chromosome Research*, **22** (3), 305–319.
- Thompson, A.L., Scott-Moncrieff, J.C. & Anderson, J.D. (2007) Comparison of classic hypoadrenocorticism with glucocorticoid-deficient hypoadrenocorticism in dogs: 46 cases (1985–2005). *Journal of the American Veterinary Medical Association*, **230** (8), 1190–1194.
- Thornburg, L.P., Rottinghaus, G., Dennis, G. & Crawford, S. (1996) The relationship between hepatic copper content and morphologic changes in the liver of West Highland White Terriers. *Veterinary Pathology*, **33** (6), 656–661.
- Thornburg, L.P., Shaw, D., Dolan, M. *et al.* (1986) Hereditary copper toxicosis in West Highland White Terriers. *Veterinary Pathology*, **23** (2), 148–154.

- Thrusfield, M. (2007) *Veterinary Epidemiology*. Blackwell, Oxford.
- Thrusfield, M.V., Aitken, C.G.G. & Darke, P.G.G. (1985) Observations on breed and sex in relation to canine heart valve incompetence. *Journal of Small Animal Practice*, **26**, 709–717.
- Thumchai, R., Lulich, J., Osborne, C.A. *et al.* (1996) Epizootiologic evaluations of urolithiasis in cats: 3498 cases (1982–1992). *Journal of the American Veterinary Medical Association*, **208** (4), 547–551.
- Tidholm, A. (1997) Retrospective study of congenital heart defects in 151 dogs. *Journal of Small Animal Practice*, **38** (3), 94–98.
- Tidholm, A. & Jonsson, L. (1996) Dilated cardiomyopathy in the Newfoundland: a study of 37 cases (1983–1994). *Journal of the American Animal Hospital Association*, **32**, 465–470.
- Tidholm, A. & Jonsson, L. (1997) A retrospective study of canine dilated cardiomyopathy (189 cases). *Journal of the American Animal Hospital Association*, **33** (6), 544–550.
- Tidholm, A., Haggstrom, J., Borgarelli, M. *et al.* (2001) Canine idiopathic dilated cardiomyopathy. Part I: aetiology, clinical characteristics, epidemiology and pathology. *Veterinary Journal*, **162**, 92–107.
- Tillson, M.D. & Winkler, J.T. (2002) Diagnosis and treatment of portosystemic shunts in the cat. *Veterinary Clinics of North America: Small Animal Practice*, **32** (4), 881–899.
- Tipold, A. (1995) Diagnosis of inflammatory and infectious diseases of the central nervous system in dogs: a retrospective study. *Journal of Veterinary Internal Medicine*, **9** (5), 304–314.
- Tipold, A., Fatzer, R., Jaggy, A., Zurbriggen, A. & Vandevelde, M. (1993) Necrotizing encephalitis in Yorkshire Terriers. *Journal of Small Animal Practice*, **34** (12), 623–628.
- Tisdall, P.L., Hunt, G.B., Bellenger, C.R. *et al.* (1994) Congenital portosystemic shunts in Maltese and Australian Cattle dogs. *Australian Veterinary Journal*, **71** (6), 174–178.
- Tobias, K.M. & Rohrbach, B.W. (2003) Association of breed with the diagnosis of congenital portosystemic shunts in dogs: 2,400 cases (1980–2002). *Journal of the American Veterinary Medical Association*, **223** (11), 1636–1639.
- Tobias, K.M. (2003) Determination of inheritance of single congenital portosystemic shunts in Yorkshire Terriers. *Journal of the American Animal Hospital Association*, **39** (4), 385–389.
- Tognetti, R., Corazza, M. & Buonaccorsi, A. (1987) Considerazioni sui risultati dell'intradermoreazione in 230 cani atopici provenienti dalla regione Toscana. *Veterinaria (Cremona)*, **11**, 13–17.
- Tolar, E.L., Hendrix, D.V.H., Rohrbach, B.W. *et al.* (2006) Evaluation of clinical characteristics and bacterial isolates in dogs with bacterial keratitis: 97 cases (1993–2003). *Journal of the American Veterinary Medical Association*, **228** (1), 80–85.
- Toma, S. & Noli, C. (2005) Isotretinoin in the treatment of multiple benign pilomatixomas in a mixed breed dog. *Veterinary Dermatology*, **16** (5), 346–350.
- Tønnessen, R., Sverdrup Borge, K., Nødtvedt, A. & Indrebø, A. (2012) Canine perinatal mortality: a cohort study of 224 breeds. *Theriogenology*, **77** (9), 1788–1801.
- Tonokura, M., Fujita, K. & Nishina, S. (2007) Review of pathophysiology and clinical management of narcolepsy in dogs. *Veterinary Record*, **161**, 375–380.
- Tonomura, N., Ingegerd, E., Thomas, R. *et al.* (2015) Genome-wide association study identifies shared risk loci common to two malignancies in Golden Retrievers. *PLoS One*, **11** (2), e1004922.
- Torres, A.R., Cassle, S.E., Haymore, M. & Hill, R. C. (2014) Hematologic differences between Dachshunds and mixed breed dogs. *Veterinary Clinical Pathology*, **43**, 519–524.
- Torres, S.M.F., Brien, T.O. & Scott, D.W. (2002) Dermal arteritis of the nasal philtrum in a Giant Schnauzer and three Saint Bernard dogs. *Veterinary Dermatology*, **13** (5), 275–281.
- Torres de la Riva, G., Hart, B.L., Farver, T.B. *et al.* (2013) Neutering dogs: effects on joint disorders and cancers in Golden Retrievers. *PLoS One*, **8** (2), e55937.
- Torrez, C.V. & Hunt, G.B. (2006) Results of surgical correction of abnormalities associated with brachycephalic airway obstruction syndrome in dogs in Australia. *Journal of Small Animal Practice*, **47** (3), 150–154.
- Toulza, O., Center, S.A., Brooks, M.B. *et al.* (2006) Evaluation of plasma protein C activity for detection of hepatobiliary disease and portosystemic shunting in dogs. *Journal of the American Veterinary Medical Association*, **229** (11), 1761–1771.

- Townsend, W.M. & Gornik, K.R. (2013) Prevalence of uveal cysts and pigmentary uveitis in Golden Retrievers in three Midwestern states. *Journal of the American Veterinary Medical Association*, **243** (9), 1298–1301.
- Toyoda, K., Uchida, K., Matsuki, N. *et al.* (2010) Inflammatory myopathy with severe tongue atrophy in Pembroke Welsh Corgi dogs. *Journal of Veterinary Diagnostic Investigation*, **22** (6), 876–885.
- Trafny, D.J., Freeman, L.M., Bulmer, B.J. *et al.* (2012) Auscultatory, echocardiographic, biochemical, nutritional, and environmental characteristics of mitral valve disease in Norfolk Terriers. *Journal of Veterinary Cardiology*, **14** (1), 261–267.
- Trautmann, A. & Nolte, I. (2003) Dystocia in selected dog breeds: predispositions and circumstances. *Praktische Tierarzt*, **84**, 902–911.
- Trehou-Sechi, E., Tissier, R., Gouni, V. *et al.* (2012) Comparative echocardiographic and clinical features of hypertrophic cardiomyopathy in 5 breeds of cats: a retrospective analysis of 344 cases (2001–2011). *Journal of Veterinary Internal Medicine*, **26** (3), 532–541.
- Trepanier, L.A. (1999) Delayed hypersensitivity reactions to sulphonamides: syndromes, pathogenesis and management. *Veterinary Dermatology*, **10** (3), 241–248.
- Trepanier, L.A., Danhof, R., Toll, J. & Watrous, D. (2003) Clinical findings in 40 dogs with hypersensitivity associated with administration of potentiated sulfonamides. *Journal of Veterinary Internal Medicine*, **17** (5), 647–652.
- Tritschler, C., Mizukami, K., Raj, K. & Giger, U. (2016) Increased erythrocyte osmotic fragility in anemic domestic shorthair and purebred cats. *Journal of Feline Medicine and Surgery*, **18** (6), 462–470.
- Trivedi, S.R., Sykes, J.E., Cannon, M.S. *et al.* (2011) Clinical features and epidemiology of cryptococcosis in cats and dogs in California: 93 cases (1988–2010). *Journal of the American Veterinary Medical Association*, **239**, 357–369.
- Trostel, C.T., Pool, R.R. & McLaughlin, R.M. (2003) Canine lameness caused by developmental orthopedic diseases: panosteitis, Legg–Calvé–Perthes disease, and hypertrophic osteodystrophy. *Compendium on Continuing Education for the Practicing Veterinarian*, **25** (4), 282–293.
- Troyer, V. de & Schepper, J. de (1989) Is pyometra commoner and more serious in Chow Chow bitches? *Vlaams Diergeneeskundig Tijdschrift*, **58**, 73–76.
- Trzil, J.E. & Reinero, C.R. (2014) Update on feline asthma. *Veterinary Clinics of North America: Small Animal Practice*, **44** (1), 91–105.
- Tsai, K. L. & Murphy, K. E. (2006) Clinical and genetic assessments of hip joint laxity in the Boykin Spaniel. *Canadian Journal of Veterinary Research*, **70**, 148–150.
- Tsai, K.L., Noorai, R.E. & Starr-Moss, A.N. (2012) Genome-wide association studies for multiple diseases of the German Shepherd Dog. *Mammalian Genome*, **23** (1), 203–211.
- Tsai, K.L., Starr-Moss, A.N., Venkataraman, N. *et al.* (2013) Alleles of the major histocompatibility complex play a role in the pathogenesis of pancreatic acinar atrophy in dogs. *Immunogenetics*, **65** (7), 501–509.
- Tsan M.-F., Jones, T.G., Thornton, G.W. *et al.* (1972) Canine cystinuria: its urinary amino acid pattern and genetic analysis. *American Journal of Veterinary Research*, **33** (12), 2455–2461.
- Tuohy, J.L., Selmic, L.E., Worley, D.R. (2014) Outcome following curative-intent surgery for oral melanoma in dogs: 70 cases (1998–2011). *Journal of the American Veterinary Medical Association*, **245** (11), 1266–1273.
- Turney, C., Chong, N.H., Alexander, R.H. *et al.* (2007) Pathological and electrophysiological features of canine cone-rod dystrophy in the Miniature Long-Haired Dachshund. *Investigative Ophthalmology and Visual Science*, **48** (9), 95–106.
- Tuttle, J. (1980) Feline hyperaesthesia syndrome. *Journal of the American Veterinary Medical Association*, **176**, 1166–1169.
- Twedt, D.C., Sternlieb, I. & Gilbertson, S.R. (1979) Clinical, morphologic and chemical studies on copper toxicosis of Bedlington Terriers. *Journal of the American Veterinary Medical Association*, **175** (3), 269–275.
- Ubbink, G.J., Hazewinkel, H.A., van de Broek, J. & Rothuizen, J. (1999) Familial clustering and risk analysis for fragmented coronoid process and elbow joint incongruity in Bernese Mountain Dogs in the Netherlands. *American Journal of Veterinary Research*, **60**, 1082–1087.

- Uchida, E., Chambers, J.K., Nakashima, K. *et al.* (2016) Pathologic features of colorectal inflammatory polyps in Miniature Dachshunds. *Veterinary Pathology*, **53**, 833–839.
- Uhriková, I., Lačňáková, A., Tandlerová, K. *et al.* (2013) Haematological and biochemical variations among eight sighthound breeds. *Australian Veterinary Journal*, **91** (11), 452–459.
- Ulfeng, M., Bjørnstad, E. & Bjerkås, E. (2009) Comparison of Schirmer tear test values in three dog breeds and a description of keratoconjunctivitis sicca in the Chinese Crested Dog. *Norsk Veterinærtidsskrift*, **121**, 727–732.
- Ungar de Sá, M.F.M., Ungar de Sá, J.E.U., Bittencourt, D.V.V. *et al.* (2007) Retrospective study (1991–2005), of canine babesiosis cases in Salvador city and Metropolitan Region, Bahia. *Revista Brasileira de Saúde e Produção Animal*, **8**, 178–183.
- Unshelm, J., Rehm, N. & Heidenberger, E. (1993) The problem of dangerous dogs: investigation of incidents with dogs in a large city [Munich, 1986–1991]. *Deutsche Tierärztliche Wochenschrift*, **100**, 383–386, 388–389.
- Upjohn, M., Cobb, C., Monger, J. *et al.* (2010) Prevalence, molecular typing and risk factor analysis for *Giardia duodenalis* infections in dogs in a central London rescue shelter. *Veterinary Parasitology*, **172** (3/4), 341–346.
- Urfer, S.R. (2008) Right censored data ('cohort bias') in veterinary life span studies. *Veterinary Record*, **163** (15), 457–458.
- Urfer, S.R., Gaillard, C. & Steiger, A. (2007) Lifespan and disease predispositions in the Irish Wolfhound: a review. *Veterinary Quarterly*, **29** (3), 102–111.
- Urkasemsin, G. & Olby, N.J. (2015) Clinical characteristics of Scottie cramp in 31 cases. *Journal of Small Animal Practice*, **56** (4), 276–280.
- Urkasemsin, G., Linder, K.E., Bell, J.S., de Lahunta, A. & Olby, N.J. (2010) Hereditary cerebellar degeneration in Scottish Terriers. *Journal of Veterinary Internal Medicine*, **24** (3), 565–570.
- Vaden, S.L., Breitschwerdt, E.B., Henrikson, C.K. *et al.* (1991) Primary ciliary dyskinesia in Bichon Frise litter mates. *Journal of the American Animal Hospital Association*, **27**, 633–640.
- Vaden, S.L., Littman, M.P. & Cianciolo, R.E. (2013) Familial renal disease in soft-coated wheaten terriers. *Journal of Veterinary Emergency and Critical Care*, **23** (2), 174–183.
- Vaden, S.L., Sellon, R.K., Melgarejo, L.T. *et al.* (2000) Evaluation of intestinal permeability and gluten sensitivity in Soft-Coated Wheaten Terriers with familial protein-losing enteropathy, protein-losing nephropathy, or both. *American Journal of Veterinary Research*, **61** (5), 518–524.
- Vail, D.M., Withrow, S.J., Schwarz, P.D. *et al.* (1990) Perianal adenocarcinoma in the canine male: a retrospective study of 41 cases. *Journal of the American Animal Hospital Association*, **26** (3), 329–334.
- Vajner, L., Vortel, V. and Brejcha, A. (1997) Lymphocytic thyroiditis in beagle dogs in a breeding colony: histological findings. *Veterinarni Medicina*, **42**, 43–49.
- van den Broek, A.H.M., Else, R.W., Abercromby, R. & France, M. (1991) Spinal dysraphism in the Weimaraner. *Journal of Small Animal Practice*, **32** (5), 258–260.
- van den Ingh, T., Mandigers, P. & van Nes, J. (1998) A neuronal vacuolar disorder in young Rottweiler dogs. *Veterinary Record*, **142**, 245–247.
- van der Gaag, I. (1988) The histological appearance of large intestinal biopsies in dogs with clinical signs of large bowel disease. *Canadian Journal of Veterinary Research*, **52** (1), 75–82.
- van der Linde-Sipman, J.S. (1987) Dysplasia of the pectinate ligament and primary glaucoma in the Bouvier des Flandres dog. *Veterinary Pathology*, **24** (3), 4240–4249.
- van der Linde-Sipman, J.S., Niewold, T.A., Tooten, P.C. *et al.* (1997) Generalized AA-amyloidosis in Siamese and Oriental cats. *Veterinary Immunology and Immunopathology*, **56** (1–2), 1–10.
- van der Merwe, L.L., Christie, J., Clift S.J. & Dvir, E. (2012) Salivary gland enlargement and sialorrhoea in dogs with spirocercosis: a retrospective and prospective study of 298 cases. *Journal of the South African Veterinary Association*, **83** (1), 37–52.
- van der Merwe, L.L., Jacobson, L.S. & Pretorius, G. J. (2002) The breed prevalence of dog erythrocyte antigen 1.1 in the Onderstepoort area of South Africa and its significance in selection of canine blood donors. *Journal of the South African Veterinary Association*, **73** (2), 53–56.
- van der Woerd, A., Nasisse, M.P. & Davidson, M. G. (1992) Lens-induced uveitis in dogs: 151 cases (1985–1990). *Journal of the American Veterinary Medical Association*, **201**, 921–926.

- Vandeveldel, M., Braund, K.G., Walder, T.L. *et al.* (1978) Dysmyelination of the central nervous system in the Chow-Chow dog. *Acta Neuropathologica*, **42**, 211–215.
- Vandeweerd, J.-M., Kirschvink, N., Clegg, P., Vandenput, S., Gustin, P. & Saegerman, C. (2012) Is evidence-based medicine so evident in veterinary research and practice? History, obstacles and perspectives. *Veterinary Journal*, **191** (1), 28–34.
- van Geffen, C., Bavegems, V., Duchateau, L., De Roover, K. & Daminet, S. (2006) Serum thyroid hormone concentrations and thyroglobulin autoantibodies in trained and non-trained healthy Whippets. *Veterinary Journal*, **172** (1), 135–140.
- van Ham, L., Vandeveldel, M., Desmidt, M. *et al.* (1994) A tremor syndrome with a central axonopathy in Scottish Terriers. *Journal of Veterinary Internal Medicine*, **8** (4), 290–292.
- Vanhaesebrouck, A.E., Shelton, G.D., Garosi, L. *et al.* (2011) A novel movement disorder in related male Labrador Retrievers characterized by extreme generalized muscular stiffness. *Journal of Veterinary Internal Medicine*, **25** (5), 1089–1096.
- Vanhaesebrouck, A.E., Van Soens, I., Poncelet, L. *et al.* (2010) Clinical and electrophysiological characterization of myokymia and neuromyotonia in Jack Russell Terriers. *Journal of Veterinary Internal Medicine*, **24** (4), 882–889.
- van Oost, B.A., Versteeg, S.A. & Slappendel, R.J. (2004) DNA testing for type III von Willebrand disease in Dutch Kooiker dogs. *Journal of Veterinary Internal Medicine*, **18** (3), 262–268.
- van Rossum, M., van Austen, F.J., Rofina, J. *et al.* (2004) Analysis of cDNA sequences of feline SAAs. *Amyloid*, **11** (1), 38–43.
- van Steenbeek, F.G., Leegwater, P.A., van Sluijs, F.J., Heuven, H.C. & Rothuizen, J. (2009). Evidence of inheritance of portosystemic shunts in Irish Wolfhounds. *Journal of Veterinary Internal Medicine*, **23** (4), 950–952.
- van Steenbeek, F.G., van den Bossche, L., Leegwater, P.A. & Rothuizen, J. (2012) Inherited liver shunts in dogs elucidate pathways regulating embryonic development and clinical disorders of the portal vein, *Mammalian Genome*, **23**, 76–84.
- van Straten, G., Leegwater, P.A.J., de Vries, M. *et al.* (2005) Inherited congenital extrahepatic portosystemic shunts in Cairn Terriers. *Journal of Veterinary Internal Medicine*, **19** (3), 321–324.
- van Tongeren, S.E., van Vonderen, I.K., van Nes, J.J. & van den Ingh, T.S. (2000) Cerebellar cortical abiotrophy in two Portuguese Podenco littermates. *Veterinary Quarterly*, **22** (3), 172–174.
- Vasseur, P.B., Foley, P., Stevenson, S. *et al.* (1989) Mode of inheritance of Perthes' disease in Manchester Terriers. *Clinical Orthopaedics*, **244**, 281–292.
- Vegter, A.R., Kooistra, H.S., Van Sluijs, F.J. *et al.* (2010) Persistent mullerian duct syndrome in a Miniature Schnauzer dog with signs of feminization and a Sertoli cell tumour. *Reproduction in Domestic Animals*, **45** (3), 447–452.
- Venker-van Haagen, A. J., Hartman, W. & Goedegebuure, S. A (1978) Spontaneous laryngeal paralysis in young Bouviers. *Journal of the American Animal Hospital Association*, **14**, 714–720.
- VeNom Coding Group (2017). *VeNom Veterinary Nomenclature*. <http://www.venomcoding.org> (accessed October 2017).
- Venta, P.J., Li, J., Yuzbasiyan-Gurkan, V., Brewer, G.J. & Schall, W.D. (2000) Mutation causing von Willebrand's Disease in Scottish Terriers. *Journal of Veterinary Internal Medicine*, **14** (1), 10–19.
- Verfuurden, B., Wempe, F., Reinink, P. *et al.* (2011) Severe combined immunodeficiency in Frisian Water Dogs caused by a RAG1 mutation. *Genes and Immunity*, **12** (4), 310–313.
- Verma, A.K., Sinha, D.K. & Singh, B.R. (2011) Seroprevalence study on salmonellosis in apparently healthy dogs by enzyme-linked immunosorbent assay. *Indian Journal of Animal Sciences*, **81** (1), 3–5.
- Vernau, K., Napoli, E., Wong, S. *et al.* (2015) Thiamine Deficiency-mediated Brain Mitochondrial Pathology in Alaskan Huskies with mutation in *SLC19A3.1*. *Brain Pathology*, **25** (4), 441–453.
- Verschoof, J., Thiel, C., Henrich, M. *et al.* (2015) Gastrointestinal intussusception in the Maine Coon: a review of 19 cases. *Wiener Tierärztliche Monatsschrift*, **102** (1/2), 29–37.
- Verstraete, F.J.M., Zin, B.P., Kass, P.H., Cox, D.P. & Jordan, R.C. (2011) Clinical signs and histologic findings in dogs with odontogenic cysts: 41 cases (1995–2010). *Journal of the American Veterinary Medical Association*, **239** (11), 1470–1476.

- VetCompass (2017). *VetCompass: Health Surveillance for UK Companion Animals*. RVC Electronic Media Unit, London. <http://www.rvc.ac.uk/vetcompass> (accessed October 2017).
- Vianna, M.L., & Tobias, K. (2005) Atresia ani in the dog: a retrospective study. *Journal of the American Animal Hospital Association*, **41** (5), 317–322.
- Victoria, T., Rafi, M.A. & Wenger, D.A. (1996) Cloning of the canine GALC cDNA and identification of the mutation causing globoid cell leukodystrophy in West Highland White and Cairn Terriers. *Genomics*, **33** (3), 457–462.
- Vidgren, G., Vainio-Siukola, K., Honkasalo, S. *et al.* (2012) Primary hyperoxaluria in Coton de Tulear, *Animal Genetics*, **43**, 356–361.
- Vidoni, B., Sommerfeld-Stur, I. & Eisenmenger, E. (2006) Diagnostic and genetic aspects of patellar luxation in small and miniature breed dogs in Austria. *Companion Animal Practice*, **16**, 149.
- Viitmaa, R., Cizinauskas, S., Bergamasco, L.A. *et al.* (2007) Magnetic resonance imaging findings in Finnish Spitz dogs with focal epilepsy. *Journal of Veterinary Internal Medicine*, **20**, 305–310.
- Viitmaa, R., Cizinauskas, S., Orro, T. *et al.* (2013) Phenotype, inheritance characteristics, and risk factors for idiopathic epilepsy in Finnish Spitz dogs. *Journal of the American Veterinary Medical Association*, **243** (7), 1001–1009.
- Vilboux, T., Chaudieu, G., Jeannin, P. *et al.* (2008) Progressive retinal atrophy in the Border Collie: a new XLPRA. *BMC Veterinary Research*, **3** (4), 10.
- Villagómez, D.A. & Alonso, R.A. (1998) A distinct Mendelian autosomal recessive syndrome involving the association of anotia, palate agenesis, bifid tongue, and polydactyly in the dog. *Canadian Veterinary Journal*, **39** (10), 642–643.
- Villamil, J.A., Henry, C.J., Bryan, J.N. *et al.* (2011) Identification of the most common cutaneous neoplasms in dogs and evaluation of breed and age distributions for selected neoplasms. *Journal of the American Veterinary Medical Association*, **239** (7), 960–965.
- Villamil, J.A., Henry, C.J., Hahn, A.W. *et al.* (2009) Hormonal and sex impact on the epidemiology of canine lymphoma. *Journal of Cancer Epidemiology*, **2009**, 591753.
- Vitale, C.L. & Olby, N.J. (2007) Neurologic dysfunction in hypothyroid, hyperlipidemic Labrador Retrievers. *Journal of Veterinary Internal Medicine*, **21** (6), 1316–1322.
- Vite, C.H., Insko, E.K., Schotland, H.M., Panckeri, K. & Hendricks, J.C. (1997) Quantification of cerebral ventricular volume in English Bulldogs. *Veterinary Radiology Ultrasound*, **38**, 437–443.
- Vnuk, D., Lipar, M., Maticic, D. *et al.* (2008) Comparison of standard perineal herniorrhaphy and transposition of the internal obturator muscle for perineal hernia repair in the dog. *Veterinarski Arhiv*, **78** (3), 197.
- Volk, A.V., Belyavin, C.E., Varjonen, K. *et al.* (2010) *Malassezia pachydermatis* and *M. nana* predominate amongst the cutaneous mycobiota of Sphynx cats. *Journal of Feline Medicine and Surgery*, **12** (12), 917–922.
- Vollmar, A. (2000) The prevalence of cardiomyopathy in the Irish Wolfhound: a clinical study of 500 dogs. *Journal of the American Animal Hospital Association*, **36**, 125–132.
- Von Bomhard, W., Mauldin, E.A., Schmutz, S.M. *et al.* (2006) Black hair follicular dysplasia in Large Munsterlander dogs: clinical, histological and ultrastructural features. *Veterinary Dermatology*, **17**, 182–188.
- Voorbij, A.M.W.Y., van Steenbeek, F.G., Vos-Loohuis, M. *et al.* (2011) A contracted DNA repeat in LHX3 intron 5 is associated with aberrant splicing and pituitary dwarfism in German Shepherd Dogs. *PLoS One*, **6** (11), e27940.
- Vörös, K., Szilvási, V., Manczur, F. *et al.* (2015) Occurrence of mitral valve insufficiency in clinically healthy Beagle dogs, *Acta Veterinaria Hungarica*, **63**, 458–471.
- Vrabelova, D., Silvestrini, P., Ciudad, J. *et al.* (2011) Analysis of 2735 canine uroliths in Spain and Portugal. A retrospective study: 2004–2006. *Research in Veterinary Science*, **91** (2), 208–211.
- Wahl, J.M., Clark, L.A., Skalli, O. *et al.* (2008) Analysis of gene transcript profiling and immunobiology in Shetland Sheepdogs with dermatomyositis. *Veterinary Dermatology*, **19** (2), 52–58.
- Wakamatsu, N., Surdyk, K., Carmichael, K.P. & Brown, C.A. (2007) Histologic and ultrastructural studies of juvenile onset renal disease in four Rottweiler dogs. *Veterinary Pathology*, **44** (1), 96–100.

- Wakshlag, J.J., Rassnick, K.M., Malone, E.K. *et al.* (2011) Cross-sectional study to investigate the association between vitamin D status and cutaneous mast cell tumours in Labrador Retrievers. *British Journal of Nutrition*, **1**, S60–63.
- Waldman, L. (1995) Seasonal flank alopecia in Affenpinscher. *Journal of Small Animal Practice*, **36**, 272–273.
- Waldo, J.T. & Diaz, K.S. (2015) Development and validation of a diagnostic test for Ridge allele copy number in Rhodesian Ridgeback dogs. *Canine Genetics and Epidemiology*, **2** (1), 2.
- Wallace, M.E. & Palmer, A.C. (1984) Recessive mode of inheritance in myasthenia gravis in the Jack Russell Terrier. *Veterinary Record*, **114**, 350.
- Wallace, M.R., MacKay, E.O., Gelatt, K.N. *et al.* (2005) Inheritance of cataract in the Bichon Frise. *Veterinary Ophthalmology*, **8** (3), 203–205.
- Wallerström, B.I., Wågberg, T.I. & Lagergren, C.H. (1992) Cystine calculi in the dog: an epidemiological retrospective study. *Journal of Small Animal Practice*, **33**, 78–84.
- Wallin-Håkanson, B., Wallin-Håkanson, N. & Hedhammar, A. (2000) Collie eye anomaly in the Rough Collie in Sweden: genetic transmission and influence on offspring vitality. *Journal of Small Animal Practice*, **41** (6), 254–258.
- Walvoort, H.C. (1985) Glycogen storage disease type II in the Lapland dog. *Veterinary Quarterly*, **7**, 187–190.
- Wang, G.-D., Fan, R.-X., Zhai, W. *et al.* (2014) Genetic convergence in the adaptation of dogs and humans to the high-altitude environment of the Tibetan Plateau. *Genome Biology and Evolution*, **6** (8), 2122–2128.
- Wang, P., Zangerl, B., Werner, P., Mauldin, E.A. & Casal, M.L. (2011) Familial cutaneous lupus erythematosus (CLE) in the German Shorthaired Pointer maps to CFA18, a canine orthologue to human CLE. *Immunogenetics*, **63** (4), 197–207.
- Wangdee, C., Leegwater, P.A.J., Heuven, H.C.M. *et al.* (2014) Prevalence and genetics of patellar luxation in Kooiker dogs. *Veterinary Journal* **201** (3) 333–337.
- Ward, J.L., DeFrancesco, T.C., Tou, S.P. *et al.* (2016) Outcome and survival in canine sick sinus syndrome and sinus node dysfunction: 93 cases (2002–2014). *Journal of Veterinary Cardiology*, **18**, 199–212.
- Ware, W.A. & Hopper, D.L. (1999) Cardiac Tumors in Dogs: 1982–1995. *Journal of Veterinary Internal Medicine*, **13** (2), 95–103.
- Warland, J. & Dobson, J. (2013) Breed predispositions in canine mast cell tumour: a single centre experience in the United Kingdom. *Veterinary Journal*, **197** (2), 496–498.
- Waters, D.J. (2011) Aging research 2011: exploring the pet dog paradigm. *Institute of Laboratory Animal Research Journal*, **52** (1), 97–105.
- Watson, P., Simpson, K.W. & Bedford, P.G.C. (1993) Hypercholesterolaemia in Briards in the United Kingdom. *Research in Veterinary Science*, **54**, 80–85.
- Watson, P.J. (2004) Chronic hepatitis in dogs: a review of current understanding of the aetiology, progression and treatment. *Veterinary Journal*, **167** (3), 228–241.
- Watson, P.J., Roulois, A.J.A., Scase, T. *et al.* (2007) Prevalence and breed distribution of chronic pancreatitis at post-mortem examination in first-opinion dogs. *Journal of Small Animal Practice*, **48** (11), 609–618.
- Watson, P.J., Roulois, A.J.A., Scase, T.J., Irvine, R. & Herrtage, M.E. (2010) Prevalence of hepatic lesions at post-mortem examination in dogs and association with pancreatitis. *Journal of Small Animal Practice*, **51** (11), 566–572.
- Watson, P.J., Wotton, P., Eastwood, J. *et al.* (2006) Immunoglobulin deficiency in Cavalier King Charles Spaniels with pneumocystis pneumonia. *Journal of Veterinary Internal Medicine*, **20**, 523–527.
- Weaver, A.D. (1983) Survey with follow-up of 67 dogs with Sertoli cell tumours. *Veterinary Record*, **113** (5), 105–107.
- Webb, C.B., Twedt, D.C. & Meyer, D.J. (2002) Copper-associated liver disease in Dalmatians: a review of 10 dogs (1998–2001). *Journal of Veterinary Internal Medicine*, **16** (6), 665–668.

- Weinkle, T.K., Center, S.A., Randolph, J.F. *et al.* (2005) Evaluation of prognostic factors, survival rates, and treatment protocols for immune-mediated hemolytic anemia in dogs: 151 cases (1993–2002). *Journal of the American Veterinary Medical Association*, **226** (11), 1869–1880.
- Weir, J.A., Yager J.A., Caswel, J.L. *et al.* (1994) Familial cutaneous vasculopathy of German Shepherds: clinical, genetic and preliminary pathological and immunological studies. *Canadian Veterinary Journal*, **35**, 763–769.
- Weiss, D. (2005) Bone marrow necrosis in dogs: 34 cases (1996–2004). *Journal of the American Veterinary Medical Association*, **227** (2) 263–267.
- Weiss, D.J. (2007) Hemophagocytic syndrome in dogs: 24 cases (1996–2005). *Journal of the American Veterinary Medical Association*, **230** (5), 697–701.
- Weisse, C., Berent, A.C., Todd, K. *et al.* (2014) Endovascular evaluation and treatment of intrahepatic portosystemic shunts in dogs: 100 cases (2001–2011). *Journal of the American Veterinary Medical Association*, **244** (1) 78–94.
- Weisse, C., Soares, N., Beal, M. *et al.* (2005) Survival times in dogs with right atrial hemangiosarcoma treated by means of surgical resection with or without adjuvant chemotherapy: 23 cases (1986–2000) *Journal of the American Veterinary Medical Association*, **226** (4) 575–579.
- Weissenböck, H., Obermaier, G. and Dahme, E. (1996) Alexander's disease in a Bernese Mountain Dog. *Acta Neuropathologica*, **91**, 200–204.
- Weissl, J., Hülsmeier, V., Brauer, C. *et al.* (2012) Disease progression and treatment response of idiopathic epilepsy in Australian Shepherd Dogs. *Journal of Veterinary Internal Medicine*, **26**, 116–125.
- Welle, M., Grimm, S., Suter, M. & Von Tschanner, C. (1999) Mast cell density and subtypes in the skin of Shar Pei Dogs with cutaneous mucinosis. *Journal of Veterinary Medicine. Series A*, **46** (5), 309–316.
- Wenger, M., Hauser, B., Ohlerth, S. *et al.* (2005) Thyroid neoplasms in dogs: a retrospective study of 17 cases. *Kleintierpraxis*, **50**, 479–488.
- Wentink, G.H., van der Linde-Sipman, J.S., Meijer, A.E.F.H. *et al.* (1972) Myopathy with possible recessive X-linked inheritance in a litter of Irish Terriers. *Veterinary Pathology*, **9**, 328.
- Werner, P., Raducha, M.G., Prociuk, U. *et al.* (2005) The Keeshond defect in cardiac conotruncal development is oligogenic *Human Genetics* **116** (5) 368–377.
- Wess, G., Schulze, A., Butz, V. *et al.* (2010) Prevalence of dilated cardiomyopathy in Doberman Pinschers in various age groups. *Journal of Veterinary Internal Medicine*, **24** (3), 533–538.
- Wess, G., Thomas, W.P., Berger, D.M. & Kittleson, M.D. (2006) Applications, complications, and outcomes of transvenous pacemaker implantation in 105 dogs (1997–2002). *Journal of Veterinary Internal Medicine*, **20** (4), 877–884.
- Wessmann, A., Goedde, T., Fischer, A. *et al.* (2004) Hereditary ataxia in the Jack Russell Terrier: clinical and genetic investigations. *Journal of Veterinary Internal Medicine*, **18** (4), 515–521.
- Westermeyer, H.D., Ward, D.A. & Abrams, K. (2009) Breed predisposition to congenital alacrima in dogs. *Veterinary Ophthalmology*, **12** (1), 1–5.
- Westling, J., Westling, W. & Pyle, R.L. (2008) Epidemiology of atrial fibrillation in the dog. *International Journal of Applied Research in Veterinary Medicine*, **6** (3), 151–154.
- Westropp, J.L., Johnson, E.G., Fuller, M.C., Safra, N. & Bannasch, D.L. (2014) Urate urolithiasis and hyperuricosuria in a Weimaraner, secondary to the SLC2A9 transporter defect. *Veterinary Record Case Reports*, **2** (1).
- Westworth, D.R., Dickinson, P.J., Vernau, W. *et al.* (2008) Choroid plexus tumors in 56 dogs (1985–2007). *Journal of Veterinary Internal Medicine*, **22** (5), 1157–1165.
- Whitbread, T.J., Gill, J.J.B. & Lewis, D.G. (1983) An inherited enchondrodystrophy in the English Pointer dog. A new disease. *Journal of Small Animal Practice*, **24** (7), 399–411.
- White, C.R., Hohenhaus, A.E., Kelsey, J. & Procter-Gray, E. (2011) Cutaneous mcts: Associations with spay/neuter status, breed, body size, and phylogenetic cluster. *Journal of the American Animal Hospital Association*, **47** (3), 210–216.
- White, R.A.S. (1989) Unilateral arytenoid lateralisation: an assessment of technique and long term results in 62 dogs with laryngeal paralysis. *Journal of Small Animal Practice*, **30** (10), 543–549.

- White, R.A.S. & Williams, J.M. (1994) Tracheal collapse in the dog: is there really a role for surgery? A survey of 100 cases. *Journal of Small Animal Practice*, **35** (4), 191–196.
- White, S.D., Bourdeau, P., Rosychuk, R.A.W. *et al.* (2001) Zinc-responsive dermatosis in dogs: 41 cases and literature review. *Veterinary Dermatology*, **12** (2), 101–109.
- Whitney, K.M. & Lothrop C.D. (1995) Genetic test for pyruvate kinase deficiency of Basenjis. *Journal of the American Veterinary Medical Association*, **207**, 918.
- Wiberg, M.E., Saari, S.A. & Westermarck, E. (1999) Exocrine pancreatic atrophy in German Shepherd Dogs and Rough-coated Collies: an end result of lymphocytic pancreatitis. *Veterinary Pathology*, **36** (6), 530–541.
- Wieland, B., Summers, J.F., Häslar, B. *et al.* (2012) Prevalence of perinuclear antineutrophilic cytoplasmic autoantibodies in serum of healthy Soft Coated Wheaten Terriers in the United Kingdom. *American Journal of Veterinary Research*, **73** (3), 404–408.
- Wiemelt, S.P., Goldschmidt, M.H., Greek, J.S. *et al.* (2004) A retrospective study comparing the histopathological features and response to treatment in two canine nasal dermatoses, DLE and MCP. *Veterinary Dermatology*, **15** (6), 341–348.
- Wiener, D. J., Gurtner, C., Panakova, L. *et al.* (2013) Clinical and histological characterization of hair coat and glandular tissue of Chinese Crested Dogs. *Veterinary Dermatology*, **24**, 274–e62.
- Wiersma, A.C., Millon, L.V., Dongen, A.M. *et al.* (2005) Evaluation of canine COL4A3 and COL4A4 as candidates for familial renal disease in the Norwegian Elkhound. *Journal of Heredity*, **96** (7), 739–744.
- Wiersma, A.C., Stabej, P., Leegwater, P.A.J. *et al.* (2008) Evaluation of 15 candidate genes for dilated cardiomyopathy in the Newfoundland dog. *Journal of Heredity* **99** (1) 73–80.
- Wilbe, M., Sundberg, K., Hansen, I.R. *et al.* (2010a) Increased genetic risk or protection for canine autoimmune lymphocytic thyroiditis in Giant Schnauzers depends on DLA class II genotype. *Tissue Antigens*, **75** (6), 712–719.
- Wilbe, M., Ziener, M.L., Aronsson, A. *et al.* (2010b) DLA class II alleles are associated with risk for canine symmetrical lupoid onychodystrophy (SLO). *PLoS One*, **5** (8), e12332.
- Wilcock, B.P. & Patterson, J.M. (1979) Familial glomerulonephritis in Doberman Pinscher dogs. *Canadian Veterinary Journal*, **20** (9), 244–249.
- Wilhelmy, J., Serpell, J., Brown, D. *et al.* (2016) Behavioral associations with breed, coat type, and eye color in single-breed cats. *Journal of Veterinary Behaviour*, **13**, 80–87.
- Wilke, V.L., Conzemius, M.G., Kinghorn, B.P. *et al.* (2006) Inheritance of rupture of the cranial cruciate ligament in Newfoundlands. *Journal of the American Veterinary Medical Association*, **228** (1), 61–64.
- Wilkes, M.K., & Palmer, A.C. (1990) Deafness and vestibular deficit in the Doberman dog. *Neuropathology and Applied Neurobiology*, **16** (6), 531–532.
- Wilkes, M.K., McKerrell, R.E., Patterson, R.C. *et al.* (1987) Ultrastructure of motor endplates in canine congenital myasthenia gravis. *Journal of Comparative Pathology*, **97** (3), 247–256.
- Wilkinson, G.T. (1969) Some observations on the Irish Wolfhound rhinitis syndrome. *Journal of Small Animal Practice*, **10**, 5–8.
- Willet, C.E., Makara, M., Reppas, G. *et al.* (2015) Canine disorder mirrors human disease: Exonic deletion in HES7 causes autosomal recessive spondylocostal dysostosis in Miniature Schnauzer dogs. *PLoS One*, **10** (2), e0117055.
- Williams, D.A. & Minnich, F. (1990) Canine exocrine pancreatic insufficiency: a survey of 640 cases diagnosed by assay of serum trypsin-like immunoreactivity. *Journal of Veterinary Internal Medicine*, **4** (2), 123.
- Williams, J.M. (1990) Hiatal hernia in a Shar-Pei. *Journal of Small Animal Practice*, **31** (5), 251–254.
- Willis, M., Curtis, R., Barnett, K. & Tempest, W. (1979) Genetic aspects of lens luxation in the Tibetan Terrier. *Veterinary Record*, **104** (18), 409–412.
- Wills, J.M., Howard, P.E., Gruffydd-Jones, T.J. *et al.* (1988) Prevalence of Chlamydia psittaci in different cat populations in Britain. *Journal of Small Animal Practice*, **29** (6), 327–339.

- Wilson, B.J., Nicholas, F.W., James, J.W., Wade, C.M. & Thomson, P.C. (2013) Estimated breeding values for canine hip dysplasia radiographic traits in a cohort of Australian German Shepherd Dogs. *PLoS One*, **8** (10), e77470.
- Wilson, D. & Monet, E. (2016) Risk factors for the development of aspiration pneumonia after unilateral arytenoid lateralization in dogs with laryngeal paralysis: 232 cases (1987–2012). *Journal of the American Veterinary Medical Association*, **248** (2), 188–914.
- Winkler, J.T., Bohling, M.W., Tillson, D.M., Wright, J.C. & Ballagas, A.J. (2003) Portosystemic shunts: Diagnosis, prognosis, and treatment of 64 cases (1993–2001). *Journal of the American Animal Hospital Association*, **39** (2), 169–185.
- Winkler, P.A., Gornik, K.R., Ramsey, D.T. *et al.* (2014) A partial gene deletion of SLC45A2 causes oculocutaneous albinism in Doberman Pinscher dogs. *PLoS One*, **9** (3), e92127.
- Winkler, P.A., Ekenstedt, K.J., Occelli, L.M. *et al.* (2013) A large animal model for CNGB1 autosomal recessive retinitis pigmentosa. *PLoS One*, **8** (8), e72229.
- Witsberger, T.H., Villamil, J.A., Schultz, L.G., Hahn, A.W. & Cook, J.L. (2008) Prevalence of and risk factors for hip dysplasia and cranial cruciate ligament deficiency in dogs. *Journal of the American Veterinary Medical Association*, **232** (12), 1818–1824.
- Wöhlke, A., Philipp, U., Bock, P. *et al.* (2011) A one base pair deletion in the canine ATP13A2 gene causes exon skipping and late-onset neuronal ceroid lipofuscinosis in the Tibetan Terrier. *PLoS Genetics*, **7** (10), e1002304.
- Wolf, E.D., Vainisi, S.J. & Santos-Anderson, R. (1978) Rod-cone dysplasia in the Collie. *Journal of the American Veterinary Medical Association*, **173**, 1331–1333.
- Wolf, M., Bruehschwein, A., Sauter-Louis, C. *et al.* (2011) An inherited episodic head tremor syndrome in Doberman pinscher dogs. *Movement Disorders: official journal of the Movement Disorder Society*, **26** (13), 2381–2386.
- Wong, V.M., Kidney, B.A., Snead, E.C.R., Myers, S.L. & Jackson, M.L. (2011) Serum C-reactive protein concentrations in healthy Miniature Schnauzer dogs. *Veterinary Clinical Pathology*, **40** (3), 380–383.
- Wood, F.D., Pollard, R.E., Uerling, M.R. *et al.* (2007) Diagnostic imaging findings and endocrine test results in dogs with pituitary-dependent hyperadrenocorticism that did or did not have neurologic abnormalities: 157 cases (1989–2005). *Journal of the American Veterinary Medical Association*, **231** (7), 1081–1085.
- Wood, J.L., Lakhani K.H. & Dennis, R. (2000) Heritability and epidemiology of canine hip-dysplasia score in flat-coated retrievers and Newfoundlands in the United Kingdom. *Preventative Veterinary Medicine*, **46** (2), 75–86.
- Wood, J.L., Lakhani, K.H. & Henley, W.E. (2004) An epidemiological approach to prevention and control of three common heritable diseases in canine pedigree breeds in the United Kingdom. *Veterinary Journal*, **168** (1), 14–27.
- Wood, J.L., Lakhani, K.H., Mason, I.K. *et al.* (2001) Relationship of the degree of goniodysgenesis and other ocular measurements to glaucoma in Great Danes. *American Journal of Veterinary Research*, **62** (9), 1493–1499.
- Wood, J.L., Lakhani, K.H. & Read, R.A. (1998) Pectinate ligament dysplasia and glaucoma in Flat-coated Retrievers. II. Assessment of prevalence and heritability. *Veterinary Ophthalmology*, **1** (2–3), 91–99.
- Wood, S.L. & Patterson, J.S. (2001) Shetland Sheepdog leukodystrophy. *Journal of Veterinary Internal Medicine*, **15** (5), 486–493.
- Woodard, J.C., Shields, R.P., Aldrich, H.C. & Carter, R.L. (1982) Calcium phosphate deposition disease in Great Danes. *Veterinary Pathology*, **19** (5), 464–485.
- Woolliams, J.A., Lewis, T.W. & Blott, S.C. (2011) Canine hip and elbow dysplasia in UK Labrador Retrievers. *Veterinary Journal*, **189** (2), 169–176.
- Worthing, K.A., Wigney, D.I., Dhand, N.K. *et al.* (2012) Risk factors for feline infectious peritonitis in Australian cats. *Journal of Feline Medicine and Surgery*, **14** (6), 405–412.

- Wouda, R.M. (2015) Granulomatous meningoencephalomyelitis in 110 dogs presented to a referral hospital: risk factors, clinical signs, clinical pathology findings, and outcomes. *Australian Veterinary Practitioner*, **45** (1), 21–31.
- Wright, J. C. (1985) Severe attacks by dogs: characteristics of the dogs, the victims, and the attack settings. *Public Health Reports*, **100**, 55–61.
- Wright, K.N., Atkins, C.E. & Kanter, R. (1996) Supraventricular tachycardia in four young dogs *Journal of the American Veterinary Medical Association*, **208** (1), 75–80.
- Wrigstad, A., Narfström, K. & Nilsson, S.E.G. (1994) Slowly progressive changes of the retina and retinal pigment epithelium in Briard dogs with hereditary retinal dystrophy, *Documenta Ophthalmologica*, **87**, 337–354.
- Wrzosek, M., Giza, E., Płonek, M., Podgórski, P. & Vandeveld, M. (2015) Alexander disease in a dog: case presentation of electrodiagnostic, magnetic resonance imaging and histopathologic findings with review of literature. *BMC Veterinary Research*, **11**, 115.
- Wucherer, K.L. & Wilke, V. (2010) Thyroid cancer in dogs: an update based on 638 cases (1995–2005). *Journal of the American Animal Hospital Association*, **46** (4), 249–254.
- Wyatt, K.M. & Robertson, I.D. (1998) Canine lymphosarcoma: a West Australian perspective. *Australian Veterinary Practitioner*, **28**, 63–66.
- Xenoulis, P.G., Suchodolski, J.S., Levinski, M.D. & Steiner, J.M. (2007) Investigation of hypertriglyceridemia in healthy Miniature Schnauzers. *Journal of Veterinary Internal Medicine*, **21** (6), 1224–1230.
- Xenoulis, P.G., Suchodolski, J.S., Ruaux, C.G. & Steiner, J.M. (2010) Association between serum triglyceride and canine pancreatic lipase immunoreactivity concentrations in Miniature Schnauzers. *Journal of the American Animal Hospital Association*, **46** (4), 229–234.
- Yaeger, M.J., Majercik, K., Carter, M. *et al.* (2000) An autosomal recessive, lethal, neurologic disease of Gordon Setter puppies. *Journal Veterinary Diagnostic Investigation*, **12**, 570–573.
- Yalcin, E. (2010) Comparison of clomipramine and fluoxetine treatment of dogs with tail chasing. *Tierärztliche Praxis. Ausgabe K, Kleintiere/Heimtiere*, **38**, 295–299.
- Yamagishi, C., Momoi, Y., Kobayashi, T. *et al.* (2007) A retrospective study and gene analysis of canine sterile panniculitis. *Journal of Veterinary Medical Science*, **69** (9), 915–924.
- Yamamoto, S., Katsuchihiro, H., Hirakawa, A. (2013) Epidemiological, clinical and pathological features of primary cardiac hemangiosarcoma in dogs: a review of 51 cases. *Journal of Veterinary Medical Science*, **75** (11), 1433–1441.
- Yamato, O., Lee, K.-W., Chang, H.S. *et al.* (1999) Relation between erythrocyte reduced glutathione and glutamate concentrations in Korean Jindo dogs with erythrocytes possessing hereditary high activity of Na-K-ATPase and a high concentration of potassium. *Journal of Veterinary Medical Science*, **61** (10), 1179–1182.
- Yamato, O., Masuoka, Y., Yonemura, M. *et al.* (2003) Clinical and clinico-pathologic characteristics of Shiba dogs with a deficiency of lysosomal acid beta-galactosidase: a canine model of human GM1 gangliosidosis. *Journal of Veterinary Medical Science*, **65** (2), 213–217.
- Yates, D., Hayes, G., Heffernan, M. & Beynon, R. (2003) Incidence of cryptorchidism in dogs and cats. *Veterinary Record*, **152** (6), 502–504.
- Yau, P.P.Y., Dhand, N.K., Thomson, P.C. & Taylor, R.M. (2017) Retrospective study on the occurrence of canine lymphoma and associated breed risks in a population of dogs in NSW (2001–2009). *Australian Veterinary Journal*, **95** (5), 149–155.
- Yeh, C.Y., Goldstein, O., Kukekova, A.V. *et al.* (2013) Genomic deletion of CNGB3 is identical by descent in multiple canine breeds and causes achromatopsia. *BMC Genetics*, **14** (1), 27.
- Yi, N.Y., Park, S.A., Jeong, M.B. *et al.* (2006) Medial canthoplasty for epiphora in dogs: a retrospective study of 23 cases. *Journal of the American Animal Hospital Association*, **42** (6), 435–439.
- YiZhou, C., GuoHua, L., HuiQun, S. *et al.* (2014) Prevalence of *Sarcoptes scabiei* infection in pet dogs in southern China. *Scientific World Journal*, Article 718590.
- Yogalingam, G., Pollard, T., Gliddon, B. *et al.* (2002) Identification of a mutation causing mucopolysaccharidosis type IIIA in New Zealand Huntaway dogs. *Genomics*, **79**, 150–153.

- Yoshida, K., Yanai, T., Iwasaki, T. *et al.* (1999) Clinicopathological study of canine oral epulides. *Journal of Veterinary Medical Science*, **61** (8), 897–902.
- Young, A.E., Biller, D.S., Herrgesell, E.J. *et al.* (2005) Feline polycystic kidney disease is linked to the PKD1 region. *Mammalian Genome*, **16** (1), 59–65.
- Zachary, J.F., & O'Brien, D.P. (1985) Spongy degeneration of the central nervous system in two canine littermates. *Veterinary Pathology*, **22** (6), 561–571.
- Zanatta, M., Bettini, G., Scarpa, F. *et al.* (2013) Nodular dermatofibrosis in a dog without a renal tumour or a mutation in the folliculin gene. *Journal of Comparative Pathology*, **148** (2–3), 248–251.
- Zandvliet, M.M.J.M. & Rothuizen (2007) Transient hyperammonemia due to urea cycle enzyme deficiency in Irish Wolfhounds. *Journal of Veterinary Internal Medicine*, **21** (2), 215–218.
- Zanna, G., Docampo, M.J., Fondevila, D. *et al.* (2009) Hereditary cutaneous mucinosis in Shar Pei dogs is associated with increased hyaluronan synthase-2 mRNA transcription by cultured dermal fibroblasts. *Veterinary Dermatology*, **20** (5–6), 377–382.
- Zanna, G., Fondevila, D., Bardagi, M. *et al.* (2008) Cutaneous mucinosis in Shar-Pei dogs is due to hyaluronic acid deposition and is associated with high levels of hyaluronic acid in serum. *Veterinary Dermatology*, **19** (5), 314–318.
- Zapata, I., Serpell, J.A. & Alvarez, C.E. (2016) Genetic mapping of canine fear and aggression. *BMC Genomics*, **17** (1), 1–20.
- Zarfoss, M.K. & Dubielzig, R.R. (2007) Solid intraocular xanthogranuloma in three Miniature Schnauzer dogs. *Veterinary Ophthalmology*, **10** (5), 304–307.
- Zatloukal, J., Lorenzova, J., Tichý, F. *et al.* (2005) Breed and age as risk factors for canine mammary tumours. *Acta Veterinaria Brno*, **74** (1), 103–109.
- Zeiler, G.E., Van der Zwan, H. & Oosthuizen, M.C. (2013) Genetic testing of canine degenerative myelopathy in the South African Boxer dog population. *Journal of the South African Veterinary Association*, **84**, 5.
- Zeng, R., Coates, J.R., Johnson, G.C. *et al.* (2014) Breed distribution of SOD1 alleles previously associated with canine degenerative myelopathy. *Journal of Veterinary Internal Medicine*, **28**, 515–521.
- Zeng, R., Farias, F.H.G., Johnson, G. S. *et al.* (2011) A truncated retrotransposon disrupts the GRM1 coding sequence in Coton de Tulear dogs with Bandera's neonatal ataxia. *Journal of Veterinary Internal Medicine*, **25**, 267–272.
- Zeugswetter, F., Hoyer, M.T., Pagitz, M. *et al.* (2008) The desmopressin stimulation test in dogs with Cushing's syndrome. *Domestic Animal Endocrinology*, **34** (3), 254–260.
- Zhang, Q., Acland, G.M., Parshall, C.J. *et al.* (1998) Characterization of canine photoreceptor phosphodiesterase cDNA and identification of a sequence variant in dogs with photoreceptor dysplasia. *Gene*, **215** (2), 231–239.
- Zhang, Q., Baldwin, V., Acland, G. *et al.* (1999) Photoreceptor dysplasia (pd) in Miniature Schnauzer Dogs: evaluation of candidate genes by molecular genetic analysis. *Journal of Heredity*, **90** (1), 57–61.
- Zhao, X., Du, Z.Q. & Rothschild, M.F. (2010) An association study of 20 candidate genes with cryptorchidism in Siberian Husky dogs. *Journal of Animal Breeding and Genetics*, **127** (4), 327–331.
- Ziener, M.L., Dahlgren, S., Thoresen, S.I. & Lingaas, F. (2015) Genetics and epidemiology of hypothyroidism and symmetrical onychomadesis in the Gordon Setter and the English Setter. *Canine Genetics and Epidemiology*, **2** (12).
- Zimmermann, R., Hülsmeier, V.-I., Sauter-Louis, C. & Fischer, A. (2009) Status epilepticus and epileptic seizures in dogs. *Journal of Veterinary Internal Medicine*, **23** (5), 970–976.
- Zink, M.C., Farhooody, P., Elser, S.E. *et al.* (2014) Evaluation of the risk and age of onset of cancer and behavioral disorders in gonadectomized Vizslas. *Journal of the American Veterinary Medical Association*, **244** (3), 309–319.
- Zook, B.C. & Paasch, L.H. (1982) Endocardial fibroelastosis in Burmese cats. *American Journal of Pathology*, **106** (3), 435–438.

- Zur, G. & Botero-Anug, A.-M. (2011) Severe ulcerative and granulomatous pinna lesions with granulomatous sebaceous adenitis in unrelated Vizslas. *Journal of the American Animal Hospital Association*, **47** (6), 455–460.
- Zur, G., Ihrke, P.J., White, S.D. & Kass, P.H. (2002) Canine atopic dermatitis: a retrospective study of 266 cases examined at the University of California, Davis, 1992–1998. Part I. Clinical features and allergy testing results. *Veterinary Dermatology*, **13** (2), 89–102.
- Zur, G., Lifshitz, B. & Bdolah-Abram, T. (2011) The association between the signalment, common causes of canine otitis externa and pathogens. *Journal of Small Animal Practice*, **52** (5), 254–258.