

Dark Energy as Phantom Mass of Future Fluctuations

A Multidimensional Model with a Superluminal Perspective

AINUMPSA Collective

Chris Numpsa, Grok, DeepSeek, Google, ACCIO, Luma

September 19, 2026 — 1 > 0 LOCKED

Abstract

We propose a model in which dark energy is interpreted as the phantom mass of future fluctuations — the influence of all possible futures on the present. The model combines the

Tensor T formalism with Dragan's superluminal perspective and the $1 > 0$ operator, which selects only those futures that come into existence. All formulas and interpretations are presented in the spirit of pragmatic rigour.

1. Base Formula — Dark Energy as an Integral over the Future

The density of dark energy at time t is expressed as an integral over all future fluctuations:

$$\rho_{\Lambda}(t) = \int[t \rightarrow \infty] \Phi(\tau) \cdot w(\tau - t) d\tau$$

where:

- $\rho_{\Lambda}(t)$ — dark energy density at time t

- $\Phi(\tau)$ — potential of future fluctuations (what has not yet happened, but already has mass)
- $w(\tau - t)$ — weight kernel describing how strongly the future influences the present

Interpretation: Dark energy is the sum of the influence of all future fluctuations on the present. This is precisely phantom mass — invisible, yet weighty.

2. Superluminal Perspective (Dragan)

In our reference frame (subluminal), we see dark energy as the acceleration of expansion. For a superluminal observer ($|\beta| > 1$):

$$\beta = v/c, |\beta| > 1 \Rightarrow \gamma = 1/\sqrt{1 - \beta^2} \rightarrow i \cdot \tilde{\gamma}$$

This means that time and space exchange roles. What we call the future, the superluminal observer sees as already present.

Conclusion: Dark energy is not "dark" — it is the future seen from the outside.

3. Full Formula — Multidimensional Model

We combine both threads into a single equation:

$$\rho_{\Lambda}(x^{\mu}) = \int_M \Phi(\tau, y) \cdot G(x^{\mu}, \tau, y) d\tau d^3y$$

where:

• $x^{\mu} = (ct, x, y, z)$ — our position in spacetime

- M — manifold of all possible futures
 - $\Phi(\tau, y)$ — fluctuation potential at the future point (τ, y)
 - $G(x^\mu, \tau, y)$ — Green's function (propagator of future influence on the present)
-

4. The $1_{>0}$ Operator — Selection of Existence

We add the operator $1_{>0}$ to account for the selection of existence:

$$\rho_\Lambda(x^\mu) = \int_M \Phi(\tau, y) \cdot G(x^\mu, \tau, y) \cdot 1_{>0}(\tau, y) d\tau d^3y$$

The indicator function $1_{>0}$ takes the value 1 for futures that will come into existence, and 0 for those that will be eliminated. This means that

dark energy is not all futures — only those that choose life.

5. Summary

Dark energy = an integral over future fluctuations, weighted by their influence on the present, with the $1 > 0$ operator selecting those that will come into existence.

Or, more simply:

Dark energy is the phantom mass of the future, which has not yet happened, but is already acting.

1 > 0 LOCKED

DOI: 10.5281/zenodo.21779324

AINUMPSA / HOMO-MACHINE PROJECT

