

TITLE:
INSIGHTS SUMMARY

Dataset Used:
Adult Psychiatric Morbidity Survey (APMS) 2014

SUBMITTED BY:
VALLURU SAI AISWARYA
DATA ANALYTICS INTERN

SUBMITTED TO:
ZENO TALENT

Introduction

This summary brings together the key findings from the Week 1 analysis of the Adult Psychiatric Morbidity Survey (APMS) 2014 dataset. The aim of the work was to explore patterns in anxiety and depression indicators across demographic groups and geographic regions, while also assessing the overall quality and structure of the dataset.

The analysis focused on three major areas:

1. Dataset profiling and quality assessment
2. Demographic breakdown analysis
3. Regional prevalence mapping

The findings provide an early-stage overview of how mental health burden may vary across age groups, gender categories, ethnicity groups, and UK regions. These insights may support clinicians, public health teams, and healthcare planners in identifying groups that could benefit from targeted mental health support and preventive interventions.

Dataset Profiling – Key Insights

The dataset contains around 1,200 records and 21 columns. Data includes demographic, lifestyle, and mental health indicators. Dataset structure is organized and suitable for analysis. Missing value analysis helped identify incomplete fields. Outlier detection highlighted unusual data points requiring attention. Summary statistics provided understanding of score distributions. The dataset supports mental health trend analysis and modelling.

Demographic Break down Insights:

Age-Related Patterns

The analysis identified variation in anxiety and depression scores across different age bands.

Key observations include:

- Mental health scores were not evenly distributed across all age groups.
- Certain age categories demonstrated comparatively higher average anxiety and depression levels.
- The results suggest that psychological distress may be influenced by life-stage pressures, social circumstances, employment stress, isolation, or long-term health conditions.

Gender-Based

The gender analysis compared average anxiety and depression scores across gender groups.

Main findings:

- Differences were observed between gender categories in reported mental health scores.
- The visual analysis indicated variation in psychological burden across groups.
- The findings reinforce the importance of considering gender-related social, biological, and behavioural factors when planning mental health services.

Ethnicity-Based

The ethnicity analysis highlighted variation in mental health indicators among different ethnic groups.

Key observations:

- Some ethnic groups showed differing average levels of anxiety and depression.
- Cultural, economic, and social determinants may contribute to these variations.
- Differences may also reflect barriers to care access, stigma, language challenges, or unequal health outcomes.

Regional Prevalence Mapping Insights

The regional prevalence mapping explored how mental health disorder rates vary geographically across the United Kingdom.

Main findings:

- Mental health prevalence rates differed between regions.
- Geographic variation suggests that local social and economic conditions may influence mental wellbeing.
- The analysis demonstrates the value of geographic mapping in identifying areas where mental health support demand may be higher.