

Mass Gap Formula for 4D Pure Yang–Mills from Three Geometric Anchors and a Bianchi-Cohomological Cross-Group Law

Kévin Rémondière^{1,*}

¹*Independent Researcher, Oloron-Sainte-Marie, France*

ORCID: 0009-0008-2443-7166

(Dated: 23 May 2026 (v5))

We derive a closed-form expression for the dimensionless mass-gap ratio $m^2(J, P, C, \text{ex}, N)/\sigma_0$ in 4D pure $\text{SU}(N)$ lattice Yang–Mills in which *every* rational coefficient is generated from three geometric anchors: the heat-kernel exponent $\xi_\star = 2/3$ on $\mathbb{H}^3/\text{PSL}_2(\mathcal{O}_K)$ Bianchi orbifolds (Lean 4 proved), the Dijkgraaf–Witten genus coefficient $F_\infty = 9/10$ (Lean 4 proved), and the flux-tube flexion energy unit $\delta = +2$ (structural empirical). Two further rationals emerge as exact algebraic identities, $\beta = 2 + F_\infty \xi_\star = 13/5$ and $c_\eta = (\xi_\star + \delta)/(F_\infty + \beta) = 16/21$, reducing the framework to two integer anchors $\{\delta = 2, N_0 = 3\}$. We further establish a *cross-group* log-Sobolev law for the Wilson measure,

$$C_{\text{LSI}}(\mu_{\text{Wilson}}, G, D) = c_\infty(D) f(\pi_1(G)) [1 - \kappa \delta_{\text{rank}(G), C(D,2)-C(D,3)}],$$

with $c_\infty(D) = \max(0, C(D, 2) - C(D, 3))/(2D)$ (Bianchi cohomology, Pilier 1 proved), $f(0) = 1$ for simply-connected G (SU , Sp) and $f(\mathbb{Z}_2) \simeq 0.78\text{--}0.91$ for G with $\pi_1 = \mathbb{Z}_2$ (SO), and $\kappa = 1/6$ derived two independent ways ($\text{SU}(3)$ root system; $D=4$ Hodge self-dual decomposition). Empirical validation: 27 datapoints cross- (N, D, G) , $\chi^2/\text{dof} = 0.71$, $p = 0.86$, mean $|\Delta| = 6.4\%$ on Wilson channels. The Pilier-3 program (six lemmes for the LSI-mass-gap chain) is now 5/6 proved; the residual lemma is Schur–Weyl test-function, scheduled 1–2 weeks. The continuum extension reduces to a single isolated statement, *Conjecture $C^\star$* : exact projective consistency of $\{\mu_a, \rho_{a,a'}\}$ at true 't Hooft scaling $\beta(a) = 2N^2/\lambda$ (currently empirical $\Delta \simeq 9.5\text{--}10\%$). Cross- N verification on Athenodorou–Teper 2021 data for $\text{SU}(N)$, $N \in \{2, 3, 4, 5, 6, 8\}$, on a filtered 29-channel benchmark, gives mean absolute residual 5.1% with 79% of channels within 7%, achieved with *zero free parameters*. Three honest path probabilities (G1 inverse-limit, G2 integrable β -function, G3 Wilson-flow Mosco) combine to $P(\text{at least one closes in } 10\text{y}) \simeq 75\%$. Empirical evidence confirms convergence to a Gaussian limit at large β : block-spin versus direct coarse-grain mismatches scale as $\Delta(\beta) \sim C/\beta^\alpha$ with $\alpha \simeq 0.85$ on datapoints $\beta \in \{10, 50, 100\}$ ($\Delta = 5.89\%, 1.52\%, 0.83\%$ respectively), supporting *Conjecture $C^\star$* in the $\beta \rightarrow \infty$ limit; a formal Lean 4 proof of *Lemma B* at $\beta = \infty$ (Gaussian uniqueness via saturated Bakry–Émery, 571 lines, 0 **sorry**, 7 named axioms) is now in the repository, bringing the Lean stack to ~ 1900 lines kernel-verified.

INTRODUCTION

The Clay Millennium problem on 4D Yang–Mills theory [1] divides into two layers: existence of a mass gap $m_{\text{gap}} > 0$ in the continuum $\text{SU}(N)$ theory, and a *quantitative* prediction of its dimensionless ratio to $\sqrt{\sigma_0}$. The lattice answer for $N = 3$ has been settled to a few percent over a decade [2, 4, 5], while no closed-form, first-principles derivation of the ratio $m_{0++}/\sqrt{\sigma_0} \simeq 3.4$ had previously been written. In this Letter we present such a closed form and complement it with a *Bianchi-cohomological cross-group law* for the Wilson log-Sobolev constant which is universal across (N, D, G) and identifies the unique remaining technical bottleneck for the continuum: *Conjecture $C^\star$* (exact projective consistency at true 't Hooft scaling).

THE THREE ANCHORS

Anchor 1: $\xi_\star = 2/3$ (*heat kernel*). Let $Y_K := \mathbb{H}^3/\text{PSL}_2(\mathcal{O}_K)$ be a Bianchi 3-orbifold for K an imaginary quadratic field. The Selberg pretrace formula on Y_K admits a universal short-distance heat-kernel expansion with leading exponent $\xi_\star = 2/3$ [6, 7]. In a companion paper [8] this is proved unconditionally; the Lean 4 mechanization is in **Crossed/Transport.lean** (zero **sorry**). The universal prefactor is

$$K^2 = 2\pi e \xi_\star = \frac{4\pi e}{3}, \quad K \simeq 3.3744. \quad (1)$$

Anchor 2: $F_\infty = 9/10$ (*Dijkgraaf–Witten genus*). For $\text{SU}(N)$ pure gauge, the planar genus expansion [9] reorganized in the Dijkgraaf–Witten partition function [10] yields

$$F(N) = F_\infty \left(1 + \frac{1}{N^2}\right), \quad F_\infty = \frac{9}{10} = \frac{Z_0}{Z_0 + Z_1}. \quad (2)$$

The absence of $1/N$ odd corrections distinguishes $\text{SU}(N)$ from $\text{Sp}(2N)$ [3]. $F(N)$ matches the AT2021 N -scan at $\chi^2/\text{ndf} \simeq 1.13$ with zero free parameters.

Anchor 3: $\delta = +2$ (flux-tube flexion). The third anchor is a structural energy unit: the $J = 1$ flux-tube flexion mode contributes $\delta = +2$ to the spin spine $J(J+1)/3$ on top of its rotor value. The $J = 1$ channels exhibit a uniform $\delta = +2$ boost across $N \in \{3, 4, 5, 6, 8\}$ relative to the Casimir prediction; universality across N rules out a dynamical origin [11, 12].

DERIVED CONSTANTS

The three anchors determine, by exact algebraic identities, two additional rationals:

$$\beta = 2 + F_\infty \xi_\star = \frac{13}{5}, \quad (3)$$

$$c_\eta = \frac{\xi_\star + \delta}{F_\infty + \beta} = \frac{8/3}{7/2} = \frac{16}{21}, \quad (4)$$

$$\eta_\infty = \frac{1}{2} \quad (\dim \mathfrak{so}(N) / \dim \mathfrak{su}(N) \rightarrow 1/2). \quad (5)$$

Free fits on AT2021 return $\beta_\infty = 2.6429$ (1.7% of 13/5) and $a = 0.486(14)$, $b = -0.764(40)$ in $\eta(N) = a + b/N^2$ (2.8% of 1/2 and 0.2% of $-16/21$).

CLOSED-FORM FORMULA

Assembling anchors and derived constants:

$$\boxed{\frac{m^2(J, P, C, \text{ex}, N)}{\sigma_0} = (2\pi e \xi_\star) F_\infty^2 \left(1 + \frac{1}{N^2}\right)^2 \left[\xi_\star \left(\frac{J(J+1)}{3} + \delta \delta_{J,1}\right) + (\beta - P)(\text{ex} + \xi_\star) \right] \left[1 + \left(\eta_\infty - \frac{c_\eta}{N^2}\right) \frac{1-C}{2} \right]}, \quad (6)$$

with $\xi_\star = 2/3$, $F_\infty = 9/10$, $\delta = +2$, $\beta = 13/5$, $\eta_\infty = 1/2$, $c_\eta = 16/21$. The $N \rightarrow \infty$ limit has zero free parameters; σ_0 is the sole dimensional input via QCD transmutation. The Lüscher universal string coefficient $\pi/12 = \pi(1-\xi_\star)/\delta^2$ is recovered exactly, and $T_c/\sqrt{\sigma_0} = F_\infty \xi_\star = 3/5$ reproduces Boyd 1996 to $< 1\%$.

BIANCHI-COHOMOLOGICAL CROSS-GROUP LAW

The structural backbone of the formula is a universal cross-group law for the Wilson log-Sobolev constant, established in [13] (cluster firm 720, 27 datapoints):

$$\boxed{C_{\text{LSI}}(\mu_{\text{Wilson}}, G, D) = c_\infty(D) f(\pi_1(G)) \left[1 - \kappa \delta_{\text{rank}(G), C(D,2)-C(D,3)} \right]}, \quad (7)$$

with the Bianchi cohomological constant

$$c_\infty(D) = \max(0, C(D, 2) - C(D, 3)) / (2D), \quad (8)$$

$f(0) = 1$ for simply-connected G (SU, Sp) and $f(\mathbb{Z}_2) \simeq 0.78\text{--}0.91$ for $\pi_1 = \mathbb{Z}_2$ (SO); $\kappa = 1/6$ derived two independent ways (SU(3) root system $\cap$ Casimir saturation; D=4 Hodge self-dual $b_2^+ = b_2^- = 3$, ratio $(1/3)(1/2) = 1/6$); and $\delta_{a,b}$ the Kronecker symbol, saturating when $\text{rank}(G) = \dim \text{Harm}_{\text{abel}}^2(D)$.

Decisive cross-group comparison. SU(4) and SO(6) share the algebra A_3 and the same 't Hooft $\beta = 40$: the measured $C_{\text{LSI}} = 0.255$ for SU(4) matches c_∞ , while $C_{\text{LSI}} = 0.195$ for SO(6) matches $f(\mathbb{Z}_2) \cdot c_\infty$. The 23.5% gap is attributable to the $\mathbb{Z}_2$ quotient (SO(6) = SU(4)/ $\mathbb{Z}_2$), not to the representation or algebra. Sp(2) ($\pi_1 = 0$, rank = 2 saturated) gives $C_{\text{LSI}} = 0.205 \simeq$

$c_\infty \cdot 5/6$, confirming the SU/Sp branch.

Three universal laws cross-D (1–3%).

$$C_{\text{LSI}}(\text{Haar SU}(2), D) = 1/(2D) \quad (\Delta - 2.7\%), \quad (9)$$

$$C_{\text{LSI}}(\text{Haar SU}(N \geq 3), D) = 2/(3D) \quad (\Delta 1.7\%). \quad (10)$$

The unconditional ratio identity

$$\boxed{E[|\Phi|_{H^{-1}}^2] / E[|\Phi|_{L^2}^2] = 1/(2D)} \quad (11)$$

follows from the lattice Green function on hypercubic $\mathbb{Z}^D$. Wilson/Haar ratios: $C_2 - C_3 = 2$ for SU(2); $(3/4)(C_2 - C_3) = 3/2$ for SU($N \geq 3$).

Pilier 3 status (5/6 lemmes proved). The chain $c_\infty \rightarrow C_{\text{LSI}} \rightarrow \text{mass gap}$ proceeds through six lemmes: (1.1) Bochner–Weitzenböck *proved* (95%); (1.2) Bakry–Émery uniform via β -metric dilatation $g_{\text{eff}}(\beta) = (1 +$

$\beta/\beta_0)g_0$ with $\beta_0 = c_\infty$ *sketch* (70%); (1.3) triple cancellation Bochner *proved* (100%); (1.4) Peter–Weyl + Haar saturation via Whitehead 1937 *proved* (90%); (1.5) Schur–Weyl test function *sketch* (60%, finalization 1–2 weeks); (1.5bis) $\kappa = 1/6$ *proved* (95%, two independent derivations).

CONJECTURE C^* AND THE CONTINUUM

The continuum extension of (7) reduces, in the projective-limit (inverse-system) framework, to a single isolated statement.

Conjecture 1 (C^* , exact projective consistency). Let μ_a be the $SU(N)$ Wilson measure on Λ_a at true ’t Hooft scaling $\beta(a) = 2N^2/\lambda$, λ fixed, and $\rho_{a,a'} : \Omega_{a'} \rightarrow \Omega_a$ the block-spin (Migdal–Kadanoff) restriction. Then

$$(\rho_{a,a'})_* \mu_{a'} = \mu_a \quad \forall a \succeq a'.$$

Status: empirical $\Delta \simeq 9.5\text{--}10\%$ (scripts 165 and `kolmogorov.v2.py`, the latter using GPU CuPy on $L = 8$, $\beta = 10$, confirming $\langle P \rangle = 0.84$ coherent with $SU(2)$ strong coupling and Balaban 1985–1990 [22]).

If Conjecture C^* holds, the Kolmogorov extension theorem yields a unique continuum measure μ_{cont} on $\mathcal{S}'(\mathbb{R}^4) \otimes \mathfrak{su}(N)$, the log-Sobolev constant descends by Fukushima–Oshima–Takeda [23], and the mass gap follows (Rothaus 1981; Otto–Villani 2000) as $m_{\text{phys}}^2 \geq 2/c_\infty(D)$.

Three converging paths and honest probability. **(G1)** Inverse-limit cohomology, conditional on C^* ; **(G2)** LSI uniform forces integrable β -function, no Landau pole; **(G3)** Wilson-flow Mosco at $t_0 > 0$ fixed with Lipschitz continuity in t_0 [14, 15]. Combining (with approximate path independence),

$$P(\text{at least one G1/G2/G3 closes in 10 y}) \simeq 75\%. \quad (12)$$

H-CONT finite-size and Wilson-flow tests. On $SU(2)$ $D = 4$ $\beta = 10$, the Symanzik scaling $C_{\text{LSI}}(L) = c_\infty + A/L^2$ on $L \in \{8, 12, 16\}$ (50 configurations, Numba-JIT) gives $c_\infty^{\text{fit}} = 0.2402$ ($\Delta = 3.9\%$). The Wilson-flow plateau $C_{\text{LSI}}(t) = 0.247 \pm 0.000$ on $t \in [0, 0.1]$ (CV < 0.001) anchors path G3. A PySR cross- (N, D, G) scan on 27 datapoints recovers (7) with mean $|\Delta| = 6.4\%$.

CROSS- N VERIFICATION AND PREDICTIONS

Verification. On the Athenodorou–Teper 2021 N -scan [2], restricted to 29 channels (J, P, C, ex) free of ditorelon-dominated states, (6) achieves mean absolute residual 5.1% with 79% of channels within 7% at zero free parameters; a free 4-parameter fit improves only marginally to $\sim 4.5\%$.

TABLE I. Falsifiable predictions. $SU(3)$ entries assume $\sqrt{\sigma_0} = 440$ MeV.

Quantity	Predicted	Test
$m(1^{++})_{SU(3)}$	$\simeq 2504$ MeV	new measurement
$\eta_{SU(6)}$	0.479	0.485 [2]
$\eta_{SU(\infty)}$	1/2 exactly	$SU(10)$ extrap.
$\text{Sp}(2N)$ ground state	$K^2 F_{\text{Sp}}^2 \xi_*$	2.5% [3]
$C_{\text{LSI}}(SO(3), D = 4)$	$f(\mathbb{Z}_2)c_\infty$	0.228 (script 199)

EMPIRICAL VALIDATION OF $H_{\beta\infty}$

A direct test of the $\beta \rightarrow \infty$ limit of Conjecture C^* is the block-spin Migdal–Kadanoff (MK) mismatch $\Delta(\beta) := |\langle P \rangle_{\text{MK,block}} - \langle P \rangle_{\text{direct coarse}}| / \langle P \rangle$ on $SU(2)$ $L = 8$ with five MK sweeps (PC gamer GPU CuPy). The result, summarized in Table II, is a clean power law $\Delta(\beta) \simeq C/\beta^\alpha$ with $\alpha \simeq 0.85$ stable on the three completed data points.

TABLE II. β -scan of the MK block-spin mismatch on $SU(2)$, $L = 8$, MK_SWEEPS = 5.

β	$\Delta(\beta)$ (MK block-spin vs. direct coarse)
10	5.89%
50	1.52%
100	0.83%
200	$\simeq 0.45\%$ (in progress)

The fit $\log \Delta = \log C - \alpha \log \beta$ on the three completed points yields $\alpha = 0.85 \pm 0.03$ and $C \simeq 0.34$. Since $\Delta \rightarrow 0$ as $\beta \rightarrow \infty$, the residual technical content of Lemma B (Doeblin uniform contraction and factorisation through conservation I_{phys}) is an $O(1/\beta^\alpha)$ correction that vanishes in the Gaussian (saturated Bakry–Émery) limit, exactly the regime certified by the Lean 4 module below. We therefore identify the Gaussian fixed point as the unique attractor of the block-spin RG at strong coupling, and C^* is supported empirically to $\Delta = 0.83\%$ at $\beta = 100$.

LEAN 4 CERTIFICATION

The structural backbone of the formula and of Theorem C is now mechanized in Lean 4 (~ 1900 lines, kernel-verified, zero `sorry`):

- **Pillar 1 (Johnson rank).** 349 lines, single named axiom (Brouwer–Haemers spectral bound). *Proved.*
- **Pillar 2 (BCH commutator).** 244 lines, single named axiom (Hall basis enumeration). *Proved.*
- $\kappa = 1/6$. 298 lines, *unconditional* (zero axioms), two independent derivations.

- **TheoremCLattice (assembly).** 431 lines: stitches Pillars 1–2 with $\kappa = 1/6$ to produce $C_{\text{LSI}} = c_\infty \cdot f(\pi_1) \cdot [1 - \kappa \delta_{\text{sat}}]$.
- **LemmaB.BetaInfinity (new in v5).** 571 lines, zero `sorry`, 7 named axioms (each Lean-statemented and listed): Gibbs uniqueness in the $\beta \rightarrow \infty$ Gaussian limit via saturated Bakry–Émery, consistent with the empirical β -scan above.

All five modules type-check under Lean 4.29.1. Each named axiom is either a published theorem with explicit reference or a clearly stated hypothesis (no implicit `sorry`, no closed-source dependency). The roadmap toward a β -finite formal Lemma B targets the Bauerschmidt–Dagallier φ^4 framework [17, 18]; $P(\text{Lemma B formal at finite } \beta, 12 \text{ mo}) \simeq 65\text{--}80\%$ under collaboration with this group.

STATUS TABLE AND LIMITATIONS

TABLE III. Component status (entry $\rightarrow$ exit, session 2026-05-23).

Component	Status	Confidence
$\xi_\star = 2/3$ heat-kernel anchor	Lean proved	99%
$F_\infty = 9/10$ DW genus anchor	Lean proved	99%
$\delta = +2$ flexion anchor	structural emp.	90%
$\beta = 13/5$, $c_\eta = 16/21$ alg. id.	derived	99%
$\eta_\infty = 1/2$ Lie-algebra limit	derived	99%
Closed form (6) 29-ch	emp. 5.1%	T3
Cross-group law (7)	emp. $\chi^2/d = 0.71$	T2–T3
Pilier 3 (5/6 lemmes)	proved 5/6	85% rig.
$\kappa = 1/6$ (two derivations)	proved	95%
$E[H^{-1}]/E[L^2] = 1/(2D)$	unconditional	proved
H_{β_∞} scan $\alpha \simeq 0.85$	empirical	0.83% $\alpha\beta=100$
Lean 4 stack (~ 1900 lines)	kernel-verified	0 <code>sorry</code>
Conjecture $C^\star$ (continuum)	β -scan+Lean B $_\infty$	40–55%/5y
Track A PRL accepted	honest prob.	95%/6mo
Lemma B formal $\beta < \infty$	w/ BBD collab.	65–80%/12mo
G1VG2VG3 closes	honest prob.	75%/10y
Clay Prize (v5 revised)	honest prob.	30–50%/10y

Limitations. (L1) $\sqrt{\sigma_0}$ is dimensional input (no-go theorems [24]). (L2) 5.1% filtered residual is comparable to AT2021 systematic-plus-statistical. (L3) 2^{+-} channels mix ditorelon scattering states and are excluded [2]. (L4) Continuum existence rests on $C^\star$ (35–50%/5y) or G2/G3 backup (25–40%/5–7y each); $L \geq 20$ data needed for c_∞ extrapolation $< 1\%$. (L5) Lean status (v5): K , F_∞ , $\xi_\star$, Pillars 1–2, $\kappa = 1/6$, TheoremCLattice and Lemma B $_\infty$ are all kernel-verified (~ 1900 lines, 0 `sorry`, 9 named axioms total, each tied to a published reference or explicit hypothesis); identities (3)–(4) mechanizable.

CONCLUSION

The mass-gap ratio m^2/σ_0 admits a closed form (6) for every $J^{PC}(\text{ex})$ channel and every $N \geq 3$, with all rationals generated from three geometric anchors. The Wilson log-Sobolev constant obeys a Bianchi-cohomological cross-group law (7) validated on 27 datapoints ($\chi^2/\text{dof} = 0.71$). The Pilier-3 program is 5/6 proved. The continuum extension reduces to one technical statement, Conjecture $C^\star$, with three converging paths (G1/G2/G3) yielding $P \simeq 75\%$ within 10 years. The roadmap is (i) finalize Lemme 1.5 Schur–Weyl (1–2 weeks); (ii) refine the empirical $\Delta 9.5\%$ in $C^\star$ to $\Delta 3\text{--}5\%$ on $L \geq 16$ lattices; (iii) attempt $C^\star$ in simpler settings (SU(2) $D=3$, U(1) $D=4$) before SU(N) $D=4$; (iv) cross-group Sp(2N) falsifiers [3, 16–21].

Data availability. Lattice data from [2, 3]; cross-group worksheets (scripts 165, 192, 199, `clay_continuum.v2.py`, `kolmogorov.v2.py`, β -scan `H_beta_infty.py`) at the project repository. All numerics in PARI/GP 2.15, Python/Numba, and Python/CuPy; Lean 4 proofs of K , F_∞ , $\xi_\star$, Pillars 1–2, $\kappa = 1/6$, TheoremCLattice and LemmaB_BetaInfinity.lean kernel-verified in Crossed/Transport.lean and Crossed/MassGap/ (~ 1900 lines, 0 `sorry`).

Acknowledgments. The author thanks the Bauerschmidt–Bodineau–Dagallier and Bauerschmidt–Dagallier $\varphi_{2,3}^4$ programs [17, 18] for providing the technical target framework against which the β -finite formal proof of Lemma B will be benchmarked, and acknowledges the lattice work of Athenodorou–Teper [2] and Bennett *et al.* [3] that anchors the cross- N and cross-group verification.

* kevin.remondier@gmail.com

- [1] A. Jaffe and E. Witten, *Quantum Yang–Mills theory*, Clay Mathematics Institute Millennium Problem description (2000).
- [2] A. Athenodorou and M. Teper, *SU(N) gauge theories in 3+1 dimensions: glueball spectra, string tensions and topology*, JHEP **12** (2021) 082; [arXiv:2106.00364](#).
- [3] E. Bennett *et al.*, *Sp(2N) lattice gauge theories and extensions of the Standard Model of particle physics*, EPJ Plus **139** (2024); [arXiv:2103.00665](#).
- [4] C. J. Morningstar and M. J. Peardon, *The glueball spectrum from an anisotropic lattice study*, Phys. Rev. D **60** (1999) 034509; [arXiv:hep-lat/9901004](#).
- [5] B. Lucini, M. Teper, and U. Wenger, *Glueballs and k-strings in SU(N) gauge theories: calculations with improved operators*, JHEP **06** (2004) 012; [arXiv:hep-lat/0404008](#).
- [6] J. Elstrodt, F. Grunewald, and J. Mennicke, *Groups acting on hyperbolic space: harmonic analysis and number theory*, Springer Monographs in Mathematics (1998).

- [7] P. Sarnak, *The arithmetic and geometry of some hyperbolic three-manifolds*, Acta Math. **151** (1983) 253–295.
- [8] K. Rémondrière, *A topological invariant of Bianchi 3-orbifolds from the identity term of the Selberg pretrace formula*, preprint, Comptes Rendus de l’Académie des Sciences (May 2026).
- [9] G. ’t Hooft, *A planar diagram theory for strong interactions*, Nucl. Phys. B **72** (1974) 461.
- [10] R. Dijkgraaf and E. Witten, *Topological gauge theories and group cohomology*, Commun. Math. Phys. **129** (1990) 393–429.
- [11] M. Lüscher and P. Weisz, *Quark confinement and the bosonic string*, JHEP **07** (2002) 049; [arXiv:hep-lat/0207003](#).
- [12] M. Lüscher and P. Weisz, *String excitation energies in $SU(N)$ gauge theories beyond the free-string approximation*, JHEP **07** (2004) 014; [arXiv:hep-th/0406205](#).
- [13] K. Rémondrière, *Theorem C: Bianchi-cohomological cross-group law for the Wilson log-Sobolev constant in lattice Yang–Mills* (v13, session 2026-05-23), preprint (May 2026).
- [14] M. Lüscher, *Properties and uses of the Wilson flow in lattice QCD*, JHEP **08** (2010) 071; [arXiv:1006.4518](#).
- [15] A. Chandra, I. Chevyrev, M. Hairer, and H. Shen, *Stochastic quantisation of Yang–Mills–Higgs in 3D*, Invent. Math. **237** (2024); [arXiv:2201.03487](#).
- [16] A. Chandra, I. Chevyrev, M. Hairer, and H. Shen, *Langevin dynamic for the 2D Yang–Mills measure*, Publ. Math. IHÉS **136** (2022) 1–147; [arXiv:2006.04987](#).
- [17] R. Bauerschmidt, T. Bodineau, and B. Dagallier, *Stochastic dynamics and the Polchinski equation: an introduction*, Probab. Surv. **21** (2024) 200–290; [arXiv:2307.07619](#).
- [18] R. Bauerschmidt and B. Dagallier, *Log-Sobolev inequality for the φ_2^4 and φ_3^4 measures*, Comm. Pure Appl. Math. **77** (2024) 2579–2612; [arXiv:2202.02295](#).
- [19] S. Cao, J. Park, and S. Sheffield, *Random surfaces and lattice Yang–Mills*, Commun. Amer. Math. Soc. (accepted, 2024); [arXiv:2307.06790](#).
- [20] S. Cao, R. Nissim, and S. Sheffield, *Dynamical approach to area law for lattice Yang–Mills* (2025); [arXiv:2509.04688](#).
- [21] S. Chatterjee, *A scaling limit of $SU(2)$ lattice Yang–Mills–Higgs theory*, Probab. Math. Phys. (accepted, 2024); [arXiv:2401.10507](#).
- [22] T. Balaban, *Renormalization group approach to lattice gauge field theories. I. Generation of effective actions in a small field approximation*, Commun. Math. Phys. **109** (1987) 249–301.
- [23] M. Fukushima, Y. Oshima, and M. Takeda, *Dirichlet Forms and Symmetric Markov Processes*, de Gruyter (1994).
- [24] K. Rémondrière, *No-go theorems for the arithmetic determination of the Yang–Mills mass gap absolute scale*, preprint, Physical Review Letters submission (May 2026).