A scalar finite-volume criterion for a continuum mass gap in $SU(2)$ lattice gauge theory: the doubling test in dimensions four and three, its continuum assembly and its converse

Kévin Rémondière*

12 November 2026 (version 1.2)

Abstract

This technical note transposes the two-dimensional finite-volume doubling criterion of OpenAI (Proposition 2.2 of *Sharp mass bounds for the two-dimensional $O(4)$ model*, 2026) to lattice gauge theories in dimensions four and three. It is predominantly a transposition, not a new mass-gap mechanism. For the $SU(2)$ Wilson theory on periodic tori set $\Delta_\beta(n) = 16 \log Z_\beta(n^4) - \log Z_\beta((2n)^4)$. (i) Under hypercubic symmetry and a positive trace-class transfer representation (a standard hypothesis, not formalised), $\Delta_\beta(n) \le 2^{-9}$ implies $\Delta_\beta(2n) \le 256\,\Delta_\beta(n)^2$ and $(\lambda_i/\lambda_0)^n \le 256\,\Delta_\beta(n) \le \frac{1}{2}$ for every excited eigenvalue of the transfer operator on every spatial torus $(2^j n)^3$; in dimension three the threshold is 2^{-7} and the constant 64. (ii) Fix $\ell > 0$ and $0 < \eta \le 2^{-9}$. Assume (W1)–(W3) and $\Delta_{\beta_k}(n_k) \le \eta$ for every k, at spacing $a_k = \ell/n_k$, and assume (W4): the lattice Osterwalder–Schrader forms converge, which is the existence of the continuum limit and is *assumed*. Then the limit satisfies $\langle v, e^{-\ell H} v\rangle \le 256\eta\|v\|^2$ on the complement of the vacuum, with the same constant for every k; given the Osterwalder–Schrader reconstruction (also assumed), it has a mass gap at least $-\log(256\eta)/\ell \ge (\log 2)/\ell$ and a unique vacuum. This implication is formalised in Lean 4 against our own Lean statement of the problem (ClayYM v1.1), with the transfer representation, the convergence and the reconstruction as named hypotheses (in the Lean statement the hypothesis on the transfer structure, `SpecLink`, is weaker than (W3), and the specialisation to Wilson's partition function requires a symmetric extension to periods 1 satisfying the spectral identities, which is an additional assumption, as stated under Table 1); the Lean sources were rebuilt on 11 November 2026 by non-authors on a clean machine, first from hashed copies of the sources and then from the git archive of the package tree (Appendix A). For square tori the passage from cylinders uses a further hypothesis (W2$^+$) and is proved on paper for one diagonal choice of the volumes only. (iii) Δ is bracketed by excited-state traces on spatial tori with sides in $\{\ell, 2\ell\}$; a finite-volume gap including the electric-flux sectors, together with a trace bound, gives an explicit sufficient inequality for the criterion; examples show that the criterion can fail when either ingredient is missing, but the example for the flux sectors has a trivial local limit, and the example for the trace bound consists of decoupled copies of an abstract theory, so that strictness within non-trivial theories, or within pure $SU(2)$, is not established. (iv) For a stipulated Maxwell spectrum (whose identification with a normalised Villain measure is not derived here) the defect exceeds the threshold at every size $n \ge 3$ and every coupling; a massive free scalar passes numerically for $M\ell \ge 11.47$. (v) A single implementation reports the sufficient bound $\Delta_\beta(2) \le 1.741 \cdot 10^{-3}$ for $\beta \le 0.005$. This interval is conditional on the finite enumeration (ENC) and rational comparison (RAT), neither independently certified here.

Relation to earlier public work. Version 5.0.0 of the series of Bhattacharjee et al. [3], whose Zenodo record was created on 11 November 2026, before the release of version 1.0 of this note, and which we read after that version was written, contains for four-dimensional tori the same

*Independent researcher, Oloron-Sainte-Marie, France. ORCID 0009-0008-2443-7166. The proofs, the programs, the Lean files and this note were prepared with the assistance of AI models (Claude Opus 5.5, Claude Sonnet 5.5 and Claude Fable 5.2, Anthropic; Qwen3.8, Qwen team; GPT-6 Astra and GPT-6.1 Sol, OpenAI; DeepSeek V4 Pro, DeepSeek); their roles are described in the statement on the use of AI at the end of the note.

doubling defect, its bracket by thermal traces, the purity reading, and a one-scale criterion whose hypothesis (B*), a family of inequalities between partition functions of finite tori, implies ours (§1.4). *Version 1.2 proves the reverse implication* (Theorem 4.1): under symmetry and a positive transfer representation, $\Delta_\beta(n) \le 2^{-9}$ implies (B*) at the scale $16n$, and the gap $(\log 2)/n$ on every integer cube of side $N \ge n$. The two criteria are therefore equivalent up to a change of scale: the reverse implication loses a factor 16 (not optimal), and a round trip, (B*) at $T_0 \Rightarrow \Delta(2T_0) \le 2^{-9} \Rightarrow$ (B*) at $32T_0$, a factor 32. The proof is paper-only for the aspect-ratio statement, and the main steps are checked by the Lean kernel in an abstract form (Appendix A). This does not answer a question left open in [3], whose data differ from ours, and it is not a proof of the mass gap. A corollary linking our criterion to their Theorem F is stated as conditional on their hypothesis (W) and their Theorem F as we read them. No priority is claimed for the one-scale criterion, the defect or the monotonicity of the free energy in a period (§1.4).

What the note does not do. It gives no estimate of Δ at weak coupling, no existence or non-triviality of the continuum limit, and no reconstruction. It does not apply as it stands to the formalised construction of OpenAI (Section 10). Reading Δ as a combination of Rényi-2 entropies of the thermal weights is a reformulation, not a result. The criterion is a sufficient condition; its equivalence with the Yang–Mills mass-gap problem is not claimed. The proofs have been checked by AI models only.

Version 1.1 (12 November 2026): literature corrections only. The principle that an inequality between partition functions of finite periodic tori (twisted against untwisted) implies decay is classical (Tomboulis–Yaffe, Kovács–Tomboulis, Kanazawa), and the comparison with these papers is recorded in §9.2.

Version 1.2 (12 November 2026). Adds Section 4 (Theorem 4.1, its remarks and the conditional Corollary 4.6), the second Lean file `lean/Reciproque.lean` and its combination with the published file `lean/main_v1_2.lean`, an example showing that positivity of the transfer operator cannot be dropped (Remark 4.2), and rewrites §1.4 and the related statements. The statements, proofs and Lean files of versions 1.0 and 1.1 are not changed: `lean/CritereScalaire.lean` and `lean/main.lean` are byte for byte those of version 1.0.

Status labels. [proved]: complete argument written in this note; [Lean]: also checked by the Lean 4 kernel (Appendix A), modulo the named hypotheses listed there; [computed]: floating-point or exact computation in the sandbox described in §11, never used in a proof unless stated; [read]: statement read in the cited source, location given; [hypothesis]: a named assumption, not proved here; [heuristic].

Contents

1 Introduction **3**

2 Setting, hypotheses and main theorem **7**

3 The lattice part **8**

4 The reverse implication: from the doubling test to a bound at one scale **10**

5 Assembly **13**

6 The converse and why it needs a finite-volume gap **14**

7 Witnesses **16**

8 Strong coupling and a reported computational instance **17**

9 Discussion **19**

10 Relation to the OpenAI construction **21**

11 Reproducibility **21**

A The Lean formalisation **22**

1 Introduction

1.1 The doubling test and where it comes from

Let $Z(n_1, n_2)$ be the periodic partition function of a two-dimensional spin model whose transfer operator is self-adjoint, positive and trace class. OpenAI [14, §2] introduce the *square-box test* $\Delta(n) = 4 \log Z(n, n) - \log Z(2n, 2n)$, in which the coefficients cancel every term proportional to the area, and prove [14, Prop. 2.2] [read]: if $n \geq 4$ is even and $\Delta(n) \leq \eta_* := 2^{-10}$, then the transfer operators on circumferences $2^k n$ satisfy $\lambda_2/\lambda_1 \leq e^{-(\log 2)/n}$, every local limit of the square tori of sides $2^k n$ has full gap at least $(\log 2)/n$, and $m_{\mathrm{lat}}(\beta) \geq (\log 2)/n$ if the periodic infinite-plane state is unique. The mechanism has four parts: the identity $e^{-p} = \sum_i t_i^2$ for the normalised trace weights and the squaring lemma [14, Lemma 2.1]; the rectangle identity [14, (2.7)]; the quadratic recursion $\Delta(2r) \leq 44\Delta(r)^2$ [14, (2.12)]; and the comparison of square tori with infinite cylinders.

The present work is predominantly a transposition of this mechanism to lattice gauge theories in dimensions $d = 4$ and $d = 3$, followed by a conditional continuum statement; it is not a new mass-gap mechanism. We claim no priority for the mechanism. The transposition to $d = 4$ was first written in internal, unpublished work of the same project (25 October 2026: threshold 2^{-12}, constant 512, gap $(\log 8)/n$; and 10 November 2026: all dimensions, constants in Table 2); the present note supersedes those notes.

The identity $e^{-p} = \sum_i t_i^2$ says that p is the Rényi-2 entropy of the normalised thermal weights t_i; OpenAI already read it as a purity [14, §2]. Reading Δ_β as a combination of such Rényi-2 entropies of the transfer spectrum is a reformulation of the same quantities: it adds no hypothesis and no result (it was written in our internal notes of 25 October 2026), and nothing below depends on it. The same reading, as the purity $\mathrm{Tr}\,\rho^2$ of a thermal state, is made for four-dimensional tori in [3, after Proposition 7.57].

1.2 Results

Let $F_\beta = \log Z_\beta$ on rectangular tori $\Lambda(\vec{n}) = \prod_{\mu=1}^4 \mathbb{Z}/n_\mu\mathbb{Z}$ (§2) and

$$\Delta_\beta(n) = 16\, F_\beta(n, n, n, n) - F_\beta(2n, 2n, 2n, 2n), \qquad \Delta_\beta^{(3)}(n) = 8\, F_\beta(n, n, n) - F_\beta(2n, 2n, 2n). \quad (1.1)$$

- *Lattice part* (Theorem 3.4, [proved], [Lean]). Under (W1)–(W2): $\Delta_\beta(n) \leq 2^{-9}$ implies $\Delta_\beta(2n) \leq 256\,\Delta_\beta(n)^2$ and, on every spatial torus $(2^j n)^3$, $(\lambda_i/\lambda_0)^n \leq 256\,\Delta_\beta(n)$ for every excited eigenvalue of the transfer operator. In $d = 3$: threshold 2^{-7}, constant 64 (Theorem 3.5).

- *Reverse implication* (Theorem 4.1, [proved], [Lean] for its main steps in abstract form). $\Delta_\beta(n) \leq 2^{-9}$ implies the one-scale bound (B*) of [3] at the scale $16n$, and the gap $(\log 2)/n$ on every integer cube of side $N \geq n$ (hence not only the dyadic cubes). Positivity of the transfer operator is used and cannot be dropped. A corollary linking the criterion to Theorem F of [3] is *conditional on their hypothesis (W) and their Theorem F as we read them, and not verified by us.*

- *Conditional assembly* (Theorem 2.2, [proved], [Lean] modulo (W2)–(W4), (W6)). If $0 < \eta \leq 2^{-9}$ and $\Delta_{\beta_k}(n_k) \leq \eta$ uniformly in k, with $a_k = \ell/n_k$, then every limit of the cylinder OS forms, in the sense of (W3)–(W4), has the q-form with $q = 256\eta$. Under the reconstruction hypothesis (W6) its mass gap is at least $-\log(256\eta)/\ell \geq (\log 2)/\ell$ and its vacuum is unique. For square tori the same conclusion needs (W5) for the family actually considered; Corollary 5.4 proves it for one diagonal choice of volumes, not for an arbitrary sequence $j_k \to \infty$.

- *Thermal-trace converse* (Proposition 6.1, Corollary 6.2, [proved]). Δ is bracketed, up to the factor $2(2^d-1)$, by the excited trace at time ℓ on d spatial tori with sides in $\{\ell, 2\ell\}$. A finite-volume gap (flux sectors included) and a trace bound give an explicit sufficient inequality for the criterion at a fixed physical size; eventual smallness as $\ell \to \infty$ needs control of the dependence of the trace bound on ℓ. Examples 6.3 and 6.4 show that neither the finite-volume gap nor the trace bound can be dropped.

- *Spectral witnesses* (§7). For the stipulated Maxwell (photon, rotor, flux) spectrum, whose identification with a normalised monopole-free Villain measure is not derived here, the defect exceeds the threshold by a factor ≥ 90 at every size $n \geq 3$ and every coupling [proved] given that spectrum (numerically by $\geq 1.7 \cdot 10^4$ [computed]). A massive free scalar passes numerically for $M\ell \geq 11.47$ ($d = 4$) and $M\ell \geq 8.47$ ($d = 3$) [computed]; these crossings come from a truncated spectrum and carry no tail certificate.

- *Reported computational instance* (Proposition 8.1). Conditional on (ENC) and (RAT), $\Delta_\beta(2) \leq 1.741 \cdot 10^{-3} < 2^{-9}$ for the $SU(2)$ Wilson theory and $0 \leq \beta \leq 0.005$. The analytic implication is proved on paper; the two computational inputs are reported from one implementation and remain uncertified here. The gap is already known in this regime.

1.3 What is new and what is not

Not new. The doubling mechanism, in $d = 2$ for spins [14]. The positive, self-adjoint transfer operator of lattice gauge theories [13, 17]. The closure lemma "same q along any filter" and the passage from the q-form to a gap through a named reconstruction hypothesis, which are part of our archived Lean statement ClayYM v1.1 [20]. The form of the result, "a local hypothesis at a fixed physical scale ℓ, uniform in the lattice spacing, plus existence of a limit, implies a gap of every limit", which is that of Theorem B of Bhattacharjee et al. [2] and of Theorem 5.3 of Rajabu [18], with different hypotheses (§9.2). The doubling defect of four-dimensional tori, its decomposition over the four directions, its bracket by thermal traces and a one-scale criterion with a gap conclusion, which are in version 5.0.0 of Bhattacharjee et al. [3] (§1.4). The unconditional squaring bound $p(2m) \leq 2(e^{p(m)}-1)^2$ (Lemma 3.1) is implicit in the chain of inequalities of [14, proof of Lemma 2.1]. Step 1 of the proof of Theorem 4.1 rests on the elementary monotonicity of the counting-measure ℓ^p norms: for $0 < L \leq L'$, whenever the traces are finite, $(\operatorname{Tr} T^{L'})^{1/L'} \leq (\operatorname{Tr} T^L)^{1/L}$ for a positive operator T. Lemmas 7.12 and 7.44 of [3] contain related monotonicity statements (of the thermal trace in the time, and of the tube energy in a side); the precise statement on the pressure used here follows from the spectral representation. The same concavity argument, with dyadic enlargement in every direction, is used long before by Brydges, Fröhlich and Seiler [4, proof of (2), p. 38 of the preprint], to bound a lattice gauge partition function below by a fractional power of its value on larger periodic rectangles: $\operatorname{Tr}(T^{L_0}) \geq (\operatorname{Tr} T^{2^{m_0}})^{L_0/2^{m_0}}$ for a positive semi-definite transfer matrix T and $2^{m_0} \geq L_0$, then the same in the other directions, which gives $\langle e^{-A_\Lambda}\rangle_0 \geq \langle e^{-A_{\widetilde{\Lambda}}}\rangle_0^{|\Lambda|/|\widetilde{\Lambda}|}$ in their notation (read on the image of the page; the optical character recognition of our copy garbles the exponents). We claim no novelty for the monotonicity.

The principle that an inequality between the partition functions of finite periodic tori, twisted against untwisted, implies a decay: for Wilson loops of $SU(2)$, $\langle W[C]\rangle \leq 2(e^{-F_{\mathrm{el}}})^{|A_C|/|A_{12}|}$ with $e^{-F_{\mathrm{el}}} = \langle \frac{1}{2}(1 - \mathcal{O}[\mathcal{V}])\rangle$, in Kovács and Tomboulis [10, eqs. (6) and (11), pp. 4–5], and restated as Theorem 1 of Kanazawa [8, p. 5], who attributes it to Tomboulis and Yaffe [19] and to [10]; for $SU(N)$ and $\mathbb{Z}_N$ fluxes, in Kanazawa [8, Theorem 2, p. 7]; and, for correlations of classical spin systems, in Kanazawa [8, Theorem 5, p. 17], for $L_\mu = 2^k n$: $|\langle \Gamma_\mu(x; n)\rangle|/\langle \Gamma_\mu(x; 0)\rangle \leq 2\{\langle \Gamma_\mu(x; 0)^2\rangle/\langle \Gamma_\mu(x; 0)\rangle^2\}^{n/L_\mu}\{1 - \int dg\, \langle \mathcal{O}(g)[\mathcal{V}]\rangle\}^{n/L_\mu}$. His abstract (p. 1) states: "Basically the inequalities guarantee that a system in a box that is sufficiently insensitive to boundary conditions has a non-zero mass gap." The hypothesis there is a ratio of twisted to untwisted partition functions, not the doubling defect, and the conclusion is a decay in a box of period L, not a bound on the transfer matrix of every spatial torus (§9.2); the family of the argument (reflection positivity,

Schwarz inequality, doublings) is the same as ours. The absence of these papers from the list of version 1.0 was a gap of the anteriority search.

Incremental content. The contribution is predominantly a routine transposition of the doubling mechanism of [14]. It consists of the higher-dimensional bookkeeping and the explicit bounds 256 at 2^{-9} in $d = 4$ and 64 at 2^{-7} in $d = 3$; the packaging of the conditional implication against the Clay-type q-form; an elementary thermal-trace converse with counterexamples to omitted hypotheses; and the witness analysis. The torus–cylinder comparison follows the insertion argument of [14], with an additional diagonal qualification when renormalised lattice observables depend on the cutoff. The reported $SU(2)$ interval up to $\beta = 0.005$ is conditional on a finite enumeration and a rational comparison that are not independently certified. We claim no new mass-gap mechanism, no optimality of the constants, and no exhaustive priority determination: the anteriority search of §9.2 is limited and recorded. In [14] it found neither a converse nor a statement in dimension $d > 2$. In [3], read on 12 November 2026 after the first draft of version 1.0 had been written, it found the four-dimensional defect, its bracket and a one-scale criterion; the part of the content listed above that is also there, and what we did not find there, are stated in §1.4.

1.4 Relation to version 5.0.0 of Bhattacharjee et al.

Version 5.0.0 of the series of Bhattacharjee et al. [3] has a Zenodo record created on 11 November 2026 at 23:29:13 UTC, before the release of this note. We read it on 12 November 2026, after the first draft of version 1.0 had been written, in its abstract, its introduction and its Sections 7.8 and 7.11; for version 1.2 we also read the lemmas and propositions cited in Section 4 and here (7.3, 7.12, 7.27, 7.33, 7.44, 7.45, 7.48, 7.51, 7.56, 7.57, 7.59) and searched the whole text by keywords. For a spatial torus s write $\widehat{Z}_s(M) = \sum_{i\geq 0} r_i^M$, where $1 = r_0 \geq r_1 \geq \cdots \geq 0$ are the eigenvalues of the transfer operator of s divided by the largest one (their relative partition function, the thermal trace at time M), and $V(s)$ for the number of sites of s. Those passages contain the following [read].

(a) *The object and its bracket* (their Lemma 7.33 and Proposition 7.34). For a four-dimensional torus $n = (n_1, \dots, n_4)$ with all sides at least 3: the doubling defect $\delta(n) = 16 \log Z(n) - \log Z(2n)$, its invariance under $Z(n) \mapsto e^{a|n|} Z(n)$, its decomposition over the four directions with the weights $8, 4, 2, 1$, and

$$8 \log \widehat{Z}_{(n_1,n_2,n_3)}(n_4) \;\leq\; \delta(n) \;\leq\; 16 \log \widehat{Z}_{(n_1,n_2,n_3)}(n_4) + 8 \log \widehat{Z}_{(n_1,n_2,2n_4)}(n_3) + 4 \log \widehat{Z}_{(n_1,2n_3,2n_4)}(n_2) + 2 \log \widehat{Z}_{(2n_2,2n_3,2n_4)}(n_1).$$

On cubic tori $\delta(n)$ is our $\Delta_\beta(n)$, the decomposition is (3.1), the upper bound is that of Proposition 6.1, and the lower bound is the first term of the lower bound of that proposition.

(b) *A one-scale criterion* (their Theorem 7.56 and Proposition 7.57). Hypothesis (B*), for an integer $T_0 \geq 3$: $\log \widehat{Z}_s(T_0) \leq \gamma V(s)$ for every spatial torus s with all its sides in $[2T_0, 48T_0)$, where $2\gamma(96T_0)^3 \leq \frac{1}{20}$. Conclusion: a doubly exponential decrease of these thermal traces along the times $2^i T_0$, and a gap at least $0.172/T_0$ for the transfer operator of every spatial torus whose smallest side a is at least $2T_0$ and whose largest side is less than $24a$. Up to a factor 2 in γ, (B*) is the family of inequalities $Z(s, T_0)^2 \leq e^{\gamma V(s)} Z(s, 2T_0)$ between partition functions of finite tori, which they read as near purity, $\operatorname{Tr} \rho^2 \geq e^{-\gamma V(s)}$, of the thermal state.

(c) *Scope remarks and a continuum statement.* Such a hypothesis contains a lattice gap (their Proposition 7.58); a gap and the free energy per site do not control the doubling defect (their Proposition 7.35); and, for continuum families satisfying their hypothesis (W), which they describe as "the shape of the conclusion claimed by the unrefereed Lean 4 construction" (their references [45, 62]; their [45] is the Wilson Yang–Mills construction of OpenAI), (B*) at one physical time extent t_0 gives a Hamiltonian with spectrum in $\{0\} \cup [0.172/t_0, \infty)$ and a simple eigenvalue 0 (their Theorem F).

What is theirs. Items (a)–(c) are earlier than this note. They cover the object Δ, the decomposition (3.1), the bracket of Proposition 6.1 in $d = 4$, the purity reading, the doubly exponential mechanism in $d = 4$, and the observation that the criterion contains a finite-volume gap. We claim no priority for any of them; our internal notes [21] are unpublished and establish none.

Their hypothesis implies ours. Let $n = 2T_0$. The four spatial tori that enter the upper bound of (a) for the cubic torus n^4 have sides in $\{2T_0, 4T_0\}$, hence in the window of (B*), and volumes $2^j n^3$, $j = 0, \dots, 3$. Since $M \mapsto \widehat{Z}_s(M)$ is non-increasing, (B*) gives $\log \widehat{Z}_s(n) \le \log \widehat{Z}_s(T_0) \le \gamma V(s)$ for each of them, and

$$\Delta_\beta(2T_0) \;\le\; \gamma n^3 \,(16 + 8 \cdot 2 + 4 \cdot 4 + 2 \cdot 8) \;=\; 512\,\gamma\, T_0^3 \;\le\; \frac{512}{40 \cdot 96^3} \;=\; \frac{1}{69120} \;<\; 2^{-9}. \qquad (1.2)$$

[proved], from the statements of [3] as read. *Conversely*, under (W1)–(W2), $\Delta_\beta(n) \le 2^{-9}$ implies (B*) at $T_0 = 16n$ with $\gamma = 1/(31 \cdot 2^{23} T_0^3)$ (Theorem 4.1, Section 4). Version 1.0 and 1.1 of this note said that we did not know this; the proof was found on 12 November 2026. Hence, up to a change of scale (a factor 16 in the reverse direction, not optimal; 32 for a round trip), the two criteria are equivalent. Note that by Proposition 6.1 alone the defect bounds the thermal traces of four boxes only; the extension to all tori of the window is obtained from the monotonicity of the pressure in each period (Step 1 of Theorem 4.1).

What differs. Our hypothesis is one inequality between the partition functions of the two cubic tori n^4 and $(2n)^4$; theirs is a family of inequalities, one for each spatial box with sides in a window. Their conclusion is wider than that of Theorem 3.4: every torus of bounded aspect ratio above the base scale, not only the dyadic cubes $(2^j n)^3$. Theorem 4.1 shows that this difference is one of scale, not of substance: the two hypotheses are equivalent up to a change of scale (a factor 16 in the reverse direction, 32 for a round trip), and the conclusion of Theorem 3.4 extends to all integer cubes of side $\ge n$ and, with a worse constant, to all spatial tori with $\min_i t_i \ge n$ and $\max_i t_i \le 24 \min_i t_i$. For the Wilson theory in $d = 4$, the gap statement (3.2) of Theorem 3.4 is thus a variant of that of their Theorem 7.56; their recursion is on normalised bounds of the thermal traces, $y_{i+1} = 16(e^{y_i} - 1)^2$, not on the defect. In the passages read we did not find: the recursion $\Delta(2n) \le 256\,\Delta(n)^2$ for the cubic defect itself, and with it a gap on all dyadic sections from a single inequality; the three-dimensional statement (Theorem 3.5); a Lean formalisation of such a criterion (Appendix A); the witnesses of §7; the reported instance of Proposition 8.1; and, by the complete reading and keyword search of 12 November 2026, the implication from a small defect at one size to (B*) (Theorem 4.1), the comparison of $f(m^4)$ with the limiting pressure of a torus by telescoping the pressure differences along $m_j = 2^j n$ (with the constant $2/31$), and the passage from a small $\Delta(n)$ at one size n to a gap on all cubes of side $N \ge n$. This is not a proof of absence. What *is* in [3] and bears on this: the telescoping $\log \widehat{Z}(M) = \sum_{k \ge 0} 2^{-k-1} \psi(2^k M)$ for one transfer operator (their Lemma 7.33, a different telescoping, in the time), and, from Proposition 7.34(ii), the fact that a small defect $\delta(n)$ of the torus n *itself*, dyadic or not, bounds the thermal trace of its spatial torus (n_1, n_2, n_3) at the time n_4 (our bracket, Proposition 6.1). So the quantity Δ and the non-dyadic case at the size where the defect is small are theirs; the new points are the implication to (B*) (and (B**), Remark 4.4) and the extension to all sizes $N \ge n$ from one size n. Their continuum statement (c) and Theorem 2.2 have different hypotheses, and we have not compared them; Corollary 4.6 is conditional on (c) as we read it.

Limits of this comparison. It was made by AI models (see "Use of AI") from the passages cited, on 12 November 2026; the rest of the 157 pages was searched by keywords only. We have not examined in which version of that series each item first appeared, and no human expert has checked the comparison.

1.5 Status of the proof

The abstract implication from $\Delta \le \eta$ to the q-form and, under the normalised reconstruction hypothesis, to `HasMassGap` of ClayYM is formalised in Lean 4 (35 declarations, standard axioms only) under the stated positive-period spectral hypotheses. The paper proof on periods at least

2 does not require the extension to periods 1 described under Table 1; its identification with that particular Lean instance is conditional on that extension. The build was reproduced on 11 November 2026 by non-authors on a clean machine, first from hashed copies of the sources and then from the git archive of the package tree (Appendix A). The transfer-spectrum representation (W2)–(W3), its identification with the lattice OS forms, the convergence (W4) and the reconstruction (W6) are named hypotheses and are not formalised; (W4) is the existence of the continuum limit and is open. The insertion estimate and the quantitative torus–cylinder comparison are paper arguments (Lean assumes that the difference tends to zero); the thermal-trace converse of Section 6 is a paper argument. The reverse implication of Section 4 is a paper argument whose steps (a), (b) and the first half of (c) are also checked by the Lean kernel in an abstract form, with the symmetry, the spectral representation and the published doubling recursion as named hypotheses; the aspect-ratio statement (d), the logarithmic form of (c), the formula of Corollary 4.6 and its link with [3] are not formalised. The Maxwell model identification is incomplete, the scalar crossings are numerical, and the strong-coupling interval is conditional on an enumeration and a rational comparison reported by one implementation. The original formalisation and draft were prepared within one model family. Read-only reviews by Qwen3.8 and GPT-6.1 Sol concerned the draft of 11 November, not this final text: they found the core implication sound subject to corrections and left the computational inputs uncertified; Claude Sonnet applied adaptations of their proposed corrections, and some proposed additions were not adopted. A read-only review of the first version of this text by GPT-6.1 Sol required sixteen corrections, which Claude Sonnet adapted. A further read-only check of this revision identified remaining corrections, recorded in `RELECTURE_SOL_TEXTE_CORRIGE.md` (one blocking and four non-blocking: the conditional status of Proposition 8.1 in the abstract and the list of results, the description of `SpecLink`, the concordance table, the attribution of the reviews, the wording of the rebuild descriptions); they are applied in this version. These reviews do not certify the computational inputs or endorse a subsequent revision. The text–Lean concordance was prepared within the author's model family. Appendix A states, for each formalised statement, what the Lean declaration says exactly. No human has checked the proofs.

2 Setting, hypotheses and main theorem

2.1 Model

For $\vec{n} = (n_1, \dots, n_4)$ with $n_\mu \geq 2$, let $\Lambda(\vec{n}) = \prod_\mu \mathbb{Z}/n_\mu\mathbb{Z}$, with links (x, μ) and plaquettes $p = (x; \mu, \nu)$, $\mu < \nu$; for $n_\mu \geq 2$ every plaquette has four distinct links. The Wilson action and partition function are

$$S_\beta(U) = \beta \sum_p (1 - \tfrac{1}{2} \operatorname{Tr} U_p), \qquad Z_\beta(\vec{n}) = \int e^{-S_\beta(U)}\, dU, \qquad F_\beta(\vec{n}) = \log Z_\beta(\vec{n}), \tag{2.1}$$

with $U_p = U_{x,\mu} U_{x+\mu,\nu} U_{x+\nu,\mu}^{-1} U_{x,\nu}^{-1}$, dU the product of normalised Haar measures on $SU(2)$ and $\beta = 4/g^2$; $Z_0 = 1$. At lattice spacing a, a period n is a physical length na. The three-dimensional model is defined in the same way on $\prod_{\mu=1}^3 \mathbb{Z}/n_\mu\mathbb{Z}$. For a profile m (a multiset of four periods) we write $\tau_\mu m$ for the profile with the μ-th period doubled and

$$p_\mu(m) = 2F(m) - F(\tau_\mu m). \tag{2.2}$$

2.2 Hypotheses

Table 1 lists every hypothesis used, with its status. The names (W1)–(W6) are used throughout, and in the Lean files.

Under (W2), let $\lambda_0 \geq \lambda_1 \geq \lambda_2 \geq \cdots \geq 0$ be the eigenvalues of K_σ with multiplicity, $r_i = \lambda_i/\lambda_0 \in [0, 1]$ $(i \geq 1)$ and $s_\sigma(N) = \sum_{i \geq 1} r_i^N$ (finite for $N \geq 1$). Then

$$F(\sigma, N) = N \log \lambda_0 + \log(1 + s_\sigma(N)). \tag{2.3}$$

Remark 2.1 (on (W2) and (W3)). For $\beta > 0$, the character expansion $e^{(\beta/2)\operatorname{Tr} g} = \sum_j c_j(\beta)\chi_j(g)$ has $c_j(\beta) = 2(2j+1)I_{2j+1}(\beta)/\beta > 0$ (at $\beta = 0$ one uses the continuous limit); this gives the positive convolution factor of the usual temporal-gauge construction. Multiplication by the spatial plaquette weights and projection onto gauge-invariant functions (the Polyakov loops inserted by fixing the temporal gauge on $N-1$ slices) must then be combined with all per-slice scalar factors of (2.1) to give the precise identity in (W2). Reflection positivity and the positive transfer operator: [13] is cited for the standard transfer construction, not re-derived here; [17, §2.3] [read]. The present paragraph is a sketch, not a complete derivation of (W2)–(W3) with our conventions; for Lemma 5.3, (W2$^+$) must also hold after gauge projection. The centre transformations on the links crossing a spatial hyperplane commute with K, so that $\mathcal{H}_{\mathrm{inv}} = \bigoplus_{e\in\mathbb{Z}_2^3} \mathcal{H}_e$ (electric-flux sectors), and the periodic trace contains *all* sectors. (W3) is the spectral theorem for the compact positive operator $\widehat{K}_\sigma$ on $\mathcal{H}_{\mathrm{inv}}(\sigma)$ (eigenbasis $e_0 = \Omega, e_1, \dots$, kernel included), together with the formula $B(F, G, ma) = \langle \psi(F), \widehat{K}^m \psi(G)\rangle$ for the reflected two-point function of the cylinder state; in coordinates $c_i(F) = \langle e_i, \psi(F)\rangle$ it reads $B(F,F,0) = \sum_i c_i(F)^2$, $B(F,F,na) = c_0(F)^2 + \sum_{i\ge 1} r_i^n c_i(F)^2$, $B(\Omega, F, 0) = c_0(F)$, $c(\Omega) = \delta_0$. The spectrum (r_i) is the same as in (2.3).

2.3 Main theorem

ClayYM [20] writes Euclidean Schwinger kernels S for the 21 components of $\operatorname{Tr} F_{\mu\nu}F_{\rho\sigma}$, their Osterwalder–Schrader form rpForm $R\,S\,(F,G,t) = \langle F, e^{-tH} G\rangle$ on finite families F, G of positive-time test functions, the truncation $v_A = A - \langle \Omega, A\rangle \Omega$, and

$$\mathrm{QGap}(R, S, \mathcal{A}_0, \ell, q): \quad 0 < \ell,\ 0 < q < 1,\ \text{and } \langle v_A, e^{-\ell H} v_A\rangle \le q\,\langle v_A, v_A\rangle \text{ for every } A \in \mathcal{A}_0. \tag{2.4}$$

Theorem 2.2 (assembly, $d = 4$)**.** *Let $(\beta_k, n_k)_k$ be couplings and integers $n_k \ge 2$, let $\ell > 0$ and $a_k = \ell/n_k$, let (j_k) be any integers ≥ 0, and let B_k be the lattice OS form of the cylinder $\mathbb{Z}\times(2^{j_k}n_k)^3$ at spacing a_k and coupling β_k. Assume (W1)–(W3) for every k, and*

$$\Delta_{\beta_k}(n_k) \le \eta \quad \text{for every } k, \qquad 0 < \eta \le 2^{-9}. \tag{2.5}$$

If $B_k \to$ rpForm $R\,S$ along a non-trivial filter in the sense of (W4), with $\mathcal{A}_0$ the set of all families, then

1. *$\mathrm{QGap}(R, S, \text{all families}, \ell, 256\eta)$, with the same $q = 256\eta \le \frac{1}{2}$ for every k;*
2. *under (W6), S has a mass gap $-\log(256\eta)/\ell \ge (\log 2)/\ell$ (`HasMassGap` of ClayYM), and every vector fixed by $e^{-\ell H}$ is a multiple of Ω; if $\eta \le 2^{-12}$, the gap is $\ge (\log 16)/\ell$.*

The same conclusions hold for a convergent family of square-torus OS forms if (W5) holds for that family. Under (W2$^+$), Corollary 5.4 establishes (W5) for its specified diagonal choice of $N_k = 2^{j_k}n_k$; this choice does not, by itself, preserve a continuum limit previously obtained along another volume sequence.

The physical time ℓ corresponds exactly to n_k lattice steps. If $j_k \to \infty$, the spatial side $2^{j_k}\ell$ diverges, as required by the lattice sequences of ClayYM. The closure argument applies along any specified non-trivial filter for which the required convergence holds. An iterated limit ("volume, then spacing") is not in general convergence along the product filter and the two must not be identified: for an iterated limit, apply Lemma 5.2 successively, with the same q at both stages. The conclusion is compatible with a trivial limit: it proves neither the existence of a continuum limit nor a nonzero excited sector, and **non-triviality is not addressed**.

3 The lattice part

Lemma 3.1 (squaring lemma)**.** *Let $r_i \in [0,1]$ with $s(N) = \sum_i r_i^N < \infty$ for $N \ge 1$, and $p(N) = 2\log(1+s(N)) - \log(1+s(2N))$. Then (a) $0 \le \log(1+s(N)) \le p(N)$, so that $r_i^N \le s(N) \le e^{p(N)} - 1$; (b) $p(2N) \le 2\left(e^{p(N)} - 1\right)^2$, without any smallness condition. [proved] [Lean: `lemme_fin_tc`]*

Proof. (a) $s(2N) = \sum_i (r_i^N)^2 \le \sum_i r_i^N = s(N)$ since $r_i^N \le 1$; hence $\log(1+s(2N)) \le \log(1+s(N))$ and $p(N) \ge \log(1+s(N)) \ge 0$. (b) $p(2N) = 2\log(1+s(2N)) - \log(1+s(4N)) \le 2s(2N) \le 2s(N)^2 \le 2(e^{p(N)}-1)^2$, using $\log(1+x) \le x$, $\log(1+s(4N)) \ge 0$ and $\sum_i (r_i^N)^2 \le (\sum_i r_i^N)^2$. □

By (W1)–(W2) and (2.3), every $p_\mu(m)$ is the quantity $p(N)$ of Lemma 3.1 for the transfer operator in direction μ on the section formed by the other three periods of m, with N the μ-th period (the eigenvalue λ_0 cancels). In particular $p_\mu(m) \ge 0$.

Lemma 3.2 (rectangle identity and decomposition)**.** *For every function F on profiles and $\mu \ne \nu$: (R) $p_\mu(\tau_\nu m) = 2p_\mu(m) - 2p_\nu(m) + p_\nu(\tau_\mu m)$. If F is symmetric (W1), then*

$$\Delta(n) = 8a_0(n) + 4a_1(n) + 2a_2(n) + a_3(n), \tag{3.1}$$

where $a_j(n)$ is p at the profile $((2n)^j, n^{4-j})$, doubling one period n. Under (W1)–(W2), $\Delta(n) \ge 0$ and $a_j(n) \le \Delta(n)/2^{3-j}$. [proved] [Lean: `decomp_4d`, which states $\Delta \ge 0$, $a_0 \le \Delta/8$ and $a_3 \le \Delta$; the identities (R) and (3.1) are steps inside the proofs of `decomp_4d` and `doublement_fin_4d`]

Proof. (R): compute $F(\tau_\mu\tau_\nu m) = 2F(\tau_\nu m) - p_\mu(\tau_\nu m) = 2F(\tau_\mu m) - p_\nu(\tau_\mu m)$ and substitute $F(\tau_\nu m) = 2F(m) - p_\nu(m)$, $F(\tau_\mu m) = 2F(m) - p_\mu(m)$. (3.1): double the four periods one after the other; the sum telescopes, and (W1) identifies each intermediate profile. □

For $j = 0, \dots, 3$ let G_j be p at the profile $(n, (4n)^j, (2n)^{3-j})$ doubling the period n, and A_j be p at the profile $(2n, (4n)^j, (2n)^{3-j})$ doubling one period $2n$ in the same direction. Then:

1. $G_0 = a_3(n) \le \Delta(n)$;
2. $A_j \le f(G_j)$, $f(x) = 2(e^x - 1)^2$: Lemma 3.1(b) in the direction of the period n of G_j;
3. $G_{j+1} \le 2G_j + A_j$: (R) with $m = (n[\mu], 2n[\nu], (4n)^j, (2n)^{2-j})$ gives $G_{j+1} = 2G_j - 2p_\nu(m) + A_j$, and $p_\nu(m) \ge 0$;
4. $\Delta(2n) = 8A_0 + 4A_1 + 2A_2 + A_3$: (3.1) at level $2n$.

Proposition 3.3 (doubling, $d = 4$)**.** *Under (W1)–(W2), if $\Delta(n) \le 2^{-9}$ then $\Delta(2n) \le 256\,\Delta(n)^2$. [proved] [Lean: `doublement_fin_4d`, `chaine_num_fin4`; the Lean theorem takes as hypotheses the two consequences of (W2) that the proof uses (positivity of p and the squaring bound, from `hyp_of_specTC`)]*

Proof. Put $D = \Delta(n) \le 2^{-9}$. On $[0,1]$, $e^x - 1 \le x + x^2$ (Mathlib: $|e^x - 1 - x| \le x^2$ for $|x| \le 1$), so $f(x) \le 2x^2(1+x)^2$. By items 1–3, $G_0 \le D$, $G_1 \le 2D + 2D^2(1+D)^2 \le 2.004\,D$, $G_2 \le 4.024\,D$, $G_3 \le 8.113\,D$, and by item 4

$$\begin{aligned}\Delta(2n) &\le 8f(G_0) + 4f(G_1) + 2f(G_2) + f(G_3)\\ &\le (16.064 + 32.384 + 65.8 + 135.9)\,D^2 = 250.148\,D^2 \le 250.15\,D^2 \le 256\,D^2.\end{aligned}$$

□

The coefficients above are rounded upward; the exact rational polynomial expression, evaluated at $D = 2^{-9}$, is approximately 250.018080; with the exact exponential the constant is 247.23, and its limit as $D \to 0$ is $2^d(2^d - 1) = 240$ (Table 2).

Theorem 3.4 (transfer gap on all sections)**.** *Under (W1)–(W2), if $\Delta(n) \le 2^{-9}$ then, for every $k \ge 0$, $256\,\Delta(2^k n) \le (256\,\Delta(n))^{2^k}$, and for every $j \ge 0$ every excited eigenvalue of the transfer operator on the section $(2^j n)^3$ satisfies*

$$(\lambda_i/\lambda_0)^n \le 256\,\Delta(n) \le \tfrac{1}{2}. \tag{3.2}$$

If $\Delta(n) > 0$ and $\lambda_i > 0$ this gives $-\log(\lambda_i/\lambda_0) \ge -\log(256\Delta(n))/n \ge (\log 2)/n$; zero eigenvalues satisfy the ratio estimate directly, and if $\Delta(n) = 0$ every excited ratio is zero. In particular λ_0 is simple. [proved] [Lean: `iteration256`, `gap_4d_fin`, which states $r_i^n \le 256\,\Delta(n)$; the bound $\le \frac{1}{2}$, the logarithmic form and the simplicity are immediate consequences not restated in Lean]

Proof. If $d_k := \Delta(2^k n) \le 2^{-9}$ then $256 d_k \le \frac{1}{2}$ and $d_{k+1} \le 256 d_k^2 \le d_k/2$; by induction $256 d_k \le (256 d_0)^{2^k}$. On the section $(2^j n)^3$ at time $2^j n$, Lemma 3.1(a) and (3.1) give (with $a_0 \le \Delta/8 \le 2^{-12} \le 1$, so that $e^x - 1 \le 2x$ applies) $r_i^{2^j n} \le e^{a_0(2^j n)} - 1 \le 2a_0(2^j n) \le \Delta(2^j n)/4 \le (256\Delta(n))^{2^j}$; take the 2^j-th root. □

Theorem 3.5 ($d = 3$)**.** *For the three-dimensional theory under the analogues of (W1)–(W2), $\Delta^{(3)}(n) = 4a_0 + 2a_1 + a_2$; if $\Delta^{(3)}(n) \le 2^{-7}$ then $\Delta^{(3)}(2n) \le 64\,\Delta^{(3)}(n)^2$ and $(\lambda_i/\lambda_0)^n \le 64\,\Delta^{(3)}(n) \le \frac{1}{2}$ on every section $(2^j n)^2$. [proved] [Lean: `doublement_fin_3d`, `gap_3d_fin` (conclusion $r_i^n \le 64\,\Delta^{(3)}(n)$)]*

Proof. Same chain with $G_0 \le D \le 2^{-7}$, $G_1 \le 2.016D$, $G_2 \le 4.098D$ and $4f(G_0) + 2f(G_1) + f(G_2) \le 60.68\,D^2 \le 64D^2$ (the polynomial expression evaluates numerically to approximately 60.655245); on the section $(2^j n)^2$, $r_i^{2^j n} \le 2a_0(2^j n) \le \Delta^{(3)}(2^j n)/2 \le (64\Delta^{(3)}(n))^{2^j}$. □

4 The reverse implication: from the doubling test to a bound at one scale

Section 1.4 shows that the one-scale hypothesis (B*) of [3] implies $\Delta_\beta(2T_0) \le 2^{-9}$. This section proves the reverse implication, at a larger scale. We use the notation of §2. For a spatial torus $t = (t_1, t_2, t_3)$ with all sides at least 3 put $V(t) = t_1 t_2 t_3$, $s_t(M) = \sum_{i \ge 1} r_i^M$ for the excited ratios $r_i = \lambda_i/\lambda_0$ of the transfer operator K_t of (W2), and

$$h_t(M) = \log(1 + s_t(M)) = \log \frac{Z(t, M)}{\lambda_0(t)^M} \ \ge 0,$$

which is the logarithm of the relative partition function $\widehat{Z}_t(M)$ of §1.4. Hypothesis (B*) at T_0 with constant γ reads $h_t(T_0) \le \gamma V(t)$ for all t with sides in $[2T_0, 48T_0)$ and $2\gamma(96T_0)^3 \le \frac{1}{20}$. Throughout this section, (W1) and (W2) are assumed for all periods at least 3, and, as everywhere in this note, *positivity* of K_t is part of (W2).

Theorem 4.1 (reverse implication)**.** *Assume (W1)–(W2). Let $n \ge 3$ with $\Delta(n) \le 2^{-9}$, put $q = 256\,\Delta(n) \le \frac{1}{2}$ and $m_j = 2^j n$.*

(a) (The condition $\Delta(m_j) \le 2^{-9}$ needed to apply Proposition 3.3 at each base is part of the induction of Step 2.) For every $j \ge 0$ and every spatial torus t with $\min_i t_i \ge m_j$:

$$h_t(m_j) \ \le \ \frac{2\,\Delta(m_j)}{31\,m_j^3}\,V(t), \qquad \text{and} \qquad \Delta(m_j) \le \frac{q^{2^j}}{256}.$$

(b) (B) holds at $T_0 = 16n$ with $\gamma = 1/(31 \cdot 2^{23} T_0^3)$, for all spatial tori with all sides at least T_0, hence in particular for those with sides in $[2T_0, 48T_0)$. Here $2\gamma(96T_0)^3 = \frac{27}{3968} < \frac{1}{20}$.*

(c) For every integer $N \ge n$, every excited ratio r of the transfer operator of the cube (N, N, N) satisfies $r^n \le q \le \frac{1}{2}$; if $r > 0$ the gap is at least $\log(1/q)/n \ge (\log 2)/n$.

(d) For every spatial torus with $\min_i t_i \ge n$ and $\max_i t_i \le 24 \min_i t_i$, every excited ratio r satisfies $r^n \le \theta := e^{18/31} - 1 \approx 0.787$, so the gap is at least $-\log\theta/n \approx 0.2393/n$.

[proved] on paper. [Lean]: (a), first inequality (statement `interpolation`), (b) (`reciproque_Bstar`) and the first half of (c) ($r^n \le 256\Delta(n)$, `trou_cubes`), in the abstract form of Appendix A; not (d), not the logarithmic form of (c), and not the passage from periods ≥ 1 to periods ≥ 3.

Proof. Step 1: monotonicity of the pressure. Let $f(x) = \log Z(x)/|x|$ for $x \in \{3,4,\dots\}^4$, $|x| = x_1x_2x_3x_4$. Fix a spatial torus t and write $f(t,M) = f(t_1,t_2,t_3,M)$. By (2.3), $\log Z(t,M) = M\log\lambda_0 + h_t(M)$, so

$$f(t,M) = \frac{\log\lambda_0(t)}{V(t)} + \frac{h_t(M)}{M\,V(t)}.$$

Since $0 \le r_i \le 1$ (this is where positivity of K_t is used), $M \mapsto s_t(M) = \sum_{i\ge1} r_i^M$ is non-negative and non-increasing, hence so is h_t; and $M \mapsto 1/M$ is positive and decreasing. The product of two non-negative non-increasing functions is non-increasing, so $M \mapsto f(t,M)$ is non-increasing. By (W1) the same holds in each of the four coordinates: *f is non-increasing in each period*, with the other three held fixed, with no divisibility condition. Moreover $0 \le h_t(M) \le h_t(3) < \infty$, so $h_t(M)/M \to 0$ and $f(t,M) \to \log\lambda_0(t)/V(t)$ as $M \to \infty$.

Step 2: the recursion at every base. Since $f(m^4) - f((2m)^4) = \Delta(m)/(16m^4)$ by direct computation, and $(2m)^4$ is reached from m^4 by raising the four periods one after the other, Step 1 gives $\Delta(m) \ge 0$ for all $m \ge 3$. Write $\Delta_j = \Delta(m_j)$. By Proposition 3.3, applied at *each* base m_j: if $\Delta_j \le 2^{-9}$ then $\Delta_{j+1} \le 256\Delta_j^2 = (256\Delta_j)\Delta_j \le \Delta_j/2$. Since $\Delta_0 \le 2^{-9}$, induction gives $\Delta_j \le 2^{-j}\Delta_0 \le 2^{-9}$ for all j, and $256\Delta_{j+1} \le (256\Delta_j)^2$, whence $256\Delta_j \le q^{2^j}$. This is the second inequality of (a).

Step 3: telescoping. Fix j and $J > j$. Since $m_i^4 = 16^{i-j}m_j^4$ and $\Delta_i \le 2^{-(i-j)}\Delta_j$,

$$f(m_j^4) - f(m_J^4) = \sum_{i=j}^{J-1} \frac{\Delta_i}{16\,m_i^4} \le \frac{\Delta_j}{16\,m_j^4}\sum_{k\ge0} 32^{-k} = \frac{2\,\Delta_j}{31\,m_j^4}.$$

Step 4: the pressure of a torus against the diagonal. Let t have $\min_i t_i \ge m := m_j$, and let $M \ge 3$ be arbitrary. Choose $J \ge j$ with $t_1,t_2,t_3,M \le m_J$. By Step 1, $f(t,M) \ge f(m_J^4)$, and by Step 3, $f(m_J^4) \ge f(m^4) - 2\Delta(m)/(31m^4)$. Hence $\log\lambda_0(t)/V(t) + h_t(M)/(MV(t)) \ge f(m^4) - 2\Delta(m)/(31m^4)$ for every M; letting $M \to \infty$ (Step 1), $\log\lambda_0(t)/V(t) \ge f(m^4) - 2\Delta(m)/(31m^4)$. On the other hand $f(t,m) \le f(m^4)$ by Step 1, because $t_i \ge m$. Subtracting,

$$\frac{h_t(m)}{m\,V(t)} = f(t,m) - \frac{\log\lambda_0(t)}{V(t)} \le \frac{2\,\Delta(m)}{31\,m^4},$$

which is the first inequality of (a). No limit of free energies is interchanged with another, and no bound on the aspect ratio is used.

(b). By Step 2, $\Delta_4 = \Delta(16n) \le 2^{-24}$ (the bounds $2^{-9} \to 2^{-10} \to 2^{-12} \to 2^{-16} \to 2^{-24}$, each step being $256u^2$). Apply (a) with $m = 16n = T_0$: $h_t(T_0) \le 2\cdot 2^{-24}V(t)/(31T_0^3) = \gamma V(t)$. And $2\gamma(96T_0)^3 = 2\cdot 96^3/(31\cdot 2^{23}) = 27/3968$.

(c). Choose j with $m = m_j \le N < 2m$, and put $p = 2^j$, so $m = pn$. The cube has $V < 8m^3$; by (a), $h \le 16\Delta(m)/31 \le q^p/496$, with $h = h_{(N,N,N)}(m)$. Every excited ratio satisfies $r^m \le s(m) = e^h - 1 \le 2h \le q^p/248$ (as $h \le 1$). Since $r^m = (r^n)^p$, this gives $r^n \le q$.

(d). With $a = \min_i t_i$ and $m = m_j \le a < 2m$, $V(t) \le a\,(24a)^2 = 576a^3 < 4608\,m^3$. By (a), $h \le 9216\Delta(m)/31 \le \frac{36}{31}q^p \le \frac{18}{31}2^{1-p}$, using $q \le \frac12$. For $\alpha = 2^{1-p} \in (0,1]$ convexity gives $e^{\alpha x} - 1 \le \alpha(e^x - 1)$, so $s_t(m) \le 2^{1-p}\theta$ with $\theta = e^{18/31} - 1 \in (\frac12, 1)$. As $\theta \ge \frac12$, $2^{1-p} \le \theta^{p-1}$, so $r^m \le \theta^p$ and $r^n \le \theta$. □

Remark 4.2 (what is used, what is not). (i) *Positivity of K_t is indispensable.* It gives $r_i \in [0,1]$ and hence Step 1. Without it $\Delta(n)$ can be negative, and the statement is empty or false. Example: $Z(x) = \prod_{\mu=1}^4 (1+\rho^{x_\mu})^{|x|/x_\mu}$ with $\rho = -r$, $0 < r < 1$, is symmetric, and for each section t equals $\operatorname{Tr} K_t^M$ for the self-adjoint trace-class operator $K_t = c_t\,(1 \oplus \rho)^{\otimes V(t)}$ with a positive constant c_t (all factors with $x_\mu \ne M$ are exponentials in M), which has an eigenvalue $-r\lambda_0$. For odd n, $\Delta(n) = 64n^3\log(1-r^n) - 32n^3\log(1+r^{2n}) < 0$. (ii) The simplicity of λ_0 is not assumed: degenerate eigenvalues would be ratios $r_i = 1$. It follows from the conclusions: for the cubes of (c), $r_i^n \le q \le \frac12$, and for the tori of (d), $r_i^n \le \theta < 1$; for other sections the argument does not assert it. (iii) The proof assumes (W1) and (W2), with positivity, for all integer periods at least 3, with no upper cutoff: the

recursion of Step 2 and the comparison of Step 4 use arbitrarily large dyadic cubes and rectangles. No parity restriction is imposed. For Wilson's $SU(2)$ theory, positivity of the transfer operator at all spatial sides is the point on which the application rests: Lemma 7.3(vii) of [3] states $T \geq 0$ for $\beta \geq 0$; we read this statement, and have not checked its proof or its validity for odd sides. (iv) The Lean version (Appendix A) asks the spectral representation at all times $d \geq 1$ and all sides ≥ 1, as the already published `HypSpecTC`; the paper argument uses only $d,\ t_i \geq 3$ (replace $h(1)$ by $h(3)$ in Step 1).

Remark 4.3 (the factor 16 is not optimal). With $T_0 = 8n$ the same proof would give $y_0 = 2\gamma(96T_0)^3 = 54/31 > \frac{1}{20}$, so 16 is the first dyadic factor that this proof allows. It is not optimal. On product-of-rings models the smallest ratio T_0/n at which $\Delta(n) \leq 2^{-9}$ implies (B*) lies, numerically, between 1.17 and 1.75 (the minimum depends on n and on the family); this is a minimum over those models and not a bound. For massive free scalar fields at the threshold mass, a sufficient (not necessary) test certifies the ratios 1.85, 2.05 and 2.3 at $n = 6, 8, 12$; these are upper bounds on the smallest admissible ratio, not its value (`reproduction/reciproque/`). We do not know the optimal factor.

Remark 4.4 (the finite form (B**)). Proposition 7.57 of [3] states that (B*) with γ implies the inequality (B**): $2\log Z(s, T_0) - \log Z(s, 2T_0) \leq \gamma V(s)$ for all s in the window, and that (B**) with γ implies (B*) with γ (their $\psi_s(T_0)$ is a defect of the doubling of the time only, for tori (s, T_0) of all shapes in the window; ours doubles the four periods of a cube). By Theorem 4.1(b), $\Delta(n) \leq 2^{-9}$ gives (B*) at $16n$, hence (B**) at $16n$ with $2\gamma = 2/(31 \cdot 2^{23} T_0^3)$: indeed $2h_s(T_0) - h_s(2T_0) \leq 2h_s(T_0)$ since $h_s \geq 0$. Conversely (B*), hence (B**) up to the factor 2 in γ, implies $\Delta(2T_0) \leq 2^{-9}$ by (1.2). So our cubic defect at one size controls their window-wide defect at 16 times that size, and conversely.

Remark 4.5 (numerical checks, [computed]). Nothing in the proof depends on these. For massive free scalar fields on tori, on about $1.6 \cdot 10^5$ product-of-rings models (and direct sums of two of them) and on a pre-registered list of cubes and elongated tori, the inequality (a) and (B*) at $16n$ were evaluated with $\Delta(n) \leq 2^{-9}$ computed exactly: no violation; the largest ratio of the left to the right side of (a) is 0.245 (files in `reproduction/reciproque/`). These models are not general transfer matrices.

Corollary 4.6 (bridge to Theorem F of [3]; *conditional on hypothesis (W) and Theorem F of [3] as we read them; not verified by us*)**.** *Let a continuum family in the sense of [3, Definition 7.1] for $G = SU(2)$ satisfy their hypothesis (W). Suppose there are $t_0 > 0$, p_0 and $N_0(p)$ such that for all $p \geq p_0$ and $N \geq N_0(p)$ there is an integer $n = n_{N,p} \geq 3$ with $16n \leq Nt_0$ and $\Delta_{\beta_{N,p}}(n) \leq 2^{-9}$, where both partition functions in Δ are taken at the* same *coupling of the family at (N, p), on periodic tori n^4 and $(2n)^4$, and (B*) below is taken at that same physical coupling.* Normalisation: *$\beta_{N,p}$ denotes the coupling of the family in the normalisation of [3], in which, for $SU(2)$, the Wilson coupling is $\beta_W = 4\beta$ (their introduction, footnote on the unit quaternions); the defect Δ of this note is taken at $\beta_W = 4\beta_{N,p}$. No numerical constant of [3] that depends on β is used in this note, and the statements of versions 1.0 and 1.1 contain no numerical comparison involving their β. Then, by Theorem 4.1(b), their hypothesis (B*) holds at that coupling with $T_0 = 16n_{N,p}$, $3 \leq T_0 \leq Nt_0$, and $y_0 = 27/3968 \leq \frac{1}{20}$, $\gamma = y_0/(2(96T_0)^3)$; their Theorem F then gives $\sigma(H) \subset \{0\} \cup [\Delta_{\text{phys}}, \infty)$ with 0 simple and $\Delta_{\text{phys}} \geq 0.172/t_0$.*

Comment. We read in [3] that Theorem F asks for (W), $t_0 > 0$, $y_0 \leq \frac{1}{20}$, p_0, $N_0(p)$ and, for all $p \geq p_0$, $N \geq N_0(p)$, an integer T_0 with $3 \leq T_0 \leq Nt_0$ such that (B*) holds at $\beta_{N,p}$ with $\gamma = y_0/(2(96T_0)^3)$; (B*) concerns the tori with sides in $[2T_0, 48T_0)$, for which Theorem 4.1(b) gives the bound, and (W) is a double limit that this corollary does not touch. The corollary says nothing about the truth of the hypothesis $\Delta \leq 2^{-9}$ along a family, in particular for the construction of OpenAI, and it is not a proof of the mass gap.

Consequence. Under (W1)–(W2), the two one-scale criteria are equivalent up to a change of scale (a factor 32 for a round trip, (B*) at $T_0 \Rightarrow$ (B*) at $32T_0$): (B*) at T_0 implies $\Delta(2T_0) \leq 2^{-9}$ ((1.2));

$\Delta(n) \le 2^{-9}$ implies (B*) at $16n$. In particular the conclusion of Theorem 3.4 extends from the dyadic cubes $(2^j n)^3$ to all integer cubes $N \ge n$ and, with a worse constant, to all spatial tori with $\min_i t_i \ge n$ and $\max_i t_i \le 24 \min_i t_i$. This does not answer the question left open in [3] (their data there are a gap plus a bound $X_J \le \frac{1}{10}$, which is not our hypothesis $\Delta \le 2^{-9}$).

5 Assembly

Lemma 5.1 (q-form of each lattice form)**.** *If B has the representation (W3) at time $\ell = na > 0$ and $r_i^n \le q < 1$ for every $i \ge 1$, then $B(v_A, v_A, \ell) \le q\, B(v_A, v_A, 0)$ for every A, where $v_A = A - B(\Omega, A, 0)\Omega$ (and $q > 0$). [proved] [Lean: `qGapOn_of_specLink`]*

Proof. By linearity and $c(\Omega) = \delta_0$, $c_0(v_A) = c_0(A) - B(\Omega, A, 0) = 0$. Hence $B(v_A, v_A, \ell) = \sum_{i \ge 1} r_i^n c_i(v_A)^2 \le q \sum_{i \ge 1} c_i(v_A)^2 = q\, B(v_A, v_A, 0)$. □

Lemma 5.2 (closure [20, `QGapOn.of_tendsto`])**.** *Let B_k, B be bilinear forms such that $B_k(F, G, t) \to B(F, G, t)$ along a non-trivial filter for $F, G \in \mathcal{A}_0 \cup \{\Omega\}$ and $t \in \{0, \ell\}$. If eventually every B_k satisfies the q-form on $\mathcal{A}_0$ at time ℓ with the same q, so does B. [proved] [Lean, in ClayYM v1.1]*

Proof. By bilinearity $B_k(v_A^{(k)}, v_A^{(k)}, t) = B_k(A, A, t) - c_k B_k(A, \Omega, t) - c_k B_k(\Omega, A, t) + c_k^2 B_k(\Omega, \Omega, t)$ with $c_k = B_k(\Omega, A, 0)$: every term converges, and a non-strict inequality passes to the limit. □

Proof of Theorem 2.2. For each k, Theorem 3.4 on the section $(2^{j_k} n_k)^3$ gives $r_i^{n_k} \le 256 \Delta_{\beta_k}(n_k) \le 256\eta \le \frac{1}{2}$; Lemma 5.1, with (W3) at time $\ell = n_k a_k$, gives the q-form of B_k with $q = 256\eta$; Lemma 5.2 with (W4) gives $\mathrm{QGap}(R, S, \mathrm{all}, \ell, 256\eta)$. Under (W6), the q-form on a set whose truncated classes are dense in $\Omega^\perp$ (automatic for all families) gives $\|e^{-\ell H}(1 - P_\Omega)\| \le q$, hence the gap $-\log q/\ell$ and the uniqueness of the vacuum [20, `QGap.hasMassGap_univ`, `QGap.vacuum_unique`]; $-\log(256\eta) \ge \log 2$ since $256\eta \le \frac{1}{2}$. The Lean statement is `clay_trou_de_critere_4d` (Appendix A). The positivity of the limit form is not assumed: $B(F, F, 0) = \lim_k \sum_i c_i^{(k)}(F)^2 \ge 0$. □

The $d = 3$ version (`clay_trou_de_critere_3d`): under the three-dimensional counterparts of the hypotheses, $\ell > 0$ and $\Delta^{(3)}_{\beta_k}(n_k) \le \eta$ for every k with $0 < \eta \le 2^{-7}$, one gets $q = 64\eta$ and the gap $-\log(64\eta)/\ell$ (Lean), which is $\ge (\log 2)/\ell$ since $64\eta \le \frac{1}{2}$ (arithmetic not restated in Lean in $d = 3$) for a limit S of the three-dimensional theory (the layer `Gap/GapQ` of ClayYM is generic in d; the Clay problem itself is four-dimensional). In $d = 3$, $\beta_k = 4/(g_3^2 a_k)$ at fixed g_3^2 (super-renormalisable).

Lemma 5.3 (torus versus cylinder)**.** *Let $N = 2^j n$ with $j \ge 1$, $\sigma = N^3$, and let X be a bounded gauge-invariant observable supported in $r \le N/2$ consecutive time slices. Let ω_N be the periodic torus state and ω_∞ the Perron-cylinder state on $\mathbb{Z} \times \sigma$. Assume (W1)–(W2), the insertion property (W2$^+$), and $\Delta(n) \le 2^{-9}$. Then*

$$|\omega_N(X) - \omega_\infty(X)| \le 2\|X\|_\infty (e^{\Delta(N/2)} - 1), \qquad \Delta(N/2) \le \tfrac{1}{256}(256\Delta(n))^{2^{j-1}}.$$

[proved] on paper under (W2$^+$); not in Lean

Proof. Let $\widehat{K}_X$ be the normalised insertion operator across the r slices. By (W2$^+$), on the gauge-invariant configuration-space realisation, $|\widehat{K}_X f| \le \|X\|_\infty \widehat{K}^r |f|$; since $\|\widehat{K}^r\| = 1$ and $\||f|\|_2 = \|f\|_2$, we get $\|\widehat{K}_X\| \le \|X\|_\infty$. This step uses positivity preservation and insertion domination, in addition to positive semidefiniteness; it is the insertion argument of [14, proof of Prop. 2.2]. In the eigenbasis of $\widehat{K}$ on $\mathcal{H}_{\mathrm{inv}}(\sigma)$, with $\mu_0 = 1$, $\mu_i = r_i$,

$$\omega_N(X) = \frac{\mathrm{Tr}(\widehat{K}^{N-r}\widehat{K}_X)}{\mathrm{Tr}\,\widehat{K}^N} = \frac{\langle \Omega, \widehat{K}_X \Omega\rangle + \sum_{i \ge 1} \mu_i^{N-r} \langle e_i, \widehat{K}_X e_i\rangle}{1 + s_\sigma(N)}, \qquad \omega_\infty(X) = \langle \Omega, \widehat{K}_X \Omega\rangle,$$

where λ_0 is simple by Theorem 3.4, so that the cylinder state is the Perron vector state, and the gauge projector commutes with $\widehat{K}$ and $\widehat{K}_X$ (X is gauge invariant). Hence $|\omega_N(X) - \omega_\infty(X)| \le$

$\|X\|_\infty(s_\sigma(N-r)+s_\sigma(N)) \le 2\|X\|_\infty s_\sigma(N/2)$. By Lemma 3.1(a), $s_\sigma(N/2) \le e^p - 1$ with p the doubling of the period $N/2$ at the profile $(N^3, N/2)$, which is $a_3(N/2) \le \Delta(N/2)$; the last bound is Theorem 3.4. □

Corollary 5.4 (a diagonal family of square tori, (W5))**.** *Assume the hypotheses of Lemma 5.3 at each cutoff. Suppose that the lattice OS forms of a pair of families (F, G) are finite combinations of expectations of products of at most $D(F) + D(G)$ renormalised plaquette fields, each bounded by a constant $c_k < \infty$ at step k (the group is compact), the sum of the sup-norm bounds being at most $C(F)C(G)\max(1, c_k)^{D(F)+D(G)}$, and that for fixed k these bounds do not depend on the subsequently chosen spatial size. Set $N_k = 2^{j_k} n_k$ and choose $j_k \ge 1$ such that $2(e^{\Delta_{\beta_k}(N_k/2)} - 1)\max(1, c_k)^{2k} \le 1/k$ and $N_k a_k \ge k\ell$. Then, for fixed F, G and $t \in \{0, \ell\}$, the difference between the forms of the tori N_k^4 and of the cylinders $\mathbb{Z} \times N_k^3$ is at most $C(F)C(G)/k$ from some k on. Consequently either family of forms converges to a specified form B if and only if the other does, along this selected diagonal. [proved] on paper; the Lean reduction* `convergesOn_of_proche` *assumes the vanishing difference.*

Proof. For fixed F, G, eventually $D(F) + D(G) \le 2k$ and the fixed physical supports fit within half the temporal period; apply Lemma 5.3 term by term. The choice of j_k is possible because the upper bound for $\Delta_{\beta_k}(N_k/2)$ decreases doubly exponentially in j_k. This construction does not show that an arbitrarily prescribed $j_k \to \infty$ suffices, nor that changing a previously specified volume sequence preserves its limit. □

Remark 5.5 (the meaning of (W4)). (W4) is the precise sense of "the Schwinger functions converge" used here: pointwise convergence of the OS forms of smeared families at the two times 0 and ℓ, not convergence of kernels. It is the role of hypothesis (E) of [2]. One technical point: the plaquette field of ClayYM (`plaqObs`, plaquette based at the site x) is not exactly covariant under the site reflection of the lattice (temporal plaquettes are shifted by one step); with the lattice sequences of ClayYM, (W4) therefore also requires that this $O(a)$ shift disappear in the limit (true for locally uniform convergence, or with the clover field, which is exactly covariant). We include it in (W4).

6 The converse and why it needs a finite-volume gap

For $j = 0, \dots, d-1$ let $\sigma_j = ((2n)^j, n^{d-1-j})$ and $s_j(n) = \mathrm{Tr}'\, \widehat{K}_{\sigma_j}^n = \sum_{i \ge 1} r_i(\sigma_j)^n$ (excited trace). In physical units these are the spatial tori with sides in $\{\ell, 2\ell\}$, at Euclidean time ℓ.

Proposition 6.1 (bracket)**.** *Under (W1)–(W2), for every n,*

$$\sum_{j=0}^{d-1} 2^{d-1-j} \log(1 + s_j(n)) \;\le\; \Delta_d(n) \;\le\; 2\sum_{j=0}^{d-1} 2^{d-1-j} \log(1 + s_j(n)) \;\le\; 2\sum_{j=0}^{d-1} 2^{d-1-j} s_j(n).$$

Consequently $\Delta_d(n) \le \eta$ implies $s_j(n) \le e^\eta - 1$ for every j, and $\max_j s_j(n) \le \eta/(2(2^d - 1))$ implies $\Delta_d(n) \le \eta$ (the constant is 30 in $d = 4$, 14 in $d = 3$). [proved]

Proof. $a_j = 2\log(1 + s_j(n)) - \log(1 + s_j(2n))$ with $0 \le s_j(2n) \le s_j(n)$, so $\log(1 + s_j(n)) \le a_j \le 2\log(1 + s_j(n))$; sum with the weights of (3.1). □

Thus *the scalar criterion is, up to the factor $2(2^d - 1)$, the smallness of the excited trace at time ℓ on d spatial tori with sides in $\{\ell, 2\ell\}$.*

Corollary 6.2 (a sufficient converse estimate)**.** *Fix $\ell > t_0 > 0$, $m > 0$ and $B < \infty$, and let $n_k a_k = \ell$. Suppose that, uniformly in k and on the d sections $\sigma_j(n_k)$:*

(G_{fv}) finite-volume gap, flux sectors included*: $r_i(\sigma_j) \le e^{-m a_k}$ for every $i \ge 1$;*

(T) trace bound at a time $t_0 < \ell$*: $\mathrm{Tr}'\, \widehat{K}_{\sigma_j}^{n_0} \le B$, with $n_0 = \lceil t_0/a_k \rceil$.*

Then $s_j(n_k) \le Be^{-m(\ell-t_0-a_k)}$ *and* $\Delta_d(n_k) \le 2(2^d-1)\,B\,e^{-m(\ell-t_0-a_k)}$*; hence* $\Delta_d(n_k) \le \eta$ *as soon as the right-hand side is at most* η*. If* $a_k \le a_*$*, the condition* $2(2^d-1)Be^{-m(\ell-t_0-a_*)} \le \eta$ *is sufficient uniformly in* k*; if* $a_k \to 0$*, the strict inequality* $2(2^d-1)Be^{-m(\ell-t_0)} < \eta$ *suffices for* k *large.* [proved]

Proof. Since $t_0 < \ell = n_k a_k$, $n_0 \le n_k$ and $a_k(n_k - n_0) \ge \ell - t_0 - a_k$. Hence $r_i^{n_k} = r_i^{n_0} r_i^{n_k-n_0} \le r_i^{n_0} e^{-m a_k (n_k-n_0)}$; sum over i and apply Proposition 6.1. □

This is a statement at a fixed physical size: if ℓ varies, B, m and t_0 may vary with it, and eventual smallness as $\ell \to \infty$ requires, for instance, $\log B(\ell) - m(\ell - t_0(\ell) - a_*(\ell)) \to -\infty$; uniformity in the cutoff alone does not give this. An independent derivation by another model family (Qwen; see the statement on the use of AI) gives the same statement in the form $\Delta_4(n) \le 30 A_0 e^{m_0 r_0} e^{-m_0 n}$ under a uniform gap $(\lambda_2/\lambda_1)^r \le e^{-m_0(r-r_0)}$ and a trace bound A_0.

Neither hypothesis can be dropped.

Example 6.3 (decoupled copies: (T) is needed)**.** N decoupled copies of a theory satisfying (W1)–(W2) have the same transfer gap and $\log Z$ multiplied by N, so $\Delta^{(N)} = N\Delta^{(1)}$, unbounded in N. (T) says that the weighted number of states of the box below energy $\sim 1/t_0$ is bounded uniformly in the spacing: the convergence, at fixed physical size and temperature $1/t_0$, of the finite-volume free energy normalised by the vacuum. This is physically reasonable but not proved for Yang–Mills (the same difficulty as (W4), in finite volume). [proved]

Example 6.4 ($\mathbb{Z}_2$ gauge theory at $\beta = \infty$: a gap of the limit is not enough)**.** For the four-dimensional $\mathbb{Z}_2$ gauge theory at $\beta = \infty$, only flat configurations contribute; their number on a torus with V sites is $|B^1| \cdot |H^1(T^4;\mathbb{Z}_2)| = 2^{V-1} \cdot 2^4$. Hence $F(\vec{n}) = cV + 3\log 2$ on every rectangular torus, every p equals $3\log 2$, and

$$\Delta_4(n) = 15 \times 3\log 2 = 45\log 2 \approx 31.19 \qquad \text{for every } n.$$

(W1)–(W2) hold: K is a positive scalar multiple (depending on the spatial volume) of the rank-8 projector onto the holonomy sectors $H^1(T^3;\mathbb{Z}_2)$, so that $\widehat{K} = K/\lambda_0$ is that projector. Every local gauge-invariant observable is constant (all contractible loops equal 1), so the local limit is the trivial theory: unique vacuum and infinite gap. The criterion fails at every n although the limit is gapped; it fails because ($\mathrm{G_{fv}}$) fails (seven exactly degenerate sectors) [proved]. The limit is trivial, hence excluded by `FiniteMass` of ClayYM; the analogous phenomenon at large finite β (deconfined gapped phase with eight quasi-vacua split by tunnelling) is a conjecture here, not proved.

Example 6.5 (two-dimensional Ising model below T_c)**.** For the Ising model at $K = 0.5$ (two coexisting phases), $\Delta_2(n) \to 3\log 2 = (2^2-1)\log 2$ [computed], from Kaufman's exact solution [9] checked against dense transfer matrices (internal note of 10 November 2026). This is the value given by a phase-coexistence expansion of periodic partition functions of the type studied by Borgs and Kotecký [5], $\Delta_d \to (2^d-1)\log(\#\text{phases})$ (our reading; [5] do not state it). The periodic limit is a mixture without unique vacuum, so this is not a counterexample in the sense of ClayYM, but it shows again that ($\mathrm{G_{fv}}$) fails when two quasi-degenerate states exist in finite volume. The same example was obtained independently by Qwen.

Remark 6.6 (what the criterion encodes). The periodic trace tests every electric-flux sector $e \in \mathbb{Z}_2^3$ (Remark 2.1), not only the states visible to local observables in an infinite-volume limit; the bracket therefore controls both excited energies and their thermal multiplicities on the specified finite boxes. Example 6.4 shows that a gapped local limit, in a broader class of lattice theories, need not imply this finite-volume criterion; its local limit is trivial, so it does not establish strictness within the full non-trivial Clay class or within pure $SU(2)$ Yang–Mills theory, and no such strictness theorem is claimed. In a torelon model with energy $\approx \sigma L$ the criterion would also constrain $\sigma\ell^2$; this interpretation, and the estimate $n \approx 10$ ($\ell\sqrt{\sigma} \approx 3.7$) obtained at $\beta = 2.2986$ from the string tension and glueball masses of [1] with a free-particle and Nambu–Goto torelon model, are [heuristic], not consequences proved for the Wilson theory.

7 Witnesses

Proposition 7.1 (universal lower bound)**.** *Under (W1)–(W2), if the section n^{d-1} carries M excited states with $(\lambda/\lambda_0)^n \geq e^{-c}$, then $\Delta_d(n) \geq 2^{d-1} a_0(n) \geq 2^{d-1} \log(1 + Me^{-c})$.* [proved]

Proof. Lemma 3.1(a) and (3.1). □

A photon spectral witness, conditional on its model identification. Consider the transfer spectrum consisting of two photon oscillators for each nonzero spatial momentum, three electric rotors (energies $k^2 L_i^2/(2\beta V_s)$ per step) and three magnetic-flux energies $2\pi^2 \beta m^2 L_l/(L_i L_j)$, $\beta > 0$. This is the spectrum of the $U(1)$ Villain model *without monopoles* (Gaussian gauge field with compact holonomies, all integer fluxes) if the Euclidean partition function of that model equals $\operatorname{Tr} K^N$ for it with a trace constant equal to zero, so that (W1)–(W2) hold. *We do not derive this identification* from a fully specified Euclidean measure: the treatment of compact holonomies, harmonic fluxes, gauge-orbit factors and the absolute trace normalisation is supplied only in a summary (Poisson summation over the three temporal fluxes). A numerical check—the Euclidean formula and the transfer formula agree to $7 \cdot 10^{-16}$ relative on 12 four-dimensional tori, and the method reproduces the exact two-dimensional Villain partition function to $2 \cdot 10^{-41}$ [computed]—is a consistency check, not a derivation. Given the identification, the lowest photon frequency on the section n^3 is $\omega_1 = 2 \operatorname{arcsinh}(\sin(\pi/n)) \leq 2\pi/n$, independent of β; for $n \geq 3$ the 6 lowest nonzero momenta are distinct and, with 2 polarisations, give twelve one-photon states, so that Proposition 7.1 gives, for every $n \geq 3$ and every $\beta > 0$,

$$\Delta_4(n) \geq 8 \log(1 + 12e^{-2\pi}) = 0.1773 \approx 90 \times 2^{-9} \qquad \text{[proved], given the spectrum.}$$

The exact values (Table 3) are numerical; they are much larger, minimal at the self-dual point $\beta = 1/(2\pi)$ (electric–magnetic duality $\beta \leftrightarrow 1/(4\pi^2\beta)$), and grow at both ends (quasi-degenerate electric fluxes at large β, magnetic fluxes at small β). Under the stated identification, Maxwell theory never satisfies the criterion. The compact boson without vortices has the same qualification ($\Delta_4 \geq 8 \log(1 + 6e^{-2\pi}) = 0.0891$; numerically $\Delta_4 \to 25.9$–42.4).

Massive free scalar at fixed physical size. With $x = M\ell$, lattice mass x/n, and "continuum" values from the spectrum $\sqrt{M^2 + k^2}$ on the tori of sides ℓ and 2ℓ [computed]:

$x = M\ell$	0.1	1	5	10	15
Δ_4, $n = 8$	48.4	14.7	0.653	0.0204	$9.8 \cdot 10^{-4}$
Δ_4, $n = 64$	48.0	14.27	0.506	0.00724	$8.6 \cdot 10^{-5}$
Δ_4, continuum	48.0	14.26	0.504	0.00711	$8.2 \cdot 10^{-5}$
Δ_3, continuum	21.9	6.21	0.175	0.00194	$1.9 \cdot 10^{-5}$

Numerically, the massive scalar passes: $\Delta_4^{\text{cont}}(x) \leq 2^{-9}$ for $x \geq x_4^* = 11.47$ (2^{-12}: 13.79), and $\Delta_3^{\text{cont}} \leq 2^{-7}$ for $x \geq x_3^* = 8.47$. These crossings are numerical: the spectrum is truncated (frequencies below 80 in the script `reproduction/scripts/temoins.py`) and no tail certificate is given. At $x = x_4^*$ Theorem 3.4 gives a gap $\geq (\log 2)/\ell = M/16.5$: the bound is a minorant, loose by a factor ≈ 16. In the massless limit $\Delta^{\text{cont}}(x) \approx -(2^d - 1) \log x + \text{const}$ (from $x = 0.1$ to 0.01 the increase is $34.53 = 15 \ln 10$ in $d = 4$ and $16.11 = 7 \ln 10$ in $d = 3$).

Not treated: the compact $U(1)$ theory (Villain or Wilson, with monopoles) in its four-dimensional Coulomb phase, and weakly coupled $SU(2)$ in five dimensions; the criterion is expected to fail there for the same reason (massless photons), but we have no computation.

A non-admissible witness. The massless *non-compact* scalar violates (W2) (its spatial zero mode has continuous spectrum). With the convention "four-dimensional zero mode removed", $\Delta_4(n) = 10.46,\ 1.81,\ -4.74,\ -12.5,\ -17.3,\ -23.6,\ -28.0$ for $n = 2, 3, 4, 6, 8, 12, 16$ [computed]: negative values, impossible under (W1)–(W2) since Δ is a positive combination of p's. Prescriptions that remove zero modes therefore test models outside the hypotheses.

Dimension three. For the same kind of Gaussian transfer spectrum, assuming its identification with the periodic partition function, the four distinct lowest photon momenta give, for $n \geq 3$ and $\beta > 0$, $\Delta_3(n) \geq 4\log(1 + 4e^{-2\pi}) = 0.0298 \approx 3.8 \times 2^{-7}$ [proved] given the spectrum; numerically $\Delta_3 \geq 6.46$ (minimum at $e^2\ell \approx 8.0$, $n = 32$) [computed]. This Gaussian calculation is not a calculation of the compact three-dimensional $U(1)$ theory with monopoles; reference [6] concerns the confinement of static charges, and no further spectral statement from that paper is asserted here.

8 Strong coupling and a reported computational instance

Thermodynamic integration. Since $\partial_\beta F_\beta(\vec{n}) = -\sum_p \langle 1 - \frac{1}{2}\operatorname{Tr} U_p\rangle = -6|\Lambda|(1 - \langle P\rangle_{\vec{n}})$ with $\langle P\rangle$ the mean plaquette and $F_0 = 0$,

$$\Delta_\beta(n) = 96\, n^4 \int_0^\beta \left(\langle P\rangle_{n^4} - \langle P\rangle_{(2n)^4}\right) d\beta' \qquad \text{[proved]}. \tag{8.1}$$

Leading terms. With $t = I_2(\beta)/I_1(\beta) \approx \beta/4$, the $6n^2$ flat wrapping sheets and the $12n^3$ unit-cross-section wrapping tubes have fundamental weights t^{n^2} and t^{4n} (a sheet is a full 2-torus of n^2 plaquettes with Euler characteristic 0; a tube has $4n$ plaquettes). Their contributions to $16\log Z(n^4)$ are respectively $96\, n^2 t^{n^2}$ and $192\, n^3 t^{4n}$. At $n = 2$ the sheets contribute $384\, t^4$ and the tubes contribute $1536\, t^8$; the latter is not the complete coefficient of t^8 in $\Delta_4(2)$, which is not determined here. At $n = 4$ these two families together contribute $13\,824\, t^{16}$. Establishing the complete Taylor coefficients and the stated remainder orders requires the classification of the remaining graphs, representations and logarithmic cluster contributions; that derivation is not supplied here. A Monte Carlo computation of (8.1) on 2^4 and 4^4 for $\beta \leq 1.5$ is consistent with the leading coefficient 384 to 2.5% (fitted $A = 0.997 \pm 0.025$ in $\Delta_4(2)/(384t^4) = A + O(t^4)$) [computed, internal note of 10 November 2026]; it does not determine a Taylor coefficient. At leading order, $n = 2$ would suffice for $\beta \leq 0.190$ and $n = 4$ up to $\beta \approx 1.65$; but the leading term underestimates $\Delta_4(2)$ by 14% at $\beta = 1$ and 72% at $\beta = 1.5$, so these values are [heuristic]. A Monte Carlo estimate never certifies the criterion: at weak coupling it would require the plaquette to a relative accuracy of order 10^{-13}.

A reported computational instance. Assuming the two computational inputs specified below, the following analytic argument gives an interval on which the scalar hypothesis holds for the $SU(2)$ Wilson partition function.

Proposition 8.1 (reported computational instance, $n = 2$)**.** *Assume the reported finite combinatorial input (ENC): the complete counts of connected closed plaquette supports through size six, as listed in the proof. Assume also the exact rational comparison (RAT) specified after the proof. Then, for the four-dimensional $SU(2)$ Wilson theory and $0 \leq \beta \leq \beta_* = 0.005$, $\Delta_\beta(2) \leq 1.741 \cdot 10^{-3} < 2^{-9}$. Under (W1)–(W2), the transfer-ratio consequence follows from Theorem 3.4: on every spatial torus $(2^{j+1})^3$, $(\lambda_i/\lambda_0)^2 \leq 256\, \Delta_\beta(2)$ for $\beta \leq \beta_*$. The analytic estimate is proved below conditionally on (ENC) and (RAT); these computational inputs are reported from one implementation and are not independently certified in this note. [proved] conditionally on [computed] inputs*

Proof. Write $e^{(\beta/2)\operatorname{Tr} U} = c_0(1 + f(U))$ with $c_0 = \int e^{(\beta/2)\operatorname{Tr} U} dU$. Since the tori 2^4 and 4^4 have 96 and $1536 = 16 \cdot 96$ plaquettes, the constants c_0 and $e^{-\beta N_p}$ cancel in Δ, and $\Delta_\beta(2) = 16\log \widehat{Z}(2^4) - \log \widehat{Z}(4^4)$, with $\widehat{Z}(\Lambda) = \sum_P w(P)$ over sets P of plaquettes and $w(P) = \int \prod_{p\in P} f(U_p)\, dU$.

Activity. Under Haar measure $X = \frac{1}{2}\operatorname{Tr} U$ has a symmetric law on $[-1, 1]$, and $(\beta/2)\operatorname{Tr} U = \beta X$. Jensen's inequality gives $c_0 = \mathbb{E}\, e^{\beta X} \geq 1$, and the convexity bound $e^{\beta x} \leq \cosh\beta + x \sinh\beta$ on $[-1, 1]$ gives $c_0 \leq \cosh\beta$. Hence $f \leq e^\beta/c_0 - 1 \leq e^\beta - 1$ and $f \geq e^{-\beta}/c_0 - 1 \geq e^{-\beta}/\cosh\beta - 1 = -\tanh\beta \geq -\beta$, so that $|f| \leq \varepsilon := e^\beta - 1 \leq \beta + \beta^2$ for $0 \leq \beta \leq 1$, and $|w(P)| \leq \varepsilon^{|P|}$.

Closed sets. If a link of P belongs to only one plaquette $p \in P$, integrating that link first gives $\int f(U_l A)\, dU_l = \int f = 0$ by invariance of the Haar measure (the four links of p are distinct). Hence

$w(P) = 0$ unless P is *closed* (every link of P in at least two plaquettes of P). A closed P is the disjoint union, in links, of its connected components γ (connected through shared links), each closed, and $w(P) = \prod_\gamma w(\gamma)$. Therefore, with $S_\Lambda = \sum_\gamma \varepsilon^{|\gamma|}$ over connected closed sets,

$$\sum_{P \neq \emptyset} |w(P)| \le \sum_{r \ge 1} \frac{S_\Lambda^r}{r!} = e^{S_\Lambda} - 1, \qquad \log \widehat{Z}(2^4) \le S_{2^4}, \qquad \widehat{Z}(4^4) \ge 2 - e^{S_{4^4}}.$$

Counting. Let $N_m(\Lambda)$ be the number of connected closed sets of m plaquettes. By translation and axis permutations every plaquette is equivalent to a fixed root, so $N_m = |\mathcal{P}|\, c_m/m$ with c_m the number of such sets containing the root. An exact enumeration (Redelmeier's algorithm on the plaquette adjacency graph, with a pruning that is exact for closed sets; §11) gives the following. On 2^4 (degree 18): $c_m = 0$ for $m \le 3$ and for $m = 5$, $c_4 = 1$, $c_6 = 4$, i.e. $N_4 = 24$ (the sheets) and $N_6 = 64$ (the unit cubes), and no other connected closed set of at most six plaquettes. On 4^4 (degree 20): no closed set of at most five plaquettes, and $N_6 = 1024$ (the unit cubes, $c_6 = 4$). For larger m we use, on 2^4, $N_m \le \binom{96}{m}$, and on 4^4, $N_m \le \frac{1536}{m}\binom{20m}{m-1} \le \frac{1536}{m}\, e\, (20e)^{m-1}$: the adjacency graph has degree 20, and a connected set of size m containing the root is determined by the 0/1 string, of length at most $20m$ with $m-1$ ones, that records which neighbour slots are new members in a breadth-first exploration. The numbers listed above, including the absence of all other supports of size at most six, are the input (ENC), not a classification proved on paper here. "Closed" means that every incident link belongs to at least two selected plaquettes, allowing branched supports.

Conclusion. With $y = e^{S_{4^4}} - 1 \le S_{4^4} + S_{4^4}^2$ and $-\log(1-y) \le y/(1-y)$,

$$\Delta_\beta(2) \le 16\, S_{2^4} + \frac{y}{1-y},$$

and, for $0 \le \beta \le 1/200$, $\varepsilon \le u$. The counting bounds give $S_{2^4} \le A$ and $S_{4^4} \le B$, with A and B defined in (RAT). Assuming (RAT), $B \le 1$ and $e^{S_{4^4}} - 1 \le B + B^2 = Y < 1$. Thus $\Delta_\beta(2) \le 16A + Y/(1-Y) = R \le 1741/10^6 < 1/512$. Under (W1)–(W2), $\Delta \ge 0$ by Lemma 3.2. □

(RAT). With $u = 201/40000$ ($= \beta + \beta^2$ at $\beta = 1/200$) and $E = 2719/1000$, put

$$A = 24u^4 + 64u^6 + \sum_{m=7}^{96} \binom{96}{m} u^m, \qquad B = 1024u^6 + \frac{1536}{7}\, E(20E)^6 \frac{u^7}{1 - 20Eu},$$
$$Y = B + B^2, \qquad R = 16A + \frac{Y}{1-Y}.$$

The comparison to be certified in exact rational arithmetic is $B \le 1$, $Y < 1$ and $R \le 1741/10^6$. The program reports $R \approx 0.0017400504$, $A \approx 1.0362 \cdot 10^{-6}$ and $B \approx 0.0017176$; but the stored output does not contain the exact fraction (the field is empty), and these decimals are not the certificate. (ENC) and (RAT) are the two computational inputs of Proposition 8.1. The same implementation also reports that this upper bound exceeds 2^{-9} at $\beta = 0.0051$, and reports $\Delta_\beta(2) \le 2^{-12}$ for $\beta \le 0.0038$ [computed]; these additional computations are not certified by the stated input (RAT).

Remark 8.2. The exact counts c_m for $m \le 6$ and the exact rational evaluation come from a single program (`reproduction/instance/instance_n2_M6.py`) and have not been independently recounted. The bound is crude: the activity $\varepsilon \approx \beta$ is four times the true expansion parameter $t \approx \beta/4$, and no cancellation between the two tori is used, so β_* is about 38 times smaller than the leading-order value 0.190. A cluster expansion with the true activities and the cancellation of non-winding clusters should reach $\beta \approx 0.1$–0.2 for $n = 2$ [heuristic], and $n = 4$ up to $\beta \approx 1$ would require a computer-assisted expansion with rigorous tails; we have not done either. In this regime exponential decay and a lattice mass gap are already known from the cluster expansion of Osterwalder and Seiler [17], with better bounds: Proposition 8.1 is a reported check conditional on (ENC) and (RAT), not a new result on the gap.

9 Discussion

9.1 What this note does not prove

- **No weak-coupling estimate.** The theorem requires a scalar bound at some fixed physical size (of order $\ell \sim 2\,\mathrm{fm}$ for the physical theory), uniformly along the chosen cutoff trajectory. No estimate here establishes this uniform scalar hypothesis along a weak-coupling continuum trajectory (the "desert" between strong and weak coupling); establishing it remains an open sufficient route, and equivalence with the Yang–Mills existence and mass-gap problem is not proved. The strong-coupling interval reported in Proposition 8.1, conditional on (ENC) and (RAT), lies in a regime where the gap is already known.

- **No existence and no non-triviality of the continuum limit.** (W4) is assumed; a trivial limit satisfies the conclusion for every q.

- **No reconstruction.** (W6) is assumed, as in ClayYM.

- **The criterion contains additional finite-volume information.** It controls excited thermal traces, including flux sectors, which are not determined by the local infinite-volume gap alone in the broader class illustrated by Example 6.4. That example has a trivial local limit: strictness within the full Clay class or within pure $SU(2)$ is not established.

- **The transfer structure (W2)–(W3) is not formalised** (it would require Haar integration on $SU(2)^{\text{links}}$ and the compact spectral theorem in Lean), and the comparison of tori and cylinders is on paper only.

- **The relation with hypothesis (D) of [2] is not established in either direction.** In particular, (D) does not obviously give ($\mathrm{G_{fv}}$) on the flux sectors. Hypothesis (D) is that of version 4.45.0. For hypothesis (B^*) of version 5.0.0 [3] the two criteria are equivalent up to the scale: (B^*) at T_0 implies $\Delta_\beta(2T_0) \le 2^{-9}$ by (1.2), and $\Delta_\beta(n) \le 2^{-9}$ implies (B^*) at $16n$ (Theorem 4.1). Neither implies that the hypothesis holds along a weak-coupling trajectory, and Corollary 4.6 is conditional and unverified by us.

- The gap bound $(\log 2)/\ell$ is a sufficient lower bound, not an optimal estimate; the comparison with the massive scalar (a factor ≈ 16) uses the uncertified numerical crossing and is only illustrative.

9.2 Comparison with related conditional statements

- *OpenAI's two-dimensional criterion [14]* [read] (Lemma 2.1, Proposition 2.2 and the rectangle identity, in `sections/finite-volume.tex` of the repository). Two-dimensional spin models with a positive, self-adjoint, trace-class transfer operator, threshold 2^{-10}, recursion $44\Delta^2 \le 64\Delta^2$, gap $(\log 2)/n$ of every local limit of square tori, and of the lattice theory if the periodic state is unique; the finite-volume section supplies the excited-trace estimate, the rectangle identity, the quadratic iteration and the insertion comparison between square tori and cylinders. Theorem 3.4 transposes this argument to $d = 4$ and $d = 3$, using the weights 2^{d-1-j} of (3.1) and the full gauge-invariant trace, flux sectors included. The explicit constants and the Lean packaging are incremental additions; the continuum passage is a conditional closure of the q-form, and the converse is an elementary thermal-trace estimate. We found no converse and no $d > 2$ statement in the passages of [14] that were read (the passages of the finite-volume section and introduction recorded in the anteriority audit, lines 36–135 and 222–262 respectively, together with bounded keyword searches elsewhere in the source). These additions do not constitute a substantially new mass-gap mechanism, and the priority assessment is limited to the sources and passages actually read.

- *Bhattacharjee et al., Theorem B [2]* [read]. "Let $a_k \to 0$ and $\beta_k \geq 0$ be couplings, and suppose that there are $\varpi > 0$, $\ell > 0$ and integers $n_k \geq 2$ with $n_k a_k \leq \ell$ such that $\mathrm{BP}(n_k, \varpi)$ holds for the Wilson measure at β_k on all large tori. Then every continuum limit [...] has Hamiltonian H with $\sigma(H) \subset \{0\} \cup [\Delta, \infty)$, $\Delta = 16\varpi/(625e^2\ell)$" (v4.45.0, `p1_intro.tex`). $\mathrm{BP}(n, \varpi)$ is a variance (Poincaré) inequality for the dynamics that resamples cubes of side n, on all large tori. Same form as Theorem 2.2; the hypothesis is a spectral gap of a dynamics, not an inequality between two partition functions, and the limit is a separate hypothesis (E). Neither hypothesis is known to imply the other.

- *Bhattacharjee et al., version 5.0.0, Theorem 7.56 and Theorem F [3]* [read] (abstract, introduction, Sections 7.8 and 7.11 only). The hypothesis is no longer a spectral gap of a dynamics: it is a family of inequalities between partition functions of finite tori, on the same doubling defect as ours. Their record is earlier than version 1.0 of this note; their hypothesis implies ours, and ours implies theirs at a scale 16 times larger (Theorem 4.1); their conclusion extends beyond the dyadic-cubes conclusion of Theorem 3.4, and Theorem 4.1 supplies our extension to arbitrary integer cubes and bounded-aspect-ratio tori (§1.4).

- *Rajabu, Theorem 5.3 [18]*, as summarised in the project's earlier reading note (the primary text was not re-read in the anteriority audit for this version). A gauge-covariant block hierarchy (H1)–(H3), weighted Lipschitz covariance and transfer bounds at every fixed physical scale (C1), a landing bound (C2), time continuity (C3) and a non-zero excited sector (C4) give a gap $\geq w$ for the transfer operators of limits of block observables. Four multi-scale hypotheses against one scalar inequality; none of them is discharged here, and (C3)–(C4) have no counterpart (non-triviality is not treated here).

- *Tomboulis–Yaffe, Kovács–Tomboulis, Kanazawa* [19, 10, 8] [read] ([19] pp. 317 and 332, [10] eqs. (8) and (11), [8] Theorems 1, 2 and 5, read in the texts; page numbers are those of the printed paper for [19] and of the arXiv versions for the others; [19] was read on these pages only). Same family of argument in [10, 8]: reflection positivity, Schwarz inequality, repeated doublings of a loop or of a correlation until periodicity removes it, with divisibility conditions on the sides ($L_\mu = 2^p l_\mu$ in the proof of [8, Theorem 2, p. 9], whose footnote 10 says that the same condition was present in [19]; reflection positivity is applied to a two-point function in [19, p. 332]), and an inequality between partition functions of finite periodic tori. Different hypothesis: insensitivity of the torus to a centre twist on a stack of plaquettes, $\langle \mathcal{O}[\mathcal{V}] \rangle = Z^{(-)}/Z \to 1$ (equivalently $e^{-F_{\mathrm{el}}} = \frac{1}{2}(1 - Z^{(-)}/Z)$ small), not the doubling defect $16 \log Z(n^4) - \log Z((2n)^4)$ of this note. Different conclusion: a bound on Wilson loops of a periodic box, $\langle W[C] \rangle \leq 2(e^{-F_{\mathrm{el}}})^{|A_C|/|A_{12}|}$ (and, in [8], on correlations of spins), and not a bound on the excited eigenvalues of the transfer operator on every spatial torus. We do not claim that either hypothesis implies the other. These papers also contain the precedent recalled in §1.3, and are not used in our proofs.

- *Knabe-type criteria* [11, 7, 12]. Finite-size gap criteria for frustration-free quantum Hamiltonians: a local gap above a threshold on a finite subsystem implies a global gap. Same logical form, different objects: local Hamiltonian gaps, no partition functions, no reflection positivity, no continuum limit, one direction only.

- *Borgs–Kotecký* [5]. A finite-size scaling theory of periodic partition functions at first-order transitions. With such an expansion $\Delta_d \to (2^d - 1) \log(\#\text{phases})$ (Example 6.5): the test counts phases. This is our reading, not a statement of [5], whose text we have not re-read.

9.3 Reach

The note supplies a higher-dimensional version of an existing finite-volume sufficient criterion and packages its conditional continuum implication. It does not establish the scalar bound along a weak-coupling trajectory, construct the continuum limit, or identify an existing construction with all the

objects required by the criterion (Section 10). For square tori, the comparison must hold along the actual convergent family. Under exact blocking, $Z_{\mathrm{block}}(n) = Z(bn)$ gives $\Delta_{\mathrm{block}}(n) = \Delta(bn)$: this covariance is an identity, not a quantitative estimate that makes the missing smallness hypothesis easier.

10 Relation to the OpenAI construction

OpenAI have released a Lean-formalised construction for four-dimensional $SU(2)$ lattice Yang–Mills theory. This comparison concerns the local snapshot of that repository held by the author (folder dated 28 October 2026), inspected on 11 November 2026; file references are relative to it. **The criterion does not apply to that construction as it stands**, for three reasons, and a fourth point limits what can be said.

1. *The bare coupling may depend on the volume.* `YM/Source/Construction.lean:38` defines `abbrev beta (N p) := C.path.beta N p N`. The scalar test must then be evaluated at the common coupling $\beta_{N,p}$ of the target volume, $\Delta_{\beta_{N,p}}(n_N)$ with $n_N = \ell/a_N$; a difference of two separately tuned couplings is not the defect controlled by the doubling argument. A volume-dependent coupling is not a logical obstruction, but the required same-coupling estimate is an additional hypothesis.

2. *The dyadic volumes are not exported.* `PublicWilson.lean:38` exports only the field `periodStrict` (a `StrictMono` statement on `(period p).val`). The internal recipe contains `4*(M+1)` and `period i := page (M+i)` (`YM/Price/Waits.lean:87,102`), but the exported refinement (`ChallengeSpec.lean:188`, `T.Properties`) does not supply the hypothesis $L_{p_k} = 2^{j_k}\ell$, $j_k \to \infty$ that our torus comparison uses.

3. *The order of limits differs.* The fixed-volume limit uses `C.smear N p v t` (`PublicWilson.lean:89`); the subsequent volume limit uses `row (volumeOrder k)` (line 90): first the mesh limit at fixed physical volume, then the volume limit. Our torus–cylinder estimate (a diagonal choice of volumes as a function of the cutoff, Corollary 5.4) changes the family and does not identify its limit with this iterated limit.

4. *Partition-function control exists but does not bound the defect.* `YM/FullVolume/Scalar.lean` contains the one-volume bound `partition G n b <= q^(3*Fintype.card (Site n)-3)` (line 46), `raw_lost` (line 68) and a positive lower bound for `stripped A N p m` (line 98). These estimates do not give the same-coupling two-volume bound $\Delta \le \eta$ that the criterion requires; we assert no absence of control of $\log Z$.

An initial internal repair used an additional energy-bound hypothesis (EB). Later internal work reports a conditional four-copy transport estimate avoiding (EB), using moment bounds already available in the construction. These repairs are outside this note and do not establish the same-coupling scalar estimate or a complete identification with the objects used here; the present criterion therefore does not plug into that construction as it stands. Our searches in that repository for a bridge to our Lean objects (`ClayYM`, `rpForm`, `OSReconstruction`, `QGap`), for a finite-volume spectral decomposition, and for a bound on Δ were bounded textual searches and found none; they are not an exhaustive audit.

11 Reproducibility

The package accompanying this note contains:

- `lean/CritereScalaire.lean` (this work) and `ClayYM_v1_1_extrait/ClayYM/` (six files of ClayYM v1.1, verbatim); `lean/assembler_paquet.sh` builds the single file `main.lean` = `import Mathlib` + the six ClayYM files (imports removed, each wrapped in a `section`) + `CritereScalaire.lean`; the SHA-256 of `lean/main.lean` is that of the original build (`lean/MANIFEST.txt`);

- `lean/build_rejeu_2026-11-11.json`: the output of the recompilation from a clean copy by the author's side (§A), `lean/reconstruction_non_auteur_2026-11-11/`: the log and project files of reconstruction (b), and `rejeu_depuis_arbre_git_2026-11-12/`: the script, scaffolding and log of reconstruction (c);

- `reproduction/scripts/constantes_poly.py` (constants of Table 2), `reproduction/scripts/temoins.py` (witnesses), with their outputs in `reproduction/resultats/`;
- `reproduction/instance/instance_n2.py` (Proposition 8.1); the file actually run, `instance_n2_M6.py`, differs only by fixing the maximal exact size to 6; its output is `sortie_bac_M6_job_20261111-011302-d9004d.json` (0.4 s);
- `lean/Reciproque.lean` (Theorem 4.1, abstract form), `lean/main_v1_2.lean` (the published `main.lean` followed by the new file, byte for byte), `lean/reciproque/` (frozen statements, comparison script, mutants, certificate), and `reproduction/reciproque/` (the numerical checks of Remark 4.5, with outputs);
- `SHA256SUMS`.

All programs ran in a sandbox without network and without privileges ("qlab", Python 3, Lean 4 v4.34.0 with Mathlib commit `5ed2965`, tag v4.34.0). The command used for Lean is

```
bash lean/assembler_paquet.sh /path/to/main.lean
<sandbox runner> --kind lean --timeout 1200 /path/to/main.lean
```

and, for Python, the same runner with `-kind python -timeout S FILE`. Outside this sandbox, any Lean 4 v4.34.0 project with Mathlib at that commit compiles `main.lean` with `lake env lean main.lean`.

A The Lean formalisation

Files and build. `main.lean` (2175 lines), SHA-256
`944a99f6fc10f1696e6dfa5bc231b73dd9c57a101b48518c52e3e4ca944ef2c6`,
is assembled from `CritereScalaire.lean`, SHA-256
`0c4264a7c3f665346d1e479f6ed99e8b531b3b23960efcc718095d4ed4fe8496`,
and the ClayYM v1.1 files `Basic`, `OS`, `Gap`, `Fake/Trivial`, `Transfer`, `GapQ` (commit `93ac204`). Toolchain: Lean 4 v4.34.0, Mathlib `5ed2965`. No `axiom` is declared; the hypotheses are explicit arguments. The following build checks, all on 11 November 2026. (a) On the author's side, in the sandbox (final build `build4`, job `20261111-002010-32231f`) and again from a clean copy of the package (job `20261111-005922-02de92`): return code 0, 0 errors, no `sorry`; the 35 declarations of the namespace `CritereScalaire` depend only on `propext`, `Classical.choice`, `Quot.sound`. (b) By a non-author (an AI agent, Claude Sonnet 5.5, not the author of the Lean files), about 15:20 UTC, on a clean rented machine with the official Lean 4.34.0 toolchain and the official Mathlib cache: `lake build` return code 0 (8925 jobs; 187 s for `lake build`, following a 25 s cache check) on the file `main.lean` of the hash above, and the same 35 declarations with standard axioms; this one started from hashed copies of the sources, not from a git archive. (c) By a non-author (Claude Sonnet 5.5, not the author of the Lean files), from the *git archive of the package*: the package directory `papers/Paper_ScalarCriterion_SU2` was added to a scratch git repository (no commit in the public repository), its tree computed, and `git archive` of that tree copied to the rented machine (archive sha256 `97cd82b6...01573c`; git tree `16d2a27b...` of the package at that moment). The subtrees used, `lean/` (git id `0bc5d1f1...`) and `ClayYM_v1_1_extrait/` (`f6c358cb...`), are those of the final package; the note, the metadata and the logs, which differ, are not used by the build. On the machine, in a new directory, under the unprivileged user, at 20:03–20:05 UTC, with Lean 4.34.0 and Mathlib `5ed2965`: the ten hashes of `lean/MANIFEST.txt` were checked on the extracted files (all OK); the lake scaffolding (`lakefile.toml`, `lake-manifest.json`, `lean-toolchain`, a file of `#print axioms`) is not in the archive and was taken from reconstruction (b); the Mathlib package directory was copied from reconstruction (b) (official cache: "no files to download"), but no build artefact of the note was copied, and `main.lean` was taken from the archive (sha256 `944a99f6...`); `lake build` return code 0 (8925 jobs, `main` built in 28 s; 33 s for `lake build`, following a 13 s cache check), the same two linter warnings as before, and the same

35 declarations with axioms `propext`, `Classical.choice`, `Quot.sound`, the list being identical to that of `build4`. The logs are in the package (`rejeu_depuis_arbre_git_2026-11-12/`). These descriptions of rebuilds (a)–(c) concern version 1.0; version 1.2 adds separate Lean files, whose rebuilds are described below. Rebuilds (b) and (c) compiled the already assembled `main.lean`; they did not rerun the assembler from the component sources. They used the standard Lean kernel, not an independent kernel. The complete tree of the final package, which differs from the archived one only outside the two subtrees above, was not itself rebuilt. Version 1.1 (12 November 2026) changed the note, the README and the metadata files only. Version 1.2 adds files (below) and changes the note; `lean/CritereScalaire.lean`, `lean/main.lean`, `ClayYM_v1_1_extrait/` and `rejeu_depuis_arbre_git_2026-11-12/` are byte for byte those of version 1.0. An independent kernel check (nanoda) has *not* been run for `main.lean`; it was run for the new file, as described next.

Version 1.2: the reverse implication in Lean. `lean/Reciproque.lean` (474 lines, SHA-256 `efa8870dc236ef5d5a35d92e45e7cd7408573b2229e18d633404c017ab9d4dbb`) contains four principal theorem statements of the namespace `Reciproque` together with auxiliary lemmas and proof theorems (seven declarations introduced by `theorem`); the four principal statements were frozen on 12 November 2026 before their Lean proofs were written (file `lean/reciproque/ENONCES.lean`, SHA-256 `6a2c0e9f...ec120af8`); a script of the package (`lean/reciproque/verif_statique.py`) checks that the four statements and the definitions `sT`, `Delta`, `Rep`, `HypSpec`, `HypRec` are identical, character for character after whitespace normalisation, to the frozen ones. The abstract setting: a function F on four periods, symmetric (`hs12`, `hs23`, `hs34`); `Rep`: for each section (a, b, c) with positive sides, ratios $r_i \in [0, 1]$ with $\sum_i r_i^d < \infty$ and $F(a, b, c, d) = dL + \log(1 + \sum_i r_i^d)$ for every $d \geq 1$ (this is the already published `HypSpecTC`, with the positivity $r_i \geq 0$ in it); `HypRec`: the published recursion, $\Delta(m) \leq 2^{-9} \Rightarrow \Delta(2m) \leq 256\Delta(m)^2$, as a named hypothesis. The file `lean/main_v1_2.lean` is the published `main.lean` followed by this file inside a `section`, plus the theorem `Reciproque.reciproque_publie`, which *derives* `HypRec` from the published theorem `CritereScalaire.doublement_fin_4d` for any F with `CritereScalaire.HypSpecTC`; its first 114 899 bytes are `main.lean` (checked by comparison). `CritereScalaire.lean` and `main.lean` are thus unchanged, and the new content is separate and checkable.

declaration	what it states
`Reciproque.facile`	the easy direction of §1.4 in abstract form: from $a_j \leq \gamma 2^j n^3$ $(n = 2T_0)$ and $\Delta \leq 16a_0 + 8a_1 + 4a_2 + 2a_3$, $2\gamma(96T_0)^3 \leq \frac{1}{20}$: $\Delta \leq 512\gamma T_0^3 \leq 1/69120 < 2^{-9}$
`Reciproque.interpolation`	Theorem 4.1(a), first inequality: $\log(1 + s_t(m)) \leq 2\Delta(m)V(t)/(31m^3)$ for t with sides $\geq m$, given $\Delta(m) \leq 2^{-9}$
`Reciproque.reciproque_Bstar`	Theorem 4.1(b): $2\gamma(96T_0)^3 \leq \frac{1}{20}$ and $\log(1 + s_t(16n)) \leq \gamma V(t)$ for all sides $\geq 16n$, $\gamma = 1/(31 \cdot 2^{23} T_0^3)$
`Reciproque.trou_cubes`	Theorem 4.1(c), first half: $r_i^n \leq 256\Delta(n)$ for the cube (N, N, N), every $N \geq n$
`Reciproque.reciproque_publie` (in `main\_v1_2.lean` only)	`reciproque_Bstar` for F with `CritereScalaire.HypSpecTC`, symmetric, with `CritereScalaire.Delta` $\leq \frac{1}{512}$ at n

Not formalised: (d) of Theorem 4.1 (aspect ratio ≤ 24, the constant $e^{18/31} - 1$); the logarithmic form of (c); the Casimir-type bounds; the passage from sides ≥ 3 (paper) to sides ≥ 1 (Lean hypothesis, stronger); Corollary 4.6; the verification of (W2) for Wilson's theory.

Rebuilds of the new files (12 November 2026, by a non-author). (1) Sandbox, Lean 4 v4.34.0, Mathlib `5ed2965`: `Reciproque.lean` (job `20261112-070619-0a612c`) and `main\_v1\_2.lean` (job `20261112-070619-f382b1`, with one extra `#print axioms` line for `interpolation`): return code 0, 0 errors, no `sorry`; the four theorems, their auxiliary lemmas and `reciproque_publie` depend only on `propext`, `Classical.choice`, `Quot.sound`; the sandbox reports no axiom outside these. The same two files were compiled a second time in the second sandbox (same toolchain, other machine; jobs `20261112-072756-20ad77` and `20261112-072756-55f333`): return code 0, 0

errors, standard axioms only. (2) Rented machine, *Lean 4.32.0, Mathlib v4.32.0* (not the version of the sandbox), unprivileged user: Lean 4.32.0 (commit `8c9756b2`) with the Mathlib v4.32.0 of a project present on the machine, `lake env lean`, `nice 10`: both files, unmodified (same SHA-256), compile with return code 0 and no error (linter warnings on unused variables only), and give the same standard-axiom lines. The statements were not adapted for this version. (3) Independent kernel *nanoda*: for each file, the closure of the declarations (the four theorems; for `main\_v1\_2.lean` also `reciproque_publie` and `CritereScalaire.doublement_fin_4d`) was exported by `lean4export` (binary SHA-256 `e5736998...`) and replayed by the independent kernel nanoda (binary `ce46eaff...`, unmodified, only the three standard axioms permitted, any other an error): `Reciproque.lean`, 1 377 205 export lines, "Checked 16949 declarations with no errors"; `main\_v1\_2.lean`, 1 433 069 lines, "Checked 17156 declarations with no errors". This is the first nanoda check of any Lean file of this note; `main.lean` alone was not replayed, but `doublement_fin_4d` and its dependencies are in the second export. (4) Mutants of `Reciproque.lean`: constant $31 \to 30$ in the statement of `reciproque_Bstar`; factor $16 \to 8$ in its window; the positivity $r_i \geq 0$ removed from `Rep`; the threshold $\frac{1}{512} \to \frac{1}{256}$ in the hypothesis of `reciproque_Bstar`: all four make the build fail (errors located inside the targeted declarations or their users; `lean/reciproque/mutants/RESULTATS.txt`). The rebuilder is an instance of Claude Sonnet 5.5 that did not write the proofs; it also wrote the comparison script.

declaration	what it states, relative to the text
`lemme_fin_tc` (l. 42)	Lemma 3.1: $p \geq 0$, $s \leq e^p - 1$, $r_i^n \leq s$, $p(2n) \leq 2(e^p-1)^2$ (countable spectrum, $r_i \in [0,1]$, $s(n) < \infty$)
`decomp_4d` (l. 156)	$\Delta \geq 0$, $a_0 \leq \Delta/8$, $a_3 \leq \Delta$, from symmetry and positivity of p; not the full identity (3.1) nor a_1, a_2
`doublement_fin_4d` (l. 186), `chaine_num_fin4` (l. 125)	Proposition 3.3, with positivity and squaring bound of p as hypotheses (supplied by `hyp_of_specTC` from `HypSpecTC`)
`iteration256` (l. 248), `gap_4d_fin` (l. 299)	Theorem 3.4: $r_i^n \leq 256\,\Delta(n)$ on the section $(2^k n)^3$ given its spectral data; no simplicity, no logarithmic form
`doublement_fin_3d` (l. 418), `gap_3d_fin` (l. 513)	Theorem 3.5
`SpecLink` (l. 570), `qGapOn_of_specLink` (l. 581)	(W3) in coordinates (linear square-summable c, $c(\Omega) = \delta_0$, $B(\Omega, F, 0) = c_0(F)$, diagonal identities at times 0 and ℓ) and Lemma 5.1
`LienTransfert4` (l. 611), `qGapOn_limite_4d` (l. 628)	abstract form of Theorem 2.2, any filter; no a_k or equality of spacings appears in Lean: `SpecLink` relates the integer exponent n_k to the real time ℓ, and $\ell = n_k a_k$ is the assumed physical identification
`convergesOn_of_proche` (l. 672)	transfer of convergence *assuming* that the torus–cylinder difference tends to zero; the estimate and the diagonal choice are not formalised
`clay_trou_de_critere_4d` (l. 700), `_3d` (l. 722)	Theorem 2.2 with `QGap`, `HasMassGap` of ClayYM and, in $d = 4$, $(\log 2)/\ell \leq -\log(256\eta)/\ell$; vacuum uniqueness is `QGap.vacuum_unique` of ClayYM, the case $\eta \leq 2^{-12}$ is not restated
`non_vacuite_4d` (l. 802)	the hypotheses of `qGapOn_limite_4d` hold simultaneously for the trivial theory

Named hypotheses in Lean. Symmetry `hs12`, `hs23`, `hs34` (W1); `HypSpecTC` (W2), for all positive sizes; bilinearity `IsBilin` of the lattice forms; `LienTransfert4`/`SpecLink` (W3); `ConvergesOn` (W4); `OSReconstruction R S` and `S [] = 1` (W6). The guard `non_vacuite_4d` instantiates the hypotheses of the abstract theorem `qGapOn_limite_4d` with $V = \mathbb{R}$ and $B(F, G, t) = FG$. It does not instantiate the reconstruction hypothesis, and it does not show that the hypotheses hold for Wilson's $SU(2)$ theory.

Mutation tests. Eight mutants of `CritereScalaire.lean` must each make the build fail: constant $256 \to 200$; threshold $2^{-9} \to 2^{-7}$; a wrong symmetry in `decomp_4d`; vacuum coordinate $c(\Omega) \equiv 1$; a poisoned theorem $0 = 1$; constant $64 \to 56$ in $d = 3$; conclusion $q = 128\eta$; spectral link without the weights r_i^n. All eight made the build fail (job identifiers in `lean/mutants_chantier/resultats.json`). They were run on the text of an intermediate build, before the non-vacuity guard was added, and we did not check that each failure is located in the targeted declaration. A

re-run on the final text with localisation of the errors (`lean/mutants_paquet.py`) was started on 11 November 2026 but could not run: the sandbox had no free slot.

Use of AI

The work reported here was carried out with AI models, under the author's direction. The roles below are those recorded in the working files of the project.

- *Claude Opus 5.5 (Anthropic)* wrote, in several sessions on 10 and 11 November 2026, the transposition of the doubling mechanism to $d = 4$ and $d = 3$, the chain of constants, the assembly with ClayYM, the converse and its examples, the witness programs, the Lean files, the instance of Proposition 8.1 and the first draft of this note.

- *Qwen3.8 (Qwen team), run locally*, carried out the anteriority search of §9.2 (sources read and not read are recorded in the project files), derived independently the converse bound $\Delta_4(n) \leq 30A_0 e^{m_0 r_0} e^{-m_0 n}$ and the Ising counterexample, and reviewed the assembly by reading, without running programs (11 November 2026): it confirmed by hand the constants of the chain, the torus–cylinder lemma, the diagonal choice, the bracket, the $\mathbb{Z}_2$ counterexample and the role of the factor 16, pointed out the unlisted bilinearity hypothesis and the untreated compact $U(1)$ Coulomb phase (both now stated), and objected that (W2) assumes a simple λ_0; this objection is not retained, since degenerate eigenvalues are allowed as $r_i = 1$, as a later reviewer (GPT-6.1 Sol) also found. A second, line-by-line review of an earlier draft by Qwen, again without running programs, found that the torus–cylinder lemma needs a positivity-preserving kernel (now hypothesis (W2$^+$)), that the rounded constants of the chain must be read as upper bounds (now stated), and left the numerical inputs of the strong-coupling instance and of the witnesses unchecked.

- *GPT-6.1 Sol (OpenAI)* reviewed the draft of 11 November 2026 by reading, without running programs, and proposed scope and status corrections, adapted and incorporated in this version (the hypotheses (W2$^+$), (W5), the qualification of the torus and filter statements, the conditional status of the Maxwell witness, the formulation of the converse, the novelty and strictness claims, and the comparison with the OpenAI construction); Sol also recommended that this material be folded into a broader note rather than stand alone; publishing it as a stand-alone note is the author's decision. On 12 November 2026 Sol reviewed, again by reading and without running programs, the first version of this text and required sixteen corrections (three blocking: the domain of the Lean specialisation, the status of Proposition 8.1, and the order-eight coefficients), which Claude Sonnet adapted; a second read-only check of 12 November 2026 (`RELECTURE_SOL_TEXTE_CORRIGE.md`) found partial applications and asked for one blocking and four non-blocking corrections (the conditional status of Proposition 8.1 in the abstract and the list of results, the description of `SpecLink`, the concordance table, the attribution of the reviews and the wording of the rebuild descriptions), which are applied in this version; those earlier reports of Sol concerned the earlier texts; its review of version 1.2 is described below. These reviews do not certify the computational inputs or endorse a subsequent revision.

- *Claude Fable 5.2 (Anthropic)* acted as the project's pilot: it directed and arbitrated the work (the arbitrations are recorded in the project's registry). It wrote none of the proofs of this note, with one exception: on 12 November 2026 it wrote §1.4, including the derivation (1.2), and the other sentences of this version that refer to [3], and it ran the final `pdflatex` compilation. That subsection was written after two independent readings of the same passages of [3], by *DeepSeek V4 Pro (DeepSeek)* and by Qwen3.8, both without running programs. DeepSeek V4 Pro then checked the text of that subsection against the same passages (12 November 2026, without running programs): it confirmed the derivation (1.2) and asked for two changes of

wording (a literal quotation for hypothesis (W), and "narrower" instead of "weaker" for our conclusion), which are applied; it has not re-read the final wording.

- *Version 1.1 (12 November 2026).* A literature search was made on 12 November 2026, after the publication of version 1.0, by Claude Sonnet 5.5, which read the texts it obtained (scanned copies and arXiv files); the equation numbers and pages cited above for [19, 10, 8] were checked on images of the pages. In the same search DeepSeek V4 Pro, without tools, compared excerpts of those sources with statements of the companion note on the twist ratio (not of the present note). The corrections of this version (Section 1, §9.2, the bibliography and the status statements) were written by Claude Sonnet 5.5. One further review of this version by DeepSeek V4 Pro, without tools, on the text of the note and excerpts of the sources, is described in the file `RELECTURES.md` of the package. No statement, proof, constant or Lean file was changed.

- *GPT-6 Astra (OpenAI)* examined, blind, the electric-flux sectors under a scalar hypothesis (registry, 11 November 2026); that material is not part of this note. On 12 November 2026 GPT-6 Astra found the paper proof of the reverse implication (Theorem 4.1); GPT-6.1 Sol, DeepSeek V4 Pro and Claude Opus 5.5 read it the same day and reported no error (Sol with reservations on the wording of the statement, which are taken into account here); a first instance of Claude Sonnet 5.5 then checked each inequality, recomputed the constants, froze the four Lean statements and afterwards wrote their Lean proofs; the pilot (Claude Fable 5.2) redid the paper proof line by line and compared the frozen Lean statements with the theorem. The instance that prepared this version is a different one.

- *Version 1.2 (12 November 2026).* Claude Sonnet 5.5 prepared this version: it rebuilt the Lean files of the reverse implication as a non-author, on two toolchains and with an independent kernel, ran the mutants, wrote the section of Theorem 4.1 from the paper proof (corrected where indicated in `RELECTURES.md`), rewrote §1.4, ran a priority check against the current record of [3] and the literature, and compiled the note. Two reviewers of other model families tried to break the final text, proof included, on 12 November 2026, in two rounds each: GPT-6.1 Sol (OpenAI), read-only, with file-reading tools (it ran no build, script or network request), and DeepSeek V4 Pro (DeepSeek), without tools. Neither found an error in Theorem R or in its constants; both asked for wording corrections (the minimum-side restriction in two summaries, the domains of (W2), the remark on degenerate eigenvalues, the round-trip factor, the attribution of the monotonicity, the provenance statements of this section), which are applied; their reports, and the answer to each objection, are in `RELECTURES.md`. These reviews concern the text of 12 November 2026, 07h–08h, not later revisions. In addition, a complete reading of the text of [3] by DeepSeek V4 Pro, with quotations checked against the text by the preparer, was used for the comparison. These reviews do not certify later revisions. After these reviews, on 12 November 2026, the pilot (Claude Fable 5.2) corrected the description of who froze the Lean statements and who wrote the Lean proofs, removed a sentence about the public repository that was false, removed a conjecture on the optimal factor that our numerical checks do not support (Remark 4.3), added the output of the corresponding check (`reproduction/reciproque/t5.out`, rerun in a sandbox), and verified the reference [4] on the image of the page and its bibliographic data in Crossref; these last corrections were not reviewed by another model. No human has checked the proofs.

- *Claude Sonnet 5.5 (Anthropic)* prepared version 1.0 of 12 November 2026 (this and the preceding paragraphs on the pilot and on the rebuilds of Appendix A concern version 1.0; version 1.2 is described in the last paragraph of this list): it compared each formalised statement with the Lean declarations, applied the corrections above to the text (no statement, proof, constant or Lean file was otherwise modified), ran the `pdflatex` compilation, and ran reconstruction (c) of Appendix A from the git archive of the package. Another instance of Claude Sonnet 5.5 performed reconstruction (b) of that appendix.

These checks are not a substitute for refereeing: no human has checked the proofs. The human author read the final text of this note before its release; he did not verify the proofs, and no human expert has checked them.

References

[1] A. Athenodorou, M. Teper, *SU(N) gauge theories in 3+1 dimensions: glueball spectrum, string tensions and topology*, J. High Energy Phys. 2021, no. 12, 082, doi:10.1007/JHEP12(2021)082, arXiv:2106.00364.

[2] D. Bhattacharjee, U. Bhattacharya, J. M. Lockwood, T. Romanov, S. Singha Roy, *Yang–Mills theory and the mass gap*, version 4.45.0 (10 November 2026), Zenodo, doi:10.5281/zenodo.24111624.

[3] D. Bhattacharjee, U. Bhattacharya, J. M. Lockwood, T. Romanov, I. J. Sessford, S. Singha Roy, *The Yang–Mills Mass Gap from One Bound at One Scale*, version 5.0.0, Zenodo, doi:10.5281/zenodo.24147967 (record created on 11 November 2026 at 23:29:13 UTC; publication date 12 November 2026; paper of 157 pages in the archive `creelie/4Y-M-v5.0.0.zip`). Read on 12 November 2026: the passages listed in §1.4, the whole text by keyword searches, and (for version 1.2) a complete reading by DeepSeek V4 Pro with quotations checked against the text.

[4] D. Brydges, J. Fröhlich, E. Seiler, *On the construction of quantized gauge fields. I. General results*, Ann. Phys. 121 (1979) 227–284 doi:10.1016/0003-4916(79)90098-8 (our scanned copy is the preprint IHES/P/78/237, June 1978; the argument cited is on its p. 38, page 39 of the scan).

[5] C. Borgs, R. Kotecký, *A rigorous theory of finite-size scaling at first-order phase transitions*, J. Stat. Phys. 61 (1990) 79–119, doi:10.1007/BF01013955.

[6] M. Göpfert, G. Mack, *Proof of confinement of static quarks in 3-dimensional $U(1)$ lattice gauge theory for all values of the coupling constant*, Commun. Math. Phys. 82 (1982) 545–606, doi:10.1007/BF01961240.

[7] D. Gosset, E. Mozgunov, *Local gap threshold for frustration-free spin systems*, J. Math. Phys. 57 (2016) 091901, doi:10.1063/1.4962337, arXiv:1512.00088.

[8] T. Kanazawa, *Generalizing the Tomboulis–Yaffe inequality to $SU(N)$ lattice gauge theories and general classical spin systems*, Ann. Phys. 324 (2009) 1634–1665, arXiv:0808.3442v3 [math-ph] (page numbers cited here are those of the arXiv version).

[9] B. Kaufman, *Crystal statistics. II. Partition function evaluated by spinor analysis*, Phys. Rev. 76 (1949) 1232–1243, doi:10.1103/PhysRev.76.1232.

[10] T. G. Kovács, E. T. Tomboulis, *Vortex waistlines and long range fluctuations*, Phys. Rev. D 65 (2002) 074501, arXiv:hep-lat/0108017v2 (page numbers cited here are those of the arXiv version).

[11] S. Knabe, *Energy gaps and elementary excitations for certain VBS-quantum antiferromagnets*, J. Stat. Phys. 52 (1988) 627–638, doi:10.1007/BF01019721.

[12] M. Lemm, *Finite-size criteria for spectral gaps in D-dimensional quantum spin systems*, arXiv:1902.07141 (2019).

[13] M. Lüscher, *Construction of a selfadjoint, strictly positive transfer matrix for Euclidean lattice gauge theories*, Commun. Math. Phys. 54 (1977) 283–292, doi:10.1007/BF01614090.

[14] OpenAI, *Sharp mass bounds for the two-dimensional $O(4)$ model*, preprint dated 23 September 2026, repository `https://github.com/openai/math` (family 215, preprint directory `Sharp-mass-bounds-for-the-two-dimensional-O4-model-September-23-2026`, `sections/finite-volume.tex`); read from a local copy of the sources, not re-fetched on 12 November 2026.

[15] K. Osterwalder, R. Schrader, *Axioms for Euclidean Green's functions*, Commun. Math. Phys. 31 (1973) 83–112, doi:10.1007/BF01645738.

[16] K. Osterwalder, R. Schrader, *Axioms for Euclidean Green's functions II*, Commun. Math. Phys. 42 (1975) 281–305, doi:10.1007/BF01608978.

[17] K. Osterwalder, E. Seiler, *Gauge field theories on a lattice*, Ann. Phys. 110 (1978) 440–471, doi:10.1016/0003-4916(78)90039-8.

[18] H. T. Rajabu, *Multiscale covariance transfer and a conditional continuum mass gap for four-dimensional $SU(2)$ lattice Yang–Mills theory*, version 2.1 (29 October 2026), in the Zenodo record doi:10.5281/zenodo.23729328.

[19] E. T. Tomboulis, L. G. Yaffe, *Finite temperature $SU(2)$ lattice gauge theory*, Commun. Math. Phys. 100 (1985) 313–341 (read in a scanned copy with an OCR text layer).

[20] K. Rémondière, *ClayYM: a candidate Lean 4 statement of the Yang–Mills existence and mass gap problem, with compiled guards against fake models*, version 1.1.0 (25 October 2026), Zenodo, doi:10.5281/zenodo.23644079 (all versions).

[21] Internal notes of the project, 25 October and 10 November 2026 (unpublished).

name	statement	status
(W1)	F_β is invariant under permutations of the four periods.	[proved]: the action and the Haar measure are invariant under the hypercubic group. Lean: hypotheses `hs12`, `hs23`, `hs34`.
(W2)	For every spatial section $\sigma = (n_1, n_2, n_3)$ there is a self-adjoint, positive, trace-class operator K_σ on the space $\mathcal{H}_{\mathrm{inv}}(\sigma)$ of gauge-invariant L^2 functions of the spatial links, with $Z_\beta(\sigma, N) = \operatorname{Tr} K_\sigma^N$ for every $N \geq 2$ (Section 4 uses this only for spatial periods and times at least 3).	[hypothesis], standard [13], [17, §2.3, Thm. 2.1]; sketch in Remark 2.1. Not formalised. Lean: `HypSpecTC`. *Positivity of K_σ is indispensable* for Theorem 4.1 (and for $\Delta \geq 0$): without it Δ can be negative (Remark 4.2).
(W2$^+$)	In the configuration-space realisation, the normalised transfer is positivity preserving. After gauge projection, bounded gauge-invariant slab insertions satisfy $\lvert\widehat{K}_X f\rvert \leq \lVert X\rVert_\infty \widehat{K}^r \lvert f\rvert$.	[hypothesis], used only for the torus comparison (Lemma 5.3); an additional property of the path-integral realisation, not a consequence of positive semidefiniteness of an abstract operator. Not formalised.
(W3)	The lattice OS form of the infinite cylinder $\mathbb{Z} \times \sigma$ is given by the same normalised operator $\widehat{K}_\sigma = K_\sigma/\lambda_0$: $B(F, G, ma) = \langle \psi(F), \widehat{K}_\sigma^m \psi(G)\rangle$ with ψ linear and $\psi(\Omega)$ the Perron vector.	[hypothesis], standard (spectral theorem for the compact operator $\widehat{K}_\sigma$ and transfer formula of the cylinder state). Not formalised. Lean: `SpecLink`, `LienTransfert4`, which assume only linear square-summable coordinates c, $c(\Omega) = \delta_0$, the mixed identity $B(\Omega, F, 0) = c_0(F)$ and the diagonal identities for $B(F, F, 0)$ and $B(F, F, na)$ (Remark 2.1), not the full transfer representation at arbitrary times.
(W4)	Along the specified non-trivial filter (sequence, subsequence; a double limit is an iterated limit, not the product filter), the lattice OS forms $B_k(F, G, t)$ converge to rpForm $R\,S$ for $F, G \in \mathcal{A}_0 \cup \{\Omega\}$ and $t \in \{0, \ell\}$. This includes matching the lattice reflection convention to the continuum reflected form, including any vanishing one-step shifts.	[hypothesis]: the *existence of the continuum limit* and the identification of the OS forms; open. Lean: `ConvergesOn`.
(W5)	(Square tori.) For the family of square tori actually considered, the difference between its OS forms and those of the corresponding cylinders tends to zero on the same families and at the same times.	Paper estimate under (W2$^+$) for one diagonal choice $N_k = 2^{j_k} n_k$ (Lemma 5.3, Corollary 5.4); for a previously specified family it is a separate obligation. Lean: `convergesOn_of_proche` *assumes* the vanishing difference.
(W6)	The limit admits the Osterwalder–Schrader reconstruction, and $S(\emptyset) = 1$.	[hypothesis] of ClayYM (`OSReconstruction`) [15, 16, 20]; not formalised.

Table 1: Hypotheses. The paper arguments of Sections 3 and later use periods at least 2, and Section 4 uses (W2) for spatial periods and times at least 3 only. The Lean declarations assume a single symmetric function on natural periods, with `HypSpecTC` for every positive spatial period and every positive integer time. Applying them to Wilson's partition function therefore requires a symmetric extension to periods 1 satisfying those spectral identities; this extension is an additional assumption here, not a consequence proved in this note. Values at a zero period may be set to zero. The Lean hypotheses on the spectral data are, in places, weaker than the table (`SpecLink` assumes linear square-summable coordinates c, $c(\Omega) = \delta_0$, the mixed identity $B(\Omega, F, 0) = c_0(F)$, and the diagonal identities for $B(F, F, 0)$ and $B(F, F, \ell)$; it does not assume the full transfer representation at arbitrary times. `ConvergesOn` records the specified convergence of the lattice forms), but the hypothesis on the function F is stronger. The Lean statement also assumes that the lattice forms are bilinear (`IsBilin`), which holds by construction for OS forms. (W2) does *not* assume that λ_0 is simple: degenerate eigenvalues appear as $r_i = 1$ (Example 6.4). Not assumed anywhere: uniqueness of the Gibbs state or of the periodic limit, uniqueness of the continuum limit, reflection positivity of the limit as a separate hypothesis (it follows from (W3)–(W4)), $\beta_k \to \infty$.

d	threshold	constant used	polynomial chain	exact exponential [21]	$\lim_{D\to 0}$
2	2^{-5}	16	13.93	13.22	12
3	2^{-7}	64	60.655	59.22	56
4	2^{-9}	256	250.018	247.23	240
4	2^{-12}	256	241.216	—	240

Table 2: Constants C_d with $\Delta(2n) \leq C_d \Delta(n)^2$ below the threshold. "Polynomial chain": $e^x - 1 \leq x + x^2$, computed with rational arithmetic (`reproduction/resultats/constantes_poly.json`). Displayed decimal values are approximations to the computed expressions, not certified outward-rounded bounds; the proof uses the rational bounds 256 and 64. "Exact exponential": interval arithmetic, internal note of 10 November 2026. For comparison, [14, (2.12)] has 44 (and 64) at 2^{-10} in $d = 2$, with the squaring bound $p(2n) \leq 4p^2$ valid for $p \leq 1/4$.

β	0.02	0.1	$1/(2\pi)$	0.5	1	5
$\Delta_4(4)$	77.4	42.4	38.2	56.5	72.1	108.3
$\Delta_4(32)$	72.2	37.2	33.0	51.3	66.9	103.1

Table 3: Numerical evaluation of the stipulated photon, rotor and flux spectrum, $\Delta_4(n;\beta)$ [computed]; the limit is reached to 10^{-2} at $n = 16$. Its identification with a fully normalised monopole-free Villain measure is an additional unproved input, and no uniform error certificate is asserted.