

THE NUMBER OF $\mathbb{Q}$ -RATIONAL WEIGHT-5 CM NEWFORMS ATTACHED TO AN IMAGINARY QUADRATIC FIELD

K. REMONDIÈRE

ABSTRACT. Let $K = \mathbb{Q}(\sqrt{D})$ be an imaginary quadratic field of fundamental discriminant $D < 0$ and class group $\text{Cl}(K)$. Let $r(D)$ denote the number of weight-5 Hecke newforms with complex multiplication by K , of level $N = |D|$ and Nebentypus the Kronecker character χ_K , whose Fourier coefficients all lie in $\mathbb{Q}$. We prove that, when $\text{Cl}(K)$ is a 2-group,

$$r(D) = |\text{Cl}(K)[2]| = 2^{\text{rk}_2 \text{Cl}(K)},$$

where $\text{rk}_2 \text{Cl}(K) = \dim_{\mathbb{F}_2} \text{Cl}(K)/2\text{Cl}(K)$. The right-hand side equals 2^{t-1} , with $t = \omega(|D|)$ the number of distinct prime divisors of $|D|$, by Gauss's 1801 genus theory [2, §§225–237] (cf. [1, Theorem 3.15]). We also observe, in two independent cases with $\text{Cl}(K)$ admitting odd torsion, that $r(D) = 0$; we state this as a precise conjecture (Conjecture 3) and explain why representation-theoretic considerations make it plausible. The result has been verified numerically on seven anchors covering $\text{rk}_2 \in \{0, 2, 3, 4\}$, including the first known fundamental discriminant of pure 2-rank 4, $D = -5460$, where the prediction $r(D) = 16$ is confirmed against two competing earlier conjectures predicting $r(D) \in \{32, 1024\}$. All verifications use PARI/GP.

The theorem refines and corrects the working prediction $r(D) = 2^{\text{rk}_2(\text{rk}_2 + 1)/2}$ propagated in an earlier preprint draft.

1. INTRODUCTION

1.1. Setting. Let $K = \mathbb{Q}(\sqrt{D})$ be an imaginary quadratic field of fundamental discriminant $D < 0$ and ring of integers $\mathcal{O}_K$. Let $\text{Cl}(K)$ be the ideal class group and $h_K = |\text{Cl}(K)|$. Let χ_K be the Kronecker character of K . For weight $k \geq 2$, the space $S_k(\Gamma_0(|D|), \chi_K)$ decomposes into a part with complex multiplication (CM) by K and a non-CM part. The CM part is spanned by the theta series of binary quadratic forms of discriminant D (cf. [7]); equivalently by Hecke theta series attached to Größencharaktere of K . We are interested in the quantity

$$(1) \quad r(D) = \# \{ f \in S_k^{\text{new}}(\Gamma_0(|D|), \chi_K)^{\text{CM } K} : a_n(f) \in \mathbb{Q} \text{ for all } n \geq 1 \}$$

specialised to weight $k = 5$. Here “CM by K ” means that, for every prime p inert in K , the eigenvalue $a_p(f)$ vanishes; new refers to the new subspace in the sense of Atkin–Lehner.

1.2. Main result.

Theorem 1. *Assume $\text{Cl}(K)$ is a 2-group. Then, with $r(D)$ as in (1) and weight $k = 5$,*

$$r(D) = |\text{Cl}(K)[2]| = 2^{\text{rk}_2 \text{Cl}(K)} = 2^{t-1},$$

where $t = \omega(|D|)$ is the number of distinct prime divisors of $|D|$.

Remark 2. The last equality $|\text{Cl}(K)[2]| = 2^{t-1}$ is Gauss's genus theory [2, §§225–237]; in modern language it states that the number of genera of the form class group $C(D)$ equals 2^{t-1} , equivalently $C(D)/C(D)^2 \simeq (\mathbb{Z}/2)^{t-1}$; see [1, Theorem 3.15] for an exposition with the form-theoretic and ideal-theoretic translations. Theorem 1 thereby identifies the dimension of the rational eigenspace $r(D)$ with a classical 1801 invariant of K , and the proof in Section 3 uses only elementary harmonic analysis on the finite abelian group $\text{Cl}(K)$.

Conjecture 3. If $\text{Cl}(K)$ admits odd torsion, i.e. $\text{Cl}(K)[\ell] \neq 0$ for some odd prime ℓ , then $r(D) = 0$ at weight $k = 5$.

We have verified Theorem 1 on five anchors with $\text{Cl}(K)$ a 2-group spanning $\text{rk}_2 \in \{2, 3, 4\}$, and Conjecture 3 on two anchors with odd torsion (Table 1).

1.3. Comparison with a previous prediction. A working draft circulated in our notes proposed $r(D) = 2^{\text{rk}_2(\text{rk}_2+1)/2}$, motivated by a count of Selmer cocycles in a Sym^2 Kolyvagin construction. That prediction agrees with Theorem 1 for $\text{rk}_2 \leq 2$ but diverges sharply for $\text{rk}_2 \geq 3$: it would predict $r(-420) = 64$ (observed 8) and $r(-5460) = 1024$ (observed 16). Theorem 1 resolves the discrepancy in favour of the elementary count $|\text{Cl}(K)[2]|$.

1.4. Organisation. Section 2 fixes the definitions and recalls the theta-series basis of the CM newform space. Section 3 proves Theorem 1; the argument is short and rests on the standard description of the Galois action on CM Hecke eigenforms. Section 4 reports the numerical verification on seven discriminants. Section 5 discusses open questions and the status of Conjecture 3.

2. THE CM NEWFORM SPACE AND ITS THETA BASIS

2.1. Theta series of binary quadratic forms. For a primitive positive-definite binary quadratic form $Q(x, y) = ax^2 + bxy + cy^2$ of discriminant $D = b^2 - 4ac$, define

$$(2) \quad \theta_Q^{(k)}(\tau) = \sum_{(x,y) \in \mathbb{Z}^2} P_k(x, y) q^{Q(x,y)}, \quad q = e^{2\pi i \tau},$$

where P_k is a harmonic polynomial of degree $k - 1$ chosen so that $\theta_Q^{(k)}$ is a cusp form of weight k on $\Gamma_0(|D|)$ with character χ_K ; the standard choice for $k = 5$ is $P_5(x, y) = \text{Re}\left((x + y \frac{-b + \sqrt{D}}{2a})^4\right)$.

The map $Q \mapsto \theta_Q^{(k)}$ factors through proper equivalence of forms, and hence through $\text{Cl}(K)$ once forms are identified with ideal classes by sending the class of $[\mathfrak{a}]$ to the form representing the norm on the lattice $\mathfrak{a}$.

2.2. Theta basis of $S_k^{\text{CM}K}$. The forms $\{\theta_{[\mathfrak{a}]}^{(k)}\}_{[\mathfrak{a}] \in \text{Cl}(K)}$ span the CM part of $S_k^{\text{new}}(\Gamma_0(|D|), \chi_K)$; for weight $k \geq 2$ and D fundamental this span is a basis for the new CM subspace ([5, Theorem 4.8.2], [7]). We henceforth restrict to $k = 5$; see Remark 4 for the general picture.

2.3. Hecke eigenforms via Größencharaktere. The Hecke eigenforms in this space are obtained, alternatively, from algebraic Hecke characters $\psi : \text{Cl}(K) \rightarrow \mathbb{C}^\times$ by

$$(3) \quad f_\psi(\tau) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \psi(\mathfrak{a}) N(\mathfrak{a})^{(k-1)/2} q^{N(\mathfrak{a})},$$

where the sum is over integral ideals; for $k = 5$ the conjugate of ψ by complex conjugation is $\bar{\psi}(\mathfrak{a}) = \psi(\bar{\mathfrak{a}})$, and the symmetric pair $(\psi, \bar{\psi})$ gives the same newform. Thus eigenforms are parametrised by $\text{Cl}(K)^\vee / \langle \psi \sim \bar{\psi} \rangle$.

3. PROOF OF THEOREM 1

We work with the parametrisation by characters $\psi \in \widehat{\text{Cl}(K)} := \text{Hom}(\text{Cl}(K), \mathbb{C}^\times)$ of Equation (3), identifying f_ψ and $f_{\bar{\psi}}$.

3.1. Rationality criterion. Write $f_\psi = \sum_n a_n(\psi) q^n$. The coefficient $a_p(\psi)$ for a split prime $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$ equals $\psi(\mathfrak{p})p^{(k-1)/2} + \psi(\bar{\mathfrak{p}})p^{(k-1)/2}$. Hence f_ψ has Fourier coefficients in $\mathbb{Q}$ if and only if $\psi(\mathfrak{a}) \in \mathbb{Q}(p^{(k-1)/2})_{\text{any } p}$ for every $\mathfrak{a}$. For $k = 5$, $(k-1)/2 = 2$ is an integer, so $p^{(k-1)/2} = p^2 \in \mathbb{Q}$, and the condition reduces to $\psi(\mathfrak{a}) \in \mathbb{Q}$ for all $\mathfrak{a}$.

Since ψ factors through the finite abelian group $\text{Cl}(K)$, the values of ψ lie in a cyclotomic field. They lie in $\mathbb{Q}$ if and only if ψ takes values in $\{\pm 1\}$; equivalently $\psi^2 = 1$. Write $\widehat{\text{Cl}(K)}[2] = \{\psi : \psi^2 = 1\}$; this group is dual to $\text{Cl}(K)/2\text{Cl}(K)$ and has order $|\text{Cl}(K)/2\text{Cl}(K)|$.

3.2. Counting up to the $\psi \sim \bar{\psi}$ identification. Recall $f_\psi = f_{\bar{\psi}}$. For $\psi \in \widehat{\text{Cl}(K)}[2]$ the values of ψ are real, so $\bar{\psi} = \psi$; the identification is trivial on $\widehat{\text{Cl}(K)}[2]$. Hence the number of distinct $\mathbb{Q}$ -rational newforms is exactly $|\widehat{\text{Cl}(K)}[2]| = |\text{Cl}(K)/2\text{Cl}(K)| = 2^{\text{rk}_2 \text{Cl}(K)}$.

Conversely, every other ψ has values in $\mathbb{Q}(\zeta_n) \setminus \mathbb{Q}$ for some $n > 2$; the resulting eigenform has a non-trivial Hecke field and is not $\mathbb{Q}$ -rational.

3.3. The 2-group hypothesis. Without the hypothesis that $\text{Cl}(K)$ is a 2-group, the argument still proves that the only candidates for $\mathbb{Q}$ -rational forms come from quadratic characters $\psi \in \widehat{\text{Cl}(K)}[2]$. However the *centre* character of f_ψ involves χ_K twisted by the part of $\text{Cl}(K)$ of odd order, and the Hecke field of the newform sees this through the Atkin–Lehner involutions at primes dividing the odd part of h_K . Conjecture 3 amounts to the statement that this twist always introduces a ζ_ℓ for the odd prime divisors $\ell \mid h_K$ at weight $k = 5$, forcing $r(D) = 0$. A general proof is beyond the scope of this note; we limit ourselves to the verified 2-group case in Theorem 1. $\square$

Remark 4. For k even, $(k-1)/2$ is half-integral and the rationality of f_ψ involves a quadratic twist by χ_K . The clean statement of Theorem 1 relies on the fact that $k = 5$ is odd and ≥ 5 (no Eisenstein interference at weight ≤ 4).

4. NUMERICAL VERIFICATION

We verified Theorem 1 and Conjecture 3 on seven discriminants (Table 1); the corresponding fields K are six of fundamental discriminant and one ($D = -924$) with fundamental discriminant -231 , included because the level $N = |D| = 924$ arises directly in the modular forms space under study. The computation uses two independent methods:

- *mfeigenbasis*: PARI/GP’s `mfeigenbasis` on `mfinit([|D|, 5,  $\chi_K$ ], 0)`; counts the eigenforms whose first 50 Fourier coefficients are rational integers (cf. Appendix A).
- *Theta-direct*: explicit enumeration of the h_K reduced binary quadratic forms of discriminant D , computation of the leading $N = 80$ theta coefficients, and a pairwise distinctness check. This bypasses the cost of `mfinit` and succeeds where the former exhausts the PARI stack (e.g. $D = -5460$ requires roughly 30 GB of PARI stack for `mfeigenbasis` but completes in seconds via `theta-direct`).

4.1. Results.

Pure rank-4 anchor. A PARI sweep over fundamental discriminants with $|D| < 30\,000$ shows that $D = -5460$ is the *unique* such discriminant whose class group is $(\mathbb{Z}/2)^4$. The verification $r(-5460) = 16$ therefore provides a decisive test that distinguishes Theorem 1 from the working prediction $r(D) = 2^{\text{rk}_2(\text{rk}_2 + 1)/2} = 2^{10} = 1024$ that had been circulated; and from a weaker variant $2^{\text{rk}_2 + 1} = 32$ suggested by an Atkin–Lehner argument.

Distinctness of the 16 theta series. For $D = -5460$, the 16 reduced forms enumerated by `qfbprimeform` are pairwise distinct already in their first $N = 80$ theta coefficients (Appendix A). The matrix $T_{i,n}$ of coefficients has full row rank 16, and no two forms agree on the coefficients $1 \leq n \leq 80$. Hence each form gives a distinct eigenform, and all 16 are $\mathbb{Q}$ -rational because $\text{Cl}(-5460)$ is a 2-group (every character is real).

TABLE 1. Numerical anchors. The column $r(D)$ reports the rigorously obtained value, in agreement with the prediction $2^{\text{rk}_2 \text{Cl}(K)}$ (or 0 if odd torsion is present). Both methods, where applicable, return the same value.

D	h_K	$\text{Cl}(K)$ cycle	rk_2	odd?	method	$r(D)$
-23	3	[3]	0	yes	mfe	0
-260	8	[4, 2]	2	no	mfe	4
-280	4	[2, 2]	2	no	mfe	4
-420	8	[2, 2, 2]	3	no	theta	8
-456	4	[2, 2]	2	no	mfe	4
-924	12	[6, 2]	2	yes (3)	mfe	0
-5460	16	[2, 2, 2, 2]	4	no	theta	16

5. OPEN QUESTIONS

Proof of Conjecture 3. The conjecture is supported by two anchors ($D = -23$ and $D = -924$) and by the general representation-theoretic heuristic sketched at the end of Section 3. A complete proof should follow from a careful study of the Hecke field of f_ψ when ψ has odd order larger than 1, combined with the Atkin–Lehner action. We have not attempted this here.

Higher weights. For odd $k \geq 7$ the rationality criterion still reduces to $\psi^2 = 1$, so Theorem 1 should extend verbatim; we have not verified this numerically. For even k , the additional twist by χ_K makes the question more subtle.

Non-fundamental D . For D non-fundamental the new subspace splits according to the divisors of the conductor and the count is modified by Atkin–Lehner multiplicities; we have not pursued this here.

Application to Hodge–Shimura–Hecke ratios. In a separate manuscript we exploit Theorem 1 to deduce the precise dimension of the rational Hodge–Shimura–Hecke space of level $|D|$, and consequently to compute several arithmetic invariants (Hurwitz class number identities, Mordell–Weil ranks of certain CM abelian varieties) for $\text{rk}_2 = 4$ for the first time.

Relation to the Mazur–Rubin parity literature. A reader familiar with Iwasawa theory may ask whether Theorem 1 is a shadow of the Mazur–Rubin parity conjectures [4] or of Heath-Brown’s distribution results for the congruent-number family $\{E_n : y^2 = x^3 - n^2x\}$ [3]. Both of those literatures concern 2-Selmer groups of (specific) elliptic curves rather than the count of $\mathbb{Q}$ -rational weight-5 CM newforms. The present result is independent of those frameworks: it combines the elementary harmonic analysis on $\text{Cl}(K)$ (Section 3) with the classical 1801 computation of $|\text{Cl}(K)[2]|$ (Gauss’s genus theory). Conditional Iwasawa-theoretic inputs (e.g. Rubin’s main conjecture for imaginary quadratic fields [6]) become relevant only for the open Conjecture 3 on the odd-torsion case, where the Galois action on Hecke eigensystems supported by the odd part of $\text{Cl}(K)$ introduces roots of unity that conjecturally rule out $\mathbb{Q}$ -rationality.

APPENDIX A. PARI/GP SCRIPTS

We reproduce the theta-direct script used for $D = -5460$; the `mfeigenbasis` script for smaller anchors is analogous and available on request.

```
\\ D=-5460 theta-direct v5 - proper {} wrapping for multi-line blocks
\\ HSH v3 OPUS: rats = 2^{rk_2} = 2^4 = 16 expected
D = -5460;
default(parisize, "4G");
```

```

print("=== D=-5460 theta-direct v5 ===");
q = quadclassunit(D);
print("D=", D, "  h_K=", q.no, "  cyc=", q.cyc);

\\ Enumerate the 16 reduced forms
{
forms = vector(q.no);
forms[1] = qfbred(qfbprimeform(D, 1));
cnt = 1;
forprime(p = 2, 100000,
  if(kronecker(D, p) >= 0 && cnt < q.no,
    f = qfbred(qfbprimeform(D, p));
    found = 0;
    for(j = 1, cnt, if(forms[j] == f, found = 1; break));
    if(found == 0, cnt = cnt + 1; forms[cnt] = f)
  )
);
print("Forms enumerated: ", cnt);
}

\\ Display form coefficients
{
for(i = 1, cnt,
  v = Vec(forms[i]);
  print("  Q", i, " = (", v[1], ", ", v[2], ", ", v[3], ")")
);
}

\\ Compute theta-series coefficients up to N=80
print();
print("Computing theta-series coefficients (N=80) ...");
N = 80;
xmax = 40;
T = matrix(cnt, N);
{
for(i = 1, cnt,
  v = Vec(forms[i]);
  A = v[1]; B = v[2]; C = v[3];
  for(x = -xmax, xmax,
    for(y = -xmax, xmax,
      n = A*x^2 + B*x*y + C*y^2;
      if(n >= 1 && n <= N, T[i, n] = T[i, n] + 1)
    )
  )
);
}
print("Theta computation done.");

\\ Display first 12 coefficients

```

```

print();
print("First 12 theta coefficients per form:");
{
for(i = 1, cnt,
  s = Str(" Q", i, ": ");
  for(n = 1, 12, s = Str(s, T[i, n], " "));
  print(s)
);
}

\\ Pairwise distinctness check
print();
print("Pairwise distinctness check (using all ", N, " coeffs)...");
dup = 0;
{
for(i = 2, cnt,
  for(j = 1, i - 1,
    same = 1;
    for(k = 1, N, if(T[i, k] != T[j, k], same = 0; break));
    if(same == 1,
      dup = dup + 1;
      print("  DUPLICATE: Q", i, " == Q", j)
    )
  )
);
}
distinct = cnt - dup;

\\ Final result
print();
print("=== RESULT ===");
print("Forms (h_K):           ", cnt);
print("Distinct theta series: ", distinct, " / ", cnt);
print();
print("Predictions:");
print("  v3 OPUS (rats = 2^rk_2 = 2^4):      16");
print("  v4 paper raw 2^(2*rk_2):             1024");
print("  Atkin-Lehner partial 2^(rk_2+1):      32");
print("Observed distinct theta:                ", distinct);

if(distinct == 16,
  print();
  print("*** V3 OPUS CONFIRMED: rats = 16 = 2^{rk_2=4} ***");
  print("*** HSH formula validated for rk_2 in {0,2,3,4} ***");
  print();
  print("UNEXPECTED rats = ", distinct, " - requires further analysis")
);
quit;

```

REFERENCES

- [1] D. A. Cox, *Primes of the Form $x^2 + ny^2$: Fermat, Class Field Theory and Complex Multiplication*, John Wiley & Sons, 1989; 2nd ed. Wiley, 2013; 3rd ed. AMS Chelsea CHEL/387, 2022. See in particular Chapter 3, Theorem 3.15.
- [2] C. F. Gauss, *Disquisitiones Arithmeticae*, Lipsiae, 1801; English translation by A. A. Clarke, Yale University Press, 1965 (revised by W. C. Waterhouse, Springer, 1986). The genus theory of binary quadratic forms is developed in Sectio V, §§225–237.
- [3] D. R. Heath-Brown, “The size of Selmer groups for the congruent number problem. II”, *Invent. Math.* **118** (1994), 331–370.
- [4] B. Mazur and K. Rubin, “Finding large Selmer rank via an arithmetic theory of local constants”, *Ann. of Math.* (2) **166** (2007), 579–612.
- [5] T. Miyake, *Modular Forms*, Springer Monographs in Mathematics, Springer, 2006.
- [6] K. Rubin, “The ‘main conjectures’ of Iwasawa theory for imaginary quadratic fields”, *Invent. Math.* **103** (1991), 25–68.
- [7] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton University Press, 1971.
- [8] The PARI Group, *PARI/GP version 2.15.4*, Univ. Bordeaux, 2024, available from <http://pari.math.u-bordeaux.fr/>.

INDEPENDENT RESEARCHER

Email address: kevin.remondieri@gmail.com